



ELSEVIER

Available online at [www.sciencedirect.com](http://www.sciencedirect.com)

ScienceDirect

Current Opinion in  
Microbiology

# Emerging insights on intestinal dysbiosis during bacterial infections

Tu Anh N Pham and Trevor D Lawley

Infection of the gastrointestinal tract is commonly linked to pathological imbalances of the resident microbiota, termed dysbiosis. In recent years, advanced high-throughput genomic approaches have allowed us to examine the microbiota in an unprecedented manner, revealing novel biological insights about infection-associated dysbiosis at the community and individual species levels. A dysbiotic microbiota is typically reduced in taxonomic diversity and metabolic function, and can harbour pathobionts that exacerbate intestinal inflammation or manifest systemic disease. Dysbiosis can also promote pathogen genome evolution, while allowing the pathogens to persist at high density and transmit to new hosts. A deeper understanding of bacterial pathogenicity in the context of the intestinal microbiota should unveil new approaches for developing diagnostics and therapies for enteropathogens.

## Addresses

Wellcome Trust Sanger Institute, Wellcome Trust Genome Campus, Hinxton CB10 1SA, United Kingdom

Corresponding author: Lawley, Trevor D ([tl2@sanger.ac.uk](mailto:tl2@sanger.ac.uk))

Current Opinion in Microbiology 2014, 17:67–74

This review comes from a themed issue on **Host–microbe interactions: bacteria**

Edited by **Olivia Steele-Mortimer** and **Agathe Subtil**

For a complete overview see the [Issue](#) and the [Editorial](#)

Available online 29th December 2013

1369-5274 © 2013 The Authors. Published by Elsevier Ltd.

Open access under [CC BY license](#).

<http://dx.doi.org/10.1016/j.mib.2013.12.002>

## Introduction

The human intestinal microbiota is composed of 500–1000 diverse species, which together contains approximately 150 times more unique genes than our genome [1]. Often viewed as a “digestive organ”, the microbiota has co-evolved with the host to form a complex mutualistic relationship [2]: the gastrointestinal tract provides a nourishing environment for its microbial community, while the microbiota performs a wide range of essential metabolic, developmental and immune functions. A health-associated microbiota also represents the first line of defence against invading pathogens or resident

opportunists, and can facilitate pathogen clearance from the intestinal tract [3].

Over the past decade, the development of high-throughput sequencing technologies and analysis tools has enabled us to study the microbiota at an exceptional depth and resolution. At the same time, there is an increasing recognition that many pathogens such as *Clostridium difficile* and enterococci harbour potent virulence factors in their genomes, yet are commonly associated with asymptomatic carriage. Thus a pathogen’s ability to manifest virulence versus commensalism cannot be determined from the genome alone [4\*], and virulence genes (e.g. those encoding bacterial toxins, antimicrobial resistance, adhesion factors) may be essentially viewed as colonization factors [4\*,5]. Disease manifestation often depends not only on the dynamic between the pathogens and host immunity, but also on the composition and activity of the cohabiting microbiota. Recent studies monitoring the microbiota in patients or murine models of bacterial infection have indeed revealed new insights about pathogen biology during dysbiosis, including host colonization, disease, adaptation and transmission. Below, we discuss emerging concepts on infection-associated dysbiosis and their implications for host–microbe interactions.

## The intestinal microbiota during homeostasis

Without exposure to antibiotics or enteropathogens, a healthy gastrointestinal tract is home to a dense and diverse microbial community, known as the microbiota. A typical intestinal microbiota is dominated by obligate anaerobes belonging to the phyla Bacteroidetes, Firmicutes and Actinobacteria, and facultative anaerobes of the Proteobacteria phylum [6]. The microbiota assembly and structure vary widely between different individuals and at different anatomical sites along the length of the intestinal tract [7]. Nevertheless, a health-associated microbiome (that is, the collective encoding potential of the microbiota) is believed to be functionally conserved, and contains a shared gene set necessary to perform important biochemical reactions for host physiology [8\*]. These functions include the degradation of xenobiotic substances, vitamin biosynthesis and fermentation of indigestible polysaccharides into beneficial short-chain fatty acids (SCFA). Colonization by microbes also promotes our immune development, including the generation of IgA-secreting plasma cells or regulatory T cells to establish intestinal homeostasis with the commensal microbiota [9]. Finally, a healthy gut ecosystem is essential for colonization resistance [10,11,12\*], whereby both the

**Box 1 High-throughput methods to study host–microbiota interactions (Figure 1)***16S rRNA gene sequencing:*

The 16S ribosomal RNA gene is highly conserved among prokaryotic species, and can be sequenced using various platforms (e.g. Roche 454 pyrosequencing, Illumina HiSeq/Miseq). By comparing to publicly available rRNA gene databases (e.g. the Ribosomal Database Project, SILVA, Greengenes), the 16S rRNA gene sequence can be used as a surrogate marker to define the microbial lineages present in a community. This allows researchers to analyse the microbiota structure, including its taxonomic and phylogenetic diversity. To classify 16S gene sequences into bacterial taxa (often called Operational Taxonomic Units – OTUs), one commonly assumes that those with  $\geq 97\%$  nucleotide identity can be assigned to a single species. This assumption largely holds true except for a few genera (e.g. *Bacillus*) in which distinct species only differ at a few bases of their 16S sequence. Another limitation is that organisms with significant polymorphism in the regions used for primer design, such as Actinobacteria and *Bifidobacteria* spp., are poorly detected with “universal” 16S gene primers [64].

*Shotgun sequencing:*

Bacterial DNA is broken into small fragments and sequenced. Fragment sequences are assembled into contigs and aligned to construct a complete genome. Traditionally, this method was used to study organisms isolated and grown in pure cultures. Therefore, genomic sequences were only obtained from cultured species and represent merely a snapshot of one bacterial clone rather than of the community as a whole. However, shotgun sequencing is being increasingly applied directly to DNA obtained from mixed-community samples (i.e. **metagenomics**). This technique allows us to understand the community structure and also to build pathways describing its function, especially when combined with mRNA or protein-based quantification of the microbiota. Thus metagenomics is gaining popularity as a method to study microbial ecosystems.

However, there are remaining challenges. Most environmental and biological communities exhibit vast diversity and unevenness, with large variations in the relative abundance of different members. Therefore the genomes of rare species, especially those with the potential to greatly influence its community or host (i.e. keystone species), may be poorly detected or assembled even with deep sequencing. Efforts are under way to improve assembly algorithms, detection and normalization of previously under-studied phylotypes. The current databases to classify microbial genes, enzymes and pathways (e.g. COG, KEGG, CAZy) are also rapidly expanding, which will enhance our ability to assign or predict the functional capacity of the microbiome from shotgun sequences.

*Culture-based methods:*

Despite the advent of culture-independent methods to study microbial communities, culture techniques still play an essential role in defining host–microbe interactions. Although only  $<30\%$  of members of the human microbiota have been cultured to date, uncultured organisms may not be in fact unculturable. Permissive growth conditions for these organisms may be uncovered with improved, high-throughput culturing techniques [62\*\*]. Microbial culturing can allow us to study biological aspects of the organism such as its growth, metabolism or behaviour in a given host. In addition, the whole genomes of these microbes may be sequenced to analyse for genetic traits (e.g. symbiosis or virulence genes), or evidence of genomic evolution.

Finally, the functional output of the microbiota may be assessed by sequencing the mRNA content (i.e. **metatranscriptomics**), quantifying the proteome (i.e. **metaproteomics**) or the active metabolites (i.e. **metabolomics**). These approaches are especially valuable for studying the microbiota function during human clinical trials.

microbial community and the basal immune responses against resident commensals can together prevent access of pathogens.

**The intestinal microbiota during infections**

The importance of the resident microbiota during intestinal infections was highlighted by two seminal papers utilizing murine infection models with the Gram-negative pathogens *Salmonella enterica* serovar Typhimurium (or *S.* Typhimurium [13\*\*]) and *Citrobacter rodentium* [14\*\*]. In both models, pathogen-induced inflammation either led to or stabilized an imbalanced state of the microbiota community structure and function, termed intestinal dysbiosis. Advanced genomic methods have since been applied to other infection models, including Gram-positive and Gram-negative pathogens, to further define dysbiosis at both the microbial community and single species levels (Box 1 and Figure 1). Below, we summarize some of the emerging concepts from these studies.


**Exploitation of dysbiosis by enteric pathogens**

Diverse enteric pathogens often exploit dysbiosis, whether precipitated by antibiotic use or host inflammation, to outcompete resident commensals and gain access to intestinal nutrients and niches. In mice, *Salmonella* Typhimurium and *Clostridium difficile* can both colonize the gut asymptotically but only overgrow to high density and induce pathology after antibiotic treatment [13\*\*,15]. *C. difficile* is also the leading cause of antibiotic-associated diarrhea in humans, whereby disease manifestation predominantly occurs following antibiotic disruption of the microbiota, or in patients with inflammatory bowel disease [16]. When dysbiosis occurs, pathogens can rapidly outcompete commensals due to a greater resistance to host defences (e.g. antimicrobial and phagocyte killing), and better utilization of the gut nutrient environment [12\*,17]. For example, *Salmonella*'s competitive advantage is partly conferred by the ability to overcome host sequestration of iron [18] and to respire anaerobically using reactive oxygen species derived from the inflamed gut [19]. The metabolic environment during dysbiosis is also high in the SCFAs acetate and formate, which positively regulate the expression of *Salmonella* pathogenicity island-1 [20,21]. In addition, antibiotic use can lead to an increased availability of mucosal carbohydrates that are normally consumed by commensal *Bacteroides*, thus opening up new replicative niches for pathogens such as *Salmonella* and *C. difficile* [22].

**Dysbiosis is characterized by a simplified community structure and function**

Characterization of intestinal dysbiosis by different 16S rRNA gene sequencing approaches has consistently shown a reduction in taxonomic diversity and species membership of the microbiota. This observation also holds true across multiple human studies and animal

Figure 1



High-throughput genomic techniques commonly applied in microbiota research. **(a)** The functional state of the microbiota can be assessed directly by measuring its transcriptome (i.e. RNA-sequencing or metatranscriptomics), proteome (i.e. metaproteomics) or metabolites (i.e. metabolomics). Such approaches are still in their infancy but hold great promise for developing microbiota-based therapies and assessing human clinical studies. **(b)** Microbiota composition and taxonomy can be determined through directed amplicon sequencing of the 16S rRNA genes or by extracting 16S rRNA gene data from metagenomic datasets. Direct sequencing of the total DNA (i.e. shotgun metagenomics) also allows a measurement of the community function by defining the proteins and pathways (e.g. KEGG, COG, RefSeq pathways) that could potentially be active in the community to infer the overall functional capacity of the community. **(c)** Microbial species from the microbiota may be isolated and cultured by high-throughput techniques, termed “culturomics”, such as the use of barcoded plates with rich non-selective agar or liquid medium. The resulting microbes can then be whole-genome sequenced to examine their genetic traits, or analysed biologically with *in vitro* or *in vivo* assays. A combination of these complementary approaches will expand our understanding of the microbiota during health and disease and may ultimately yield microbiota-based therapeutics and diagnostics.

infection models, including *S. Typhimurium*, *C. rodentium*, *C. difficile*, and vancomycin-resistant enterococci (VRE) [14<sup>••</sup>,23–25,26<sup>••</sup>,27<sup>•</sup>,28]. The overall bacterial biomass may decrease in some cases depending on the inflammatory insult [14<sup>••</sup>,29]. In addition, dysbiosis generally leads to a depletion of obligate anaerobic bacteria such as *Bacteroides* and *Ruminococcus* spp., and conversely, a bloom in facultative anaerobes including the family Enterobacteriaceae (e.g. *E. coli*, *Klebsiella* spp., *Proteus* spp.). This shift may partly be due to the ability of Enterobacteriaceae species to respire using reactive nitrogen species — a byproduct of host inflammation, thereby outcompeting other commensals [30]. However,

the complex mechanism underlying other population-wide changes during dysbiosis (e.g. the bloom of anaerobic *Prevotella* spp. driven by NLRP6 inflammatory deficiency [31]) remains unclear.

Among the functional consequences of a simplified microbiota is a reduced metabolic capacity, often exemplified by a decline in SCFA production. This outcome may be in part due to a reduction in anaerobic bacteria, including dominant SCFA-producing genera such as *Bacteroides*, *Clostridium*, *Bifidobacterium* and *Roseburia*. SCFAs are physiological byproducts of carbohydrate fermentation by the microbiota, and serve to salvage energy for the

host as well as to enhance the mucosal barrier, inhibit intestinal inflammation and oxidative stress [32]. Dysbiosis caused by broad-spectrum antibiotics (e.g. clindamycin, cephalosporins), which can trigger opportunistic infection by *C. difficile* and enterococci, is commonly associated with low intestinal SCFA levels [23,33]. Furthermore, *C. difficile* infection may itself lead to decreased amounts of faecal acetate and butyrate, both in humans and equivalent murine models [25,33]. In the streptomycin-induced model of *S. Typhimurium* infection, butyrate level also decreases in the large intestine, which may promote bacterial invasion by stimulating expression of the *Salmonella* pathogenicity island genes [34]. As such, the microbiota's declining metabolic capacity may further impair host defence to pathogens and promote the stability of a dysbiotic community.

#### The dysbiotic microbiota acts as a pathogenic community

In *S. Typhimurium* infection, a microbiota with simplified structure (e.g. in mice treated with clinically relevant doses of antibiotics) or increased Enterobacteriaceae abundance may exacerbate disease outcome [35,36,37]. The pathogenic role of a dysbiotic microbiota is also shown in *C. difficile*-associated diarrhea, in which dysbiosis caused by an epidemic *C. difficile* strain leads to relapsing infection with more severe pathology [25]. Interestingly in some infection models such as *C. difficile* [25] and *C. rodentium* [14,28], microbiota analyses reveal that the inciting pathogens often constitute only a minor fraction of the overall microbial community. Together, these findings suggest that low-abundance pathogens could induce global changes to the microbiota structure and function, in a manner that further destabilizes the intestinal ecosystem. Such enteropathogens may be considered 'keystone species' [38], and likely influence the microbial community through a combination of their virulence expression, and of the host inflammatory and metabolic responses.

A dysbiotic microbiota may also be enriched for pathobionts — resident species with virulence potential that are normally kept at low levels. An overgrowth of commensal Enterobacteriaceae (*Klebsiella* spp. and *Proteus* spp.) or *Helicobacter typhlonius* has been shown to occur during intestinal dysbiosis, and can directly trigger colitis in mice [39]. Moreover, the depletion of anaerobic commensals during dysbiosis can lead to intestinal overgrowth of VRE, both in murine infection models or patients undergoing antibiotic therapies [26,40]. This consequently predisposes the host to invasive enterococcal infections with life-threatening sequelae [26]. Similarly, a multidrug-resistant *E. coli* pathobiont can expand in the mouse intestine following antibiotic disruption of the microbiota, causing bacteremia and sepsis [41]. Therefore during dysbiosis, the host may be increasingly susceptible to both pathogens and pathobionts, and the

microbiota may be viewed collectively as a pathogenic community.

#### An ecosystem for pathogen virulence expression and genome evolution


The microbiota often influences pathogen virulence and fitness upon passage through the gastrointestinal tract. Signaling from commensal bacteria has been shown to upregulate the virulence genes of enterohaemorrhagic *E. coli* O157:H7 and facilitates its adaptation to the host [42]. Another attaching-effacing pathogen, *C. rodentium*, also upregulates its virulence genes early during infection in a microbiota-dependent manner [43]. In both *C. rodentium* and *Vibrio cholerae*-induced diarrhea, passage through the gut allows the pathogens to efficiently colonize subsequent hosts [44,45]. "Hyperinfectious" *V. cholerae* can also persist in aquatic reservoirs — a phenotype associated with significant changes in the bacterial transcriptome, including a repression of chemotactic factors and upregulation of carbon metabolism [46].

In addition, pathogens may acquire virulence, fitness and antimicrobial resistance genes from the gut community, as they evolve under the selective pressures from host immune defence, microbial competition or antibiotic use. Transfer of antibiotic resistance genes by conjugative transposons has long been shown to occur extensively among pathogens and commensals, within the gut reservoirs of both humans and farm animals [47]. The hospital-associated pathogen *Enterococcus faecalis* V583 can also evolve in the intestinal tract by disseminating fluoroquinolone resistance and fitness-enhancing bacteriophages [48,49]. Using whole-genome sequencing and phylogenetics, He et al. recently demonstrated the rapid evolution of an epidemic *C. difficile* strain (ribotype 027), fuelled by antibiotic use and the transfer of mobile genetic elements with other intestinal bacteria [50,51]. In addition, Stecher and colleagues combined 16S rRNA gene and shotgun genome sequencing to show that during enterobacterial blooms in the inflamed gut, pathogenic *Samonella* and commensal *E. coli* can efficiently exchange fitness genes via conjugative plasmids [52]. The intestinal ecosystem represents a rich, dynamic reservoir for pathogens to intermingle and exchange genetic materials, especially during dysbiosis-induced bloom [53]. Therefore limiting dysbiosis, especially in the hospital setting, may have broad implications for the control of emerging infectious diseases.

#### Enhanced disease persistence and host-to-host transmission

Dysbiosis can promote pathogen transmission by increasing the levels of shedding and prolonging the infectious period. In both murine models of *Salmonella* and *C. difficile*-induced disease, antibiotic disruption of the microbiota leads to a remarkably high bacterial load ( $10^8$ - $10^9$  CFU/g of faeces) [15,25,54]. This phenomenon,

Figure 2



Features of intestinal dysbiosis during bacterial infections. **(a)** A healthy microbiota is typically diverse in structure and performs a wide range of functions (e.g. xenobiotic metabolism, production of SCFAs), thereby maintaining a mutualistic metabolic relationship with the host. Colonization resistance relies in part on the ability of the resident microbiota to outcompete pathogens for niches and nutrients. **(b)** During dysbiosis induced by pathogen-mediated inflammation or antibiotic perturbation, the microbiota is reduced in both taxonomic diversity and function, and intestinal colonization resistance is impaired. Diverse Gram-negative and Gram-positive pathogens can maintain dysbiosis by acting as keystone species to modulate community-wide shifts in the microbiota, possibly by orchestrating the host inflammatory response. As a result, the microbial community becomes more pathogenic, wherein pathogens and resident pathobionts may overgrow and even invade to cause systemic infection. Interactions with the gut microbiota often also allow pathogens to express their virulence factors and evolve under selective pressures. Consequently, the pathogens' increased fitness and over-colonization may exacerbate pathology and enhance host-to-host transmission.

also known as the “supershedder” phenotype, allows pathogens to transmit very effectively through direct contact or environmental contamination. For example, *C. difficile* supershedders can spread infection by releasing millions of infectious spores, which persist in the environment for long periods of months or even years [25,55]. In a clinical study involving VRE-infected patients, those shedding high bacterial levels have also been shown to contaminate their hospital surroundings [56]. In addition, the clearance of many pathogens including *Salmonella*, *C. difficile* and *C. rodentium* depends on the presence of a healthy microbiota [25,36,43]. As such, intestinal dysbiosis can also promote the spread of pathogens by allowing them to establish persistent infection within the host.

### Future perspectives on microbiota restoration

Each year, infectious diseases are increasingly difficult to treat because of rising antimicrobial resistance and a shortage in antibiotics discovery. Given the significant impact of dysbiosis on pathogen-mediated disease and transmission (Figure 2), the restitution of a healthy microbiota holds great promise as a therapy, at least for some infections. There are compelling evidences to suggest that administration of a diverse microbiota, or individual

probiotic species, can restrict or eliminate enteric pathogens. For example, microbiota transplantation may be used in mice to prevent lethal disease caused by *C. rodentium* and VRE [57,58]. Further, patients with recurrent *C. difficile* infection, or murine supershedders, can completely eliminate *C. difficile* after receiving a healthy microbiota [25,59]. Interestingly, *C. difficile* suppression may also be achieved by inoculating mice with Lachnospiraceae species or a defined group of six commensals (including the known probiotic *Lactobacillus reuteri* and previously uncharacterized Bacteroidetes and *Anaerostipes* spp.) [25,60]. This suggests that some microbial species derived from a healthy microbiota can potentially serve as standardized treatment, or “bacteriotherapy” [61]. The development of bacteriotherapy will require multiple complementary approaches, including high-throughput bacterial culturing (or “culturomics” [62,63]) and functional characterization of the human microbiota (e.g. pathway analyses of the microbiome, metatranscriptomics or metabolomics). Finally, mechanistic experiments in gnotobiotic animal models can inform us on how specific commensals influence colonization resistance and potentially be utilized as a microbiota-based therapeutic.

## Acknowledgements

We thank Alan Walker and members of the Bacterial Pathogenesis team for helpful comments of the manuscript.

## References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Sommer F, Backhed F: **The gut microbiota—masters of host development and physiology.** *Nat Rev Microbiol* 2013, **11**:227-238.
  2. Dethlefsen L, McFall-Ngai M, Relman DA: **An ecological and evolutionary perspective on human–microbe mutualism and disease.** *Nature* 2007, **449**:811-818.
  3. Stecher B, Hardt WD: **The role of microbiota in infectious disease.** *Trends Microbiol* 2008, **16**:107-114.
  4. Medini D, Serruto D, Parkhill J, Relman DA, Donati C, Moxon R, Falkow S, Rappuoli R: **Microbiology in the post-genomic era.** *Nat Rev Microbiol* 2008, **6**:419-430.
- This review describes emerging concepts in bacterial genomics and considers their application to the field of pathogen evolution and epidemiology.
5. Falkow S: **What is a pathogen?** *ASM News* 1997, **63**:359-365.
  6. Eckburg PB, Bik EM, Bernstein CN, Purdom E, Dethlefsen L, Sargent M, Gill SR, Nelson KE, Relman DA: **Diversity of the human intestinal microbial flora.** *Science* 2005, **308**:1635-1638.
  7. Lozupone CA, Stombaugh J, Gonzalez A, Ackermann G, Wendel D, Vazquez-Baeza Y, Jansson JK, Gordon JI, Knight R: **Meta-analyses of studies of the human microbiota.** *Genome Res* 2013, **23**:1704-1714.
  8. Qin J, Li R, Raes J, Arumugam M, Burgdorf KS, Manichanh C, Nielsen T, Pons N, Levenez F, Yamada T *et al.*: **A human gut microbial gene catalogue established by metagenomic sequencing.** *Nature* 2010, **464**:59-65.
- A landmark study from the MetaHIT (Metagenomics of the Human Gastrointestinal Tract) consortium characterizing the gut microbiome of 124 European individuals by metagenomic sequencing. The authors showed presence of a “core” gene set shared by most individuals, which encodes many genes necessary for beneficial metabolic functions.
9. Hooper LV, Macpherson AJ: **Immune adaptations that maintain homeostasis with the intestinal microbiota.** *Nat Rev Immunol* 2010, **10**:159-169.
  10. Kamada N, Seo SU, Chen GY, Nunez G: **Role of the gut microbiota in immunity and inflammatory disease.** *Nat Rev Immunol* 2013, **13**:321-335.
  11. Lawley TD, Walker AW: **Intestinal colonization resistance.** *Immunology* 2013, **138**:1-11.
  12. Stecher B, Hardt WD: **Mechanisms controlling pathogen colonization of the gut.** *Curr Opin Microbiol* 2011, **14**:82-91.
- This review describes our current understanding about the role of the microbiota in colonization resistance, with a wealth of experimental evidences to support different non-exclusive mechanisms.
13. Stecher B, Robbiani R, Walker AW, Westendorf AM, Barthel M, Kremer M, Chaffron S, Macpherson AJ, Buer J, Parkhill J *et al.*: **Salmonella enterica serovar Typhimurium exploits inflammation to compete with the intestinal microbiota.** *PLoS Biol* 2007, **5**:2177-2189.
- In this seminal study, the authors showed that *Salmonella* employs its virulence genes to elicit a robust host inflammatory response, and benefits from the resulting disruption of the microbiota. Accordingly, an avirulent *Salmonella* strain cannot compete with the microbiota, except in the inflamed gut environment of *IL10*<sup>-/-</sup> mice.
14. Lupp C, Robertson ML, Wickham ME, Sekirov I, Champion OL, Gaynor EC, Finlay BB: **Host-mediated inflammation disrupts the intestinal microbiota and promotes the overgrowth of Enterobacteriaceae.** *Cell Host Microbe* 2007, **2**:204.

Using whole 16S sequence analyses, the authors thoroughly examined the shift in microbiota composition during different models of intestinal infection and inflammation, including *Citrobacter rodentium*, *Campylobacter jejuni* and chemical-induced colitis. An overgrowth of Enterobacteriaceae family was identified as a common feature during intestinal inflammation.

15. Lawley TD, Bouley DM, Hoy YE, Gerke C, Relman DA, Monack DM: **Host transmission of Salmonella enterica serovar Typhimurium is controlled by virulence factors and indigenous intestinal microbiota.** *Infect Immun* 2008, **76**:403-416.
  16. Rupnik M, Wilcox MH, Gerding DN: **Clostridium difficile infection: new developments in epidemiology and pathogenesis.** *Nat Rev Microbiol* 2009, **7**:526-536.
  17. Keeney KM, Finlay BB: **Enteric pathogen exploitation of the microbiota-generated nutrient environment of the gut.** *Curr Opin Microbiol* 2011, **14**:92-98.
  18. Raffatellu M, George MD, Akiyama Y, Hornsby MJ, Nuccio SP, Paixao TA, Butler BP, Chu H, Santos RL, Berger T *et al.*: **Lipocalin-2 resistance confers an advantage to Salmonella enterica serotype Typhimurium for growth and survival in the inflamed intestine.** *Cell Host Microbe* 2009, **5**:476-486.
  19. Winter SE, Thiennimitr P, Winter MG, Butler BP, Huseby DL, Crawford RW, Russell JM, Bevins CL, Adams LG, Tsolis RM *et al.*: **Gut inflammation provides a respiratory electron acceptor for Salmonella.** *Nature* 2010, **467**:426-429.
  20. Huang Y, Suyemoto M, Garner CD, Cicconi KM, Altier C: **Formate acts as a diffusible signal to induce Salmonella invasion.** *J Bacteriol* 2008, **190**:4233-4241.
  21. Lawhon SD, Maurer R, Suyemoto M, Altier C: **Intestinal short-chain fatty acids alter Salmonella typhimurium invasion gene expression and virulence through BarA/SirA.** *Mol Microbiol* 2002, **46**:1451-1464.
  22. Ng KM, Ferreyra JA, Higginbottom SK, Lynch JB, Kashyap PC, Gopinath S, Naidu N, Choudhury B, Weimer BC, Monack DM *et al.*: **Microbiota-liberated host sugars facilitate post-antibiotic expansion of enteric pathogens.** *Nature* 2013, **502**:96-99.
  23. Buffie CG, Jarchum I, Equinda M, Lipuma L, Gobbourne A, Viale A, Ubeda C, Xavier J, Pamer EG: **Profound alterations of intestinal microbiota following a single dose of clindamycin results in sustained susceptibility to Clostridium difficile-induced colitis.** *Infect Immun* 2012, **80**:62-73.
  24. Kaiser P, Diard M, Stecher B, Hardt WD: **The streptomycin mouse model for Salmonella diarrhea: functional analysis of the microbiota, the pathogen's virulence factors, and the host's mucosal immune response.** *Immunol Rev* 2012, **245**:56-83.
  25. Lawley TD, Clare S, Walker AW, Stares MD, Connor TR, Raisen C, Goulding D, Rad R, Schreiber F, Brandt C *et al.*: **Targeted restoration of the intestinal microbiota with a simple, defined bacteriotherapy resolves relapsing Clostridium difficile disease in mice.** *PLoS Pathog* 2012, **8**:e1002995.
  26. Ubeda C, Taur Y, Jenq RR, Equinda MJ, Son T, Samstein M, Viale A, Succi ND, van den Brink MR, Kamboj M *et al.*: **Vancomycin-resistant Enterococcus domination of intestinal microbiota is enabled by antibiotic treatment in mice and precedes bloodstream invasion in humans.** *J Clin Invest* 2010, **120**:4332-4341.
- This study combines clinical evidence with a mouse model of VRE infection to demonstrate that enterococci may “dominate”, i.e. representing a major component (up to 95%) of the microbiota, following antibiotic disruption. This strongly correlated with subsequent enterococcal bacteremia in humans.
27. Chang JY, Antonopoulos DA, Kalra A, Tonelli A, Khalife WT, Schmidt TM, Young VB: **Decreased diversity of the fecal microbiome in recurrent Clostridium difficile-associated diarrhea.** *J Infect Dis* 2008, **197**:435-438.
- In this clinical study, the authors used 16S rRNA gene analysis to characterize the microbiota diversity of patients with initial and recurrent *C. difficile* infection. Those with recurrent disease had significant less diversity and a depletion in Bacteroidetes.
28. Hoffmann C, Hill DA, Minkah N, Kim T, Troy A, Artis D, Bushman F: **Community-wide response of the gut microbiota to**

- enteropathogenic *Citrobacter rodentium* infection revealed by deep sequencing.** *Infect Immun* 2009, **77**:4668-4678.
29. Willing BP, Russell SL, Finlay BB: **Shifting the balance: antibiotic effects on host-microbiota mutualism.** *Nat Rev Microbiol* 2011, **9**:233-243.
  30. Winter SE, Winter MG, Xavier MN, Thiennimitr P, Poon V, Keestra AM, Laughlin RC, Gomez G, Wu J, Lawhon SD *et al.*: **Host-derived nitrate boosts growth of *E. coli* in the inflamed gut.** *Science* 2013, **339**:708-711.
  31. Elinav E, Strowig T, Kau AL, Henao-Mejia J, Thaiss CA, Booth CJ, Peaper DR, Bertin J, Eisenbarth SC, Gordon JI *et al.*: **NLRP6 inflammasome regulates colonic microbial ecology and risk for colitis.** *Cell* 2011, **145**:745-757.
  32. Maslowski KM, Mackay CR: **Diet, gut microbiota and immune responses.** *Nat Immunol* 2011, **12**:5-9.
  33. Reeves AE, Theriot CM, Bergin IL, Huffnagle GB, Schloss PD, Young VB: **The interplay between microbiome dynamics and pathogen dynamics in a murine model of *Clostridium difficile* infection.** *Gut Microbes* 2011, **2**:145-158.
  34. Gantois I, Ducatelle R, Pasmans F, Haesebrouck F, Hautefort I, Thompson A, Hinton JC, Van Immerseel F: **Butyrate specifically down-regulates salmonella pathogenicity island 1 gene expression.** *Appl Environ Microbiol* 2006, **72**:946-949.
  35. Sekirov I, Tam NM, Jogova M, Robertson ML, Li Y, Lupp C, Finlay BB: **Antibiotic-induced perturbations of the intestinal microbiota alter host susceptibility to enteric infection.** *Infect Immun* 2008, **76**:4726-4736.
  36. Endt K, Stecher B, Chaffron S, Slack E, Tchitchek N, Benecke A, Van Maele L, Sirard JC, Heikenwalder M *et al.*: **The microbiota mediates pathogen clearance from the gut lumen after non-typhoidal *Salmonella* diarrhea.** *PLoS Pathog* 2010, **6**:e1001097.
- Here the authors showed that a diverse microbiota was necessary for clearance of *Salmonella* in the late stages of infection whereas secretory IgA was dispensable, and that mice given a low-complexity microbial mixture (the Altered Schaedler Flora) developed persistent infection.
37. Stecher B, Chaffron S, Kappeli R, Hapfelmeier S, Friedrich S, Weber TC, Kirundi J, Suar M, McCoy KD, von Mering C *et al.*: **Like will to like: abundances of closely related species can predict susceptibility to intestinal colonization by pathogenic and commensal bacteria.** *PLoS Pathog* 2010, **6**:e1000711.
  38. Hajishengallis G, Darveau RP, Curtis MA: **The keystone-pathogen hypothesis.** *Nat Rev Microbiol* 2012, **10**:717-725.
- This review proposes the ecological concept of 'keystone species', whereby low-abundance pathogens (e.g. *P. gingivalis*, *B. fragilis*) can drive community-wide effects on the microbiota.
39. Powell N, Walker AW, Stolarczyk E, Canavan JB, Gokmen MR, Marks E, Jackson I, Hashim A, Curtis MA, Jenner RG *et al.*: **The transcription factor T-bet regulates intestinal inflammation mediated by interleukin-7 receptor + innate lymphoid cells.** *Immunity* 2012, **37**:674-684.
  40. Stiefel U, Pultz NJ, Helfand MS, Donskey CJ: **Increased susceptibility to vancomycin-resistant *Enterococcus* intestinal colonization persists after completion of anti-anaerobic antibiotic treatment in mice.** *Infect Control Hosp Epidemiol* 2004, **25**:373-379.
  41. Ayres JS, Trinidad NJ, Vance RE: **Lethal inflammasome activation by a multidrug-resistant pathobiont upon antibiotic disruption of the microbiota.** *Nat Med* 2012, **18**:799-806.
  42. Hughes DT, Terekhova DA, Liou L, Hovde CJ, Sahl JW, Patankar AV, Gonzalez JE, Edrington TS, Rasko DA, Sperandio V: **Chemical sensing in mammalian host-bacterial commensal associations.** *Proc Natl Acad Sci U S A* 2010, **107**:9831-9836.
  43. Kamada N, Kim YG, Sham HP, Vallance BA, Puente JL, Martens EC, Nunez G: **Regulated virulence controls the ability of a pathogen to compete with the gut microbiota.** *Science* 2012, **336**:1325-1329.
- Using the *Citrobacter rodentium* infection model, the authors demonstrated that the microbiota is important for both virulence expression (early in infection) and for pathogen clearance (late in infection). Further, *C. rodentium* clearance is in part due to a competition for similar carbohydrates with commensal bacteria.
44. Merrell DS, Butler SM, Qadri F, Dolganov NA, Alam A, Cohen MB, Calderwood SB, Schoolnik GK, Camilli A: **Host-induced epidemic spread of the cholera bacterium.** *Nature* 2002, **417**:642-645.
  45. Wiles S, Dougan G, Frankel G: **Emergence of a 'hyperinfectious' bacterial state after passage of *Citrobacter rodentium* through the host gastrointestinal tract.** *Cell Microbiol* 2005, **7**:1163-1172.
  46. Schild S, Tamayo R, Nelson EJ, Qadri F, Calderwood SB, Camilli A: **Genes induced late in infection increase fitness of *Vibrio cholerae* after release into the environment.** *Cell Host Microbe* 2007, **2**:264-277.
  47. Salyers AA, Gupta A, Wang Y: **Human intestinal bacteria as reservoirs for antibiotic resistance genes.** *Trends Microbiol* 2004, **12**:412-416.
  48. Duerkop BA, Clements CV, Rollins D, Rodrigues JL, Hooper LV: **A composite bacteriophage alters colonization by an intestinal commensal bacterium.** *Proc Natl Acad Sci U S A* 2012, **109**:17621-17626.
- This paper showed that *Enterococcus faecalis* V583 produces a bacteriophage in a nutrient-dependent manner and outcompetes phage-sensitive *E. faecalis* *in vitro* and *in vivo*. This suggests that bacteriophage production represents a mechanism for bacteria to establish colonization and domination in the intestinal tract.
49. Matos RC, Lapaque N, Rigottier-Gois L, Debarbieux L, Meylheuc T, Gonzalez-Zorn B, Repoila F, Lopes Mde F, Serror P: ***Enterococcus faecalis* prophage dynamics and contributions to pathogenic traits.** *PLoS Genet* 2013, **9**:e1003539.
  50. He M, Miyajima F, Roberts P, Ellison L, Pickard DJ, Martin MJ, Connor TR, Harris SR, Fairley D, Bamford KB *et al.*: **Emergence and global spread of epidemic healthcare-associated *Clostridium difficile*.** *Nat Genet* 2013, **45**:109-113.
  51. He M, Sebailia M, Lawley TD, Stabler RA, Dawson LF, Martin MJ, Holt KE, Seth-Smith HM, Quail MA, Rance R *et al.*: **Evolutionary dynamics of *Clostridium difficile* over short and long time scales.** *Proc Natl Acad Sci U S A* 2010, **107**:7527-7532.
  52. Stecher B, Denzler R, Maier L, Bernet F, Sanders MJ, Pickard DJ, Barthel M, Westendorf AM, Krogfelt KA, Walker AW *et al.*: **Gut inflammation can boost horizontal gene transfer between pathogenic and commensal Enterobacteriaceae.** *Proc Natl Acad Sci U S A* 2012, **109**:1269-1274.
  53. Stecher B, Maier L, Hardt WD: **'Blooming' in the gut: how dysbiosis might contribute to pathogen evolution.** *Nat Rev Microbiol* 2013, **11**:277-284.
  54. Lawley TD, Clare S, Walker AW, Goulding D, Stabler RA, Croucher N, Mastroeni P, Scott P, Raisen C, Mottram L *et al.*: **Antibiotic treatment of *Clostridium difficile* carrier mice triggers a supershedder state, spore-mediated transmission, and severe disease in immunocompromised hosts.** *Infect Immun* 2009, **77**:3661-3669.
  55. Deakin LJ, Clare S, Fagan RP, Dawson LF, Pickard DJ, West MR, Wren BW, Fairweather NF, Dougan G, Lawley TD: **The *Clostridium difficile* spo0A gene is a persistence and transmission factor.** *Infect Immun* 2012, **80**:2704-2711.
  56. Donskey CJ, Chowdhry TK, Hecker MT, Huyen CK, Hanrahan JA, Hujer AM, Hutton-Thomas RA, Whalen CC, Bonomo RA, Rice LB: **Effect of antibiotic therapy on the density of vancomycin-resistant enterococci in the stool of colonized patients.** *N Engl J Med* 2000, **343**:1925-1932.
  57. Willing BP, Vacharaksa A, Croxen M, Thanachayanont T, Finlay BB: **Altering host resistance to infections through microbial transplantation.** *PLoS ONE* 2011, **6**:e26988.
- Using microbiota transplantation, the authors showed that a susceptible mouse strain, C3H/HeJ, may become more resistant to *C. rodentium* infection partly due to increased IL-22 signaling.
58. Ubeda C, Bucci V, Caballero S, Djukovic A, Toussaint NC, Equinda M, Lipuma L, Ling L, Gobourne A, No D *et al.*: **Intestinal microbiota containing *Barnesiella* species cures vancomycin-resistant *Enterococcus faecium* colonization.** *Infect Immun* 2013, **81**:965-973.

Here the authors found that introduction of a complete, diverse microbiota to mice heavily colonized with VRE could lead to VRE elimination. This effect was mediated by the obligate anaerobic members of the microbiota, in manner dependent on the presence of *Barnesiella* spp. (members of the Porphyromonadaceae family, *Bacteroidetes* phylum).

59. van Nood E, Vrieze A, Nieuwdorp M, Fuentes S, Zoetendal EG, de Vos WM, Visser CE, Kuijper EJ, Bartelsman JF, Tijssen JG *et al.*: **Duodenal infusion of donor feces for recurrent *Clostridium difficile***. *N Engl J Med* 2013, **368**:407-415.

A randomized controlled trial showing that fecal microbiota transplantation achieved 81% response in patients with recurrent *C. difficile* diarrhea, superior to vancomycin therapy alone or with bowel lavage.

60. Reeves AE, Koenigsknecht MJ, Bergin IL, Young VB: **Suppression of *Clostridium difficile* in the gastrointestinal tracts of germfree mice inoculated with a murine isolate from the family Lachnospiraceae**. *Infect Immun* 2012, **80**:3786-3794.
61. Adamu BO, Lawley TD: **Bacteriotherapy for the treatment of intestinal dysbiosis caused by *Clostridium difficile* infection**. *Curr Opin Microbiol* 2013.
62. Goodman AL, Kallstrom G, Faith JJ, Reyes A, Moore A, Dantas G, ●● Gordon JI: **Extensive personal human gut microbiota**

**culture collections characterized and manipulated in gnotobiotic mice**. *Proc Natl Acad Sci U S A* 2011, **108**:6252-6257.

The authors introduced the technique of culturing and archiving a large number of microbial species from human microbiota samples in bar-coded, multiwell plates. The complete microbiota or cultured microbes were then administered to germ-free mice to characterize their function *in vivo*.

63. Lagier JC, Armougom F, Million M, Hugon P, Pagnier I, Robert C, ● Bittar F, Fournous G, Gimenez G, Maraninchi M *et al.*: **Microbial culturomics: paradigm shift in the human gut microbiome study**. *Clin Microbiol Infect* 2012, **18**:1185-1193.

Here the authors used 212 different culture conditions, assisted by automatic colony picking and mass spectrometry for rapid identification of microbial species, to isolate 340 species from the microbiota of lean and obese individuals. Rare, fastidious species were recovered by co-culturing and elimination of more dominant bacteria. The authors also coined the term “culturomics” to describe large-scale, systematic culturing of microbes from a complex microbiota.

64. Farris MH, Olson JB: **Detection of *Actinobacteria* cultivated from environmental samples reveals bias in universal primers**. *Lett Appl Microbiol* 2007, **45**:376-381.