

of soluble electron carriers may have significant functional consequences since they would introduce macroscopic heterogeneities in the chain. The expected signature of such heterogeneities is essentially kinetic calling for new methods allowing the time-resolved analysis of the electron transfer sequence associated with the mitochondrial respiration *in vivo*. We are currently developing such a method based on the flash-induced photolysis of CO in the presence of O₂, as fruitfully conducted for the mechanistic dissection of complex IV. The advantages and pitfalls of the approach will be described and preliminary results will be presented and discussed.

doi:10.1016/j.bbabi.2008.05.362

S13.19 A role for sodium ions in the respiratory chain of *Rhodothermus marinus*

Andreia S. Fernandes^a, Ana P. Batista^a, Ricardo O. Louro^a, Miguel Teixeira^a, Julia Steuber^b, Manuela M. Pereira^a

^aInstituto de Tecnologia Química e Biológica, Universidade Nova de Lisboa, Oeiras, Portugal

^bDepartment of Biochemistry, University of Zurich, Switzerland

E-mail: andreaiaf@itqb.unl.pt

Rhodothermus marinus is a strictly aerobic and thermohalophilic organism isolated from submarine hot springs in Iceland and Açores. Its respiratory complexes have been studied and include a complex I (NADH:menaquinone oxidoreductase), a complex II (succinate:menaquinone oxidoreductase), a novel complex III and at least three different dioxygen reductases. Since it is a halophilic organism, and because a proton/sodium antiporter gene was found among its complex I genes, a possible role of sodium ions in *R. marinus* bioenergetics was investigated. We prepared inside-out vesicles from *R. marinus* and demonstrated that the vesicles maintained an electrochemical K⁺ potential imposed by K⁺/valinomycin. The membrane potential driven by the addition of substrates NADH and succinate to *R. marinus* membrane vesicles was followed using the sensitive dye oxonol V. It was observed that the NADH-driven membrane potential was sodium ion dependent, while the build-up of a membrane potential during succinate oxidation seems not to be influenced by Na⁺. To investigate the mode of Na⁺ transport during NADH respiration, ²³Na in membrane vesicles was followed by NMR spectroscopy.

doi:10.1016/j.bbabi.2008.05.363

S13.20 Structural characterization of respiratory complexes in potato tuber

Jelle B. Bultema^a, Roman Kouřil^a, Hans-Peter Braun^b, Egbert J. Boekema^a

^aDepartment of Biophysical Chemistry, University of Groningen, The Netherlands

^bInstitute for Plant Genetics, Faculty of natural Sciences, University of Hannover, Germany

E-mail: j.b.bultema@rug.nl

The aim of this study was to determine the structures of potato respiratory supercomplexes. Therefore, mitochondrial inner membranes from potato tuber cells were isolated, mildly solubilized with digitonin and the respiratory supercomplexes were separated by sucrose gradient ultra centrifugation. Finally, content of sucrose gradient fractions was inspected with Blue Native electrophoresis and electron microscopy. Single particle analysis of our data revealed several projection maps of complex I, monomeric and dimeric ATP synthase, supercomplex III₂+IV₁, supercomplex I+III₂ and larger

unassigned supercomplexes. In some side-view projection maps of complex I the structure of carbonic anhydrase shows its trimeric features. Furthermore, one projection map revealed an extra unknown density at the intermembrane side of complex I. Top-view projection maps of I+III₂ supercomplex showed similar features found in other plant species including the presence of carbonic anhydrase. Besides the top-views, two different side-views and several angular views of the I+III₂ supercomplex were revealed which allowed a better assignment of interaction between complex I and III₂ within the supercomplex. The side-views of the largest supercomplex most likely do not represent the structure of the I+III₂+IV₁ supercomplex, also known as the respirasome. The largest particles represent probably a supercomplex composed of two copies of complex I and one copy of complex III₂.

doi:10.1016/j.bbabi.2008.05.364

S13.21 Production, characterization, and determination of the real catalytic properties of the 'succinate dehydrogenase' from *Wolinella succinogenes*

Hanno D. Juhnke^a, Heiko Hiltcher^a, Hamid R. Nasiri^b, Harald Schwalbe^b, C. Roy D. Lancaster^{a,c}

^aCluster of Excellence "Macromolecular Complexes", Max Planck Institute of Biophysics, Department of Molecular Membrane Biology, Frankfurt am Main, Germany

^bCluster of Excellence "Macromolecular Complexes", Institut für Organische Chemie und Chemische Biologie, Center for Biomolecular Magnetic Resonance, Johann Wolfgang Goethe-Universität, Frankfurt am Main, Germany

^cChair of Structural Biology, Faculty of Medicine, Saarland University, Homburg (Saar), Germany

E-mail: Roy.Lancaster@structural-biology.eu

The genomes from both of the ϵ -proteobacteria *Wolinella succinogenes* and *Campylobacter jejuni* contain operons (*sdhABE* operons) that encode for hitherto uncharacterized enzyme complexes annotated as 'non-classical' succinate dehydrogenases. In the framework of a functional genomics project, a genetic system has been established for the homologous (over-)production and manipulation of the SdhABE complex from *W. succinogenes*. The catalytic properties of the purified enzyme were examined using various possible electron donor and acceptor substrates. Strikingly, for the SdhABE complex annotated as a 'succinate dehydrogenase', no succinate oxidation activity could be detected, neither with DCPIP, nor with methylene blue, nor with the high-potential quinone EQ-0 as electron donor. Although the complex catalyzes fumarate reduction with the menaquinol-6 analog 2,3-dimethyl-1,4-naphthoquinol (DMNH₂) the activities are very low. In addition to menaquinol-6, membranes of *C. jejuni* and of *W. succinogenes* contain a second quinol, 8-methylmenaquinol-6 (8-MMKH₂-6). Supplying an 8-MMKH₂-6 analog as a substrate increased specific quinol:fumarate reductase activity by about one order of magnitude. Furthermore, studies on variant enzymes demonstrated that the hydrophilic subunits of the complex are, in contrast to all other members of the superfamily, exported into the periplasm via the *tat*-pathway. Our studies reveal that the putative succinate dehydrogenase is in fact a novel periplasmic 8-methylmenaquinol:fumarate reductase with no detectable succinate dehydrogenase activity. These results provide an explanation for apparently puzzling previously published observations on the regulation of the *C. jejuni sdhABE* operon.

doi:10.1016/j.bbabi.2008.05.365