



King Saud University
Saudi Journal of Biological Sciences

www.ksu.edu.sa
www.sciencedirect.com



ORIGINAL ARTICLE

Selective inhibition of toxic cyanobacteria by β -carboline-containing bacterium *Bacillus flexus* isolated from Saudi freshwaters



Saad A. Alamri ^{a,*}, Zakaria A. Mohamed ^b

^a King Khalid University, Faculty of Science, Biological Science Department, P.O. Box 10255, Abha 61321, Saudi Arabia

^b Sohag University, Faculty of Science, Botany Department, Sohag 82524, Egypt

Received 9 January 2013; revised 15 March 2013; accepted 3 April 2013

Available online 18 April 2013

KEYWORDS

Antialgal bacteria;
Bacillus flexus;
Cyanobacteria;
Harmine;
Norharmane

Abstract A bacterial strain SSZ01 isolated from a eutrophic lake in Saudi Arabia dominated by cyanobacterial blooms, showed an antialgal activity against cyanobacteria species. Based on the analysis of the 16S rDNA gene sequence, the isolated strain (SSZ01) most likely belonged to the genus *Bacillus* with a 99% similarity to *Bacillus flexus* strain EMGA5. The thin layer chromatography (TLC) analysis of the ethyl acetate extract of this bacterium revealed that this strain can produce harmine and norharmane compared to different β -carboline analog standards. Harmine and norharmane were also detected in considerable amounts in bacterial growth medium, indicating a potential excretion of these compounds into the aquatic environment. The crude extract of *Bacillus flexus* as well as pure materials of harmine and norharmane inhibited the growth of tested species of cyanobacteria. However, the bacterial crude extract has a higher toxicity against tested species of cyanobacteria than harmine and norharmane. In addition, harmine was more toxic to cyanobacteria than norharmane. On the other hand, neither pure compounds of harmine and norharmane nor crude bacterial extract showed any antialgal activity against tested species of green algae. The results of the present study suggest that *B. flexus* SSZ01 or its crude extract containing harmine and norharmane could be a candidate for the selective control of cyanobacterial blooms without affecting other algal species.

© 2013 Production and hosting by Elsevier B.V. on behalf of King Saud University.

1. Introduction

Cyanobacteria usually dominate in eutrophic lakes and cause environmental problems due to the production of malodorous compounds and toxins (Oliver and Ganf, 2000). Cyanobacteria blooms are also responsible for wildlife and human health hazards, causing economic losses on fisheries, aquaculture and recreational activities (Codd, 1999; Anderson et al., 2002; Hallegraeff, 2003). Cyanobacteria may produce hepatotoxins

* Corresponding author. Tel.: +966 569010654; fax: +966 72417852.

E-mail address: amri555@yahoo.com (S.A. Alamri).

Peer review under responsibility of King Saud University.



Production and hosting by Elsevier

(microcystins, nodularins and cylindrospermopsins), neurotoxins (anatoxins and saxitoxins), cytotoxins and irritant compounds (lipopolysaccharides (LPS)) (Codd, 1999; Long and Carmichael, 2003; Apeldoorn et al., 2007). Some of these toxins can act as cancer promoters (Falconer et al., 1999; Kuiper-Goodman et al., 1999). In addition, cyanobacterial toxins significantly affect the growth of plants and accumulate in their edible tissues (Mohamed and Al-Shehri, 2009) providing an additional route of human and animal exposure to these toxins.

Much attention has been given in recent decades to strategies for cyanobacteria bloom control and management. Various chemical or synthetic agents (e.g., copper, chlorine, aluminum, calcium, and potassium permanganate) are used to control nuisance phytoplankton in aquatic ecosystems (Lam et al., 1995; Falconer et al., 1999). However, these algacides often induce the release of phytotoxins, which threaten drinking water supplies, accumulate and persist in the environment, and are toxic to fish (Lam et al., 1995; Karan et al., 1998; Boyd and Massaut, 1999; Meepagala et al., 2005). Therefore, biological control agents such as viruses (Garry et al., 1998), bacteria (Imai et al., 1995; Park et al., 2006a), and protozoa (Kim et al., 2009) are of particular interest.

Research into the relationship between bacteria and algae has resulted in the isolation of several strains of bacteria capable of inhibiting or killing harmful algal bloom species (Lovejoy et al., 1998; Yoshinaga et al., 1998; Amaro et al., 2005; Su et al., 2007). This is because these bacteria may contain natural compounds acting as algacides against algae and cyanobacteria (Kodani et al., 2002). Such algacides are likely to be specific and biodegradable, and may therefore offer an environmentally friendly method for control of algal blooms (Park et al., 2006a,b). Among antialgal substances isolated from bacteria are β -carboline (e.g. harmine, norharmine and norharmaline) (Kodani et al., 2002; Volk, 2005, 2006; Volk and Ferkert, 2006; Volk and Mundt, 2007).

In this study, different strains of bacteria, isolated from phytoplankton samples collected from a eutrophic lake in Saudi Arabia, were screened for their antialgal activity. The strongest antialgal bacterium was checked for the presence of β -carbolines. In addition, the current study investigates the effect of β -carboline compounds, isolated from such an antialgal bacterium, on the selective inhibition of cyanobacterial growth.

2. Materials and methods

2.1. Tested algae

The tested cyanobacteria (*Merismopedia tenuissima*, *Oscillatoria limnetica*) and green algae (*Chorella vulgaris*, *Ankistrodesmus falcatus*) used in the antialgal activity experiments, were isolated from the same place (Tendaha Lake), where the antialgal bacteria were isolated. All algal species were grown and maintained in BG-11 medium (Stanier et al., 1971) under the conditions of 25 °C and illumination of 30 $\mu\text{mol photon m}^{-2} \text{s}^{-1}$.

2.2. Isolation and screening of antialgal bacteria

Surface water samples were collected during May 2011 from Tendaha Lake, southwest of Saudi Arabia. Samples were serially diluted with sterile distilled water. One hundred μl aliquots

of each dilution were spread onto Nutrient Broth (NB) Agar (Difco) medium plates, containing 0.3% Beef Extract, 0.5% Peptone, 0.5% NaCl and 1.5% Agar, followed by incubation for 3 d at 30 °C. Colonies with different colony color and morphological shape were chosen for isolation. As a result, three different strains were isolated in this study. To test the antialgal activity of bacterial strains, a small sample of each colony was inoculated on plates of BG-11 medium containing a tested algal species. The plates were incubated for 7 d at 25 °C under illumination of 30 $\mu\text{mol photon m}^{-2} \text{s}^{-1}$. A bacterial strain exhibiting antialgal activity against either species of cyanobacteria or eukaryotic algae was selected for further study.

Antialgal bacteria were inoculated and precultured into three sterilized tissue culture tubes containing 10 ml of liquid NB at 30 °C, at 200 rpm for 48 h. These 10 ml cultures were then transferred into three 250 ml Erlenmeyer flasks containing 10 ml of liquid NB. Flasks were incubated for 3 d at 30 °C and 200 rpm. Bacterial cells were then removed by centrifugation (5,000g for 20 min at 4 °C) and filtration (0.2 μm pore-size membrane filter). Both the pellets and supernatants were ultrasonicated and extracted separately with ethyl acetate. Subsequently, the ethyl acetate layer was evaporated and stored at -20 °C until the next procedure.

The analysis of β -carboline compounds in bacterial extract was performed according to Kodani et al. (2002) using thin layer chromatography (TLC). Briefly, the ethyl acetate extracts were subjected to silica gel chromatography TLC (Kieselgel 60F254, 20 \times 20 cm, Merck) and eluted with chloroform:methanol (9:1) according to Kodani et al. (2002). Ten μl of test solutions (cell extract and medium extract) and reference solutions (harmine, harmine and norharmine) with concentration of 200 $\mu\text{g ml}^{-1}$ methanol were applied to the plates. β -carboline compounds (harmine, harmine and norharmine) were noticed in UV light (UV 254 nm). TLC plates were scanned, digitized and analyzed by UN-scan-it gel scanning software (Silk Scientific, Orem, Utah).

2.3. Growth inhibition experiments

As the yield of harmine and norharmine isolated from *Bacillus flexus* SSZ01 during the present study was very low, it was decided to use harmine and norharmine obtained from Fluka and Sigma companies, respectively. Different concentrations of chemical harmine and norharmine (1, 10, 20, 30 $\mu\text{g ml}^{-1}$), as well as crude bacillus extract containing the same concentrations of harmine and norharmine were added to each tested algal suspension in BG-11 medium (25 ml) in 50 ml Erlenmeyer flasks. The flasks were incubated at the same above conditions of algal growth for 7 d. The numbers of surviving cells were counted directly on a hemacytometer at a magnification of $\times 400$. The antialgal activity of antialgal bacterium was calculated using the following equation (Kim et al., 2009).

$$2.4. \text{Antialgal activity (\%)} = (1 - Tt/Ct) \times 100$$

where Tt and Ct are the cell numbers of treated and control cultures, respectively, of tested species. Algal cultures with no treatments were used as control. All experiments were carried out in triplicate, and results are given as the mean and standard deviation of raw data. The IC₅₀ values (i.e. the concentrations of tested compounds that inhibited growth (cell

numbers) by 50% relative to the controls) were calculated by those obtained from probity regression analysis.

2.5. Identification and sequence analysis of 16S rRNA gene

The antialgal bacterium was identified on the basis of an analysis of 16S rRNA gene sequences as reported in the previous study (Alamri, 2012).

2.6. Statistical analysis

Statistical analyses were performed using the Statistical Package for the Social Sciences for Windows (SPSS, version 10.0, Chicago, IL, USA). Data are presented as means with their standard deviation. Statistical evaluation of the data was performed using a one-way ANOVA test. Values were considered statistically significant when $p \leq 0.05$.

3. Results

3.1. Screening of antialgal bacteria

The preliminary screening based on agar diffusion assay for antialgal activity of three morphologically different strains of




Figure 1 TLC of standards of harmine (1), harmine (2), and norharmane (3), and of ethyl acetate extracts of *Bacillus flexus* cells (4) and growth medium (5).

bacteria isolated from phytoplankton samples collected from a eutrophic lake in Saudi Arabia, showed that only one strain (SSZ01) had antialgal activity against tested species of cyanobacteria. The bacterial 16S rRNA gene sequences showed close relationships between the strain SSZ01 and the genus *Bacillus*. The highest sequence similarity value (99%) was obtained between strain SSZ01 and *B. flexus* strain EMGA5 (Genbank accession number, EU602312). Thus, this strain was designated to be *B. flexus* and deposited in the Genbank with an accession number of GU112451.

The TLC analysis of ethyl acetate extract of antialgal bacterium (*B. flexus*), revealed that both bacterial cells and NB medium contained the β -carboline derivatives, harmine and norharmane with different concentrations compared to standard β -carbolins (Fig. 1). By digitizing the scanned TLC plates, the results showed that the extract of antialgal bacterium cells contained 10 and 15 μg harmine and norharmane ml^{-1} , respectively while growth medium of this bacterium contained 5 and 8 μg ml^{-1} harmine and norharmane, respectively.

3.2. Growth inhibition of algae β -carboline compounds produced by bacteria

The results of 7 d exposure of cyanobacteria and green algae to harmine and/or norharmane in a liquid medium showed that the crude extract of *B. flexus* cells containing harmine and norharmane inhibited the growth of cyanobacteria (*M. tenuissima* and *O. limnetica*) as determined by cell number (Fig. 2). Although there was no significant variation in the growth as affected by this extract between these two species, *Merismopedia* was more sensitive than *Oscillatoria*. IC_{50} values of bacterial crude extract varied substantially between the two species ($p < 0.05$), i.e. IC_{50} value for *Merismopedia* was less than that of *Oscillatoria* (Table 1).

The pure materials of either harmine or norharmane were less toxic to tested cyanobacteria species than crude bacterial extract containing harmine and norharmane. These β -carboline analogs decreased the cell number and chl.a contents in treated cultures of tested cyanobacteria compared to controls




Figure 2 Effect of ethyl acetate extract of *Bacillus flexus* on the cell number of tested species of cyanobacteria and green algae.

Table 1 IC₅₀ values ($\mu\text{g ml}^{-1}$) obtained for crude extract of antialgal *Bacillus* sp., pure harmine and norharmane toward cyanobacteria and green algae species screened in this study.

Tested algae	IC ₅₀ ($\mu\text{g ml}^{-1}$)		
	Crude extract	Harmine	Norharmane
<i>M. tenuisssima</i>	5.3 ± 0.9	11.62 ± 1.7	13.2 ± 1.2
<i>O. limnetica</i>	9.6 ± 1.1	11.25 ± 1.3	12.5 ± 1.4
<i>A. falcatus</i>	–	–	–
<i>C. vulgaris</i>	–	–	–

(Figs. 3 and 4). In contrast to crude extract, IC₅₀ values of either harmine or norharmane did not differ significantly between the two cyanobacteria species (*M. tenuisssima* and *O.*

limnetica) (Table 1). However, harmine was more toxic to cyanobacteria than norharmane (as shown by higher IC₅₀ values). Furthermore, the antialgal activities of either crude bacterial extract or pure harmine/norharmane toward tested cyanobacteria increased with increasing the concentration of antialgal substance. The strongest antialgal activity for both tested species of cyanobacteria was obtained at a concentration of 30 $\mu\text{g ml}^{-1}$ (Table 2). However, the antialgal activities differed significantly among crude extract, harmine and norharmane for both species of cyanobacteria at the lowest concentrations only (1, 10 $\mu\text{g ml}^{-1}$) ($p < 0.05$), but no significant variation in these activities was observed at the highest concentrations (20, 30 $\mu\text{g ml}^{-1}$) used during the present study. In contrast to cyanobacteria, green algae (*A. falcatus* and *C. vulgaris*) had not been affected by either bacterial crude extract or pure




Figure 3 Effect of a pure harmine of *Bacillus flexus* on the cell number of tested species of cyanobacteria and green algae.




Figure 4 Effect of a pure norharmane of *Bacillus flexus* on the cell number of tested species of cyanobacteria and green algae.

Table 2 Antialgal activity (%) obtained for different concentrations ($\mu\text{g ml}^{-1}$) of crude extract of antialgal *Bacillus flexus*, pure harmine and norharmine toward cyanobacteria species screened in this study.

Tested species/antialgal concentrations	Antialgal activity (%)		
	Crude extract	Harmine	Norharmine
<i>M. tenuisissima</i>			
0	0	0	0
1	38.5	15.4	4.6
10	61.5	46.2	41.6
20	86.2	83.4	83.1
30	100	96.9	98.4
<i>O. limnetica</i>			
0	0	0	0
1	45.6	24.6	3.5
10	77.2	59.6	64.9
20	96.5	86	94.7
30	100	98.2	98.2

β -carboline compounds (harmine, norharmine). The cell number of tested species of green algae did not differ significantly between treated and control cultures (Figs. 3 and 4).

4. Discussion

Bacteria with antialgal activities were reported to belong to *Cytophaga-Flavobacteria-Bacteroides* CFB group (50%) and γ -*Proteobacteria* (45%), while the remaining strains represent the Gram-positive genera *Micrococcus*, *Bacillus* and *Planomicrobium* (Fukuyo et al., 2002; Mayali and Azam, 2004; Hare et al., 2005). Some *Bacillus* strains were reported to completely inhibit the growth of bloom-forming cyanobacteria and green algae in highly eutrophic lakes (Ahn et al., 2003; Mu et al., 2007). In the present study, *B. flexus* was isolated as bacterial strain with an antialgal activity against cyanobacteria, from a Saudi eutrophic pond containing toxic cyanobacterial blooms.

Antialgal bacteria may inhibit the algal growth effectively through direct attack that is required for cell-to-cell contact (Imai et al., 1993) or indirect attack by the production of extracellular compounds (Fukami et al., 1992; Imai et al., 1993; Wang et al., 2005). In this study, *B. flexus* strain might inhibit the growth of cyanobacteria indirectly by the production of two β -carboline derivatives (harmine and norharmine). β -carboline derivatives have been described to occur naturally in some higher plants (Allen and Holmstedt, 1980; Bourke et al., 1992; Cheng and Mitchelson, 1997), and in Indonesian sponge (Rao et al., 2003). Recently, these compounds were isolated and identified in some species of fresh and marine cyanobacteria (Volk, 2005, 2006, 2008; Volk and Furkert, 2006; Volk and Mundt, 2007; Mohamed, 2013), but their production in bacteria was confined to a few species including the gliding bacterium *Myxobacter* (Böhendorf et al., 1996), *Pseudomonas* (Kodani et al., 2002) *Pseudoalteromonas piscicida* (Zheng et al., 2005) and *Enterococcus faecium* (Aassila et al., 2003).

The bacterial crude extract containing the β -carbolines, harmine and norharmine, showed antagonistic activity against cyanobacteria, affecting the cell division in these species. These

results confirmed the results of previous studies reporting the inhibition of many species of cyanobacteria by norharmine and other related β -carbolines. Kodani et al. (2002) reported the antialgal activity of β -carbolines, harmine and norharmine, produced by *Pseudomonas* sp. against the cyanobacteria *Anabaena cylindrica*, *A. variabilis*, *Anacystis marina*, *Microcystis aeruginosa*, *M. viridis* and *Oscillatoria agardhii* at a concentration of $30 \mu\text{g disk}^{-1}$. Volk (2006) also investigated the activity of three β -carbolines harmine, norharmine and norharmalane toward some species of cyanobacteria (*Arthrospira laxissima*, *Chroococcus minutus*, *Nostoc carneum*, *Nostoc insulare*, *Synechocystis aquatilis*, *Synechococcus* species), and he found that all these compounds were cytotoxic against the cyanobacterial test organisms in low quantities (0.5–18.0 μg).

The results of the present study showed that both crude bacterial extracts containing harmine, norharmine as well as synthetic harmine and norharmine exhibited an antialgal activity against cyanobacteria species at different IC_{50} s. The crude extract was more toxic to cyanobacteria ($\text{IC}_{50} = 5.3\text{--}9.6 \mu\text{g ml}^{-1}$) than harmine and norharmine ($\text{IC}_{50} = 11.25\text{--}13.2 \mu\text{g ml}^{-1}$). The high toxicity of crude extract may be due to the synergistic effect of both harmine and norharmine present together in the extract. Previously, it has been stated that the inhibitory effect of algicides is likely to be a synergistic effect of various compounds (Ball et al., 2001; Ferrier et al., 2005). On the other hand, the results of this study revealed differences in the antialgal activity between harmine and norharmine toward tested species of cyanobacteria; where harmine was of a significant higher toxicity than norharmine. These results are in conformity with those obtained by Volk (2006) reporting differences in the activity of norharmine and harmine (the 1-methyl-derivate of norharmine) toward the cyanobacteria, *Arthrospira laxissima* and *Synechococcus* sp., harmine was of a significant higher toxicity than norharmine. Based on the hypothesis of Volk (2006) that the hydrophobic alkaloid bases could easily penetrate the external plasma membrane of cells than the hydrophilic ones, a higher penetration rate of the more hydrophobic harmine could also explain the higher activity of this β -carboline in comparison to the less hydrophobic and unmethylated norharmine.

In contrast to cyanobacteria, the tested species of green algae for antialgal activity of harmine and norharmine during the present study did not show any growth inhibition toward these compounds. Accordingly, previous studies showed that β -carbolines have antialgal activity toward cyanobacteria with no remarkable effect on green algae (Kodani et al., 2002). This finding indicates that these β -carboline compounds have selective inhibition of cyanobacterial growth rather than green algae. The potential for selectively controlling cyanobacteria without affecting other algal species could be useful for the preservation of ecosystems with the least impact on the environment.

The results of the present study also showed that *B. flexus* not only contains the β -carbolines, harmine and norharmine, within the cells, but also can release considerable amounts of these compounds into culture medium. This study is not the first to demonstrate that bacteria and cyanobacteria can excrete such antialgal metabolites into the culture media. Previous studies detected β -carboline derivatives in culture media of bacteria (Kodani et al., 2002) and cyanobacteria (Volk, 2005, 2006, 2008; Volk and Furkert, 2006). These results indicate the potential excretion of these antialgal compounds into

the aquatic environment and thus function naturally as allelopathic substances.

5. Conclusion

This study is the first to report that *B. flexus* can produce anti-algal β -carboline (harmine and norharmine). This bacterium and its β -carboline inhibited the growth of cyanobacteria only, but not green algae, which is an advantage of bacterial agents in removing undesirable algal species but providing niches for other beneficial species to colonize and thrive.

Acknowledgment

The author would like to acknowledge and thank King Khalid University for supporting this study.

References

- Aassila, H., Bourguet-Kondracki, M.L., Rifai, S., Fassouane, A., Guyot, M., 2003. Identification of harmine as the antibiotic compound produced by a tunicate-associated bacterium. *Marine Biotechnol.* 5, 163–166.
- Ahn, C.Y., Joung, S.H., Jeon, J.W., Kim, H.S., Yoon, B.D., Oh, H.M., 2003. Selective control of cyanobacteria by surfactin-containing culture broth of *Bacillus subtilis* C1. *Biotechnol. Lett.* 25, 1137–1142.
- Alamri, S., 2012. Biodegradation of microcystin-RR by *Bacillus flexus* isolated from a Saudi freshwater lake. *Saudi J. Biol. Sci.* 19, 435–440.
- Allen, J.R., Holmstedt, B.R., 1980. The simple-carboline alkaloids. *Phytochem.* 19, 1573–1582.
- Amaro, A.M., Fuentes, M.S., Ogalde, S.R., Venegas, J.A., Suarez-Isla, B.A., 2005. Identification and characterization of potentially algalytic marine bacteria strongly associated with the toxic dinoflagellate *Alexandrium catenella*. *J. Eukaryot. Microbiol.* 52, 191–200.
- Anderson, D., Glibert, P., Burkholder, J., 2002. Harmful algal blooms and eutrophication: nutrient sources, composition, and consequences. *Estuaries* 25, 704–726.
- Apeldoorn, M., Egmond, H., Speijers, G., Bakker, G., 2007. Toxins of cyanobacteria. *Mol. Nutr. Food Res.* 51, 7–60.
- Ball, A., Williams, M., Vincent, D., Robinson, J., 2001. Algal growth control by a barley straw extract. *Bioresour. Technol.* 77, 177–181.
- Böhlendorf, B., Bedorf, N., Jansen, R., Trowitzsch-Kienast, W., Höfle, G., Forche, E., Gerth, K., Irschik, H., Kunze, B., Reichenbach, H., 1996. Antibiotics from gliding bacteria, LXXIII indole and quinoline derivatives as metabolites of tryptophan in myxobacteria. *Liebigs Ann.* 1, 49–53.
- Bourke, C.A., Stevens, G.R., Carrigan, M.J., 1992. Locomotor effects in sheep of alkaloids identified in Australia *Tribulus terrestris*. *Aust. Vet. J.* 69, 163–165.
- Boyd, C., Massaut, L., 1999. Risks associated with the use of chemicals in pond aquaculture. *Aquaculture Eng.* 20, 113–132.
- Cheng, J., Mitchelson, K.R., 1997. Improved separation of six harmine alkaloids by high-performance capillary electrophoresis. *J. Chromatogr.* 761, 297–305.
- Codd, G.A., 1999. Cyanobacterial toxins: their occurrence in aquatic environments and significance to health. *Bull. Inst. Oceanogr. Monaco* 19, 483–500.
- Falconer, I., Bartram, J., Chorus, I., Kuiper-Goodman, T., Utkilen, H., Burch, M., Codd, G., 1999. Safe levels and safe practices. In: Chorus, I., Bartram, J. (Eds.), *Toxic Cyanobacteria in Water: a Guide to their Public Health Consequences, Monitoring and Management*. WHO/ Spon, London, pp. 155–178.
- Ferrier, M., Butler, B., Terlizzi, D., Lacouture, R., 2005. The effects of barley straw (*Hordeum vulgare*) on the growth of freshwater algae. *Bioresour. Technol.* 96, 1788–1795.
- Fukami, K., Yuzawa, A., Nishijima, T., Hata, Y., 1992. Isolation and properties of a bacterium inhibiting the growth of *Gymnodinium nagasakiense*. *Nippon Suisan Gakkai-shi* 58, 1073–1077.
- Fukuyo, Y., Imai, I., Kodama, M., Tamai, K., 2002. Red tides and other harmful algal blooms in Japan. In: Taylor, F.J., Trainer, V.L. (Eds.), *Harmful Algal Blooms in the PICES Region of the North Pacific*. North Pacific Marine Science Organization (PICES), Sidney, Canada, pp. 7–20.
- Garry, R.T., Hearing, P., Cosper, E.M., 1998. Characterization of a lytic virus infectious to the bloom-forming microalga *Aureococcus anophagefferens* (Pelagophyceae). *J. Phycol.* 34, 616–621.
- Hallegraeff, G.M., 2003. Harmful algal blooms: a global overview. In: Hallegraeff, G.M., Anderson, D.M., Cembella, A.D. (Eds.), *Manual on Harmful Marine Microalgae*. UNESCO Publishing, Paris, pp. 25–49.
- Hare, C.E., Demir, E., Coyne, K.J., Cary, S.C., Kirchman, D.L., Hutchins, D.A., 2005. A bacterium that inhibits the growth of *Pfiesteria piscicida* and other dinoflagellates. *Harmful Algae* 4, 221–234.
- Imai, I., Ishida, Y., Hata, Y., 1993. Killing of marine phytoplankton by a gliding bacterium *Cytophaga* sp., isolated from the coastal sea of Japan. *Marine Biol.* 116, 527–532.
- Imai, I., Ishida, Y., Sakaguchi, K., Hata, Y., 1995. Antialgal marine-bacteria isolated from northern Hiroshima Bay. *Jpn. Fish. Sci.* 61, 628–636.
- Karan, V., Vitorovic, S., Tutundzic, V., Poleksic, V., 1998. Functional enzymes activity and gill histology of carp after copper sulfate exposure and recovery. *Ecotoxicol. Environ. Saf.* 40, 49–55.
- Kim, Y.S., Lee, D.S., Jeong, S.Y., Lee, W.J., Lee, M.S., 2009. Isolation and characterization of a marine anti-algal bacterium against the harmful raphidophyceae *Chattonella marina*. *J. Microbiol.* 47, 9–18.
- Kodani, S., Imoto, A., Mitsutani, A., Murakami, M., 2002. Isolation and identification of the anti-algal compound, harmine (1-methyl- β -carboline), produced by the anti-algal bacterium, *Pseudomonas* sp. K44-1. *J. Appl. Phycol.* 14, 109–114.
- Kuiper-Goodman, T., Falconer, I., Fitzgerald, J., 1999. Human health aspects. In: Chorus, I., Bartram, J. (Eds.), *Toxic Cyanobacteria in Water: A Guide to their Public Health Consequences, Monitoring and Management*. WHO/Spon, London, pp. 113–153.
- Lam, A.K., Prepas, E.E., Spink, D., Hruday, S.E., 1995. Chemical control of hepatotoxic phytoplankton blooms: implications for human health. *Water Res.* 29, 1845–1854.
- Long, B.M., Carmichael, W., 2003. Marine cyanobacterial toxins. In: Hallegraeff, G.M., Anderson, D.M., Cembella, A.D. (Eds.), *Manual on Harmful Marine Microalgae*. UNESCO, Venice, pp. 279–296.
- Lovejoy, C., Bowman, J.P., Hallegraeff, G.M., 1998. Antialgal effects of a novel marine *Pseudoalteromonas* isolate (class Proteobacteria, gamma subdivision) on harmful algal bloom species of the genera *Chattonella*, *Gymnodinium*, and *Heterosigma*. *Appl. Environ. Microbiol.* 64, 2806–2813.
- Mayali, X., Azam, F., 2004. Antialgal bacteria in the sea and their impact on algal blooms. *J. Eukaryot. Microbiol.* 51, 139–144.
- Meepagala, K., Schader, K., Wedge, D., Duke, S., 2005. Antialgal and antifungal compounds from the roots of *Ruta graveolens* and synthesis of their analogs. *Phytochemical* 66, 2689–2695.
- Mohamed, Z.A., 2013. Allelopathic activity of the norharmine-producing cyanobacterium *Synechocystis aquatilis* against cyanobacteria and microalgae. *Oceanol. Hydrobiol. Stud.* 42, 1–7.
- Mohamed, Z.A., Al-Shehri, A.M., 2009. Microcystins in groundwater wells and their accumulation in vegetable plants irrigated with contaminated waters in Saudi Arabia. *J. Hazard. Mater.* 172, 310–315.

- Mu, R.M., Fan, Z.Q., Pei, H.Y., Yuan, X.L., Liu, S.X., Wang, X.R., 2007. Isolation and algae-lysing characteristics of the antialgal bacterium B5. *J. Environ. Sci.* 19, 1336–1340.
- Oliver, R.L., Ganf, G., 2000. Freshwater blooms. In: Whitton, B.A., Potts, M. (Eds.), *The Ecology of Cyanobacteria: their Diversity in Time and Space*. Kluwer Academic Publishers, Dordrecht, pp. 149–194.
- Park, M.H., Han, M.S., Ahn, C.Y., Kim, H.S., Yoon, B.D., Oh, H.M., 2006a. Growth inhibition of bloom-forming cyanobacterium *Microcystis aeruginosa* by rice straw extract. *Lett. Appl. Microbiol.* 43, 307–312.
- Park, M.H., Hwang, S.J., Ahn, C.Y., Kim, B.H., Oh, H.M., 2006b. Screening of seventeen oak extracts for the growth inhibition of the cyanobacterium *Microcystis aeruginosa* Kutz. em. Elenkin. *Bull. Environ. Contam. Toxicol.* 77, 9–14.
- Rao, K.V., Santarsiero, B.D., Mesecar, A.D., Schinazi, R.F., Tekwani, B.L., Hamann, M.T., 2003. New manzamine alkaloids with activity against infectious and tropical parasitic diseases from an Indonesian sponge. *J. Nat. Prod.* 66, 823–828.
- Stanier, R.Y., Kunisawa, R., Mandel, M., Cohen-Baziere, G., 1971. Purification and properties of unicellular blue-green algae (order Chroococcales). *Bacteriol. Rev.* 35, 171–205.
- Su, J., Yang, X., Zheng, T., Tian, Y., Jiao, N., Cai, L., Hong, H., 2007. Isolation and characterization of marine antialgal bacterium against the toxic dinoflagellate *Alexandrium tamarense*. *Harmful Algae* 6, 799–810.
- Volk, R., 2005. Screening of microalgal culture media for the presence of antialgal compounds and isolation and identification of two bioactive metabolites excreted by the cyanobacteria *Nostoc insulare* and *Nodularia harveyana*. *J. Appl. Phycol.* 17, 339–347.
- Volk, R., 2006. Antialgal activity of several cyanobacterial exometabolites. *J. Appl. Phycol.* 18, 145–151.
- Volk, R., 2008. Screening of microalgae for species excreting norharmane a manifold biologically active indole alkaloid. *Microbiol. Res.* 163, 307–313.
- Volk, R., Furkert, F., 2006. Antialgal, antibacterial and antifungal activity of two metabolites produced and excreted by cyanobacteria during growth. *Microbiol. Res.* 161, 180–186.
- Volk, R., Mundt, S., 2007. Cytotoxic and non-cytotoxic exometabolites of the cyanobacterium *Nostoc insulare*. *J. Appl. Phycol.* 19, 55–62.
- Wang, X.L., Gong, L.Y., Liang, S.K., Han, X.R., Zhu, C.J., Li, Y.B., 2005. Antialgal activity of rhamnolipid biosurfactants produced by *Pseudomonas aeruginosa*. *Harmful Algae* 4, 433–443.
- Yoshinaga, I., Kim, M.C., Katanozaka, N., Imai, I., Uchida, A., Ishida, Y., 1998. Population structure of antialgal marine bacteria targeting the red tide forming alga *Heterosigma akashiwo* (Raphidophyceae), determined by restriction fragment length polymorphism analysis of the bacterial 16S ribosomal RNA genes. *Mar. Ecol. Prog. Ser.* 170, 33–44.
- Zheng, L., Chen, H., Han, X., Lin, W., Yan, X., 2005. Antimicrobial screening and active compound isolation from marine bacterium NJ6-3-1 associated with the sponge *Hymeniacidon perleve*. *World J. Microbiol. Biotechnol.* 21, 201–206.