

Available online at www.sciencedirect.com

ScienceDirect

journal homepage: www.e-jds.com

ORIGINAL ARTICLE

The normal range of maximum mouth opening and its correlation with height or weight in the young adult Chinese population

Xiao-Yan Li ^{a*}, Cheng Jia ^a, Zi-Chuan Zhang ^b^a Department of Stomatology, The Third Hospital of Hebei Medical University, Shijiazhuang, P.R. China^b Department of Stomatology, Bethune International Peace Hospital, Shijiazhuang, P.R. China

Received 21 June 2016; Final revision received 11 September 2016

Available online ■ ■ ■

KEYWORDS

adults;
Chinese;
height;
maximum mouth
opening;
weight

Abstract *Background/purpose:* Maximum mouth opening (MMO) is an important diagnostic reference for dental clinicians. However, the relationship between either body height or weight of the individual and their subsequent MMO has, up to now, been unclear. The purpose of this study was to measure the MMO of healthy young Chinese adults and to analyze the possible correlation of MMO with either height or weight.

Materials and methods: A total of 452 young Chinese adults, aged 20–35 years (238 males, 214 females) were selected for this cross-sectional study. We recorded the MMO, age, sex, height, and weight of the participants. Two standardized examiners performed the clinical oral assessments. Independent sample *t* tests were used to examine the difference in MMO relative to sex. Pearson's correlation and simple linear regression were used to estimate the correlation between MMO and either height or weight.

Results: The average MMO across the 452 participants was 52.02 ± 5.09 mm, and the average MMO of males (54.18 ± 5.21 mm) was significantly larger than that of females (49.62 ± 3.69 mm; $P < 0.001$). The mean MMO was moderately positively correlated with height ($r = 0.54$; $P < 0.001$) and weight ($r = 0.50$; $P < 0.001$). In the regression model, it was estimated that, for every 10 cm or 10 kg, MMO increased by about 3.6 mm or 1.8 mm, respectively. *Conclusion:* With the limits of the present study, both height and weight were found to be significantly correlated with the MMO of Chinese young adults and may be significant predictors of MMO measurement.

Copyright © 2016, Association for Dental Sciences of the Republic of China. Published by Elsevier Taiwan LLC. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

* Corresponding author. Department of Stomatology, The Third Hospital of Hebei Medical University, 139 Zi-Qiang Road, Shijiazhuang 050051, P.R. China.

E-mail address: lixiaoyan.1208@163.com (X.-Y. Li).

<http://dx.doi.org/10.1016/j.jds.2016.09.002>

1991-7902/Copyright © 2016, Association for Dental Sciences of the Republic of China. Published by Elsevier Taiwan LLC. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Please cite this article in press as: Li X-Y, et al., The normal range of maximum mouth opening and its correlation with height or weight in the young adult Chinese population, Journal of Dental Sciences (2016), <http://dx.doi.org/10.1016/j.jds.2016.09.002>

Introduction

Mandibular function is evaluated using a series of diagnostic tests, including palpation of the masticatory muscles and temporomandibular joint (TMJ), occlusal evaluation, and radiographic examination.¹ Maximum mouth opening (MMO) is also an important diagnostic reference for dental clinicians as a preliminary evaluation. Limited mouth opening during mandibular movements may result from a temporomandibular joint disorder (TMD), oral submucous fibrosis, rheumatic disease, infection, malignancies, or facial trauma.^{2,3}

Establishing a normal range for MMO could allow dental clinicians to objectively evaluate the treatment effects and set therapeutic goals for patients performing mandibular functional exercises. Previous studies have attempted to compile an average range of physiological mouth opening capacity measurements, but the variability in MMO is quite large because it varies dramatically with age, sex, race, joint condition, mandibular size, cranial base size, body height, and weight.^{4–11} There is, as yet, no reference for clinicians to determine whether a patient has limited mouth opening capacity, as it is so dependent on other particular physiological characteristics. It is therefore necessary to investigate the correlation between MMO and any related physiological indicators.

The correlation between age and MMO in ethnic Chinese adults has previously been studied and published in the literature.⁴ However, no correlation has been established between MMO and either body height or weight. The objective of the present study was to establish a normal range of MMO in a sample of young Chinese adults and to analyze possible correlations of MMO with either height or weight.

Materials and methods

The study included 452 young ethnic Chinese adults, aged 20–35 years (238 males, 214 females), who were undergoing regular physical examinations in the Medical Examination Center of the Third Hospital of Hebei Medical University (Shijiazhuang, China). A dental examination was performed for volunteers meeting the following inclusion criteria: (1) generally healthy; (2) had signed the informed consent; (3) had full permanent dentitions and no dental prosthesis; (4) no history of oral submucous fibrosis; (5) no history of head or neck tumors; (6) no history of jaw or face pain, either at rest or during activity; (7) no history of TMD or bruxism; (8) no history of TMJ, jaw, head, or face trauma; (9) no clinical symptoms of excessive dental attrition (2nd and 3rd degrees of severity); and (10) no Class III malocclusion (anterior crossbite). The Ethics Committee of the Bethune International Peace Hospital (Protocol No. 2014-12-8) approved the protocol of this study, and informed consent was obtained from each participant.

Age, sex, MMO, height, and weight were collected from each participant. The methods for height and weight measurements have been previously reported in detail.¹¹ Height and weight were both measured to the nearest tenth in centimeters and kilograms, respectively.

The MMO measurements were taken by two previously trained examiners and were then standardized ($\kappa > 0.8$). Prior to measuring MMO, the participants were asked to rest for at least 10 minutes. They were then seated comfortably in the dental chair in an upright and relaxed position, looking straight ahead. Each participant was instructed to open his or her mouth as wide as possible. The linear distance was measured between the mesioincisal edge of the upper right central incisor and the mesioincisal edge of the lower right central incisor, using a Boley gauge (Pearson Dental Suppliers Corporation, Sylmar, CA, USA). In order to ensure the accuracy and reproducibility of the results, each participant was repeatedly measured three times within 15 minutes. The mean value of the MMO readings was recorded as the outcome for all analyses.

The collected data were entered into spreadsheets in Microsoft Excel 2007 (Microsoft Corporation, Redmond, WA, USA) and imported into SPSS software (version 19.0; SPSS Incorporated, Chicago, IL, USA) for statistical analysis. An independent sample *t* test was used to examine the differences in MMO, relative to sex. Pearson's correlation and simple linear regression were used to estimate the correlations between MMO and either height or weight. A *P* value < 0.05 was considered statistically significant.

Results

The mean age of the 452 participants was 27.8 ± 4.2 years, and the average age of females (26.9 ± 4.2 years) was significantly younger than that of males (28.6 ± 3.9 years, $P < 0.001$). The average MMO for all participants was 52.02 ± 5.09 mm, and the mean MMO of males (54.18 ± 5.21 mm) was statistically larger than that of females (49.62 ± 3.69 mm; $t = 10.63$, $P < 0.001$).

There was a moderately positive correlation between MMO and height (Pearson's correlation coefficient $r = 0.54$; $P < 0.001$), as well as between MMO and weight ($r = 0.50$; $P < 0.001$). The tendency of MMO to increase with height or weight was obvious, as shown in Figures 1 and 2, displaying scatter and linear regression diagrams. In the regression model, it was estimated that for every 10 cm or 10 kg, MMO increased by about 3.6 mm or 1.8 mm, respectively. Regression equations were deduced by calculating the regression coefficient and intercept. For the 20–35-years age group, the regression equations for height and weight were: $\text{MMO (mm)} = 0.36 \times \text{height} - 10.15$ ($F = 180.37$, $P < 0.001$), and $\text{MMO (mm)} = 0.18 \times \text{weight} + 39.87$ ($F = 151.80$, $P < 0.001$), respectively.

Discussion

The values for a restricted mouth opening have generally been reported to be < 35 mm for joint-related disorders and < 40 mm for muscular disorders.¹² These parameters, however, are not applicable to all individuals, because MMO varies considerably from one individual to another. The MMO of healthy young Chinese adults in the present study was 52.02 ± 5.09 mm (54.18 ± 5.21 and 49.62 ± 3.69 mm for males and females, respectively). The tendency for the MMO of males to be significantly greater than that of females was seen consistently across studies. This may be




Figure 1 Scatter and linear regression diagrams of maximum mouth opening (MMO) associated with height.




Figure 2 Scatter and linear regression diagrams of maximum mouth opening (MMO) associated with weight.

because the anatomical structure of the male head and face bones are generally larger than those of females. Nevertheless, there were still some differences with the results of previous studies.

Differences across studies may result from several causes. First, participants with different racial backgrounds may have different facial proportions, which could make a significant difference in MMO.¹ Studies carried out in different countries and targeting different ethnic populations have shown that wide variability exists across racial groups. It has been reported that mean MMO ranges from 43.3 mm to 59.0 mm.¹⁰ Ying et al¹¹ also demonstrated that there was a positive correlation between race and MMO. Second, age is an important predictor of MMO measurement. MMO has been shown to steadily increase after birth and into adulthood, and then to gradually decrease as aging progresses, which may be explained by the

development of the articular eminence of the TMJ,¹¹ as well as by any morphologic changes related to bruxism and trauma.¹ This trend has also been established for ethnic Chinese adults.⁴ The mean MMO in this study was almost 1 mm larger than that in the 20–39-years age group, as reported by Yao et al,⁴ which may be attributable to the younger participants in this study. Third, methodological discrepancies could result in the variation in reported MMO values. MMO can be expressed either as the interincisal distance or as the interincisal distance plus the overbite. The former refers to the vertical distance between the upper central incisor to the lower central incisor of the same side, and the latter means the vertical distance traveled by the mandible. It is thought that interincisal distance, being the actual functional opening capacity of the mouth, is the more important measurement, because its value lies in its effect on chewing and dental

treatments.¹³ Therefore, the interincisal distance was used as the MMO measurement in this study. There are other measurement methods worth considering, as well. For example, Wood and Branco¹⁴ compared direct (intraoral) and indirect (extraoral) methods, concluding that the direct method was preferable, which is the reason why we chose the intraoral measurement for this study.

Head position is also an important influencing factor in determining MMO in healthy adults.^{15,16} The vertical mandibular opening significantly and successively decreased in the forward, neutral, then retracted head positions.¹⁵ In this study, therefore, all volunteers were placed in an upright and relaxed position, in order to eliminate the possible influence of different head and neck positions.

The correlation between MMO and either height or weight is controversial. It has been demonstrated that body height was strongly correlated with MMO,¹⁷ and Placko et al⁶ found that MMO was larger in tall patients, regardless of sex or age. Nevertheless, Ying et al¹¹ found a significant correlation between MMO and weight, but not between MMO and height. Gallagher et al⁸ and Reicheneder et al⁹ failed to find an association between weight and MMO. Chen et al¹⁸ examined the interincisal distance in 518 preschool children, ages 3–5 years, and investigated the factors correlated with MMO in that sample population. They found that MMO was correlated with weight and increased by 0.19 mm/increase in kg. In our study, a moderately positive correlation was found between MMO and height ($r = 0.54$) and between MMO and weight ($r = 0.50$).

There are two limitations to this study. First, participants were recruited from a pool of people who were undergoing regular medical examinations in our hospital. Most of these volunteers came from satellite cities and may or may not be representative of the province's young adults. Second, radiographic and magnetic resonance images of the TMJ were not collected or taken into consideration in the inclusion criteria, which may have resulted in the unintentional inclusion of some asymptomatic participants with limitations in mouth opening. However, the absence of any history of jaw pain and the lack of TMD or trauma history should have minimized the number of participants with undetected mouth opening anomalies.

Even within the limits of this study, we concluded that both height and weight have significant influences on the MMO of healthy young Chinese adults. An increased sample size and a wider range of ages should be examined in future research, including other underlying factors, such as the length, width, and angle of the mandible, in order to confirm and elaborate on these findings.

Conflicts of interest

The authors have no conflicts of interest relevant to this article.

References

- Juan FC, Carlo EM, Alejandro JC, et al. Clinical characterization of mouth opening among Mexican adolescents and young adults. *J Dent Sci* 2012;7:81–4.
- Fukui T, Tsuruta M, Murata K, Wakimoto Y, Tokiwa H, Kuwahara Y. Correlation between facial morphology, mouth opening ability, and condylar movement during opening–closing jaw movements in female adults with normal occlusion. *Eur J Orthod* 2002;24:327–36.
- Sari S, Kucukesmen C, Sonmez H. Evaluation of the applicability of temporomandibular opening index in Turkish children with and without signs and symptoms of temporomandibular joint disorders. *Cranio* 2008;26:197–201.
- Yao KT, Lin CC, Hung CH. Maximum mouth opening of ethnic Chinese in Taiwan. *J Dent Sci* 2009;4:40–4.
- Agerberg G. Maximal mandibular movements in young men and women. *Sven Tandlak Tidskr* 1974;67:81–100.
- Placko G, Bellot-Samson V, Brunet S, et al. Normal mouth opening in the adult French population. *Rev Stomatol Chir Maxillofac* 2005;106:267–71 [In French, English abstract].
- Khare N, Patil SB, Kale SM, Sumeet J, Sonali I, Sumeet B. Normal mouth opening in an adult Indian population. *J Maxillofac Oral Surg* 2012;11:309–13.
- Gallagher C, Gallagher V, Whelton H, Cronin M. The normal range of mouth opening in an Irish population. *J Oral Rehabil* 2004;31:110–6.
- Reicheneder C, Kardari Z, Proff P, Fanghaenel J, Faltermeier A, Römer P. Correlation of condylar kinematics in children with gender, facial type and weight. *Ann Anat* 2013;195:243–7.
- Zawawi KH, Al-Badawi EA, Lobo-Lobo S, Melis M, Mehta NR. An index for the measurement of normal maximum mouth opening. *J Can Dent Assoc* 2003;69:737–41.
- Ying QV, Bacic J, Abramowicz S, Sonis A. Cross sectional: normal maximal incisal opening and associations with physical variables in children. *Pediatr Dent* 2013;35:61–6.
- Dworkin SF, LeResche L. Research diagnostic criteria for temporomandibular disorders: review, criteria, examinations and specifications, critique. *J Craniomandib Disord* 1992;6:301–55.
- Mezitis M, Rallis G, Zachariades N. The normal range of mouth opening. *J Oral Maxillofac Surg* 1989;47:1028–9.
- Wood GD, Branco JA. A comparison of three methods of measuring maximal opening of the mouth. *J Oral Surg* 1979;37:175–7.
- Higbie EJ, Seidel-Cobb D, Taylor LF, Cummings GS. Effect of head position on vertical mandibular opening. *J Orthop Sports Phys Ther* 1999;29:127–30.
- Visscher CM, Huddleston Slater JJ, Lobbezoo F, Naeije M. Kinematics of the human mandible for different head postures. *J Oral Rehabil* 2000;27:299–305.
- Landtwing K. Evaluation of the normal range of vertical mandibular opening in children and adolescents with special reference to age and stature. *J Maxillofac Surg* 1978;6:157–62.
- Chen HS, Yang PL, Lee CY, Chen KK, Lee KT. Analysis of maximum mouth opening and its related factors in 3- to 5-year-old Taiwanese children. *Odontology* 2015;103:84–8.