

Available online at www.sciencedirect.com**ScienceDirect**

Procedia Technology 22 (2016) 465 – 470

Procedia
Technology

9th International Conference Interdisciplinarity in Engineering, INTER-ENG 2015, 8-9 October
2015, Tirgu-Mures, Romania

Effect of Metal Ions, Chemical Agents, and Organic Solvent on *Bacillus Sp.R2* Chitinase Activity

Ben Amar Cheba^{a,*}, Taha Ibrahim Zaghoul^b, Mohamad Hisham EL-Massry^b, Ahmad
Rafik EL-Mahdy^c

^aDepartment of Biotechnology, Faculty of Nature and Life Sciences, University of Sciences and Technology of Oran -Mohamed Boudiaf
(USTOMB), BP 1505 Al Mnaouar, Oran 31000,Algeria

^bDepartment of Biotechnology, Institute of Post Graduate Studies and Research, University of Alexandria,Alexandria,Egypt

^cDepartment of Food Science and Technology-Faculty of Agriculture-University of Alexandria- Alexandria-Egypt

Abstract

Bacillus sp. R₂ purified chitinase was subjected to the effect of metal ions, chemical agents, and organic solvent on its activity ,the results showed that K⁺, Mn²⁺, Na⁺, Mg²⁺ and Ca²⁺ stimulated chitinase activity by 2,4,7,15 and 21% respectively whereas Cu²⁺, Fe²⁺, Zn²⁺, Ag⁺ and Hg²⁺ inhibited the enzyme. 10mM β-mercaptoethanol and EDTA decreased the activity by 38.7 and 78.4% respectively whereas 1% SDS inactivated the enzyme completely, also 1mM galactose, glucose and N-acetyl glucosamine inactivated the enzyme by about 14.4, 25.2 and 38.7% respectively ethanol, acetone and isopropanol inhibited about 30% of chitinase activity however full or nearly full inhibition was observed with chloroform diethylether and ethylacetate. Furthermore the chitinase shelf life was 30 day at -20C and survived 4 freezing and thawing with lost of about 30% of its original activity. This is the first contribution about marine chitinase characterization study from novel gram variable *Bacillus* isolated from the Red sea.

© 2016 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license

(<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Peer-review under responsibility of the “Petru Maior” University of Tirgu Mures, Faculty of Engineering

Keywords: *Bacillus sp. R₂*; chitinase; characterization; metal ions; chemical agents; organic solvent; activity.

* Corresponding author. Tel.: +213-699-15-67-95.

E-mail address: benach57@yahoo.com

1. Introduction

Chitinases (E.C. 3.2.1.14) hydrolyze the linear polymer chitin, a polysaccharide of β -(1,4) linked *N*-acetylglucosamine (GlcNAc; 2-acetamino-2-deoxy- β -d-glucose) units, have gained intensive research interest for their versatile biotechnological applications that ranging from chitinous wastes recycling and phytopathogens biocontrol [1,2] to the preparation of pharmaceutically important chito oligosaccharides [3,4]. A wide range of microorganisms secrete chitinases for multiple purposes such as nutrition, morphogenesis, defence, antagonism, parasitism and pathogenesis [5–6].

Due to their diverse properties and technological characteristics, bacterial chitinases particularly were produced, purified, and characterized from many Gram negative and positive genera including, *Serratia*, *Aeromonas*, *Pseudomonas*, *Alteromonas*, *Enterobacter*, *Bacillus*, *Clostridium*, *Micromonospora* and *actinomyces* [7, 8, 9, 10, 11], however no reports or papers were found about marine chitinases from Gram variable Bacilli, for this reason in previous papers [12, 13] we described chitinase production and purification from marine Gram variable *Bacillus* sp.R2, whereas the present work was a successful contribution to characterize the enzyme via investigating the effect of certain factors such as metal ions, chemical agents, and organic solvent on chitinase activity.

2. Materials and methods

2.1. Chemicals

Yeast extract and peptone were obtained from (Oxoid Hampshire, England). While trypton and agar from (Difco Michigan, USA). Beef extract from (BBL-Maryland, USA), crab shell chitin flakes (Win-lab, UK). 2-Hydroxy 3,5-dinitrosalselic acid (DNSA) obtained from (Merck, Darmstadt- Germany). Crab shells chitosan, galactose, glucose, *N*-acetyl glucosamine and bovine serum albumin (BSA) were from (Sigma -USA). All of the other chemicals and reagents that were used were of highest grade commercially available.

2.2. Microorganism and purified chitinase characterization

Bacillus sp.R2 marine bacterial strain isolated and identified biochemically and molecularly by Cheba et al 2006 (strain accession number in NCBI Gen Bank was: DQ923161). The chitinase enzyme was produced and purified to homogeneity as reported in [12, 13], moreover the enzyme was subjected to characterization.

2.2.1. Effect of metal ions on chitinase activity

The enzyme was preincubated separately with 1 mM of various metal ions dissolved in 50 mM Tris-HCl buffer, pH 7.5 at 37°C for 30 min. Enzyme activities without metal ions addition (controls) were measured simultaneously and the residual activity was calculated.

2.2.2. Effect of major sea water cations concentration on chitinase activity

The reaction mixture of enzyme and substrate was preincubated separately for 30 min at 37°C with different concentration (1, 5, 10, 15, 25, 50 and 100 mM) of cations dissolved in 50 mM Tris-HCl, pH 7.5. The tested major cations were chloride form of Na^+ , K^+ , Ca^{+2} and Mg^{+2} .

2.2.3. Effect of chemical reagent and some monosugars on chitinase activity

The enzyme solution was incubated separately in 50 mM Tris-HCl, pH 7.5 containing various chemicals for 10 min at 37°C. Enzyme activities without adding chemicals (controls) were measured in parallel. The residual activity was determined by adding the substrate and carrying out the enzyme assay under the optimum condition.

2.2.4. Effect of organic solvents on chitinase activity

The enzyme was preincubated separately for 10 min at 37°C with 50% of the organic solvents. Reaction mixtures were carried out as above.

2.3. Analytical procedures

2.3.1. Chitinase assay

Chitinase activity was analyzed according to the method of Miller (1959) [14] by estimating the released reducing sugars spectrophotometrically at 540nm. A standard curve was established prepared with a series of dilutions of N-acetyl – D-glucosamine (NAG) and DNSA .One unit of chitinase activity was defined as the amount of enzyme required to release 1 μ mol of NAG per minute during reaction conditions.

2.3.2. Protein content assay

The amounts of protein in crude as well as purified enzyme were determined by the methods of Bradford (1976) [15] and Lowry et al. (1951) [16] respectively, using bovine serum albumin as standard.

3. Results and discussion

3.1. Effect of metal ions on chitinase activity

Data given in Table 1 revealed an increase in the chitinase activity 2, 4, 7, 15 and 21% by the addition of 1mM of K⁺, Mn²⁺, Na⁺, Mg²⁺ and Ca²⁺, respectively, where as other ions had a drastic effect on the enzyme activity and their inhibitory effect increased in the following order: Cu²⁺ < Fe²⁺ < Zn²⁺ < Ag⁺ < Hg²⁺. The chitinase activation by Ca²⁺, Mg²⁺, Mn²⁺, Na⁺ and K⁺ was reported by many authors[17,18,19] and the inhibition by Hg²⁺, Ag²⁺, Fe²⁺, Cu²⁺ and Zn²⁺ also was noticed in many literatures[17,18,19,20] However, certain ions such Cu²⁺ and Zn²⁺ could sometimes act as chitinase stimulators [21,22]and sometimes it could be chitinase inhibitors [17,19,23]

Table 1.Effect of metal ions on chitinase activity.

Metal ions (1mM)	Relative activity (%)	Metal ions (1mM)	Relative activity (%)
None	100	MgCl ₂	115
KCl	102	MnCl ₂	104
NaCl	107	FeSO ₄	34.5
HgCl ₂	0	ZnSO ₄	25.3
CaCl ₂	121	MgSO ₄	98.6
CuCl ₂	64.7	AgNO ₃	12.8

3.2. Effect of major sea water cations concentration

The results registered and graphically illustrated in Figure (1) showed that the four major cations contained in sea water (Na⁺, K⁺, Mg²⁺ and Ca²⁺) activated the enzyme, among them Ca²⁺ remarkably increased the activity by about 0.6 times at the concentration 10-50 mM followed by Mg²⁺ and Na⁺ which enhanced the activity by about 1.3 fold at the concentration of 5-15 mM and 15-50 mM for Mg²⁺ and Na⁺, respectively. Similar findings were obtained by Tsujibo et al. (1992)[17]who found that the chitinase of the marine bacterium *Alteromonas* sp. strain 0-7 was stimulated by the four major seawater cations, but he found also that Mg²⁺ at the concentration of 1-40 mM enhanced the activity by 1.6 fold instead of Ca²⁺. Other investigators also discussed the role of Ca²⁺ ion for the hydrolase activity generally and for the chitinases specially. Wright and Smucker (1986)[24]reported that the chitinase activity of the American oyster in full strength sea water was strongly inhibited by Mg²⁺, slightly

inhibited by Na⁺, but highly stimulated by Ca²⁺. Our chitinase also showed higher activity and was more stable in 75% sea water containing various cations such as Na⁺, K⁺, Mg²⁺ and Ca²⁺ than in artificial sea water or in 50mM Tris - HCl buffer, pH 7.5 (Data not shown). These results collectively emphasized the fact stating that the enzyme produced from marine organisms may have properties different from those produced by terrestrial organisms. Therefore, it is interesting to investigate the chitinase structure and function relationship.




Fig. 1. Effect of sea water major salts concentration on chitinase activity.

3.3. Effect of chemical reagent and some monosugars

The response of chitinase to the action of various chemical modifying reagents was performed and designed with the aim to understand the catalytic mechanism. Table (2) indicated that the chitinase activity was completely eliminated by 1% SDS but 22.5% and 84.3% remained at the concentration of 0.1% of SDS and Triton X-100, respectively. β -mercaptoethanol and EDTA at a concentration of 1mM inactivated the enzyme by about 26% and 43.1%, respectively but the concentration of 10mM of β -mercaptoethanol and EDTA decreased the activity by 38.7% and 78.4%, respectively. The other reagents did not have a significant effect on the chitinase. These results were in accordance with those reported in the literature that showed the inhibitory effects of SDS[21,25] Triton X-100[25] β -mercaptoethanol[27,28] EDTA [26,28] on the chitinase activity. Table (2) also showed that 1mM of galactose, glucose and N-acetyl glucosamine inactivated the enzyme by about 14.4%, 25.2% and 38.7%, respectively. Sakai et al. (1994) [21] also found that galactose, glucose and N-acetyl glucosamine inhibited the enzyme of *B. stearothermophilus* by 15%, 30% and 28%, respectively and the chitinases of *Enterobacter* sp. NRG4 [26] and *Metarhizium anisopliae* [29] was inhibited by 19.22 and 81% and 28.79 and 21% after addition 10mM of glucose, D-glucosamine and N-acetyl D-glucosamine, respectively. Others also demonstrated that the chitin hydrolysis end products such as N-acetyl glucosamine, chitobioses and the other chito oligosaccharides inhibited the chitinase activity [30, 21, 28].

3.4. Effect of organic solvents

As shown in Table (3) the chitinase exhibited a moderate tolerance (retained up to 70% of its original activity) to some organic solvents such as ethanol, acetone, isopropanol. However, full or nearly full inhibition by chloroform, diethyl ether, ethylacetate and diethylene dioxide (dioxane) was established. Sakai et al. (1994) [21] also found that butanol, ethanol, propanol, acetonitrile and dioxane decreased the chitinase activity of *Bacillus stearothermophilus* CH-4.

Table 2. Effect of chemical reagents and some mono sugars on chitinase activity

Chemical reagents	Concentration	Relative activity (%)	Chemical reagents	Concentration	Relative activity (%)
None	0 - 0	100 - 100	PMSF	3 mM	96.3
SDS	0.1- 1%	22.5 - 0	Ammonium sulfate	1mM	83.69
Triton X-100	0.1- 1%	84.3 - ND	Tween 20	0.5%	94.2
Urea	0.1- 1%	92.6 - 83.5	Glycerol	0.5%	96.8
β -mercaptoethanol	1-10 mM	74 - 61.3	Glucose	1mM	74.8
EDTA	1-10 mM	56.9 - 21.6	Galactose	1mM	85.6
KCN	1mM	98	N-acetyl glucosamine	1mM	61.8

Table 3. Effect of organic solvents on chitinase activity

Organic solvent (50%)	Relative activity (%)	Organic solvent (50%)	Relative activity (%)
None	100	Acetone	77.9
Methanol	67.5	Chloroform	0
Ethanol	76	Toluene	8.3
Propanol	37.5	Ethylacetate	2
Isopropanol	70.3	Diethylether	0
Butanol	45.6	Diethylene dioxide (dioxane)	6.5

4. Conclusion

The main conclusion obtained from the characterization of marine chitinase from novel gram variable *Bacillus* sp. R₂ that the enzyme showed higher activity and stability in 75% sea water containing various cations such as Na⁺, K⁺, Mg²⁺ and Ca²⁺. These results collectively emphasized the fact stating that the enzyme produced from marine organisms may have properties different from those produced by terrestrial organisms. Therefore, further investigation is needed such as chitinase structure and function relationship site directed mutagenesis and enzyme conditions optimizations which expected to makes the enzyme industrially useful for biotechnological applications

Acknowledgements

This work was supported by the Algerian ministry of higher education and scientific research. The author also thanks Professor Almahdy R.A for kindly providing some chemicals and his useful scientific discussion. We are grateful to Dr. Zaghoul T.I. and Dr. EL-Massry M.H for their scientific assistance and helpful suggestions during the work.

References

- [1] Ghanem KM, Al-Fassi FA, Farsi RM, Statistical optimization of cultural conditions for chitinase production from shrimp shellfish waste by *Alternaria alternata*, African Journal of Microbiology Research 2011; 5(13):1649–1659,
- [2] Cheba BA. Chitin and Chitosan: Marine Biopolymers with Unique Properties and Versatile Applications. Global Journal of Biotechnology & Biochemistry., 2011. 6 (3): 149-153.
- [3] Fernandes JC, Spindola H, De Sousa V, et al., "Anti-inflammatory activity of chitooligosaccharides in vivo," Marine Drugs, 2010 vol. 8, no. 6, pp. 1763–1768
- [4] Aam BB, Heggset EB, Norberg AL, Sorlie M, Varum KM, Eijsink VG. Production of chitooligosaccharides and their potential applications in medicine. Mar Drugs. 2010;8(5):1482–1517. doi: 10.3390/md8051482.

- [5] Ju C, Yue W, Yang Z et al. Antidiabetic effect and mechanism of chitoooligosaccharides. *Biological and Pharmaceutical Bulletin* 2010; 33, (9):1511–1516.
- [6] Lee CG, Da Silva CA, Lee JY, Hartl D, Elias JA. Chitin regulation of immune responses: an old molecule with new roles. *Curr. Opin. Immunol.* 2008;20:684–689
- [7] Gutiérrez-Román MI, Holguín-Meléndez F, Bello-Mendoza R, Guillén-Navarro K, Dunn MF, Huerta-Palacios G, Production of prodigiosin and chitinases by tropical *Serratia marcescens* strains with potential to control plant pathogens, in *World Journal of Microbiology and Biotechnology* 2012; 28:145–153.
- [8] Chernin LS, De La Fuente L, Sobolev V, Haran S, Vorigas CE, Oppenheim AB, Chet I. Molecular cloning, structural analysis, and expression in *Escherichia coli* of a chitinase gene from *Enterobacter agglomerans*. *Applied Environmental Microbiology* 1997; 63(3):834–839.
- [9] Bharat B, Gurinder Singh H. Isolation, purification and properties of a thermostable chitinase from an alkalophilic *Bacillus* sp. BG-11. *Biotechnology Letters*, February 1998, vol. 20, no. 2, p. 157-159
- [10] Kim, Kyoung Ja; Yang, Yong Joon and Kim, Jong Gi. Purification and characterization of chitinase from *Streptomyces* sp. M-20. *Journal of Biochemistry and Molecular Biology* 2003; 36(92):185-189.
- [11] Bhattacharya D, Nagpure A, Gupta RK. Bacterial chitinases: properties and potential. *Crit Rev Biotechnol.* 2007;27(1):21–28. doi: 10.1080/07388550601168223
- [12] Cheba BA, Zaghoul TI, EL-Mahdy AR, EL-Massry MH. Enhanced Production of *Bacillus* sp. R2 Chitinase through Cell Immobilization. *ACT-Biotechnology Research Communications* 2011;1(1):8-13.
- [13] Cheba BA, Zaghoul TI, El-Mahdy AR, El-Massry MH. *Bacillus* sp. R2 Chitinase: Affinity Purification and Immobilization *Procedia Technology*. 2015. 19, , Pages 958–964 doi:10.1016/j.procy.2015.02.137
- [14] Miller GR. Use of Dinitrosalicylic Acid reagent for determination of reducing sugar. *Anal. Chem.*, 1959,31 (3). : 426 – 428.
- [15] Bradford MM. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* 1976; 72: 248 – 254.
- [16] Lowery OH, Rosebrough NJ, Farr AL, Randell RJ. Protein measurement with the Folin phenol reaction. *J. Biol. Chem.* 1951; 193: 265 – 275.
- [17] Yuli PE, Suhartono MT, Rukayadi Y, Hwanj JK, Pyun YR. Characteristic of thermostable chitinase enzymes from the Indonesian *Bacillus* sp. 13.26. *Enz. Microb. Technol.* 2004. 35: 147-153.
- [18] Aunpad R, Panbamgrad W. Cloning and characterization of the constitutively expressed chitinase C gene from a marine bacterium *Salinivibrio costicola* strain 5SM-1. *J. Biosc. Bioeng.* 2003. 96:529-536. 42
- [19] Tsujibo H, Minoura K, Miyamoto K, Endo H, Moriwaki M, Inamori Y. Purification and properties of a thermostable chitinase from *Streptomyces thermoviolaceus* OPC-520. *Appl. Environ. Microbiol.* 1993. 59 (2): 620-622
- [20] Donderski W, Brzezinska MS. The influence of heavy metals on the activity of chitinases produced by planktonic benthic and epiphytic bacteria. *Polish J. Enviro. Studies.* 2005.14 (6):851-859
- [21] Sakai K, Narihara M, Kasama Y, Wakayama M, Moriguchi M. Purification and characterization of thermostable β -N-acetyl hexosaminidase of *Bacillus stearothermophilus* CH-4 isolated from chitin-containing compost. *Appl. Environ. Microbiol.* 1994.60 (8): 2911-2915.
- [22] Tang Y, Zhao J, Ding S, Liv S, Yang Z. Purification and properties of chitinase from *Enterobacter aerogenes*. *Wei Sheng Wu Xue Bao. Acta Microbiologica Sinica*: 2001. 41 (1): 82-86.
- [23] Wen C, Teseng C, Cheng C, Li Y. Purification, characterization and cloning of a chitinase from *Bacillus* sp. NTU2. *Biotechnol. App. Biochem.* 2002; 35: 213-219
- [24] Wright DA, Smucker RA. Ionic requirements for chitinase/chitobiase activity in the oyster, *Crassostrea virginica*. *Comp. Biochem. Physiol.* A. 1986. 84: 495-497
- [25] Joshi S, Kozlowski M, Selvaraj G, Lyer VN, Davies RW. Cloning of the genes of the chitin utilization of *Serratia liquefaciens*. *J. Bacteriol.* 1988. 170: 2984-2988
- [26] Dahiya N, Tewari R, Tewari RP, Hoondal GS. Chitinase from *Enterobacter* sp. NRG4: its purification, characterization and reaction pattern. *Electronic. J. Boitechnol.* 2005. 8 (2): 134-145.
- [27] Zhang Z, Yuen GY. The role of chitinase production by *Stenotrophomonas maltophilia* strain C3 in biological control of *Bipolaris sorokiniana*. *Phytopathol.* 2000.90: 384-389.
- [28] Sakai K, Yokota A, Kurokawa H, Wakayama M, Moriguchi M. Purification and characterization of three thermostable endochitinases of a noble *Bacillus* strain, MH-1 isolated from chitin-containing compost. *App. Environ. Microbiol.* 1998. 64 (9): 3397-3402
- [29] Wang SY, Moyne AL, Thottappilly G, Wu SJ, Locy RD, Singh NK. Purification and characterization of *Bacillus cereus* exochitinase. *Enz. Microb. Technol.* 2001.28: 492-498.
- [30] Shimosaka M, Nogawa M, Wang X, Kumehara M, Okazaki M. Production of two chitosanases from a chitosan-assimilating bacterium, *Acinetobacter* sp. strain CHB101. *App. Environ. Microbiol.* 1995. 61 (2): 438-442.