

Differential membrane perturbation caused by the cell penetrating peptide Tp10 depending on attached cargo

Elsa Bárány-Wallje^a, Jugnu Gaur^a, Pontus Lundberg^b, Ülo Langel^b, Astrid Gräslund^{a,*}

^a Department of Biochemistry and Biophysics, The Arrhenius Laboratories, Stockholm University, S-10691 Stockholm, Sweden

^b Department of Neurochemistry, The Arrhenius Laboratories, Stockholm University, S-10691 Stockholm, Sweden

Received 16 February 2007; revised 18 April 2007; accepted 19 April 2007

Available online 26 April 2007

Edited by Maurice Montal

Abstract The membrane leakage caused by the cell penetrating peptide Tp10, a variant of transportan, was studied in large unilamellar vesicles with the entrapped fluorophore calcein. The vesicles were composed of zwitterionic 1-palmitoyl-2-oleoyl-sn-glycero-3-phosphocholine. A significant decrease in membrane leakage was found when the 55 kDa streptavidin protein was attached to Tp10. When a 5.4 kDa peptide nucleic acid molecule was attached, the membrane leakage was comparable to that caused by Tp10 alone. The results suggest that direct membrane effects may cause membrane translocation of Tp10 alone and of smaller complexes, whereas these effects do not contribute for larger cargoes.

© 2007 Federation of European Biochemical Societies. Published by Elsevier B.V. All rights reserved.

Keywords: Cell-penetrating peptides; Transportan 10; Cell membrane perturbation; Streptavidin; Peptide nucleic acid

1. Introduction

Cell-penetrating peptides (CPPs) have the ability to translocate through cell membranes with high efficiency [1–3]. When they are covalently linked to a larger cargo, such as polypeptides, oligonucleotides or proteins, they still retain their translocation properties [4]. These short, synthetic peptide based vectors are considered as biocompatible and economical candidates for delivery of hydrophilic drugs (especially in connection with gene therapy) [1,2,5,6]. Transportan is a chimeric CPP constructed from the (1–12) amino acid residues from the N-terminal part of the neuropeptide galanin linked with a lysine residue to the full length wasp venom mastoparan [7]. Transportan 10 (Tp10) was developed as an analog to transportan by deleting the first six N-terminal amino acids, in order to reduce toxicity [8]. Tp10 has the sequence AGYLLGKINLKALAALAKKIL [8]. Tp10 and transportan have remarkably good translocation capabilities and has successfully been used as delivery vectors for peptide nucleic acid

(PNA) [9,10], peptides and proteins [11,12] but they have been shown to induce membrane leakage in model systems [13]. Previous studies have also shown that Tp10 is toxic to cancer cells (assayed as leakage of lactate dehydrogenase) and that the peptide induces hemolysis [14].

The CPPs are a variable group of peptides, and their mechanisms of action have been under intense debate. Only relatively recently [15] some consensus has appeared: Cargo delivery is mostly endocytosis-driven and very likely dependent on endosomal escape. Direct membrane interaction mechanisms seem to operate in parallel in some cases, as has been suggested for Tp10 [16]. Such interactions have been mainly associated with highly efficient CPPs, which however also display cytotoxic activities [14]. The modifying effects of different cargoes on the uptake mechanisms have also been reported [17,18].

Whereas it seems clear that cargoes affect the CPP uptake mechanisms in cells, less work has been done on how attachment of cargo may affect the membrane perturbations induced by CPPs. Of particular interest is then the group of highly efficient CPPs which are considered to be somewhat cytotoxic. The aim of the present work is to evaluate the membrane interactions of Tp10 carrying different biologically relevant cargoes. Tp10 and transportan are structurally similar to active antimicrobial peptides, since they contain the mastoparan sequence [19,20]. In this respect similar to the antimicrobial peptides magainin and buforin, Tp10 may also have more than one uptake mechanism, and the relative importance of these mechanisms may depend on the cargo [17,18].

We have compared the membrane perturbation effects of Tp10_{bt} alone and in complex with a large and hydrophilic cargo such as streptavidin (55 kDa), or a smaller and hydrophobic cargo such as 18 bases long PNA (5.4 kDa). Streptavidin [21,22] was coupled to Tp10 via a biotin molecule attached to the Lys 7 of Tp10. The PNA was covalently attached to Tp10 by a disulfide bond. The membrane perturbation by the different constructs was studied by calcein leakage experiments in large unilamellar vesicles (LUVs) composed of phospholipids. Our hypothesis was that membrane perturbation (manifested as calcein leakage in a vesicle model system) should be strongly connected to CPP activity due to direct membrane interactions, as well as to the endosomal escape of efficient CPPs.

Phospholipid vesicles are artificial membrane systems which are suitable mimics of the properties of biological membranes. With phospholipid vesicles it is possible to manipulate the various parameters of the membrane system such as size or

*Corresponding author. Fax: +46 8 155597.
E-mail address: astrid@dbb.su.se (A. Gräslund).

Abbreviations: CPP, cell-penetrating peptide; Tp10, transportan 10; PNA, peptide nucleic acid; LUVs, large unilamellar vesicles; POPC, 1-palmitoyl-2-oleoyl-sn-glycero-3-phosphocholine; POPG, 1-palmitoyl-2-oleoyl-sn-glycero-3-[phospho-rac-(1-glycerol)]; bt, biotin labeled

charge, etc. 1-Palmitoyl-2-oleoyl-sn-glycero-3-phosphocholine (POPC) was used to create neutral vesicles, whereas POPC:POPG mixtures were used in order to create vesicles with net negative surface charge. The membrane perturbing effect of Tp10_{bt} and transportan on neutral vesicles was compared to the effect that they had on vesicles including 25% negatively charged phospholipids. We also compared the membrane leakage caused by transportan, Tp10 and mastoparan, respectively.

2. Materials and methods

2.1. Materials

Transportan was purchased from Neosystems (Strasbourg, France). Biotin labeled Tp10 was synthesized at the Department of Neurochemistry, Stockholm University. Mastoparan from the venom of wasp (zoological name) was obtained from NeoMPS (Strasbourg, France). The identity and purity were controlled by amino acid, mass spectrometric and HPLC analysis. 1-Palmitoyl-2-oleoyl-phosphocholine (POPC) and 1-palmitoyl-2-oleoyl-sn-glycero-3-[phospho-*rac*-(1-glycerol)] (POPG) were purchased from Avanti Polar Lipids (Alabaster, Alabama), of the best quality, and were used without further purification. Calcein, a fluorescein derivative (C₃₀H₂₆N₂O₁₃, Mw = 622.5 g/mole), was purchased from Sigma-Aldrich (Stockholm, Sweden). Streptavidin was obtained from MP Biomedicals (Illkirch, France), and was used as purchased. The Tp10-PNA construct was prepared as previously described [23]. In brief, the PNA was attached with a disulphide bridge to the ε-amino group of Lys 7. The single stranded PNA used was an 18 bases long sequence (underlined), C- and N-terminally flanked with two lysines (K) on either side: (Cys)-KKCCTCTTACCTCAGTTACAKK (Mw = 5380.4 Da).

2.2. Determination of peptide concentrations

The peptide concentrations in the stock solutions were determined by light absorption on a Cary 4 spectrophotometer using cuvettes with a 1 cm light path. All the spectra were baseline corrected. A molar absorptivity of 5600 M⁻¹ cm⁻¹ and 1280 M⁻¹ cm⁻¹, at 280 nm, for tryptophan and tyrosine, respectively, were applied in the calculations. Mastoparan does not include any tryptophans or tyrosines and its concentration was therefore determined by amino acid analysis at Protein Analysis Center, Department of Medical Biochemistry and Biophysics, Karolinska Institute, Stockholm.

2.3. Determination of PNA concentration

PNA concentration in the stock solution was determined by light absorption on a Cary 4 spectrophotometer using cuvettes with 1 cm light path. The spectra were baseline corrected. A molar absorptivity of 164.3 mM⁻¹ cm⁻¹ at 260 nm was used for this PNA.

2.4. Sample preparations

Large unilamellar vesicles were prepared by initially dissolving the phospholipids (POPC or POPC:POPG 3:1 mixtures, respectively) at the desired concentration in methanol, to ensure complete mixing of the components, and then removing the solvent by placing the sample under a stream of nitrogen or argon to create a thin lipid film. Further solvent was removed by vacuum drying for 3 h. The dried lipids were then dispersed in 50 mM potassium phosphate buffer (pH 7.0). The solution was vortexed to form large multilamellar vesicles (LMVs). The dispersion was run through a repeated freeze-thaw cycle 10 times, which resulted in decreased lamellarity, and then extruded through two polycarbonate filters (0.1 μm pore size) 20 times in an Avanti manual extruder. This process gave unilamellar vesicles with a size of 100 nm.

2.5. Fluorescence spectroscopy

Fluorescence was measured on a Perkin-Elmer LS 50B luminescence spectrometer with FLWINLAB software. All measurements were made in 4 ml plastic cuvettes (1 × 1 cm) at an ambient temperature of approximately 20 °C. The fluorescence intensity of calcein was measured at 512 nm (excited at 490 nm). Scans were recorded with 2.5 nm excitation and emission bandwidths and a scan speed of 800 nm/min.

2.6. Peptide-induced calcein release

LUVs with entrapped calcein were prepared by hydrating a lipid film of desired composition with 70 mM calcein present in the buffer (the final pH was adjusted to 7.0 by addition of NaOH from a 1 M stock solution). Calcein gives low fluorescence intensity at 70 mM because of self-quenching and the intensity increases considerably upon dilution. Free calcein was separated from the LUVs on a Sephadex-G25 column purchased from Amersham Biosciences (Uppsala, Sweden). Increasing concentration of peptides was added to LUVs composed of 100 μM phospholipids. After 5 min incubation, release of calcein from the LUVs was monitored by an increase in the fluorescence intensity at 512 nm.

The maximum fluorescence intensity corresponding to 100% leakage was determined by lysing the vesicles with 10% triton X-100. The % leakage was then calculated according to the following equation:

$$\% \text{Leakage} = 100 \{ (F - F_0) / (F_{\text{max}} - F_0) \}$$

where F_0 represents the fluorescence intensity of the intact vesicles, F and F_{max} , the intensity before and after the addition of the detergent, respectively.

3. Results

3.1. Peptide-induced calcein release from LUVs

The membrane perturbing effect of Tp10_{bt} was investigated through calcein-release experiments in LUVs. Fig. 1 depicts the % leakage induced by Tp10_{bt} with neutral POPC LUVs, after 5 min of incubation, at room temperature. The results show that Tp10_{bt} causes membrane leakage at quite low concentrations.

3.2. Membrane perturbations with streptavidin as cargo

Fig. 1 also depicts the calcein leakage effect of Tp10_{bt} when attached to streptavidin (55 kDa). Streptavidin can bind up to four molecules of Tp10_{bt}. Two and four times excess concentration of streptavidin to Tp10_{bt} was used, respectively. For both cases a similar and drastic decrease in the membrane leakage could be seen compared to Tp10_{bt} alone. Streptavidin was used in excess to ensure complete utilization of Tp10_{bt} in complex formation. This complex was found much less membrane perturbing than Tp10_{bt} alone. The leakage was reduced almost




Fig. 1. Calcein leakage from LUVs induced by Tp10_{bt} (black squares), Tp10_{bt} pre-incubated with two times excess of streptavidin (black circles) and Tp10_{bt} pre-incubated with four times excess of streptavidin (open circles) respectively. Increasing concentrations of the peptide and peptide + streptavidin solutions were added to LUVs composed of 100 μM POPC with 70 mM calcein entrapped. The % calcein release after 5 min at room temperature and pH 7.0 was plotted as a function of peptide concentration.

500 times when a Tp10_{bt}:streptavidin ratio of 1:2 or 1:4 was used (Fig. 1).

3.3. Membrane perturbations effects with PNA

Peptide nucleic acid (PNA, 5.4 kDa) was covalently coupled to Tp10. Fig. 2 shows the membrane leakage caused by Tp10_{bt}, and the Tp10-PNA construct. In this case the added cargo had only a small effect on the membrane perturbation and leakage caused by Tp10.

3.4. Comparison of activities of peptides

The membrane perturbation effect of Tp10_{bt} was compared with that of its parent peptide transportan and with the potent antimicrobial peptide mastoparan which has the same sequence as the C-terminus of Tp10 and transportan. The results in Fig. 3 show that mastoparan was most potent and caused severe membrane leakage at very low concentrations. Transportan was less membrane perturbing than mastoparan, but Tp10_{bt} was found to be far less membrane disturbing than both the other peptides. The kinetics of calcein release from 100 μ M LUVs was studied for mastoparan, transportan and Tp10, respectively. For the three peptides the release appeared to be biphasic with a rapid initial phase followed by a much slower process. When peptide concentrations were adjusted




Fig. 2. Calcein leakage from POPC LUVs caused by addition of Tp10_{bt} (black squares) and Tp10-PNA (grey triangles) as a function of peptide concentration.




Fig. 3. Calcein leakage caused by the three different peptides; mastoparan (grey diamonds), transportan (black circles) and Tp10_{bt} (black squares). LUVs consisting of 100 μ M POPC with 70 mM calcein enclosed were titrated with the different peptides, respectively. The experiments were done at room temperature at pH 7.0 with an incubation time of 5 min.




Fig. 4. Comparison between the calcein leakage caused by Tp10_{bt} and transportan from LUVs consisting of POPC or POPC:POPG 3:1 mixtures, respectively. (a) Calcein release when Tp10_{bt} was added to 100 μ M POPC vesicles (black squares) and 100 μ M POPC:POPG 3:1 vesicles (grey diamonds), respectively. (b) Calcein release when transportan was added to 25 μ M POPC vesicles (black circles) and 25 μ M POPC:POPG 3:1 vesicles (grey triangles), respectively.

to correspond to a similar amount of leakage (60% after 20 min), the kinetics for the different peptides turned out to be very similar (data not shown).

3.5. Membrane perturbing effects on vesicles including negatively charged phospholipids

The membrane perturbing effect of Tp10_{bt} (Fig. 4a) and transportan (Fig. 4b) on neutral vesicles was compared to the effect that they had on vesicles including 25% negatively charged phospholipids. The negatively charged vesicles contained POPC:POPG mixtures (molar ratio 3:1). Both peptides had a much stronger effect on the neutral vesicles than on the charged vesicles at the lowest peptide concentrations.

4. Discussion

Tp10 structurally resembles antimicrobial, pore forming peptides. It interacts with the membrane in a partly hydrophobic manner and exhibits strong effects on the membrane integrity. These observations provide a link between observed translocation across cell membranes and the induction of leakage in the LUV model systems, similar to what has been reported for magainin and buforin [17,18].

We have now studied the extent of membrane leakage when different cargo molecules are attached to Tp10. To use streptavidin as a large and hydrophilic cargo, Tp10 was biotinylated at position Lys 7. Streptavidin has a high affinity for biotin. It

has been shown previously that Tp10_{bt} is able to translocate through membranes when bound to streptavidin [24]. The large reduction of leakage for the Tp10-streptavidin complex suggests that the use of a suitable cargo can reduce the toxicity of Tp10 to some extent.

In other experiments PNA was used as a cargo. The Tp10-PNA construct, which is relatively hydrophobic and with additional positive charges, turned out to be only slightly less membrane perturbing than Tp10_{bt} alone. PNA are useful tools for antisense gene therapy, but they suffer from poor uptake since they are membrane impermeable [25]. Tp10-mediated uptake should be quite efficient, given that the cytotoxicity can be tolerated. In this study the PNA was conjugated with the CPP by a disulphide bond, which has proven to be an efficient way to do the linkage [26]. Inside the cell, this bond may be reduced by cytosolic components and free PNA can move to its designated target [27].

Tp10 and mastoparan are both fragments of transportan. Mastoparan, a potent wasp venom peptide, caused maximal membrane leakage. It is suggested to have a transmembrane orientation, favoring pore formation [28–31]. A pore may be defined either as a well defined structure, e.g. composed of a number of helices in parallel alignment, or as a less well-defined peptide-induced structural defect in the membrane through which ions or molecules can pass. The latter definition may be most suitable for CPPs like Tp10 and transportan.

The membrane perturbing effect of Tp10 as well as transportan is significantly stronger on neutral vesicles consisting of POPC than on negatively charged vesicles consisting of POPC:POPG mixtures (see Fig. 4). The differences observed in the present study are surprisingly big at the lowest measured peptide concentrations. Although these effects may partly be due to specific interactions between the peptides and the phospholipid headgroups, they also suggest a competition between electrostatic forces and hydrophobic forces [32]. For a bilayer consisting of zwitterionic phospholipids these peptides could possibly be inserted more deeply into the hydrophobic part of the bilayer causing leakage, whereas for negatively charged lipids the electrostatic attraction could keep the peptide in the headgroup region. Similar observations have previously been reported for the bee venom melittin [28,29,31,33].

The varying induction of membrane leakage by Tp10 with and without cargoes indicates that the mechanism by which this class of CPPs translocates through the membrane may depend on whether a large cargo is attached to the peptide or not. In the absence of large hydrophilic cargoes the peptide may induce membrane disturbances that allows for direct translocation of peptide into the cell. In the presence of a large hydrophilic cargo, such as streptavidin, the membrane disturbance is small and endocytosis is probably the only translocation mechanism [24,34–36]. For another class of CPPs (e.g. penetratin) that are generally less toxic, there is no evidence of membrane leakage in model systems [37].

The present results suggest a further step forward in understanding the mechanisms of CPPs. Given the right conditions CPPs use endocytosis (including endosomal escape) when entering cells, without or with cargo. Certain peptides like transportan or Tp10 which cause leakage in membrane model systems, can in addition enter cells via direct membrane interactions (transient pore formation). Our present results suggest that this additional pathway is strongly dependent on cargo. The non-endocytotic pathway may be inhibited by large

hydrophilic cargoes, but allowed with smaller hydrophobic or positively charged cargoes.

Acknowledgements: This work was supported by the Swedish Research Council, the Swedish Cancer Society and the Foundation for Strategic Research (Bio-X, SSF A3 05:175 and Center for Biomembrane Research).

References

- [1] Derossi, D., Chassaing, G. and Prochiantz, A. (1998) Trojan peptides: the penetrating system for intracellular delivery. *Trends Cell Biol.* 8, 84–87.
- [2] Järver, P. and Langel, Ü. (2004) The use of cell-penetrating peptides as a tool for gene regulation. *Drug Discovery Today* 9, 395–402.
- [3] Langel, Ü. (2002) *Cell-penetrating Peptides. Processes and Applications*, CRC Press, Boca Raton.
- [4] Dietz, G.P. and Bähr, M. (2004) Delivery of bioactive molecules into the cell: the Trojan horse approach. *Mol. Cell. Neurosci.* 27, 85–131.
- [5] Deshayes, S., Morris, M.C., Divita, G. and Heitz, F. (2005) Cell-penetrating peptides: tools for intracellular delivery of therapeutics. *Cell Mol. Life Sci.* 62, 1839–1849.
- [6] Magzoub, M. and Gräslund, A. (2004) Cell-penetrating peptides: from inception to application. *Q. Rev. Biophys.* 37, 147–195.
- [7] Langel, Ü., Pooga, M., Kairane, C., Zilmer, M. and Bartfai, T. (1996) A galanin-mastoparan chimeric peptide activates the Na⁺,K⁺-ATPase and reverses its inhibition by ouabain. *Regul. Pept.* 62, 47–52.
- [8] Soomets, U., Lindgren, M., Gallet, X., Hällbrink, M., Elmquist, A., Balaspiri, L., Zorko, M., Pooga, M., Brasseur, R. and Langel, Ü. (2000) Deletion analogues of transportan. *Biochim. Biophys. Acta* 1467, 165–176.
- [9] El-Andaloussi, S., Johansson, H., Magnusdottir, A., Järver, P., Lundberg, P. and Langel, Ü. (2005) TP10, a delivery vector for decoy oligonucleotides targeting the Myc protein. *J. Contr. Rel.* 110, 189–201.
- [10] Pooga, M., Soomets, U., Hällbrink, M., Valkna, A., Saar, K., Rezaei, K., Kahl, U., Hao, J.X., Xu, X.J., Wiesenfeld-Hallin, Z., Hökfelt, T., Bartfai, T. and Langel, Ü. (1998) Cell penetrating PNA constructs regulate galanin receptor levels and modify pain transmission *in vivo*. *Nat. Biotechnol.* 16, 857–861.
- [11] Pooga, M., Kut, C., Kihlmark, M., Hällbrink, M., Fernaeus, S., Raid, R., Land, T., Hallberg, E., Bartfai, T. and Langel, Ü. (2001) Cellular translocation of proteins by transportan. *FASEB J.* 15, 1451–1453.
- [12] Säälik, P., Elmquist, A., Hansen, M., Padari, K., Saar, K., Viht, K., Langel, Ü. and Pooga, M. (2004) Protein cargo delivery properties of cell-penetrating peptides. A comparative study. *Bioconjug. Chem.* 15, 1246–1253.
- [13] Magzoub, M., Eriksson, L.E.G. and Gräslund, A. (2003) Comparison of the interaction, positioning, structure induction and membrane perturbation of cell-penetrating peptides and non-translocating variants with phospholipid vesicles. *Biophys. Chem.* 103, 271–288.
- [14] Saar, K., Lindgren, M., Hansen, M., Eiriksdóttir, E., Jiang, Y., Rosenthal-Aizman, K., Sassian, M. and Langel, Ü. (2005) Cell-penetrating peptides: a comparative membrane toxicity study. *Anal. Biochem.* 345, 55–65.
- [15] Mäe, M. and Langel, Ü. (2006) Cell-penetrating peptides as vectors for peptide, protein and oligonucleotide delivery. *Curr. Opin. Pharmacol.* 6, 509–514.
- [16] Yandek, L.E., Pokorny, A., Florén, A., Knoelke, K., Langel, Ü. and Almeida, P.F.F. (2007) Mechanism of the cell-penetrating peptide Tp10 permeation of lipid bilayers. *Biophys. J.* 92, 2434–2444.
- [17] Takeshima, K., Chikushi, A., Lee, K.K., Yonehara, S. and Matsuzaki, K. (2003) Translocation of analogues of the antimicrobial peptides magainin and buforin across human cell membranes. *J. Biol. Chem.* 278, 1310–1315.
- [18] Tünnemann, G., Martin, R.M., Haupt, S., Patsch, C., Edenhofer, F. and Cardoso, M.C. (2006) Cargo-dependent mode of uptake

- and bioavailability of TAT-containing proteins and peptides in living cells. *FASEB J.* 20, 1775–1784.
- [19] Bárány-Wallje, E., Andersson, A., Gräslund, A. and Mäler, L. (2004) NMR solution structure and position of transportan in neutral phospholipid bicelles. *FEBS Lett.* 567, 265–269.
- [20] Lindberg, M., Jarvet, J., Langel, Ü. and Gräslund, A. (2001) Secondary structure and position of the cell-penetrating peptide transportan in SDS micelles as determined by NMR. *Biochemistry* 40, 3141–3149.
- [21] Agarana, C.E., Kuntz, I.D., Birken, S., Axel, R. and Cantor, C.R. (1986) Molecular cloning and nucleotide sequence of the streptavidin gene. *Nucleic Acids Res.* 14, 871–882.
- [22] Weber, P.C., Ohlendorf, D.H., Wendoloski, J.J. and Salem, F.R. (1989) Structural origins of high affinity biotin binding to streptavidin. *Science* 243, 85–88.
- [23] El-Andaloussi, S., Johansson, H., Lundberg, P. and Langel, Ü. (2006) Induction of splice correction by cell-penetrating peptide nucleic acids. *J. Gene Med.* 8, 1262–1273.
- [24] Padari, K., Säälik, P., Hansen, M., Koppel, K., Raid, R., Langel, Ü. and Pooga, M. (2005) Cell transduction pathways of transportans. *Bioconjug. Chem.* 16, 1399–1410.
- [25] Gait, M.J. (2003) Peptide-mediated cellular delivery of antisense oligonucleotides and their analogues. *Cell Mol. Life Sci.* 60, 844–853.
- [26] Bendifallah, N., Rasmussen, F.W., Zachar, V., Ebbesen, P., Nielsen, P.E. and Koppelhus, U. (2006) Evaluation of cell-penetrating peptides (CPPs) as vehicles for intracellular delivery of antisense peptide nucleic acid (PNA). *Bioconjug. Chem.* 17, 750–758.
- [27] Hällbrink, M., Florén, A., Elmquist, A., Pooga, M., Bartfai, T. and Langel, Ü. (2001) Cargo delivery kinetics of cell-penetrating peptides. *Biochim. Biophys. Acta* 1515, 101–109.
- [28] Duforc, E.J., Smith, I.C.P. and Dufourcq, J. (1986) Molecular details of melittin-induced lysis of phospholipid membranes as revealed by deuterium and phosphorus NMR. *Biochemistry* 25, 6448–6455.
- [29] Hinch, D.K. and Crowe, J.H. (1996) The lytic activity of the bee venom peptide melittin is strongly reduced by the presence of negatively charged phospholipids or chloroplast galactolipids in the membranes of phosphatidylcholine large unilamellar vesicles. *Biochim. Biophys. Acta* 1284, 162–170.
- [30] Hori, Y., Demura, M., Iwadate, M., Ulrich, A.S., Niidome, T., Aoyagi, H. and Asakura, T. (2001) Interaction of mastoparan with membranes studied by ¹H-NMR spectroscopy in detergent micelles and by solid-state ²H-NMR and ¹⁵N-NMR spectroscopy in oriented lipid bilayers. *Eur. J. Biochem.* 268, 302–309.
- [31] Ladokhin, A.S. and White, S.H. (2001) “Detergent like” permeabilization of anionic lipid vesicles by melittin. *Biochim. Biophys. Acta* 1514, 253–260.
- [32] Dathe, M., Schumann, M., Wieprecht, T., Winkler, A., Beyersmann, M., Krause, E., Matsuzaki, K., Murase, O. and Bienert, M. (1996) Peptide helicity and membrane surface charge modulate the balance of electrostatic and hydrophobic interactions with lipid bilayers and biological membranes. *Biochemistry* 35, 12612–12622.
- [33] Matsuzaki, K., Yoneyama, S. and Miyajima, K. (1997) Pore formation and translocation of melittin. *Biophys. J.* 73, 831–838.
- [34] Console, S., Marty, C., Garcia-Echeverria, C., Schwendener, R. and Ballmer-Hofer, K. (2003) Antennapedia and HIV transactivator of transcription (TAT) “protein transduction domains” promote endocytosis of high mol. wt. cargo upon binding to cell surface glycosaminoglycans. *J. Biol. Chem.* 278, 35109–35114.
- [35] Drin, G., Cottin, S., Blanc, E., Rees, A.R. and Temsamani, J. (2003) Studies on the internalization mechanism of cationic cell-penetrating peptides. *J. Biol. Chem.* 278, 31192–31201.
- [36] Fischer, R., Köhler, K., Fotin-Mleczek, M. and Brock, R. (2004) A stepwise dissection of the intracellular fate of cationic cell-penetrating peptides. *J. Biol. Chem.* 279, 12625–12635.
- [37] Bárány-Wallje, E., Keller, S., Serowy, S., Geibel, S., Pohl, P., Bienert, M. and Dathe, M. (2005) A critical reassessment of penetratin translocation across lipid membranes. *Biophys. J.* 89, 2513–2521.