

NCoR Repression of LXRs Restricts Macrophage Biosynthesis of Insulin-Sensitizing Omega 3 Fatty Acids

Pingping Li,^{1,8} Nathanael J. Spann,^{2,8} Minna U. Kaikkonen,^{2,5} Min Lu,¹ Da Young Oh,¹ Jesse N. Fox,² Gautam Bandyopadhyay,¹ Saswata Talukdar,¹ Jianfeng Xu,¹ William S. Lagakos,¹ David Patsouris,¹ Aaron Armando,³ Oswald Quehenberger,^{1,3} Edward A. Dennis,^{3,4} Steven M. Watkins,⁶ Johan Auwerx,⁷ Christopher K. Glass,^{1,2,*} and Jerrold M. Olefsky^{1,*}

¹Department of Medicine

²Department of Cellular and Molecular Medicine

³Department of Chemistry and Biochemistry

⁴Department of Pharmacology

University of California, San Diego, 9500 Gilman Drive, La Jolla, CA 92093, USA

⁵A.I. Virtanen Institute, Department of Biotechnology and Molecular Medicine, University of Eastern Finland, P.O. Box 1627, 70211 Kuopio, Finland

⁶Lipomics Technologies, Inc., West Sacramento, CA 95691, USA

⁷Laboratory of Integrative and Systems Physiology, Ecole Polytechnique Fédérale de Lausanne, 1015 Lausanne, Switzerland

⁸These authors contributed equally to this work

*Correspondence: cglass@ucsd.edu (C.K.G.), jolefsky@ucsd.edu (J.M.O.)

<http://dx.doi.org/10.1016/j.cell.2013.08.054>

SUMMARY

Macrophage-mediated inflammation is a major contributor to obesity-associated insulin resistance. The corepressor NCoR interacts with inflammatory pathway genes in macrophages, suggesting that its removal would result in increased activity of inflammatory responses. Surprisingly, we find that macrophage-specific deletion of NCoR instead results in an anti-inflammatory phenotype along with robust systemic insulin sensitization in obese mice. We present evidence that derepression of LXRs contributes to this paradoxical anti-inflammatory phenotype by causing increased expression of genes that direct biosynthesis of palmitoleic acid and ω 3 fatty acids. Remarkably, the increased ω 3 fatty acid levels primarily inhibit NF- κ B-dependent inflammatory responses by uncoupling NF- κ B binding and enhancer/promoter histone acetylation from subsequent steps required for proinflammatory gene activation. This provides a mechanism for the *in vivo* anti-inflammatory insulin-sensitive phenotype observed in mice with macrophage-specific deletion of NCoR. Therapeutic methods to harness this mechanism could lead to a new approach to insulin-sensitizing therapies.

INTRODUCTION

Insulin resistance is a characteristic pathophysiologic defect in the great majority of patients with type 2 diabetes mellitus (Haffner and Taegtmeier, 2003; Olefsky and Glass, 2010; Reaven,

2005). Obesity is the most common cause of insulin resistance, and the current obesity epidemic in Westernized countries is driving the parallel type 2 diabetes epidemic (Shoelson et al., 2007). Obesity-associated chronic tissue inflammation is a key mechanism for decreased insulin sensitivity (Olefsky and Glass, 2010), and in obesity, excessive numbers of proinflammatory, M1-like macrophages accumulate in adipose tissue (Xu et al., 2003) and liver (Lanthier et al., 2010), where they locally release a variety of cytokines that act on insulin target cells to impair insulin signaling. Although other immune cell types such as lymphocytes, eosinophils, and neutrophils can also contribute to the tissue inflammatory state in obesity (Kintscher et al., 2008; Wu et al., 2011; Yang et al., 2010), their function in this context is largely to regulate macrophage migration and activation.

Inflammatory pathways in macrophages are under stringent control by a variety of transcription factors and coregulatory molecules. Among these are NF- κ B, AP1, the PPAR family, LXR, and their associated coactivators and corepressors. As a general rule, coactivators are recruited to nuclear receptors (NR) by receptor agonists, whereas corepressors are resident at promoter regions in the absence of ligands (Glass and Ogawa, 2006). The major corepressors, nuclear receptor corepressor (NCoR) and silencing mediator of retinoid and thyroid hormone receptors (SMRT), are important in the regulation of metabolic processes. In the basal state, NCoR is resident on a number of inflammatory pathway genes, keeping them in the repressed state (Medzhitov and Horng, 2009). Upon signal-dependent inflammatory pathway activation, NCoR dissociates from the promoter complex, allowing proinflammatory transcription factors, such as NF- κ B and AP1, to induce increased expression of these genes (Glass and Saijo, 2010; Ogawa et al., 2004).

Although NCoR can be detected on the promoter regions of genes that are targets of a variety of transcription factors, our

previous studies (Li et al., 2011), as well as those of others (Yamamoto et al., 2011), have shown that deletion of NCoR does not lead to broad derepression of a large number of transcriptional programs. For example, in adipocytes, the dominant effect of NCoR knockout (KO) is PPAR γ derepression (Li et al., 2011), whereas, in skeletal muscle, PPAR δ and myocyte enhancer factor (MEF) activation are the major effects of NCoR gene deletion (Yamamoto et al., 2011). This leads to the concept that NCoR has a restricted set of functions within the context of a particular cell type. Based on these ideas, we used Lys M-Cre-mediated excision (Clausen et al., 1999) of NCoR to generate macrophage/neutrophil NCoR KO mice. Although NCoR can cause basal repression of inflammatory pathways (Glass and Saijo, 2010), we unexpectedly found that macrophage NCoR KO mice displayed an anti-inflammatory, insulin-sensitive phenotype. Mechanistic studies traced this surprising phenotype to LXR derepression. Thus, a major effect of NCoR KO in macrophages is to derepress LXR, which leads to the induction of lipogenic pathway genes. This causes increased biosynthesis of palmitoleic acid (POA) and ω 3 fatty acids within macrophages, which exert robust local anti-inflammatory effects and are associated with the *in vivo* insulin sensitive phenotype.

RESULTS

Creation of Macrophage/Neutrophil-Specific NCoR Knockout Mice

Mice carrying “floxed” NCoR alleles, with LoxP sites flanking exon 11 of the NCoR gene, were crossbred with mice transgenic for Lys M-driven Cre expression. Both of these mouse lines have been extensively bred (>10 generations) to the C57B/6J background. Because Lys M is highly expressed in macrophage/neutrophil lineages, the resulting NCoR^{F/F}·Lys M-Cre mice exhibit NCoR deletion in these cell types. Consistent with our previous studies (Hevener et al., 2007), this system is highly efficient for generation of tissue-specific NCoR KO mice and resulted in 80%–90% deletion of NCoR from macrophages (Figures 1A and 1B). For convenience, these animals are hereafter referred to as MNKO mice.

Metabolic Studies in Wild-Type and MNKO Mice

MNKO mice exhibited normal fertility, and their offspring followed the predicted Mendelian pattern with no developmental abnormalities noted. On normal chow diet, wild-type (WT) and MNKO mice reached the same body weight, whereas the MNKO group was modestly more insulin sensitive during insulin tolerance test (ITT) testing (data not shown). When placed on 60% high-fat diet (HFD) for 12 weeks, WT and MNKO mice exhibited similar increases in body weight (Figure 1C) and fat mass (Figure 1D). Given the known function of NCoR to repress basal expression of macrophage inflammatory pathway genes, we predicted increased inflammation and insulin resistance in the HFD MNKO mice. Surprisingly, insulin and free fatty acid (FFA) levels were significantly lower in HFD KO mice compared to WT littermates (Figures 1E and 1F), indicating an improved overall metabolic phenotype. This phenotype was more robust following oral glucose administration, which revealed enhanced glucose tolerance (Figure 1G) in MNKO animals. Together, with

the reduced basal insulin levels, these results indicate that deletion of NCoR from macrophages confers a systemic insulin-sensitive phenotype.

To quantitate this effect and to identify tissue-specific responses, we performed hyperinsulinemic-euglycemic clamp studies (Li et al., 2011), which quantitatively assess insulin action in target tissues *in vivo*. The HFD MNKO mice were markedly more insulin sensitive compared to HFD WT mice. This was manifested by an increased overall glucose infusion rate (GIR) (Figure 1H) and glucose disposal rate (GDR) (Figure 1I). In addition, HFD MNKO mice exhibited enhanced insulin-stimulated glucose disposal rate (IS-GDR, a measure of skeletal muscle insulin sensitivity) (Figure 1J) and an increased ability of insulin to suppress hepatic glucose production (HGP, Figure 1K) and circulating FFA levels (Figure 1L). Thus, the MNKO mice displayed greater *in vivo* insulin sensitivity in all three major insulin target tissues, muscle, liver, and fat. Interestingly, treatment of HFD WT mice with rosiglitazone (3 weeks) led to the expected increase in systemic insulin sensitivity; however, the untreated MNKO mice were already insulin sensitive, and rosiglitazone treatment had little, if any, additive beneficial effect (Figures 1H–1L).

Inflammatory Signaling and Macrophage Function

Given the unexpected insulin-sensitive phenotype in the MNKO mice, we characterized inflammatory pathway signaling in WT and NCoR-depleted macrophages. We measured messenger RNA (mRNA) expression levels of a variety of inflammatory genes in the basal state and after TLR4 (KLA), TLR3 (Poly(I:C)), and TLR2 (Pam3CSK4) stimulation in thioglycollate-elicited intraperitoneal macrophages (*i.p.* Macs) obtained from WT and MNKO mice (Figure 2). Consistent with the *in vivo* results for glucose homeostasis and insulin sensitivity, basal gene expression was decreased in the MNKO macrophages (Figure 2A), and, more importantly, NCoR deletion led to broadly impaired inflammatory responses to TLR4, TLR2, and TLR3 agonists (Figures 2B–2D). Complementing the results in macrophages from MNKO mice, RNAi-mediated depletion of NCoR in WT macrophages results in the same attenuation of inflammatory gene responses (Figure 2E). Interestingly, a subset of genes characteristic of alternatively activated, M2-like, macrophages exhibited increased expression in NCoR KO cells in the basal state and after IL-4 stimulation (Figures 2F and 2G).

In light of the reduced inflammatory state in the MNKO mice, we went on to measure systemic aspects of HFD-mediated inflammation (Figure 3). As can be seen in Figures 3A–3D, circulating levels of multiple cytokines, typically released by macrophages, were lower in the MNKO mice compared to WT, which is fully consistent with the gene expression data in Figure 2. In addition, the circulating TNF α levels were lower in MNKO mice as compared to WT mice following lipopolysaccharide (LPS) administration *in vivo* (Figure S1A available online). Immunohistochemical analyses of adipose tissue sections were conducted by staining of adipose tissue sections for the macrophage marker F4/80 (Figure 3E), which revealed that adipose tissue from MNKO mice contained fewer macrophages compared to WT mice. We extended this finding by performing fluorescence-activated cell sorting (FACS) studies of adipose tissue




Figure 1. Improved Glucose Metabolism and Insulin Sensitivity of MNKO Mice

(A) Shown (top to bottom) are WT, floxed, and deleted NCoR gene loci. (B) Relative messenger RNA levels of NCoR in macrophages. Values are fold induction of gene expression normalized to the housekeeping gene 36B4 and expressed as mean \pm SEM. (C) Body weight of WT and KO mice on 60% HFD. (D) Epi-WAT mass. (E) Fasting blood insulin levels. (F) Fasting FFA levels. (G) Glucose tolerance tests. (H) GIR during hyperinsulinemic euglycemic clamp. (I) GDR. (J) IS-GDR. (K) Percent suppression of HGP by insulin (HGP suppression). (L) Percent suppression of free FA levels (FFA suppression). Values are expressed as mean \pm SEM. * $p < 0.05$, ** $p < 0.01$ for KO versus WT, or for comparisons as indicated.

NCoR KO i.p. Macs to migrate toward adipocyte-conditioned medium (CM) was substantially impaired compared to WT cells (Figure 3G), providing an additional mechanism for reduced ATMs and decreased inflammatory status in the MNKO mice.

Consistent with the reduced intrinsic-inflammatory state of NCoR-depleted macrophages and the decreased ATM accumulation in HFD MNKO mice, we also found decreased epididymal white adipose tissue (eWAT) expression of a number of proinflammatory genes typically associated with the inflammatory/insulin-resistant state (Figure 3H). A similar decrease in inflammatory gene expression was observed in perirenal, subcutaneous, and brown adipose tissue (BAT) from HFD MNKO mice (Figure S1B). Interestingly, although the overall degree of inflammation and ATM accumulation was less in subcutaneous WAT and BAT

stromal vascular cells (SVCs) (Figure 3F), which demonstrated reduced macrophage content, as well as decreased CD11c⁺ proinflammatory macrophages in the MNKO mice.

Macrophages and adipocytes are the cellular sources of chemokines in adipose tissue of obese mice, and the decreased expression of MCP-1 and keratinocyte-derived chemokine (KC) (Figure 3A) is one mechanism for decreased adipose tissue macrophage (ATM) accumulation. However, we considered whether the intrinsic chemotactic properties of NCoR-depleted macrophages were impaired. To this end, we conducted in vitro chemotaxis experiments. We found that the ability of

relative to eWAT and perirenal fat, the decreased inflammation in the MNKO fat depots was still evident. Similar to adipose tissue, hepatic inflammation also occurs in obesity with increased numbers of recruited hepatic macrophages (RHMs), which are distinct from Kupffer cells, and increased inflammatory gene expression. As seen in Figure S1C, macrophage markers (F4/80 and CD11c) and inflammatory gene expression are reduced in liver tissue from HFD MNKO mice compared to WT controls.

The mechanism of inflammation-induced insulin resistance largely involves local release of inflammatory mediators, which exert paracrine effects to directly diminish insulin signaling in

insulin target cells (Jager et al., 2007; Rotter et al., 2003). To assess this mechanism in MNKO and WT mice, CM was harvested from basal and LPS-treated WT and MNKO macrophages. L6 myocytes were then treated with CM, followed by measurements of insulin-stimulated glucose transport (Figure 3I). After treatment with basal CM, myocyte glucose transport was substantially higher with MNKO CM administration. LPS stimulation leads to a marked increase in cytokine release by macrophages, and consequently, CM harvested from the LPS-treated WT cells robustly decreased the ability of insulin to stimulate transport in myocytes (Figure 3I). In contrast, CM from LPS-treated MNKO cells was without effect to inhibit insulin action. These findings indicate that CM from WT macrophages contains factors (most likely cytokines), which attenuate insulin signaling, and that this is more pronounced after LPS stimulation. CM from the MNKO cells contains less insulin-resistance-promoting cytokines, which is consistent with the results in Figures 1, 2, 3, and S1. It is also possible that CM from MNKO cells contains factors that potentiate insulin-stimulated glucose transport.

Genome-wide Consequences of NCoR Deletion on Inflammatory Response Genes

The unexpected insulin-sensitive phenotype of MNKO mice and the hyporesponsiveness of the corresponding macrophages to Toll-like receptor (TLR) agonists prompted us to evaluate the consequences of NCoR deletion in macrophages on a genome-wide scale. We next used global run-on sequencing (GRO-seq) analysis to quantify nascent RNAs that are actively being transcribed in macrophages from WT and MNKO mice under basal conditions or upon activation with the TLR4 agonist KLA. GRO-seq thereby provides a more direct measure of transcription rates than total mRNA levels. Using this approach, we identified 3,397 expressed transcripts with RefSeq annotations that were >1.5-fold increased in MNKO versus WT macrophages under basal conditions, exemplified by *Mmp12* (Figure 4A). Conversely, 1,746 expressed transcripts were reduced >1.5-fold in MNKO compared to WT macrophages. Unexpectedly, the overall pattern of differential gene regulation under basal conditions is consistent with roles of NCoR in both positive and negative regulation of innate immune functions. In particular, downregulated genes were significantly enriched for functional annotations linked to immune response, defense response, and inflammatory response (Figure 4B). Furthermore, a number of the RefSeq transcripts induced more than 2-fold in WT macrophages following 1 hr of KLA treatment exhibited <50% of maximal activation in MNKO macrophages, as exemplified by *Nos2* and *Cxcl10* (Figures 4C and 4D). Indeed, a direct comparison of the KLA-stimulated induction of these induced genes in WT macrophages showed that ~26% of these genes achieved levels of induction in MNKO macrophages that were 66% or less than in WT macrophages (indicated by red data points in Figure 4E and by box plot in Figure S2). We define this subset of KLA-induced genes as hypo-responsive, and collectively, they specify a genome-wide signature of the hyporesponsive MNKO phenotype. Visualization of GRO-seq tag density for the hyporesponsive set of genes indicated no changes in levels of transcripts from paused Pol II at the transcriptional start site but indicated uniform reduction in tag

densities throughout the gene bodies under KLA-stimulated conditions (Figure 4F).

To investigate consequences of NCoR deletion on deposition of a histone acetylation mark associated with transcriptional activation (Hargreaves et al., 2009), we performed ChIP-seq analysis of histone H4 lysine 5 acetylation (H4K5ac) and of NCoR in WT and MNKO macrophages. As expected, H4K5ac was increased in the vicinity of NCoR binding sites in MNKO macrophages (Figure 4G), which is consistent with a role of NCoR in recruiting histone deacetylase 3 (HDAC3) to these locations (Figure S2B). Evaluation of H4K5ac at specific loci illustrated the expected increase at genes exhibiting upregulation in MNKO cells, exemplified by the *Mmp12* locus (Figure 4A). However, H4K5ac was also increased at genes exhibiting hyporesponsiveness to KLA stimulation, exemplified by the *Nos2* and *Cxcl10* genes (Figures 4C and 4D). As shown in Figure 4H, this pattern was observed for the entire set of hyporesponsive genes in MNKO cells. ChIP analysis confirmed that, in MNKO macrophages, HDAC3 binding is significantly reduced at NCoR sites on inflammatory loci demonstrating both hyperacetylation and TLR4 hyporesponsiveness, as exemplified by *Nos2*, *IL-1 β* , and *Cxcl10* (Figure S2C).

NCoR Deficiency Induces an LXR Program Leading to Altered Fatty Acid Metabolism

Based on these results, we evaluated genes that were derepressed in MNKO macrophages for functions that might negatively regulate the TLR4 signaling pathway. Interestingly, derepressed genes were significantly enriched for functional annotations linked to fatty acid metabolism, including the terms linoleic acid metabolism, biosynthesis of unsaturated fatty acids, and α linoleic acid metabolism (Figure 4B). Further, several of the derepressed genes encode enzymes involved in the synthesis of monounsaturated (MUFA) and polyunsaturated fatty acids (PUFA) that have been documented to have anti-inflammatory and/or insulin-sensitizing effects (Oh et al., 2010; Spann et al., 2012). These included *Elovl5*, *Fads1*, and *Fads2*, involved in the elongation and desaturation of long-chain fatty acids to generate anti-inflammatory PUFAs, and *Scd1/Scd2*, which catalyze the synthesis of 16:1 POA, suggested to be an insulin-sensitizing adipokine (Cao et al., 2008). Quantitative PCR assays confirmed significant upregulation of *Fads2*, *Elovl5*, and *Scd2* (Figure 5A).

Prior ChIP-seq studies of the LXR β cistrome in macrophages indicate that several of these genes are direct targets of liver X receptors (LXRs) (Heinz et al., 2010). Studies in LXR-deficient mice demonstrated that, in the unliganded state, LXRs mediate active repression at select LXR target genes dependent on interaction with NCoR (Wagner et al., 2003). Consistent with these prior findings, a significant fraction of known LXR target genes is derepressed in MNKO cells, including *Abca1* and *Fasn* (Figure 5A), suggesting that upregulation of LXR target genes involved in fatty acid metabolism in MNKO macrophages could be a consequence of locus-specific derepression of LXR. In agreement with this possibility, overlap of ChIP-seq data revealed a high degree of colocalization of NCoR with LXR β binding sites in the macrophage (Figure 5B). Given that motif analysis verifies that these sites as likely bona fide LXR binding sites




Figure 2. Reduced Inflammation in i.p. Macrophages of MNKO Mice

(A) Relative mRNA levels of inflammatory cytokines in i.p. macrophages without treatment.

(B) Relative mRNA levels of inflammatory cytokines in i.p. macrophages with TLR4 agonist treatment.

(C and D) (C) Relative mRNA levels of inflammatory cytokines in i.p. macrophages after TLR2 agonist treatment or (D) TLR3 agonist treatment.

(legend continued on next page)




Figure 3. Hypoinflammatory Phenotype of MNKO Mice

(A–D) Inflammatory cytokine expression in serum, including IL-6 (A), KC (B), MIP-1 α (C), and MCP-1 (D). (E) F4/80 staining in Epi-WAT of WT and MNKO mice. (F) FACS analysis of macrophages and CD11c-positive macrophage content in Epi-WAT. (G) Chemotaxis assay on the i.p. macrophages from WT and MNKO mice. (H) Proinflammatory cytokine expression in Epi-WAT from WT and MNKO. (I) Effect of conditioned medium from i.p. macrophages treated with LPS on glucose uptake in L6 cells. Values are expressed as mean \pm SEM. * $p < 0.05$, ** $p < 0.01$. See also [Figure S1](#).

(Figure 5C), and that NCoR tag densities demonstrating most significant enrichment precisely centered on LXR β peaks (Figure 5D), it is likely that loss of NCoR would have a significant

role at many of these loci containing LXR-NCoR complexes normally poised for active repression under basal conditions. Consistent with this, direct measure of LXR complex activity by

(E) Relative mRNA levels of NCoR and indicated inflammatory cytokines in control (siCTL) versus siNCoR-treated i.p. macrophages after TLR4 agonist treatment.

(F and G) Relative mRNA levels of M2-like cytokines in i.p. macrophages without IL-4 treatment or (G) with IL-4 treatment.

Values are relative to GAPDH and are expressed as mean \pm SEM. * $p < 0.05$, ** $p < 0.01$ for KO versus WT, or siCTL versus siNCoR in 2E.




Figure 4. Genome-wide Impact of NCoR Deletion

(A) UCSC genome browser image illustrating normalized tag counts for H4K5ac ChIP-seq, GRO-seq, and NCoR ChIP-seq (Barish et al., 2012) in WT and MNKO macrophages at the *Mmp12* locus.

(B) Functional KEGG annotations associated with genes demonstrating increased or reduced expression in MNKO macrophages by more than 1.5-fold.

(C and D) UCSC genome browser images corresponding to (A) for the *Nos2* and *Cxcl10* genomic loci.

(E) Scatter plot of fold change of genes induced >2-fold by 1 hr KLA treatment in WT macrophages versus fold change of the same genes in MNKO macrophages. Fold activation is plotted as normalized GRO-seq tag counts for genes comparing log₂ values of KLA versus vehicle treated. MNKO hypo-responsive genes are highlighted in red. The Pearson's correlation coefficient is reported in the figure ($p < 0.0001$).

(F) Normalized distribution of GRO-seq tag counts in the vicinity of the transcriptional start sites of TLR4-responsive genes exhibiting compromised activation in MNKO macrophages.

(G) Normalized distribution of H4K5ac ChIP-seq tag counts in the vicinity of NCoR binding sites under basal conditions.

(legend continued on next page)




Figure 5. Impact of NCoR KO on LXR Target Gene Expression in Macrophages

(A) qPCR analysis of the indicated LXR target genes in WT and MNKO macrophages. (B) Venn diagram of overlap between LXR β and NCoR peaks in macrophages. (C) Sequence motifs associated with LXR β and NCoR cobound sites. (D) Distribution of NCoR ChIP-seq tag density in the vicinity of LXR β binding sites. (E) LXR-dependent *Abca1*-luciferase reporter assays in WT and MNKO macrophages. (F) Distribution of H4K5ac ChIP-seq tag density in the vicinity of LXR β binding sites. (G) UCSC genome browser image illustrating normalized tag counts for H4K5ac ChIP-seq, GRO-seq, LXR β ChIP-seq, and NCoR ChIP-seq (Barish et al., 2012) at the *Scd2* locus. Values are expressed as mean \pm SEM. * $p < 0.05$ and ** $p < 0.01$ for KO versus WT.

a marked increase in H4K5ac (Figure 5G, upper track). Collectively, these data indicate that one significant consequence of NCoR deletion in the macrophage is derepression of LXR target genes, including those that control fatty acid metabolism.


These findings suggested that NCoR KO macrophages might exhibit reprogramming of fatty acid metabolism, generating lipid products with anti-inflammatory effects. To assess this, we performed a lipidomic analysis on WT and NCoR KO i.p. Macs (Figure 6A). 18:3n3 linolenic acid (ALA) is a precursor for the biosynthetic pathway, containing the gene products of *Fads1*, *Fads2*, and *Elovl5*, whose enzymatic activities enable production of the long chain ω 3 FAs 20:5n3(EPA) and 22:6n3(DHA). Importantly, ALA concentrations are markedly decreased in the NCoR KO macrophages, whereas levels of EPA and DHA

were 4- to 6-fold increased (Figure 6A). The potent anti-inflammatory effects of ω 3 FAs are well known (Mori and Beilin, 2004; Oh et al., 2010), providing a potential mechanistic explanation for the decreased overall inflammatory tone of the MNKO mice. Furthermore, Cao et al. (2008) identified palmitoleic acid (16:1n7, POA) as a “lipokine” that can exert insulin-like actions. In addition, recent studies have demonstrated that POA exhibits potent anti-inflammatory activity in macrophages (Spann et al.,

use of luciferase reporter constructs revealed that MNKO macrophages demonstrated significantly higher LXR reporter activity from a construct driven by LXR-binding sites from the *Abca1* locus (Figure 5E). Further, LXR binding sites were associated with increased H4K5ac in MNKO macrophages (Figure 5F) and increased expression of nearby genes, exemplified by *Scd2* (Figure 5G, track 2). Accordingly, the *Scd2* locus enhancer region occupied by LXR and NCoR (Figure 5G, tracks 3 and 4) indicates

(H) Normalized distribution of H4K5ac ChIP-seq tag counts in the vicinity of transcriptional start sites (TSS) under basal conditions. For Figures 4G and 4H, H4K5ac tag counts are presented as averages at indicated positions of KLA-activated genes demonstrating hyporesponsiveness in MNKO macrophages. TLR4 MNKO hyporesponsive loci are chosen according to those genes demonstrating 1.5-fold decrease in KLA response of MNKO relative to WT, as measured by comparison of normalized GRO-seq tag counts of MNKO versus WT macrophages.

See also Figure S2.



(legend on next page)

2012). In line with increased expression of *Scd1* and *Scd2*, the levels of POA were also 2- to 3-fold increased in the MNKO macrophages (Figure 6A). We also measured EPA, DHA, and POA concentrations within each of the major lipid classes, finding increases in the cholesterol ester and triacylglycerol fractions, but not in total phospholipids (Figures 6B–6D).

No appreciable differences in circulating EPA, DHA, or POA levels were observed in MNKO mice compared to WT mice (data not shown), suggesting that the biological consequences of increased production of these lipid species would most likely result from autocrine or paracrine mechanisms. To investigate this possibility, we first evaluated the impact of EPA on TLR responses in macrophages derived from WT and MNKO animals. As expected, EPA treatment strongly suppressed the KLA response of *Cxcl10*, *Nos2*, *Il1b*, *Tnf*, and *Irfg* in WT macrophages (Figure 6E). In MNKO macrophages, responses of *Cxcl10*, *Nos2*, *Il1b*, *Tnf*, and *Irfg* to TLR4 ligation were reduced, as previously noted. With the exception of *IL6*, the inhibitory effects of EPA were much less robust in MNKO versus WT macrophages, which is consistent with increased endogenous levels of ω 3 FAs in the MNKO macrophages.

To further investigate roles of endogenous POA and ω 3 FAs in determining the MNKO phenotype, we used an RNAi approach to knock down *Fads2*, *Elovl5*, and *Scd2* in both WT and MNKO macrophages. Knockdown of these enzymes responsible for the production of POA and ω 3 FAs (Figure S3) led to increased KLA-stimulated inflammatory gene expression of MNKO hyporesponsive genes, exemplified by *Cxcl10* and *Nos2* (Figures 6F–6H). The knockdowns demonstrated direct target specificity, as known LXR target genes such as *Abca1* and *Abcg1* were not reduced by the siRNAs, relative to the specific target genes *Fads2*, *Elovl5*, and *Scd2* (Figures S3A–S3C). Further, the siRNA-mediated derepression of TLR target genes was mainly selective for the hyporesponsive genes in the MNKO cells. Thus, many of the KLA-stimulated genes, either hyperresponsive or unaffected by loss of NCoR, failed to exhibit the same increases with *Fads* or *Scd2* knockdown (Figures S3B and S3C). Taken together, these data are consistent with the hypothesis that NCoR depletion derepresses LXR, leading to increased expression of select lipid metabolic target genes in the POA and ω 3 FA biosynthetic pathways.

ω 3 FAs Suppress Transcriptional Outputs of NF- κ B Regulatory Elements

To specifically address the relationship of the hyporesponsive phenotype of MNKO macrophages to NF- κ B-dependent gene expression, we assessed NF- κ B-driven reporter gene activity in MNKO and WT macrophages. Reporter activity of constructs, driven by either NF- κ B-bound enhancer regions from MNKO hyporesponsive genes (*Nos2* and *Cxcl10*) or a consensus NF-

κ B response element, exhibited significantly reduced activity in MNKO versus WT macrophages (Figure S4A). Next, we performed ChIP-seq measurements of the p65 component of NF- κ B to determine whether reduced NF- κ B activity was due to a reduction in DNA binding. Unexpectedly, ChIP-seq analysis revealed a very similar pattern of p65 binding following KLA stimulation in both WT and MNKO macrophages. Motif analysis confirmed these as bona fide p65 binding sites, with highest enrichment for NF- κ B motifs, as well as motifs for macrophage lineage-determining factors PU.1, AP-1, and C/EBP (Figure S4B). The similarity in p65 binding was observed not only for the entire set of p65 binding sites associated with TLR4-responsive genes induced > 2-fold in WT macrophages (Figure 7A) but also for the subset of p65 binding sites associated with MNKO hyporesponsive loci (Figures 7B–7D), exemplified by the *Cxcl10* locus (Figure 7E). These sites were also associated with increased H4K5ac following TLR4 stimulation (Figure 7F), indicating that binding of NF- κ B resulted in recruitment of histone acetyltransferases. These findings indicate that much of the reduction in expression of NF- κ B target genes in MNKO macrophages is independent of changes in NF- κ B binding and occurs at a step subsequent to histone acetylation. Recent studies have implicated histone H3 lysine 4 mono- and dimethylation (H3K4me1 and H3K4me2) as additional indicators of active local transcription (Ng et al., 2003; Ostuni et al., 2013). In an effort to resolve the paradox of increased histone acetylation and undiminished p65 binding at hyporesponsive genes in the MNKO cells, we performed ChIP-seq analysis of H3K4me2 in WT and MNKO macrophages. These studies revealed that MNKO macrophages demonstrated significantly reduced levels of TLR4-dependent H3K4me2 deposition at hyporesponsive genes, as shown for the *Cxcl10* locus (Figure 7E). This pattern of reduced H3K4me2 deposition, at both p65-bound enhancer elements and within the associated gene bodies, was observed for the entire set of hyporesponsive genes (Figures 7G and S4C). Conversely, genes that are derepressed in MNKO macrophages exhibit increased H3K4me2 levels, exemplified by the *Mmp12* locus (Figure S4D).

In addition to H3K4 methylation, recent studies have demonstrated that enhancers direct the expression of eRNAs that are linked to enhancer activity (Lai et al., 2013). Because GRO-seq analysis not only identifies nascent mRNA transcripts but also enables quantification of eRNAs, we also evaluated eRNA expression at p65 binding sites associated with hyporesponsive genes. This analysis indicated significantly reduced eRNA production in MNKO-derived macrophages following TLR4 activation (Figure 7H). Interestingly, in contrast to these defects in the NF- κ B transcriptional pathway, we found no differences in JNK activity between WT and MNKO macrophages (Figure S4E), which is consistent with both the transcriptomic results and the view that the primary effects of ω 3 FAs in the MNKO cells are

Figure 6. Increased ω 3 Fatty Acid in i.p. Macrophages from MNKO Mice

(A) ALA, EPA, DHA, and Pamitoleic acid content in vehicle or KLA-treated i.p. macrophages from WT and MNKO mice.

(B–D) (B) EPA level, (C) DHA level, and (D) pamitoleate level in different lipid fractions in i.p. macrophages from WT and MNKO mice.

(E) Effects of ω 3 fatty acid treatment on inflammatory gene expression in KLA-treated WT and MNKO i.p. macrophages.

(F–H) (F) KLA-induced inflammatory gene expression in i.p. macrophages after siRNA knockdown of *Elovl5*, *Fads2* (G), or *Scd2* (H).

Values are expressed as mean \pm SEM. * $p < 0.05$, ** $p < 0.01$, † $p < 0.001$ for KO versus WT, or EPA-KLA versus KLA in (E), ‡ or KLA-treated target-specific siRNA (*Scd2*, *Elovl5*, or *Fads2*) versus siCTL in (F)–(H). See also Figure S3.




Figure 7. Increased ω 3 FA Production Leads to Decreased NF- κ B Activity in i.p. Macrophages of MNO Mice

(A) Venn diagram of overlap between p65 peaks within 50 kb of TLR4-responsive genes (WT fold change > 2, RPKM > 0.1, and FDR < 0.05) in KLA-treated WT and MNO macrophages.

(B) Venn diagram of overlap between p65 peaks within 50 kb of MNO hypo-responsive genes in KLA-treated WT and MNO macrophages.

(C) Normalized distribution of for p65 ChIP-seq tag counts at enhancer-like regions associated with MNO hypo-responsive genes (defined in B) in vehicle or KLA-treated WT and MNO macrophages.

(D) Scatterplot of normalized tag counts for p65 at enhancer-like regions associated with TLR4-responsive genes (defined in A) in KLA-treated MNO and WT macrophages. The p65 peaks corresponding to hypo-responsive genes in MNO macrophages (defined in B) are highlighted in red. The Pearson's correlation coefficient is reported in the figure ($p < 0.0001$).

(legend continued on next page)

intracellular. Together, these findings provide evidence that loss of NCoR leads to uncoupling of p65 binding and local TLR-dependent histone acetylation from histone methylation and eRNA production, resulting in reduced signal-responsive gene expression.

To directly link these observations to effects of ω 3 FAs on NF- κ B activity, we evaluated effects of EPA and DHA on NF- κ B-driven reporter genes in primary WT macrophages. These experiments demonstrated the expected decrease in reporter gene activity (Figure S4F). Importantly, ChIP assays indicated that ω 3 treatment of WT macrophages resulted in decreased TLR-dependent deposition of H3K4me2 (Figure 7I) with no changes in p65 binding (Figure 7J) at p65 binding sites in enhancers associated with the *Nos2* and *Cxcl10* genes. These observations suggest that ω 3 fatty acids suppress the transcriptional functions of p65 at a step downstream of NF- κ B binding.

DISCUSSION

Here, we present evidence that deletion of NCoR from macrophages results in an anti-inflammatory phenotype with protection from systemic insulin resistance in the setting of HFD-induced obesity. This surprising phenotype in the MNKO mice was manifested by decreased macrophage accumulation in adipose tissue, decreased expression of inflammatory pathway genes, and reduced cytokine protein levels. In addition, we observed improved glucose tolerance in these animals, as well as systemic insulin sensitivity, as measured by the hyperinsulinemic euglycemic clamp method. Interestingly, treatment with rosiglitazone had little further effect to augment insulin sensitivity in the MNKO mice, possibly because they were already insulin sensitive in the untreated state or because the anti-inflammatory effects attributable to rosiglitazone (Bouhrel et al., 2007; Odegaard et al., 2007) require NCoR (Pascual et al., 2005). This improvement in inflammatory status and insulin sensitivity reinforces the connection between macrophage-mediated tissue inflammation and insulin resistance. In vitro studies further demonstrated that, although some inflammatory response genes were upregulated in MNKO cells, a substantially larger number exhibited hyporesponsiveness to TLR4 ligation.

NCoR and the related protein SMRT were initially discovered based on their functions as corepressors of retinoic acid and thyroid hormone receptors (Chen and Evans, 1995; Hörlein et al., 1995). Subsequent studies extended these corepressor functions to additional nuclear receptors, including LXRs (Hu et al., 2003), as well as a number of signal-dependent transcription factors such as AP-1, NF- κ B, and Ets (Mottis et al., 2013).

The corepressor activities of NCoR and SMRT are based on their ability to serve as platforms for the assembly of large corepressor complexes, with a core NCoR/SMRT complex containing Tbl1, TblR1, HDAC3, and GPS2 (Rosenfeld et al., 2006). HDAC3 contributes directly to the corepressor activity of NCoR/SMRT complexes by catalyzing the removal of histone acetylation marks that are involved in transcriptional activation (Codina et al., 2005).

Genome-wide analysis confirmed increased H4K5 acetylation in MNKO cells at genome sites where NCoR is bound in WT cells, which is consistent with loss of recruitment of HDAC3 due to the NCoR deletion. At these sites, acetylation of histone H4 at lysine residues is linked to recruitment of pTEFb, which is associated with increased transcriptional elongation (Yang et al., 2005). However, increased H4K5ac was observed not only at genes exhibiting increased expression in MNKO cells but also at the large subset of TLR4-responsive genes exhibiting impaired KLA stimulation (hyporesponsive gene set, Figure 4E). In addition, our p65 ChIP-seq results showed unimpaired p65 occupancy of the promoters/enhancers for these hyporesponsive genes. Thus, even when NF- κ B binds to its target genes at hyperacetylated sites, its transcriptional functions are compromised in NCoR-deficient macrophages. These observations indicate a paradoxical uncoupling of histone acetylation and p65 DNA binding from transcriptional activation.

Investigation of possible causes of the hyporesponsive phenotype of MNKO macrophages led to the finding of derepression of LXR target genes involved in generation of anti-inflammatory fatty acids, including long-chain ω 3-FAs and POA. Thus, NCoR can bind to LXR, mediating basal transcription repression at NCoR/LXR binding sites. In the absence of NCoR, LXR is derepressed, leading to increased transcription of these targets, which include the lipogenic genes *Elovl5*, *Fads1*, *Fads2*, *Scd1*, and *Scd2*. These lipogenic genes drive the biosynthesis of ω 3 FAs and POA within macrophages, effectively reprogramming macrophage lipid metabolism. Derepression of these genes was not observed in fetal liver-derived macrophages from systemic NCoR KO mice (Ghisletti et al., 2009), which may account for the relatively activated phenotype reported for those cells. Administration of these ω 3 FAs was less effective at inhibiting TLR4 responses in MNKO cells, consistent with the elevated endogenous levels. Conversely, knockdown of the lipogenic enzymes responsible for their production partially restored inflammatory responses in NCoR KO cells. These results support the conclusion that increased expression of anti-inflammatory FAs contribute to the hyporesponsive phenotype of MNKO cells. The relative importance of anti-inflammatory

(E) UCSC genome browser image illustrating normalized tag counts for H3K4me2 ChIP-seq, GRO-seq, p65 ChIP-seq, and NCoR ChIP-seq (from Barish et al., 2012) in WT and MNKO macrophages for the *Cxcl10* genomic locus as indicated.

(F) Normalized distribution of H4K5Ac ChIP-seq tag density in the vicinity of p65 binding sites at enhancers associated with MNKO hyporesponsive genes (defined in B) in vehicle or KLA-treated WT and MNKO macrophages.

(G) Normalized distribution of H3K4me2 ChIP-seq tag density in the vicinity of p65 binding sites at enhancers associated with MNKO hyporesponsive genes (defined in B) in KLA-treated cells.

(H) Normalized distribution of GRO-seq tag counts in the vicinity of p65 peaks associated with MNKO hyporesponsive genes (defined in B) in vehicle or KLA-treated cells.

(I) ChIP for H3K4me2 at *Nos2* and *Cxcl10* loci in WT macrophages treated with KLA in the presence or absence of EPA.

(J) ChIP for p65 at *Nos2* and *Cxcl10* loci in WT macrophages treated with KLA in the presence or absence of EPA.

Values are expressed as mean \pm SEM. * $p < 0.05$ and ** $p < 0.01$ for KLA versus EPA-KLA. See also Figure S4.

FAs appears to vary in a gene-specific manner, indicating that additional mechanisms contribute.

Genome-wide and locus-specific analyses indicate that ω 3 FAs provide at least two levels of suppression of inflammatory response genes. ω 3 FAs exert anti-inflammatory effects by serving as agonists for GPR120 on macrophages (Oh et al., 2010). GPR120 ligation inhibits proinflammatory signaling by blocking activation of TAK1 with reduced downstream signaling to IKK β /NF- κ B (Oh et al., 2010). This, in turn, would result in reduced translocation of NF- κ B to the nucleus. Consistent with this prediction, reduced occupancy of p65 was observed on a subset of inflammatory gene promoters and enhancers in the NCoR-deficient macrophages and in WT macrophages treated with ω 3 FAs. However, this effect was quantitative and locus specific, with the great majority of TLR4 hyporesponsive genes remaining largely or fully occupied by p65. Consistent with this, we found no decrease in JNK activity in the MNKO macrophages compared to WT. This indicates that, in MNKO cells, the major effects of endogenously produced ω 3 FAs are intracellular and largely independent of GPR120. Furthermore, these sites of p65 binding exhibited increased H4K5ac, which requires the active recruitment of histone acetyltransferases such as CREB-binding protein (CBP) and p300 by active NF- κ B. In contrast, H3K4 dimethylation and enhancer RNA production were consistently inhibited at genomic regulatory elements associated with the hyporesponsive genes. Previous studies have demonstrated local deposition of H3K4me2 as a valid measure of signal-responsive transcription factor activity (He et al., 2010). Further, signal-dependent inductions of H3K4me2 and transcriptional output from local transcription-factor-activated enhancers reflect the increased induction of associated genes (He et al., 2010; Ostuni et al., 2013). Reduced H3K4me2 levels were further correlated with reduced eRNA production at p65 enhancers associated with hyporesponsive genes, which is further indicative of a loss of transcriptional output of NF- κ B at these locations. Thus, although histone acetylation and p65 DNA binding are required for activation, they are not sufficient to promote transcriptional activation of the hyporesponsive inflammatory gene set in the context of NCoR deletion or ω 3 FA treatment.

Together, the present studies provide evidence in support of two main conclusions. First, these observations suggest that macrophages can be a source of sufficient ω 3 FAs and POA to exert autocrine and/or paracrine effects that result in improved insulin sensitivity. This concept is also exemplified by our findings that the normal effects of CM harvested from WT macrophages to inhibit insulin-stimulated glucose transport do not occur using CM from MNKO macrophages. That these ω 3 FAs effects are local is supported by our data demonstrating increased levels of ω 3 FAs in macrophages without changes in systemic concentrations. This concept raises the possibility that treatments, which could reprogram macrophage lipid metabolism into the more metabolically favorable state characteristic of NCoR macrophage KO mice, could have important therapeutic effects to reduce tissue inflammation and promote insulin sensitivity. Second, these findings reveal a molecular function of unsaturated ω 3 FAs by which they reduce H3K4 methylation, enhancer transcription, and mRNA expression at regulatory ele-

ments that control inflammatory response genes and thereby establish a hyporesponsive state. These effects are due to reduced NF- κ B complex activity independent of effects on NF- κ B localization itself. Identification of the molecular targets and mechanisms responsible for these effects could also lead to development of new approaches for diseases characterized by pathogenic forms of inflammation.

EXPERIMENTAL PROCEDURES

Animal Care and Use

Animals were maintained on a 12 hr/12 hr light/dark cycle with free access to food and water. All animal procedures were in accordance with University of California, San Diego research guidelines for the care and use of laboratory animals.

Creation of Control and Monocyte/Neutrophil-Specific NCoR KO Mice

NCoRfl/fl mice were generated as described previously (Li et al., 2011). To generate macrophage/neutrophil-specific NCoR null mice, NCoRfl/fl were bred with transgenic mice harboring Cre recombinase driven by myeloid-specific lysozyme M promoter (Clausen et al., 1999) to create the following genotypes: NCoRfl/fl (control) and NCoRfl/fl-LysMCre (MNKO). These mice were backcrossed to the C57BL/6J strain for more than ten generations.

Lipid Measurements

Thioglycollate-elicited peritoneal macrophages were plated at 2.0×10^6 cells per well of a 6-well plate in growth medium containing RPMI 1640 (with L-glutamine) (Invitrogen) plus 10% heat-inactivated FBS (Hyclone). After 24 hr, cells were washed twice with PBS and switched to culture medium containing RPMI 1640 (Invitrogen) without serum and supplemented with FA-free BSA (sigma) for 6 hr. Media and cells were collected for detection and quantitation of lipids by mass spectrometric methods as described previously (Brown et al., 2007).

Lipid Fraction Analyses in Macrophages

The lipids from thioglycollate-elicited macrophages were extracted in the presence of authentic internal standards by the method of Folch et al. (1957). Using chloroform:methanol (2:1 v/v), individual lipid classes within each extract were separated by liquid chromatography (Agilent Technologies model 1100 series). Each lipid class was transesterified in 1% sulfuric acid in methanol in a sealed vial under a nitrogen atmosphere at 100°C for 45 min. The resulting fatty acid methyl esters were extracted from the mixture with hexane containing 0.05% butylated hydroxytoluene and prepared for gas chromatography by sealing the hexane extracts under nitrogen. Fatty acid methyl esters were separated and quantified by capillary gas chromatography (Agilent Technologies model 6890) equipped with a 30 m DB-88MS capillary column (Agilent Technologies) and a flame-ionization detector.

High-Throughput Sequencing and Data Analysis

ChIP fragments were sequenced for 36 or 50 cycles on an Illumina Genome Analyzer or HiSeq 2000, respectively, according to the manufacturer's instructions. GRO-seq results were trimmed to remove A-stretches originating from the library preparation. Each sequence tag returned by the Illumina Pipeline was aligned to the mm9 assembly using ELAND, allowing up to two mismatches. Only tags that mapped uniquely to the genome were considered for further analysis.

Data analysis was performed using HOMER, and the detailed instructions for analysis can be found at <http://biowhat.ucsd.edu/homer/> or as previously described in Heinz et al. (2010). Each sequencing experiment was normalized to a total of 10^7 uniquely mapped tags by adjusting the number of tags at each position in the genome to the correct fractional amount given the total tags mapped. Sequencing experiments were visualized by preparing custom tracks for the UCSC genome browser. GRO-seq density within gene bodies was determined by adding the number of tags within the gene body defined

by RefSeq and normalizing by the length of the gene. Differentially expressed genes were identified as described previously (Escoubet-Lozach et al., 2011), using log values of gene body tag densities to calculate indicated fold change thresholds. For gene ontology analysis, DAVID Bioinformatics Resources 6.7 was used.

For additional methods, see the [Extended Experimental Procedures](#).

Statistical Analyses

Statistical analyses were performed using Excel or GraphPad Prism 5. The images were prepared using Photoshop CS5.1 or GraphPad Prism 5. Data are presented as the mean \pm SEM. For experiments involving two factors, data were analyzed by two-way ANOVA followed by Bonferroni post tests. Individual pair-wise comparisons were performed using Student's *t* test. $p < 0.05$ was considered significant.

ACCESSION NUMBERS

Sequencing data supporting these studies can be found at the Gene Expression Omnibus database under accession number GSE50944. Data from previously published ChIP-seq experiments can be found under accession numbers GSE27060 and GSE33596.

SUPPLEMENTAL INFORMATION

Supplemental Information includes Extended Experimental Procedures, four figures, and one table and can be found with this article online at <http://dx.doi.org/10.1016/j.cell.2013.08.054>.

ACKNOWLEDGMENTS

We thank the Flow Cytometry Resource (Dennis Young) for FACS analysis at the Rebecca and John Moores Cancer Center and thank Jachelle M. Ofrecio for genotyping at UCSD. This study was funded in part by grants to J.M.O. (DK033651, DK074868, DK-063491, and DK-09062), C.K.G. (DK074868, GM069338, and DK091183), and J.A. (the Ecole Polytechnique Federale de Lausanne, the Swiss National Science Foundation, and the EU Ideas program [ERC-AdG-23118]). This study was also funded by the Eunice Kennedy Shriver NICHD/NIH through a Cooperative Centers Program in Reproduction and Infertility Research to J.M.O.

Received: August 2, 2012

Revised: June 18, 2013

Accepted: August 27, 2013

Published: September 26, 2013

REFERENCES

- Barish, G.D., Yu, R.T., Karunasiri, M.S., Becerra, D., Kim, J., Tseng, T.W., Tai, L.J., Leblanc, M., Diehl, C., Cerchiotti, L., et al. (2012). The Bcl6-SMRT/NCoR2 cistrome represses inflammation to attenuate atherosclerosis. *Cell Metab.* **15**, 554–562.
- Bouhlef, M.A., Derudas, B., Rigamonti, E., Dièvert, R., Brozek, J., Haulon, S., Zawadzki, C., Jude, B., Torpier, G., Marx, N., et al. (2007). PPAR γ activation primes human monocytes into alternative M2 macrophages with anti-inflammatory properties. *Cell Metab.* **6**, 137–143.
- Brown, H.A., Henage, L.G., Preininger, A.M., Xiang, Y., and Exton, J.H. (2007). Biochemical analysis of phospholipase D. *Methods Enzymol.* **434**, 49–87.
- Cao, H., Gerhold, K., Mayers, J.R., Wiest, M.M., Watkins, S.M., and Hotamisligil, G.S. (2008). Identification of a lipokine, a lipid hormone linking adipose tissue to systemic metabolism. *Cell* **134**, 933–944.
- Chen, J.D., and Evans, R.M. (1995). A transcriptional co-repressor that interacts with nuclear hormone receptors. *Nature* **377**, 454–457.
- Clausen, B.E., Burkhardt, C., Reith, W., Renkawitz, R., and Förster, I. (1999). Conditional gene targeting in macrophages and granulocytes using LysMCre mice. *Transgenic Res.* **8**, 265–277.
- Codina, A., Love, J.D., Li, Y., Lazar, M.A., Neuhaus, D., and Schwabe, J.W. (2005). Structural insights into the interaction and activation of histone deacetylase 3 by nuclear receptor corepressors. *Proc. Natl. Acad. Sci. USA* **102**, 6009–6014.
- Escoubet-Lozach, L., Benner, C., Kaikkonen, M.U., Lozach, J., Heinz, S., Spann, N.J., Crotti, A., Stender, J., Ghisletti, S., Reichart, D., et al. (2011). Mechanisms establishing TLR4-responsive activation states of inflammatory response genes. *PLoS Genet.* **7**, e1002401.
- Folch, J., Lees, M., and Sloane Stanley, G.H. (1957). A simple method for the isolation and purification of total lipides from animal tissues. *J. Biol. Chem.* **226**, 497–509.
- Ghisletti, S., Huang, W., Jepsen, K., Benner, C., Hardiman, G., Rosenfeld, M.G., and Glass, C.K. (2009). Cooperative NCoR/SMRT interactions establish a corepressor-based strategy for integration of inflammatory and anti-inflammatory signaling pathways. *Genes Dev.* **23**, 681–693.
- Glass, C.K., and Ogawa, S. (2006). Combinatorial roles of nuclear receptors in inflammation and immunity. *Nat. Rev. Immunol.* **6**, 44–55.
- Glass, C.K., and Saijo, K. (2010). Nuclear receptor transrepression pathways that regulate inflammation in macrophages and T cells. *Nat. Rev. Immunol.* **10**, 365–376.
- Haffner, S., and Taegtmeier, H. (2003). Epidemic obesity and the metabolic syndrome. *Circulation* **108**, 1541–1545.
- Hargreaves, D.C., Horng, T., and Medzhitov, R. (2009). Control of inducible gene expression by signal-dependent transcriptional elongation. *Cell* **138**, 129–145.
- He, H.H., Meyer, C.A., Shin, H., Bailey, S.T., Wei, G., Wang, Q., Zhang, Y., Xu, K., Ni, M., Lupien, M., et al. (2010). Nucleosome dynamics define transcriptional enhancers. *Nat. Genet.* **42**, 343–347.
- Heinz, S., Benner, C., Spann, N., Bertolino, E., Lin, Y.C., Laslo, P., Cheng, J.X., Murre, C., Singh, H., and Glass, C.K. (2010). Simple combinations of lineage-determining transcription factors prime cis-regulatory elements required for macrophage and B cell identities. *Mol. Cell* **38**, 576–589.
- Hevener, A.L., Olefsky, J.M., Reichart, D., Nguyen, M.T., Bandyopadhyay, G., Leung, H.Y., Watt, M.J., Benner, C., Febbraio, M.A., Nguyen, A.K., et al. (2007). Macrophage PPAR γ is required for normal skeletal muscle and hepatic insulin sensitivity and full antidiabetic effects of thiazolidinediones. *J. Clin. Invest.* **117**, 1658–1669.
- Hörlein, A.J., Näär, A.M., Heinzel, T., Torchia, J., Gloss, B., Kurokawa, R., Ryan, A., Kamei, Y., Söderström, M., Glass, C.K., et al. (1995). Ligand-independent repression by the thyroid hormone receptor mediated by a nuclear receptor co-repressor. *Nature* **377**, 397–404.
- Hu, X., Li, S., Wu, J., Xia, C., and Lala, D.S. (2003). Liver X receptors interact with corepressors to regulate gene expression. *Mol. Endocrinol.* **17**, 1019–1026.
- Jager, J., Grémeaux, T., Cormont, M., Le Marchand-Brustel, Y., and Tanti, J.F. (2007). Interleukin-1 β -induced insulin resistance in adipocytes through down-regulation of insulin receptor substrate-1 expression. *Endocrinology* **148**, 241–251.
- Kintscher, U., Hartge, M., Hess, K., Foryst-Ludwig, A., Clemenz, M., Wabitsch, M., Fischer-Posovszky, P., Barth, T.F., Dragun, D., Skurk, T., et al. (2008). T-lymphocyte infiltration in visceral adipose tissue: a primary event in adipose tissue inflammation and the development of obesity-mediated insulin resistance. *Arterioscler. Thromb. Vasc. Biol.* **28**, 1304–1310.
- Lai, F., Orom, U.A., Cesaroni, M., Beringer, M., Taatjes, D.J., Blobel, G.A., and Shiekhattar, R. (2013). Activating RNAs associate with Mediator to enhance chromatin architecture and transcription. *Nature* **494**, 497–501.
- Lanthier, N., Molendi-Coste, O., Horsmans, Y., van Rooijen, N., Cani, P.D., and Leclercq, I.A. (2010). Kupffer cell activation is a causal factor for hepatic insulin resistance. *Am. J. Physiol. Gastrointest. Liver Physiol.* **298**, G107–G116.
- Li, P., Fan, W., Xu, J., Lu, M., Yamamoto, H., Auwerx, J., Sears, D.D., Talukdar, S., Oh, D., Chen, A., et al. (2011). Adipocyte NCoR2 knockout decreases PPAR γ phosphorylation and enhances PPAR γ activity and insulin sensitivity. *Cell* **147**, 815–826.

- Medzhitov, R., and Horng, T. (2009). Transcriptional control of the inflammatory response. *Nat. Rev. Immunol.* *9*, 692–703.
- Mori, T.A., and Beilin, L.J. (2004). Omega-3 fatty acids and inflammation. *Curr. Atheroscler. Rep.* *6*, 461–467.
- Mottis, A., Mouchiroud, L., and Auwerx, J. (2013). Emerging roles of the corepressors NCoR1 and SMRT in homeostasis. *Genes Dev.* *27*, 819–835.
- Mullican, S.E., Gaddis, C.A., Alenghat, T., Nair, M.G., Giacomin, P.R., Everett, L.J., Feng, D., Steger, D.J., Schug, J., Artis, D., and Lazar, M.A. (2011). Histone deacetylase 3 is an epigenomic brake in macrophage alternative activation. *Genes Dev.* *25*, 2480–2488.
- Ng, H.H., Robert, F., Young, R.A., and Struhl, K. (2003). Targeted recruitment of Set1 histone methylase by elongating Pol II provides a localized mark and memory of recent transcriptional activity. *Mol. Cell* *11*, 709–719.
- Odegaard, J.I., Ricardo-Gonzalez, R.R., Goforth, M.H., Morel, C.R., Subramanian, V., Mukundan, L., Red Eagle, A., Vats, D., Brombacher, F., Ferrante, A.W., and Chawla, A. (2007). Macrophage-specific PPARgamma controls alternative activation and improves insulin resistance. *Nature* *447*, 1116–1120.
- Ogawa, S., Lozach, J., Jepsen, K., Sawka-Verhelle, D., Perissi, V., Sasik, R., Rose, D.W., Johnson, R.S., Rosenfeld, M.G., and Glass, C.K. (2004). A nuclear receptor corepressor transcriptional checkpoint controlling activator protein 1-dependent gene networks required for macrophage activation. *Proc. Natl. Acad. Sci. USA* *101*, 14461–14466.
- Oh, D.Y., Talukdar, S., Bae, E.J., Imamura, T., Morinaga, H., Fan, W., Li, P., Lu, W.J., Watkins, S.M., and Olefsky, J.M. (2010). GPR120 is an omega-3 fatty acid receptor mediating potent anti-inflammatory and insulin-sensitizing effects. *Cell* *142*, 687–698.
- Olefsky, J.M., and Glass, C.K. (2010). Macrophages, inflammation, and insulin resistance. *Annu. Rev. Physiol.* *72*, 219–246.
- Ostuni, R., Piccolo, V., Barozzi, I., Polletti, S., Termanini, A., Bonifacio, S., Curina, A., Prosperini, E., Ghisletti, S., and Natoli, G. (2013). Latent enhancers activated by stimulation in differentiated cells. *Cell* *152*, 157–171.
- Pascual, G., Fong, A.L., Ogawa, S., Gamliel, A., Li, A.C., Perissi, V., Rose, D.W., Willson, T.M., Rosenfeld, M.G., and Glass, C.K. (2005). A SUMOylation-dependent pathway mediates transrepression of inflammatory response genes by PPAR-gamma. *Nature* *437*, 759–763.
- Reaven, G.M. (2005). The insulin resistance syndrome: definition and dietary approaches to treatment. *Annu. Rev. Nutr.* *25*, 391–406.
- Rosenfeld, M.G., Lunyak, V.V., and Glass, C.K. (2006). Sensors and signals: a coactivator/corepressor/epigenetic code for integrating signal-dependent programs of transcriptional response. *Genes Dev.* *20*, 1405–1428.
- Rotter, V., Nagaev, I., and Smith, U. (2003). Interleukin-6 (IL-6) induces insulin resistance in 3T3-L1 adipocytes and is, like IL-8 and tumor necrosis factor-alpha, overexpressed in human fat cells from insulin-resistant subjects. *J. Biol. Chem.* *278*, 45777–45784.
- Shoelson, S.E., Herrero, L., and Naaz, A. (2007). Obesity, inflammation, and insulin resistance. *Gastroenterology* *132*, 2169–2180.
- Spann, N.J., Garmire, L.X., McDonald, J.G., Myers, D.S., Milne, S.B., Shibata, N., Reichart, D., Fox, J.N., Shaked, I., Heudobler, D., et al. (2012). Regulated accumulation of desmosterol integrates macrophage lipid metabolism and inflammatory responses. *Cell* *151*, 138–152.
- Wagner, B.L., Valledor, A.F., Shao, G., Daige, C.L., Bischoff, E.D., Petrowski, M., Jepsen, K., Baek, S.H., Heyman, R.A., Rosenfeld, M.G., et al. (2003). Promoter-specific roles for liver X receptor/corepressor complexes in the regulation of ABCA1 and SREBP1 gene expression. *Mol. Cell Biol.* *23*, 5780–5789.
- Wu, D., Molofsky, A.B., Liang, H.E., Ricardo-Gonzalez, R.R., Jouihan, H.A., Bando, J.K., Chawla, A., and Locksley, R.M. (2011). Eosinophils sustain adipose alternatively activated macrophages associated with glucose homeostasis. *Science* *332*, 243–247.
- Xu, H., Barnes, G.T., Yang, Q., Tan, G., Yang, D., Chou, C.J., Sole, J., Nichols, A., Ross, J.S., Tartaglia, L.A., and Chen, H. (2003). Chronic inflammation in fat plays a crucial role in the development of obesity-related insulin resistance. *J. Clin. Invest.* *112*, 1821–1830.
- Yamamoto, H., Williams, E.G., Mouchiroud, L., Cantó, C., Fan, W., Downes, M., Héligon, C., Barish, G.D., Desvergne, B., Evans, R.M., et al. (2011). NCoR1 is a conserved physiological modulator of muscle mass and oxidative function. *Cell* *147*, 827–839.
- Yang, Z., Yik, J.H., Chen, R., He, N., Jang, M.K., Ozato, K., and Zhou, Q. (2005). Recruitment of P-TEFb for stimulation of transcriptional elongation by the bromodomain protein Brd4. *Mol. Cell* *19*, 535–545.
- Yang, H., Youm, Y.H., Vandanmagsar, B., Ravussin, A., Gimble, J.M., Greenway, F., Stephens, J.M., Mynatt, R.L., and Dixit, V.D. (2010). Obesity increases the production of proinflammatory mediators from adipose tissue T cells and compromises TCR repertoire diversity: implications for systemic inflammation and insulin resistance. *J. Immunol.* *185*, 1836–1845.