

telling the whole story. Under physiological conditions a cell membrane operates very close to the melting transition, i.e., to the point where the fluid membrane becomes a solid gel. Recent theoretical work indicates that signal propagation in a nerve cell may also involve a thermo-acoustic pulse of partial gellification. Natural selection should have led to optimal propagation under physiological conditions.

When apolar molecules are dissolved in the apolar membrane of the nerve cell, the freezing temperature of the membrane is lowered. This would interfere with pulse propagation and thus lead to anesthesia. However, if the theory is right, the effect should be reversed if we let the propagation take place at lower temperature. This is because the lower temperature would bring us closer again to the freezing transition.

We experimentally test this idea on the sciatic nerve of frogs. We follow the propagation of a signal with different concentrations of Argon in the medium and at different temperatures. Argon is an anesthetic that is chemically inert and it is expected to have its anesthetic effect just through interfering with the fluidity of the membrane. As a control we also perform the same experiments with Lidocain as the involved anesthetic. Lidocain is an anesthetic that is well known to work through interfering with voltage gated sodium channels.

#### 494-Pos Board B373

##### Ci-VSP Is A Depolarization-Activated PI(4,5)P<sub>2</sub> And PI(3,4,5)P<sub>3</sub> 5' Phosphatase

Christian R. Halaszovich, Dominik Oliver.

University Marburg, Marburg, Germany.

Phosphoinositides are membrane-delimited regulators of protein function and control many different cellular targets. The differentially phosphorylated isoforms have distinct concentrations in various subcellular membranes, which can change dynamically in response to cellular signaling events. Maintenance and dynamics of phosphoinositide levels involve a complex set of enzymes, among them phospholipases and lipid kinases and phosphatases. Recently, a novel type of phosphoinositide-converting protein, termed Ci-VSP, was isolated, which contains a voltage sensor domain. It was already shown that Ci-VSP can alter phosphoinositide levels in a voltage-dependent manner. However, the exact enzymatic reaction catalyzed by Ci-VSP is not known. We used fluorescent phosphoinositide-binding probes and total internal reflection microscopy together with patch-clamp measurements from living cells to delineate substrates and products of Ci-VSP. Upon activation of Ci-VSP by membrane depolarization, membrane association of PI(4,5)P<sub>2</sub> and PI(3,4,5)P<sub>3</sub>-specific binding domains decreased, revealing consumption of these phosphoinositides by Ci-VSP. Depletion of PI(4,5)P<sub>2</sub> was coincident with an increase in membrane PI(4)P. Similarly, PI(3,4)P<sub>2</sub> was generated during depletion of PI(3,4,5)P<sub>3</sub>. These results suggest that Ci-VSP acts as a 5'-phosphatase of PI(4,5)P<sub>2</sub> and PI(3,4,5)P<sub>3</sub>.

## IP<sub>3</sub> Receptors

#### 495-Pos Board B374

##### Toward A Computational Model Of IP<sub>3</sub>R1-associated Ataxia

Sherry-Ann Brown, Leslie M. Loew.

University of Connecticut Health Center, Farmington, CT, USA.

Individuals with ataxia suffer impaired imbalance and incoordination of motor functions. Approximately 150,000 Americans are afflicted with ataxia, as are thousands of individuals worldwide. Among these are families with reduced levels of IP<sub>3</sub>R1 protein, the primary receptor for IP<sub>3</sub> in cerebellar Purkinje neurons. Mice with reduced levels of IP<sub>3</sub>R1 are also ataxic; cerebellar microsomes from IP<sub>3</sub>R1 knockout mice exhibit little calcium release when probed with IP<sub>3</sub>. This suggests that altered calcium response to IP<sub>3</sub> may mediate the pathophysiology of cerebellar ataxia associated with reduced IP<sub>3</sub>R1. Currently, there are no direct therapeutics for hereditary ataxias. We hypothesized that adjusting IP<sub>3</sub>R1 sensitivity to IP<sub>3</sub> in the context of reduced IP<sub>3</sub>R1 could restore normal calcium response. To investigate our hypothesis, we adapted a computational compartmental model of a cerebellar Purkinje neuron previously published by our laboratory, using optimal parameters for calcium release. These parameters were dependent on the shape of the IP<sub>3</sub> signal produced from PIP<sub>2</sub> hydrolysis, determined in a recent study published by our group. In our optimized model, we reduced the value of  $J_{max}$ , the variable representing IP<sub>3</sub>R1 abundance in Purkinje spines, to 50%, 40%, 30%, 20%, and 10% of the normal level of IP<sub>3</sub>R1 found in mouse cerebellum. Next, we adjusted the sensitivity of IP<sub>3</sub>R1 to IP<sub>3</sub> in a similar cumulative fashion to see whether increasing sensitivity could rescue low abundance. We did this by varying values for  $d_{IP_3}$ , the dissociation constant for IP<sub>3</sub> from the receptor. We found that corresponding increases in IP<sub>3</sub>R1 sensitivity to IP<sub>3</sub> restored normal calcium response when IP<sub>3</sub>R1 abundance was reduced to as low as 30% of its normal value.

This promises significant therapeutic benefit for individuals with 'IP<sub>3</sub>R1-associated ataxia', as the phosphorylation status of IP<sub>3</sub>R1 can be regulated experimentally to adjust its sensitivity. (Supported by NIH RR013186)

#### 496-Pos Board B375

##### Electron Cryomicroscopy of IP<sub>3</sub>R1 Calcium Release Channel

Que T. Ngo<sup>1</sup>, Joshua T. Maxwell<sup>2</sup>, Gregory A. Mignery<sup>2</sup>, Wah Chiu<sup>3</sup>, Steven J. Ludtke<sup>3</sup>, Irina I. Serysheva<sup>1</sup>.

<sup>1</sup>The University of Texas Medical School, Houston, TX, USA, <sup>2</sup>Stritch School of Medicine Loyola University at Chicago, Maywood, IL, USA,

<sup>3</sup>Baylor College of Medicine, Houston, TX, USA.

The inositol 1, 4, 5 - trisphosphate receptor (IP<sub>3</sub>R) is an intracellular Ca<sup>2+</sup> release channel that mediates ligand-gated release of Ca<sup>2+</sup> from the endoplasmic reticulum (ER) into the cytoplasm. IP<sub>3</sub>R1 is the predominant type in the cerebellar ER membrane where it forms homotetramers with a  $M_r$  over 1.2 MDa. The gating of IP<sub>3</sub>R1 channel is still poorly understood due to the lack of high-resolution structure of the channel complex. Although several low-resolution 3D structures of the IP<sub>3</sub>R1 were reported, these 3D maps are broadly consistent in the overall size and shape. To achieve a reliable structure of IP<sub>3</sub>R1 channel at higher resolution, substantial improvements were made to cryo-specimen preparations that allowed acquiring electron images of ice-embedded channel protein, which exhibit substantially improved contrast and image quality. The structure of IP<sub>3</sub>R1 was analyzed under conditions favoring the closed channel conformation, i.e. in the absence of the two co-agonists, Ca<sup>2+</sup> and IP<sub>3</sub>. Ice-embedded IP<sub>3</sub>R1 particles were imaged at 60,000X magnification on a JEOL 2010F electron cryomicroscope with a Gatan 4k x 4k CCD camera. Image processing and the reconstruction were performed using EMAN. The improved map clearly exhibits more structural detail in both the cytoplasmic and membrane-spanning regions of the channel, connected through the stalk-like region. Available x-ray structures of the IP<sub>3</sub>-binding core region (pdb code: 1N4K) and the ligand binding suppressor domain (pdb code: 1XZZ) were docked into the cryo-EM density map to interpret visualized structural domains. Currently, structural analysis of IP<sub>3</sub>R1 in other physiologically relevant functional states is being performed to reveal the gating mechanism of the IP<sub>3</sub>R1 channel.

This research is supported by grants from NIH (R01GM072804, P41RR02250).

#### 497-Pos Board B376

##### The Amplification Of InsP<sub>3</sub>R Activity By NCS-1 Is Attenuated By Medications Used In The Treatment Of Bipolar Disorder

Christin Schulze<sup>1,2</sup>, Jessica Olofsson<sup>1,3</sup>, Barbara E. Ehrlich<sup>1</sup>.

<sup>1</sup>Yale University, New Haven, CT, USA, <sup>2</sup>Friedrich-Schiller University, Jena, Germany, <sup>3</sup>Chalmers University of Technology, Gothenburg, Sweden.

Neuronal Calcium Sensor-1 (NCS-1) is a high-affinity, low-capacity calcium-binding protein abundantly expressed in neuronal and neuroendocrine cells. We previously showed that NCS-1 interacts with the inositol 1,4,5-trisphosphate receptor (InsP<sub>3</sub>R) and modulates calcium signaling by enhancing InsP<sub>3</sub>-dependent InsP<sub>3</sub>R channel activity and intracellular calcium transients. Furthermore, it is known that NCS-1 is overexpressed in the prefrontal cortex of bipolar disorders and schizophrenic patients. Because we had reported that addition of lithium, a compound used for treatment of bipolar disorders, attenuates the NCS-1/InsP<sub>3</sub>R association, we hypothesized that other medications used for these disorders also might target the interaction between NCS-1 and the InsP<sub>3</sub>R. After overexpressing NCS-1 in a human neuroblastoma cell line to simulate the situation in the prefrontal cortex of bipolar patients, and using calcium sensitive dyes, we assessed the effect of the three main categories of medications used in bipolar disease on InsP<sub>3</sub>R-dependent intracellular calcium transients. We found that long-term treatment (8h) of cells overexpressing NCS-1 with therapeutic concentrations of chlorpromazine (CPZ) or valproic acid (VPA) attenuate the amplification effect of NCS-1 on InsP<sub>3</sub>-mediated Ca<sup>2+</sup> release. This finding is dependent on NCS-1 overexpression and was not observed in cells with reduced NCS-1 levels due to shRNA mediated NCS-1 knockdown. Furthermore, no alterations due to treatment were observed in either the calcium loading of the intracellular stores or in the expression level of NCS-1 or InsP<sub>3</sub>R. Therefore, the treatment with all three main categories of bipolar medications - lithium, anti-convulsants like VPA and antipsychotics like CPZ - appear to target the interaction between NCS-1 and the InsP<sub>3</sub>R. This study suggests a new approach to investigating and understanding the etiology and treatment of bipolar disorder.

#### 498-Pos Board B377

##### The Role of the Pore-forming Region in the Regulation of IP<sub>3</sub> Receptor by Luminal Ca<sup>2+</sup>

Shitian Cai, Wenqian Chen, Lin Zhang, Wayne S.R. Chen.

University of Calgary, Calgary, AB, Canada.

It is well known that submaximal concentrations of IP<sub>3</sub> release only a portion of the intracellular Ca<sup>2+</sup> store via the IP<sub>3</sub> receptor (IP<sub>3</sub>R), a phenomenon known

as "quantal" Ca<sup>2+</sup> release. Such quantal behavior of IP<sub>3</sub>R is thought to be due to the feedback regulation of the channel by luminal Ca<sup>2+</sup>. A high level of luminal Ca<sup>2+</sup> enhances the sensitivity of IP<sub>3</sub>R to IP<sub>3</sub>, while a reduced luminal Ca<sup>2+</sup> level desensitizes IP<sub>3</sub>R. Despite its importance, the molecular basis underlying the regulation of IP<sub>3</sub>R by luminal Ca<sup>2+</sup> is unknown. Ryanodine receptors (RyRs), another family of intracellular Ca<sup>2+</sup> release channels, also exhibit quantal Ca<sup>2+</sup> release in response to agonists, and are regulated by luminal Ca<sup>2+</sup>. We have recently demonstrated that mutations in the TM10 helix (the pore inner helix) of the RyR2 channel markedly alter the sensitivity of the channel to activation by luminal Ca<sup>2+</sup>. Given the high degree of sequence homology in the channel pore-forming region between RyR and IP<sub>3</sub>R, we hypothesize that the TM6 helix in IP<sub>3</sub>R, corresponding to TM10 in RyR, is also important for luminal Ca<sup>2+</sup> regulation of IP<sub>3</sub>R. To test this hypothesis, we have generated a number of mutations in the TM6 of IP<sub>3</sub>R and established stable, inducible HEK293 cell lines expressing these mutants. By monitoring the ER luminal Ca<sup>2+</sup> level using a fluorescent ER Ca<sup>2+</sup> sensor protein, D1ER, we found that mutations in TM6 either increase or decrease the rate of IP<sub>3</sub>-induced Ca<sup>2+</sup> release in permeabilized mutant cells. These mutations also affect the sensitivity of ATP-triggered Ca<sup>2+</sup> release in intact cells. Further studies at the single channel level should provide new insights into the role of the pore-forming region in the luminal Ca<sup>2+</sup> regulation of IP<sub>3</sub>R.

#### 499-Pos Board B378

##### A C-terminal Fragment of Chromogranin B Amplifies Inositol (1,4,5)-Triphosphate Receptor Mediated Signaling

Stefan Schmidt<sup>1,2</sup>, Felix M. Heidrich<sup>1,3</sup>, Michelle Mo<sup>1</sup>, Barbara E. Ehrlich<sup>1</sup>.

<sup>1</sup>Yale University, New Haven, CT, USA, <sup>2</sup>University of Goettingen, Goettingen, Germany, <sup>3</sup>Dresden University of Technology, Dresden, Germany.

Chromogranin B (CGB) is a low affinity, high capacity calcium binding protein belonging to the granin family. It is located in the lumen of the endoplasmic reticulum (ER) and is also found in secretory granules. Addition of CGB amplifies calcium release from ER stores and increases the activity of the inositol (1,4,5)-triphosphate receptor (InsP<sub>3</sub>R). We previously demonstrated that CGB is non-uniformly distributed within neurons, and its spatial localization is cell type specific. We also showed that stimulation of the InsP<sub>3</sub>R in neurons leads to initiation of intracellular calcium release where the concentration of CGB is highest. When we expressed the N-terminal region of CGB, which binds to the third intraluminal loop of the InsP<sub>3</sub>R, the functional interaction between CGB and the InsP<sub>3</sub>R was disrupted and the initiation site of calcium release was altered. We now report that a 20 amino acid fragment of the C-terminal region plays a critical role in regulating calcium transients from the InsP<sub>3</sub>R. Addition of the C-terminal region of CGB increased the activity of single InsP<sub>3</sub>R currents in lipid bilayers. When intracellular calcium transients were monitored in 3T3 cells lacking CGB, InsP<sub>3</sub>R dependent calcium release was markedly amplified after expression of full length CGB or expression of the C-terminal region. In contrast, expression of the N-terminal region was unable to amplify the intracellular calcium transients. In SHSY5Y cells with endogenous CGB, expression of the C-terminal region induced a prolonged response to extracellular agonists compared to native cells whereas expression of the N-terminal region depressed calcium signaling and altered the signal initiation site. These effects of CGB on calcium transients in neuronal cells indicate its importance in physiological processes and will guide investigation into pathophysiological processes.

#### 500-Pos Board B379

##### Comparison of IP<sub>3</sub>R and RyR Expression and Ca<sup>2+</sup> Release Characteristics in Isolated Cardiac Nuclei

Susan Currie<sup>1</sup>, Richard D. Rainbow<sup>2</sup>, Marie-ann Ewart<sup>1</sup>, John G. McCarron<sup>1</sup>.

<sup>1</sup>University of Strathclyde, Glasgow, United Kingdom, <sup>2</sup>University of Leicester, Leicester, United Kingdom.

In cardiac muscle, the role of the inositol triphosphate receptor (IP<sub>3</sub>R) and its regulation is not fully understood. A contribution to nuclear Ca<sup>2+</sup> signalling has been proposed. This study compares expression and Ca<sup>2+</sup> release characteristics of the IP<sub>3</sub>R and the ryanodine receptor (RyR) in purified functional cardiac nuclei. It also examines whether the IP<sub>3</sub>R may exist as a multi-protein complex in these preparations. Quantitative immunoblotting of IP<sub>3</sub>R and RyR protein levels in isolated nuclei demonstrated greater expression of the IP<sub>3</sub>R; nucleolin was used as an internal control for quantification. Ca<sup>2+</sup> release in response to IP<sub>3</sub> and caffeine from single isolated nuclei was used to compare IP<sub>3</sub>R and RyR activity. Changes in nuclear [Ca<sup>2+</sup>]<sub>Nuc</sub> were measured as fluorescence signals from nuclei loaded with 10µM Fluo 5N-AM. IP<sub>3</sub> or caffeine was applied by hydrostatic pressure ejection and signals expressed as ratios (F/F<sub>0</sub>) of fluorescence counts relative to baseline. Ca<sup>2+</sup> release in response to IP<sub>3</sub> (10µM) was signif-

icantly greater than that released in response to caffeine (10mM) (0.12 ± 0.02 v's 0.017 ± 0.002 [Ca<sup>2+</sup>]<sub>Nuc</sub> (F/F<sub>0</sub>) for IP<sub>3</sub> and caffeine respectively, n=6). When tetracaine (100µM) was applied to the nuclei, IP<sub>3</sub>-mediated Ca<sup>2+</sup> release was unaffected but the response to caffeine was abolished, suggesting RyR activation does not contribute to IP<sub>3</sub>-mediated nuclear Ca<sup>2+</sup> release. The potential for other nuclear proteins interacting with the nuclear IP<sub>3</sub>R was also investigated. Immunoblot analysis demonstrated expression of both FKBP12 and calcineurin in cardiac nuclei. These proteins are known to interact with the IP<sub>3</sub>R in other tissue types. Co-immunoprecipitation experiments using an anti-IP<sub>3</sub>R (type II) antibody suggest IP<sub>3</sub>R/calcineurin/FKBP12 interaction specifically at the nucleus. These results highlight the existence of a nuclear multi-protein IP<sub>3</sub>R complex, providing further scope for regulation of cardiac nuclear Ca<sup>2+</sup> release.

#### 501-Pos Board B380

##### Type 2 Inositol 1,4,5-trisphosphate Receptor Phosphorylation and Modulation by Ca<sup>2+</sup>/Calmodulin-dependent Protein Kinase II

Joshua T. Maxwell, Ademuyiwa S. Aromolaran, Gregory A. Mignery.

Loyola University Medical Center, Maywood, IL, USA.

InsP<sub>3</sub>-mediated intracellular Ca transients can activate Ca<sup>2+</sup>-calmodulin-dependent protein kinase II (CaMKII), a multifunctional Serine/Threonine protein kinase involved in many signaling pathways. Recent results show that InsP<sub>3</sub>Rs in the heart (InsP<sub>3</sub>R2) are primarily targeted to the nuclear envelope in ventricular cardiac myocytes. Here it forms a macromolecular complex with CaMKIIδ (Bare et al, 2005, *JBC*). Upon stimulation of InsP<sub>3</sub> production, Ca<sup>2+</sup> released through the InsP<sub>3</sub>R2 activates CaMKIIδ, allowing it to act on downstream targets, such as histone deacetylases 4 & 5 (HDAC4 & HDAC5) (Zhang et al, 2007, *JBC*). Additionally, CaMKII activity feedback modulates InsP<sub>3</sub>R2 function by direct phosphorylation and results in a decrease in the channel's open probability. The results of this study show that in planar lipid bilayers the channel activity of InsP<sub>3</sub>Rs can be inhibited by CaMKII-mediated phosphorylation, and that effect can be reversed by addition of protein phosphatases. Furthermore, the N-terminal 1078 amino acids of the rat InsP<sub>3</sub>R2 have been shown to interact with, as well as be phosphorylated by CaMKII in *in vitro* kinase assays. A smaller fragment spanning amino acids 1-708 of the InsP<sub>3</sub>R2 has been shown to be phosphorylated in a CaMKII-dependent manner. Conversely, C-terminal regions were not phosphorylated by CaMKII *in vitro*. We have also shown that the N-terminal region of the rat InsP<sub>3</sub>R1 spanning amino acids 1-1081 can be phosphorylated by CaMKII. Our results from mass spectrometry and *in vitro* kinase assays indicate that the putative CaMKII regulatory phosphorylation site lies within amino acids 150-340 of the InsP<sub>3</sub>R2. This work was supported by National Institutes of Health Grant HL-80101

#### 502-Pos Board B381

##### Regulation Of Inositol 1,4,5-Trisphosphate Receptor Isoforms By O-Linked Glycosylation

Patricia Bimboese<sup>1,2</sup>, Craig J. Gibson<sup>1</sup>, Stefan Schmidt<sup>1,3</sup>, Jere Paavola<sup>1,4</sup>, Barbara E. Ehrlich<sup>1</sup>.

<sup>1</sup>Yale University, New Haven, CT, USA, <sup>2</sup>Friedrich-Schiller University, Jena, Germany, <sup>3</sup>University of Goettingen, Goettingen, Germany,

<sup>4</sup>University of Helsinki, Helsinki, Finland.

The inositol 1,4,5 trisphosphate receptor (InsP<sub>3</sub>R), an intracellular calcium channel, is a family of three isoforms. All three isoforms display a significant level of sequence identity yet they differ in expression level, localization and many functional aspects. We previously showed that InsP<sub>3</sub>R type 1 is modified by O-linked β-N-acetylglucosamine glycosylation (O-GlcNAcylated). Through this dynamic and inducible modification a single monosaccharide is covalently attached to serine and threonine residues of the protein backbone, providing protein regulation similar to O-phosphorylation. We also reported that increased O-GlcNAcylation of the InsP<sub>3</sub>R type 1 reduced the percent of cells that responded to addition of extracellular agonists and those that did respond had a decreased InsP<sub>3</sub> dependent calcium release from the endoplasmic reticulum (ER). We now report that the InsP<sub>3</sub>R type 3 is also O-GlcNAcylated. Interestingly, the functional impact of O-GlcNAcylation on InsP<sub>3</sub>R type 3 channel is opposite to the effect measured with the InsP<sub>3</sub>R type 1. Human cholangiocytoma cells (MzChA-1) contain >90% InsP<sub>3</sub>R type 3. When these cells were incubated in hyperglycemic media there was an increase in the percent cells responding to InsP<sub>3</sub> generating stimuli and there was an increase in the InsP<sub>3</sub> dependent calcium release from the ER. A difference in functional response between InsP<sub>3</sub>R isoforms was reported previously for phosphorylation by cyclic AMP dependent protein kinase (PKA). In contrast, the InsP<sub>3</sub>R type 2 showed no detectable O-GlcNAc glycosylation and no significant functional changes even though the enzymes necessary for both the addition and removal of the monosaccharide are present in all cell types tested. The dynamic and inducible nature of O-GlcNAcylation and the isoform specificity suggests that this form of modification of the InsP<sub>3</sub>R and subsequent changes in intracellular