

increased from  $-0.42/-0.28$  to  $-0.74$  for a one- vs. four year observation period. When not including a common time points in the analysis, correlations of short- and long-term changes were weak to moderate. The application of the SDC approach showed a moderate consistency between the knees identified as progressors over short- and long-term periods. Longer observation periods may therefore be preferable to achieve robust results in individual knees.

### 368 FEASIBILITY OF BONE DENSITY EVALUATION USING PLAIN DIGITAL RADIOGRAPHY

M.B. Kinds<sup>1,2</sup>, A.C. Marijnissen<sup>1</sup>, K.L. Vincken<sup>2</sup>, L.W. Bartels<sup>2</sup>, M.A. Viergever<sup>2</sup>, H.W. de Jong<sup>3</sup>, F.P. Lafeber<sup>1</sup>. <sup>1</sup>Rheumatology & Clinical Immunology, Univ. Med. Ctr. Utrecht, Utrecht, Netherlands; <sup>2</sup>Image Sci. Inst., Univ. Med. Ctr. Utrecht, Utrecht, Netherlands; <sup>3</sup>Radiology, Univ. Med. Ctr. Utrecht, Utrecht, Netherlands

**Purpose:** For the evaluation of subchondral bone density (BD) changes due to OA, Dual Energy X-ray Absorptiometry (DEXA) is the most validated method. The need for DEXA might be reduced if the quantitative measurement of clinically relevant BD changes on radiographs is proved feasible, since radiographs are commonly acquired to evaluate structural changes due to OA. Precision of BD evaluation might be influenced by variations in acquisition settings that commonly occur in clinical practice and by post-processing (PP) that was introduced with the transition from conventional film-screen radiography to digital radiography. The objective of this study was to evaluate the effects of PP and acquisition settings on the precision and with that feasibility of BD measurement using plain digital radiography.

**Methods:** A bone density standard (BDS) was created consisting of eight cups with hydroxyapatite (HA: range 1.0-5.75 g/cm<sup>2</sup>). Digital radiographs of the BDS were taken (Philips Digital Diagnost), with variations in the acquisition and PP settings. Tube voltage (in kilovolt: kV), exposure (in milliamperere seconds: mAs), tube added filtering, and BDS position in the field-of-view were systematically varied and the default clinical PP was compared with minimal PP (at minimal strength). An aluminum step wedge served as an internal reference to express gray values of the BDS in mm aluminum equivalents (mmAl), by use of custom made software. In all cases a human (cadaver) knee joint was added to simulate clinical conditions. The relation (R<sup>2</sup>) between the BD values normalized to the reference wedge (in mmAl equivalents) and actual BD (HA in g/cm<sup>2</sup>), with variations in acquisition and PP settings was evaluated with linear regression analyses. Precision of BD measurement of the BDS was calculated in early OA (Cohort Hip & Cohort Knee: CHECK) to evaluate the relevance for clinical (research) practice.

**Results:** The BDS was validated by DEXA scanning and the relation between actual HA (g/cm<sup>2</sup>) and DEXA values was strongly linear: R<sup>2</sup>=0.99. In general for digital radiographs a strong correlation between actual BD and BD in mmAl was found for all settings. The correlation improved by changing PP from clinical (R<sup>2</sup>=0.96) to minimal (R<sup>2</sup>=0.98). Higher kV improved the correlation further. Even, for clinical PP mean SD was 0.97 mmAl, much smaller than the change of 2.51 mmAl clinically observed during two-year follow-up in early OA which implies the feasibility of BD measurements using plain digital radiography.

**Conclusions:** Accurate bone density measurement using digital radiography is feasible in a clinically relevant range, which removes the need for additional DEXA scans since plain radiographs are acquired for to evaluate structural changes due to osteoarthritis. Care should be taken in changing post-processing and acquisition settings, which can have profound effect on outcome.

Sponsor: Dutch Arthritis Association.

Acknowledgements: We thank the radiology department for their cooperation, and Paco Welsing for methodological support.

### 369 THE OCCURRENCE OF CAM IMPINGEMENT IN YOUNG MALE SOCCER PLAYERS

R. Agricola<sup>1</sup>, M. Heijboer<sup>1</sup>, A. Ginai<sup>1</sup>, R. van der Heijden<sup>1</sup>, J. Verhaar<sup>1</sup>, H. Weinans<sup>1,2</sup>, J. Waarsing<sup>1</sup>. <sup>1</sup>Erasmus Med. Ctr., Rotterdam, Netherlands; <sup>2</sup>Delft Univ. of Technology, Delft, Netherlands

**Purpose:** Femoroacetabular Impingement (FAI) is a cause of hip pain and might cause osteoarthritis (OA) of the hip due to abnormal contact between the femoral neck and acetabulum. Cam impingement is a

subtype of FAI, in which the abnormal contact is caused by a cam-type deformity in the femoral head-neck junction. Cam impingement is mostly seen in young active males. However, no studies have focused on the presence of cam-type deformities during skeletal development. This study aimed to determine the age of onset and prevalence of cam-type deformities in young male soccer players versus non-athletic controls.

**Methods:** 89 elite pre-professional soccer players and 90 controls aged 12-19 years were included in this study. In the soccer players group, both an anteroposterior (AP) and a Lauenstein radiograph of the hip were obtained according to a standardized protocol. Controls with both an AP and a Lauenstein radiograph with no signs of hip pathology were obtained from radiology databases. The alpha angle was calculated in all radiographs using semi-automatized software. An alpha angle larger than 60° was considered to define a cam-type deformity. All radiographs were scored by an orthopedic surgeon and a radiologist, using a three-point scoring system. The anterosuperior head-neck junction was classified as normal (1), flattened (2) or having a prominence (3). The soccer players completed a questionnaire, and range of motion (ROM) and impingement tests were performed. Differences in prevalence were tested using logistic regression, corrected for age. Differences between the mean alpha angle in soccer players and the control group, and differences in the ROM between cam-type deformity cases and normal soccer players were tested using Generalized Estimating Equations, corrected for age.

**Results:** The mean age was 14.8 years for the soccer players and 13.7 years for the controls. An alpha angle >60° was first found at the age of 12 in some soccer players and controls. A cam-type deformity defined by alpha angle was more prevalent in soccer players (26%) than in controls (18%), though not significantly when corrected for age (Figure 1).




Fig. 1. The prevalence of a cam-type deformity (alpha angle >60°) per age in soccer players and controls. In both soccer players and controls an alpha angle >60° was found starting at the age of 12 years. Total number (n) per age is given on top of each bar.

The mean alpha angle in the soccer players was larger than in controls in both the Lauenstein view (50.8 vs 46.8, p=0.002), and the AP view (50.9° vs 48.0° p=0.079). A prominence in the anterosuperior head-neck junction was first found at the age of 13 years and the prevalence was 13.5% in the soccer players. No prominences were found in the controls. A flattening of the head-neck junction was also more frequently found in the soccer players (53% vs 19%, p=0.0001) (Figure 2).




Fig. 2. Both a flattening and a prominence in the anterior head-neck junction were significantly more frequent in soccer players than in controls. Aspherical hips were especially found from the age of 14 years, starting with a flattening and with increasing age more prominences were found in the young soccer players.

Internal rotation was significantly reduced in soccer players with a cam-type deformity defined by alpha angle (19.7 vs 26.2, p=0.002), whereas a positive impingement test did not associate with the presence of a cam-type deformity.

**Conclusions:** A cam-type deformity can be present and recognizable from the age of 13 years. Cam-type deformities are more prevalent, and more pronounced in young soccer players than in their non-athletic