

Available online at www.sciencedirect.com

ScienceDirect

Procedia Food Science 7 (2016) 93 – 96

Procedia
Food Science

9th International Conference on Predictive Modelling in Food

Modeling *Penicillium expansum* growth response to thyme essential oil at selected water activities and pH values using surface response methodology

A. Rosas-Gallo, N. Ramírez-Corona, E. Palou, and A. López-Malo*

Departamento de Ingeniería Química, Alimentos y Ambiental, Universidad de las Américas Puebla, Ex hacienda Sta. Catarina Mártir, San Andrés Cholula, Puebla 72810, México

Abstract

Our objective was to evaluate, using a full factorial design, the effects of selected water activities (0.990, 0.945, or 0.900), pHs (5, 4, or 3), and thyme essential concentration (TEO, 0, 25, 50, or 100 ppm) oil on *Penicillium expansum* lag time (λ) and radial growth rate (μ_m) obtained by modeling mold response using Gompertz equation, and corresponding polynomial quadratic models. Potato-dextrose agar formulated with every studied factor combination was inoculated with 10^3 spores/ml, and incubated at 25°C up to 30 days. Mold colony diameter was periodically measured during incubation and adjusted with Gompertz equation to determine λ and μ_m . Decreasing a_w and pH, and increasing TEO concentration decreased μ_m and increased λ . At low a_w and pH, the increase in TEO concentration had a dramatic effect on *P. expansum* response since 25 ppm of TEO inhibited its growth for 30 days at 25°C. Gompertz parameters exhibited that *P. expansum* was sensitive to the evaluated combined factors, allowing us to construct a secondary predictive growth model. TEO in combination with a_w and pH reduction effectively inhibited *P. expansum* growth.

© 2016 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license

(<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Peer-review under responsibility of Department of Food Science, Faculty of Food Engineering, University of Campinas.

Keywords: *Penicillium expansum*; kinetic model; thyme essential oil; predictive models

1. Introduction

Antimicrobial agents are chemical additives or naturally-present compounds that delay or inactivate microbial growth in food products, and therefore slow down quality deterioration and help maintain food safety. Many researchers have concluded that the evolution of food additives should be based on a balance of risks and benefits,

* Corresponding author. Tel.: +52 (222) 2292126; fax: +52 (222) 2292729.

E-mail address: aurelio.lopezm@udlap.mx

so that in the future the most beneficial will be those that fulfill several functions, depending on the food to which they are added¹. Antimicrobials are currently produced synthetically, but these are also present naturally in various food components². Thus, they can be classified as antimicrobial compounds naturally present in the food or added intentionally. Among antimicrobials approved for direct use are organic acids and their derivatives, some esters, nitrites, nitrates, and several others³. Thyme is used as a flavoring and as a medicinal plant; the main components of its essential oil are thymol, anethole, and borneol (in its leaves), as well as carvacrol and cineole (in the whole plant). Thyme is produced in great amounts in some states of Mexico; so far, it is primarily used to enhance the flavor of dishes and for culinary preparations. However, few investigations have been made on the effect of its essential oil to inhibit fungal growth⁴. Today's consumers associate safe foods with fresh or least-processed and are demanding foods with characteristics similar to those of their fresh counterparts: without the addition of chemical additives, including those that serve as antimicrobial agents. This has led to a search for new forms of conservation, and to study the effect that different natural products in combination with some traditional forms may have on food preservation. The use of natural products to achieve an extended shelf-life while maintaining microbiological stability and safety is a current trend in alternative in the search for antimicrobial agents². Therefore, it is important to study the essential oil of thyme used as a natural antimicrobial agent to inhibit mold growth, and also evaluate the effect of changing pH and/or a_w on the antimicrobial activity of this essential oil. The antimicrobial activity of essential oils from several plants and spices has been recognized for many years. However, data on the effect of essential oils in combination with other factors on mold growth is still scarce. Additionally, there are few models to predict performance when natural preservatives are used in combination with other factors. Our objective was to evaluate, using a full factorial design, the effects of selected a_w (0.990, 0.945, or 0.900), pHs (5, 4, or 3), and concentration of TEO (0, 25, 50, or 100 ppm) on *Penicillium expansum* lag time (λ) and radial growth rate (μ_m) obtained by modeling mold response using Gompertz's equation, and corresponding polynomial quadratic models.

2 Materials and Methods

Thyme was acquired from a local market in San Francisco Tepeyecac (Puebla, Mexico). The essential oil was obtained by steam-distillation in a Clevenger-type apparatus. The oil was analyzed using a gas chromatograph (Agilent Technologies 6850N) coupled to a mass spectrometer detector (MSD, Agilent 5975C). A capillary column (Agilent, 5% phenyl methyl polysiloxane, 30 m length, 250 μ m diameter and 0.25 μ m thick) was used. 1 μ l of oil was injected using a 10:1 split at a temperature of 300°C. The ramp column temperature started at 60°C for 2 min, increasing to 250°C at 10°C/min and was maintained for 10 min. The carrier gas was helium at a constant flow rate of 1.1 mL/min. The mass spectrometer was operated at 70 eV and maintained at 200°C^{5, 6}. *P. expansum* was obtained from the Food Microbiology collection – Universidad de las Américas Puebla and maintained in PDA slants at 25°C for 10 days⁷. Spores were harvested washing the culture surface with 10 ml of sterile Tween (0.1%) aqueous solution and the spore suspension was used as inocula the same day^{5, 7}. Solid model systems were prepared using a PDA base; following a full factorial design including studied factors and levels. Agar was adjusted to different a_w values with sodium chloride, the amount needed was calculated by Chirife and Favetto's equation⁸:

$$a_w = 1 - K * m$$

where: K is a constant characteristic of the solute (0.03710 for NaCl) and m is solute molality in the aqueous phase. Model systems with the adjusted a_w were sterilized at 121°C for 15 min, cooled and the pH (5.0, 4.0 or 3.0) was adjusted with citric acid (10%) previously sterilized. The amounts of acid needed to achieve the desired pH were previously determined by titration curves. TEO (0, 25, 50, or 100 ppm) was thoroughly mixed with agar and poured into Petri dishes and allowed to solidify. Each of the prepared systems was inoculated (10^6 spores/plate). Three replicates of each combination were performed. Inoculated systems were kept at 25°C in airtight plastic chambers. The diameter of the mold colony was measured with a digital caliper every 24 h starting from the day when growth became visible. Mold growth response was adjusted to Gompertz's equation^{9, 10}:

$$\ln\left(\frac{D_t}{D_0}\right) = A \exp\left[-\exp\left[\frac{\mu_m e}{A}\right](\lambda - t) + 1\right]$$

where: D_t (cm) is the average colony diameter at time t (h), D_0 (mm) is the average initial diameter, A is the maximum growth, μ_m is the maximum growth rate (1/h), and λ is the lag phase (h). With the parameters obtained from Gompertz model, an analysis of the experimental surface-response design was carried out:

$$f(x) = \beta_0 + \beta_1 x_1 + \dots + \beta_i x_i + \dots + \beta_i^2 x_i^2 + \dots + \beta_i x_i x_j$$

obtaining the coefficients for the polynomial models using Minitab 16 (Minitab Inc., State College, PA,) software. Where: $f(x)$ is the parameter defined as the dependent variable (μ_m or λ), $x_{i,j}$ are the independent variables, and β_i are the polynomial model coefficients.

3. Results

The yield of the extraction of thyme essential oil was 1.02 ± 0.14 (ml/100g), identified compounds in the essential oil were p-cymene (53.29%), thymol (12.95%), carvacrol (10.25%), β -caryophyllene (8.65%), γ -terpinene (4.90%), and Bornyl acetate (4.24%) were the major compounds. Cosentino et al.¹¹ reported that these compounds in different thyme phenotypes make up 53.6-96.9% of the total composition of the essential oil used. p-cymene is in greater proportion than γ -terpinene as reported by Burt³, who similarly reported that TEO has a greater proportion of thymol than carvacrol which agrees with our analysis. Figure 1 presents *P. expansum* growth response at a_w of 0.99 in systems with pH 4 or 3 with and without 25 ppm of TEO. It can be observed that without essential oil, growth begins after 25-30 h. Figure 1 also shows that Gompertz's equation adequately fits the experimental data. Table 2 presents the constants of Gompertz's equation; in mediums formulated with a_w of 0.99 and pH 5, μ_m is higher compared to that found at pH 3, in the same way λ is longer at pH 3 compared to pH 5 and when concentration of TEO increases from 0 to 25 ppm. Similarly, reduction in a_w caused a dramatic effect on mold growth responses; the addition of 25 ppm of TEO was inhibitory for many of the tested combinations. The addition of 50 ppm allows mold growth only at pH 5. Thus, for concentrations of 100 ppm no growth was detected even after 2,160 h.




Fig. 1. *Penicillium expansum* growth response in model systems formulated with a_w of 0.99, pH 4 or 3, and 0 or 25 ppm of thyme essential oil. Lines represent the adjustment of Gompertz's equation.

Table 2 . Average Gompertz parameters that describe *Penicillium expansum* growth under studied conditions.

pH	Thyme essential oil concentration (ppm)	a_w 0.99		a_w 0.95		a_w 0.90	
		Growth rate (h^{-1})	Lag time (h)	Growth rate (h^{-1})	Lag time (h)	Growth rate (h^{-1})	Lag time (h)
3	0	0.0948	26.1	0.0072	124.0	0.0022	707.9
	25	0.0064	122.6	--	>2,160	--	>2,160
	50	--	>2,160	--	>2,160	--	>2,160
4	0	0.1347	27.3	0.0389	62.9	0.0042	507.5
	25	0.0085	62.6	--	>2,160	--	>2,160
	50	--	>2,160	--	>2,160	--	>2,160
5	0	0.1331	27.2	0.0414	21.7	0.0052	190.8
	25	0.1184	27.0	0.0063	46.0	--	>2,160
	50	0.0071	121.2	--	>2,160	--	>2,160

According to Kumar et al.¹², the mycelial growth of *Penicillium* spp. was inhibited with 0.7 mM solutions of TEO and exhibited a broad fungicidal spectrum in eight different foods. Analyzing Gompertz's growth parameters using surface-response methodology, polynomial coefficients that describe the effects of the studied variables were calculated.




Fig. 2. Surface plots of *Penicillium expansum* growth rate (GR, h⁻¹) in media formulated without thyme essential oil (Right), and lag time (h) in media with 50 ppm of thyme essential oil (Left).

The obtained polynomial models adequately described (R^2 , 0.86-0.95) the effects of a_w , pH, and essential oil concentration on mold growth rate and lag phase.

$$GR = 5.841 - 12.491 * a_w - 0.191 * pH + 0.0101 * TEO + 6.683 * a_w^2 + 0.211 * a_w * pH - 0.012 * a_w * TEO$$

$$\lambda = 89844.9 - 174449.0 * a_w - 2454.4 * pH + 101.5 * TEO + 84243.0 * a_w^2 - 14.9 * pH^2 + 0.1038 * TEO^2 + 2622.8 * a_w * pH - 91.9 * a_w * TEO - 2.82 * pH * TEO$$

Surface plots showing the effects of the valuated factors are presented in Figure 2. As it can be observed lowering a_w and/or pH has a dramatic effect on *P. expansum* growth rate. The addition of TEO increases λ as can be observed in Figure 2 and Table 1. Table 2 also presents the minimum inhibitory concentrations for *P. expansum* under the different a_w -pH combinations tested. As can be observed, lowering a_w and pH diminished the concentration of TEO needed to cause growth inhibition.

4. Conclusions

The essential oil of thyme could be utilized as an antifungal agent considering that relatively low quantities are required to inhibit studied mold for long periods of time (60 days). Because of its flavor and aromatic contributions it could be used in the production of sausages or other compatible products. We conclude that the minimum inhibitory concentration of thyme essential oil for *Penicillium expansum*, depends on a_w and pH of the medium, while combining a_w , pH, and concentration of essential oil are essential factors affecting fungal growth.

Acknowledgements

We acknowledge financial support from the National Council for Science and Technology of Mexico (CONACyT) and Universidad de las Americas Puebla (UDLAP). Author Rosas-Gallo acknowledges financial support for his Ph.D. studies in Food Science from CONACyT and UDLAP.

References

1. Suhr KI, Nielsen PV. Antifungal activity of essential oils evaluated by two different application techniques against rye bread spoilage fungi. *J Appl Microbiol* 2003; **94**:665-674.
2. Burt SA. Essential oils: Their antibacterial properties and potential applications in foods-a review. *Int J Food Microbiol* 2004; **94**:223-253.
3. Burt SA, Reinders RD. Antibacterial activity of selected plant essential oils against *Escherichia coli* O157:H7. *Lett Appl Microbiol* 2003; **36**:162-167.
4. Inouye S, Abe S, Yamaguchi H, Asakura M. Comparative study of antimicrobial and cytotoxic effects of selected essential oils by gaseous and solution contacts. *Int J Aromatherapy* 2003; **13**:34-42.
5. Gómez-Sánchez A, Palou E, López-Malo A. Antifungal activity evaluation of Mexican oregano (*Lippia berlandieri* Schauer) essential oil on the growth of *Aspergillus flavus* by gaseous contact. *J Food Protect* 2011; **74**:2192-2198.
6. Mejía-Garibay B, Palou E, López-Malo A. Composition, diffusion, and antifungal activity of black mustard (*Brassica nigra*). *J Food Protect* 2015; **78**:843-848.
7. López-Malo A, Alzamora SM, Palou E. *Aspergillus flavus* growth in the presence of chemical preservatives and naturally occurring antimicrobial compounds. *Int J Food Microbiol* 2005; **99**:119-128.
8. Favetto GJ, Chirife J. Simplified method for the prediction of water activity in binary solutions. *J Food Technol* 1985; **20**:631-636.
9. Char CD, Guerrero SN, Alzamora SM. Growth of *Eurotium chevalieri* in milk jam: influence of pH, potassium sorbate and water activity. *J Food Safety* 2007; **27**(1):1-16.
10. Zwietering MH, Jongenburger I, Rombouts FM, Van't Riet K. Modeling of the bacterial growth curve. *App & Environ Microbiol* 1990; **56**:1875-1881.
11. Cosentino S, Tuberoso CIG, Pisano B, Satta M, Mascia V, Arzede E, Palmas F. In-vitro antimicrobial activity and chemical composition of Sardinian *Thymus* essential oils. *Lett Appl Microbiol* 1999; **29**:130-135.
12. Kumar A, Shukla R, Singh P, Prasad CS, Dubey NK. Assessment of *Thymus vulgaris* L. essential oil as a safe botanical preservative against post harvest fungal infestation of food commodities. *Inn Food Sci & Emerging Technol* 2008; **9**:575-580.