

Mechanisms of Resistance to RAF Inhibitors in Melanoma

Andrew E. Aplin^{1,2,3}, Fred M. Kaplan¹ and Yongping Shao¹

The recent RAF inhibitor trial with PLX4032/RG7204 in late-stage mutant B-RAF melanoma patients has been lauded as a success story for personalized cancer therapy since short-term clinical responses were observed in the majority of patients. However, initial responses were followed by subsequent tumor regrowth, and a subset of patients showed intrinsic resistance. Bi-directional translational efforts are now essential to determine the mechanisms underlying acquired/secondary and intrinsic resistance to RAF inhibitors.

Journal of Investigative Dermatology (2011) **131**, 1817–1820;
doi:10.1038/jid.2011.147; published online 19 May 2011

INTRODUCTION

Metastatic dissemination of melanoma occurs to multiple sites, and is associated with 2–16% 10-year survival expectancy depending on the site of metastasis. Melanoma is a paradigm for chemoresistance and treatment options have remained limited for decades. Standard chemotherapy, such as the alkylating agent and dacarbazine, elicits a response rate of only 10% (Flaherty, 2010). Immunotherapy options, such as high-dose IL 2 and anti-CTLA4 (ipilimumab), have shown improved median survival benefits, but again response rates are low and the side effects of these treatments can be severe. Recently, targeted therapies have been designed to selectively kill melanoma cells harboring mutations in the serine-threonine kinase, B-RAF. Approximately, 50–60% of melanomas harbor B-RAF mutations (Davies *et al.*, 2002); the most frequent mutation is a valine to glutamic acid substitution at codon 600 (V600E). V600E, as well as V600K/D mutations, lead to constitutive B-RAF kinase activity and elevate downstream signaling through the MEK (mitogen-activated protein kinase/extracellular

signal-regulated kinase)–ERK1/2 (extracellular signal-regulated kinase 1/2) pathway. B-RAF^{V600E} is a driver mutation that promotes melanoma growth and survival in a variety of pre-clinical models and inhibiting B-RAF expression/activity results in growth inhibition and cell death (Dhomen and Marais, 2007). However, B-RAF^{V600E} mutations are also found in benign nevi and, hence, are not sufficient for malignancy.

PLX4032/RG7204 was recently developed as a potent adenosine triphosphate-competitive inhibitor of RAFs, with modest preference *in vitro* for mutant B-RAF and C-RAF compared with wild-type B-RAF (Bollag *et al.*, 2010). However, in cells it functions as a selective inhibitor of mutant B-RAF signaling (Bollag *et al.*, 2010), probably because of the higher adenosine triphosphate $K_{m(\text{app})}$ for B-RAF^{V600E} in mM cellular concentrations of adenosine triphosphate compared with wild-type forms of B-RAF and C-RAF (Hatzivassiliou *et al.*, 2010). In a phase 1 trial, 81% of melanoma patients harboring B-RAF^{V600E} showed objective tumor regression by RECIST criteria following PLX4032 treatment (Flaherty *et al.*, 2010). In addition, a second adenosine triphosphate-competitive RAF inhibitor, GSK2118436, is showing promising results in phase 1 trials with a 63% response rate observed in mutant B-RAF^{V600E/K/D} patients (Kefford *et al.*, 2010). However, in the PLX4032 trial, the clinical effects were temporary and the length of tumor-free survival averaged 7 months (Smalley and Sondak, 2010). Furthermore, 19% of patients in the phase 1 trial did not show tumor regression >30% (Flaherty *et al.*, 2010). Thus, acquired and intrinsic modes of resistance are hampering the clinical efficacy of PLX4032. It is critical to understand the mechanisms of resistance in order to optimize PLX4032 activity and improve the response rates, as well as the duration of clinical benefit. Emerging evidence from patient-matched pre-treatment and post-relapse samples (Table 1) highlights that multiple mechanisms underlie resistance to PLX4032 and likely other RAF inhibitors (Figure 1). These mechanisms can be divided into four non-mutually exclusive categories: re-activation of RAF–MEK signaling, alterations in ERK1/2-regulated cell cycle events, activation of alternative signaling pathways, and chromatin-regulating events.

RE-ACTIVATION OF THE RAF–MEK SIGNALING PATHWAY

The most direct route for a melanoma cell to by-pass RAF inhibitor action is by re-activation of the RAF–MEK–ERK1/2 pathway. Indeed, enhanced phosphoERK1/2 levels have been observed in mutant B-RAF melanoma cell lines that have acquired resistance to RAF inhibitors (PLX4032 and its non-clinical tool compound PLX4720, AZ628, and SB-590885)

¹Department of Cancer Biology, Thomas Jefferson University, Philadelphia, Pennsylvania, USA; ²Kimmel Cancer Center, Thomas Jefferson University, Philadelphia, Pennsylvania, USA and ³Department of Dermatology and Cutaneous Biology, Thomas Jefferson University, Philadelphia, Pennsylvania, USA

Correspondence: Andrew E. Aplin, Department of Cancer Biology, Kimmel Cancer Center, Thomas Jefferson University, 233 South 10th Street, Philadelphia, Pennsylvania 19107, USA.

E-mail: Andrew.Aplin@KimmelCancerCenter.Org

Abbreviations: ERK1/2, extracellular signal-regulated kinase 1/2; MAP3K, mitogen-activated protein kinase kinase kinase; MEK, mitogen-activated protein kinase/extracellular signal-regulated kinase; PDGFR β , platelet-derived growth factor receptor β

Received 31 January 2011; revised 22 March 2011; accepted 1 April 2011; published online 19 May 2011

Table 1. Evidence from patient-matched pre-treatment and post-relapse samples

Gene	Alteration(s) in patient samples	Reference
<i>Cot1</i>	Enhanced <i>Cot1</i> mRNA levels following PLX4032 treatment in two out of three patient samples analyzed	Johannessen <i>et al.</i> (2010)
<i>IGF-1R</i>	Enhanced IGF-1R staining in relapse samples in two out of five patients compared with pre-treatment samples	Villanueva <i>et al.</i> (2010)
<i>N-RAS</i>	Two out of 16 relapse samples harbored acquired N-RAS mutations. The two tumors were independent metastases from the same patient.	Nazarian <i>et al.</i> (2010)
<i>PDGFRβ</i>	Four out of 11 PLX4032-resistant tumor samples showed elevated PDGFRβ staining compared with patient-matched samples from the pre-treatment condition.	Nazarian <i>et al.</i> (2010)

Abbreviations: IGF-1R, IGF-1 receptor; PDGFRβ, platelet-derived growth factor receptor β.




Figure 1. Multiple mechanisms of resistance to RAF inhibitors in mutant B-RAF cells. Resistance to RAF inhibitor (i) blockade of signaling through the MEK-ERK1/2 pathway can occur via acquired mutation in N-RAS (Q61K or Q61R) or upregulation of RTK. These mechanisms enhance RAS activity, which promotes C-RAF dimerization and activation. MEK-ERK1/2 pathway activation can also occur through mutations in the B-RAF target, MEK1 (P124L), and via upregulation of the MAP3K and Cot1. Activation of the parallel PI-3 kinase-Akt pathway is promoted by loss of PTEN expression/activity, often through mutation and upregulation of RTKs including IGF-1R and possibly PDGFRβ. Re-activation of the ERK1/2 pathway and PI-3K-Akt signaling promote G1/S cell cycle events including cyclin D1 upregulation and downregulation of the cyclin-dependent inhibitor, p27^{Kip1}. In addition, these pathways promote survival events by promoting expression of the anti-apoptotic protein, Mcl-1, as well as down-modulating levels of the pro-apoptotic BH3-only proteins, Bim-EL, and Bmf. Alterations in the expression of these cell cycle and survival proteins may also promote resistance to RAF inhibitors. ERK, extracellular signal-regulated kinase; IGF-1R, IGF-1 receptor; MAP3K, mitogen-activated protein kinase kinase kinase; MEK, mitogen-activated protein kinase/extracellular signal-regulated kinase; PDGFRβ, platelet-derived growth factor receptor β; PI-3K, phosphatidylinositol 3 kinase; PTEN, phosphatase and tensin homolog; RTK, receptor tyrosine kinase.

through continued culture in the presence of drug (Montagut *et al.*, 2008; Paraiso *et al.*, 2010; Tap *et al.*, 2010; Villanueva *et al.*, 2010). A notable difference (at least to date) from resistance mechanisms to Abl kinase and EGFR inhibitors in

chronic myelogenous leukemia and non-small cell lung cancer, respectively, is the lack of secondary gatekeeper mutations in the drug target. Although mutation of the gatekeeper amino acid, threonine 529, renders B-RAF^{V600E} resistant to PLX4720 in cell-based assays (Whittaker *et al.*, 2010), deep-sequencing analysis indicates that threonine 529 or other sites in B-RAF are not mutated in resistant tumor samples and cell lines (Nazarian *et al.*, 2010; Tap *et al.*, 2010). Rather upregulation/amplification of other mitogen-activated protein kinase kinase kinases (MAP3Ks) seems to be a more prominent mechanism. C-RAF and A-RAF are paralogs of B-RAF. Settleman and colleagues showed that mutant B-RAF expression was associated with mutant C-RAF expression was associated with mutant B-RAF melanoma cell resistance to AZ628 (Montagut *et al.*, 2008). Depletion of C-RAF from melanoma cells with acquired resistance to AZ628 enhanced susceptibility to RAF inhibition. Conversely, C-RAF overexpression in parental cells enhanced resistance to AZ628 (Montagut *et al.*, 2008). AZ628-resistant cells showed enhanced susceptibility to geldanamycin, an agent that reduces RAF protein levels by targeting its chaperone protein, heat shock protein 90 (Montagut *et al.*, 2008). A modest increase in C-RAF expression was also observed in melanoma cells acquiring resistance to SB-590885 (Villanueva *et al.*, 2010). In these studies, the authors propose that RAF isoform switching mediates the maintenance of elevated ERK1/2 phosphorylation. This notion is based on the requirement for molecular depletion of A-RAF and C-RAF, as well as SB-590885 (which is more potent against B-RAF compared with A- or C-RAF) to completely inhibit ERK1/2 phosphorylation in SB-590885-acquired resistant cells (Villanueva *et al.*, 2010). However, it should be noted that maintenance of phosphoERK1/2 levels in these studies co-occurs with acquisition of alternative resistance pathways (see below).

Further support for a role for C-RAF in resistance to B-RAF inhibition was provided by Garraway and colleagues using a complementary DNA screen for kinases that prevent PLX4720-mediated cell growth arrest (Johannessen *et al.*, 2010). This screen also identified a non-RAF MAP3K, Cot1/Tpl2/MAP3K8. Cot1 activated MEK-ERK1/2 signaling in PLX4720-treated mutant B-RAF cell lines and cell lines endogenously expressing elevated levels of Cot1 were intrinsically resistant to PLX4720 (Johannessen *et al.*, 2010). Importantly, Cot1 mRNA levels were increased in

two out of three relapsed tumors samples analyzed from PLX4032-treated patients, indicating relevance to the *in vivo* situation (Johannessen *et al.*, 2010). B-RAF inhibition in cell lines led to increased Cot1 levels, raising the possibility that Cot1 upregulation is an adaptive response to PLX4032. However, RAF inhibitor-induced Cot1 upregulation in cells was independent of mRNA alterations, indicating a distinct mechanism to the Cot1 upregulation observed in patients.

The above studies highlight that compensatory upregulation of MAP3Ks, such as Cot1, is likely to represent one mechanism of resistance to RAF inhibitors in patients. However, mechanisms upstream of MAP3Ks also appear likely in a subset of resistant tumors. Upstream of RAFs are the RAS guanosine triphosphatases that recruit RAFs to the membrane for activation. N-RAS is mutated in ~15% of melanomas in a manner that is mutually exclusive from B-RAF mutation. The potential role of RAS in altering the effects of RAF inhibitors was highlighted by a series of papers showing that RAF inhibitors lead to a paradoxical hyperactivation of MEK-ERK1/2 signaling in cells harboring mutant N-RAS/high RAS activity (reviewed in Kaplan *et al.* (2010). Indeed, ectopic expression of mutant N-RAS in a mutant B-RAF colorectal cancer cell line nullifies the inhibitory effects of PLX4720 (Poulikakos *et al.*, 2010), a result that is reproducible in B-RAF^{V600E} melanoma cells (Johannessen *et al.*, 2010; Nazarian *et al.*, 2010). Knockdown of mutant N-RAS reverses the insensitivity of PLX4032-resistant cell lines and, importantly, N-RAS mutations were detected in two independent progressing tumors from a PLX4032-treated patient (Nazarian *et al.*, 2010). Elevated RAS activity in the absence of RAS mutations was detected in cell lines with acquired resistance to PLX4032 indicating that events that promote elevated RAS signaling will likely elicit similar effects. Thus, the co-occurrence of B-RAF mutations and RAS mutation/activation is likely to promote resistance to PLX4032 and other RAF inhibitors.

Another possible mechanism to by-pass the activity of RAF inhibitors is through alterations in B-RAF effectors. MEK1 and 2 are phosphorylated and activated by B-RAF and a P124L mutation in MEK1 was identified through a MEK1 random mutagenesis screen that confers cellular resistance to the MEK inhibitor, AZD6244 (Emery *et al.*, 2009). MEK1^{P124L} expression conferred resistance to PLX4720 in cell-based assays and was identified in a metastasis from a patient with acquired resistance to AZD6244, indicating the clinical relevance of this mutation. Interestingly, the combination of PLX4720 plus AZD6244 overcame the resistance conferred by MEK1^{P124L} expression. These and other data have prompted the use of MEK inhibitors in clinical trials enrolling mutant B-RAF melanoma patients who were previously treated with or without a BRAF inhibitor.

ALTERATIONS IN ERK1/2-REGULATED CELL CYCLE EVENTS

Mutant B-RAF signaling via ERK1/2 promotes G1/S cell cycle progression, at least in part, through control of cyclin D1 levels, which enhance cyclin-dependent kinase activity to promote hyperphosphorylation of retinoblastoma (Bhatt

et al., 2005). Studies using retinoblastoma-null fibroblasts and tumor lines have demonstrated the requirement of an intact cyclin-cyclin-dependent kinase-retinoblastoma axis for MEK inhibitor prevention of cell cycle entry (D'Abaco *et al.*, 2002). Approximately, 25% of mutant B-RAF melanoma cell lines and tumor samples harbor amplifications in cyclin D1 (Smalley *et al.*, 2008). Furthermore, overexpression of cyclin-dependent kinase 4 and cyclin D1 in a SB-590885-sensitive cell line promotes resistance to this RAF inhibitor (Smalley *et al.*, 2008). An implication of this work is that mutant B-RAF patients with amplification of the cyclin D1 locus or other alteration on the cyclin-cyclin-dependent kinase-retinoblastoma axis will likely be intrinsically resistant to PLX4032 and, thus, patients stratified on this basis may be predictive of the response.

ACTIVATION OF ALTERNATIVE SIGNALING PATHWAYS

Multiple laboratories have generated mutant B-RAF cell lines that have acquired resistance to RAF inhibitors but not all of these lines show re-activation of the ERK1/2 pathway (Montagut *et al.*, 2008). These findings indicate that input from alternative, ERK1/2-independent pathways induce RAF inhibitor resistance mechanisms. In the Garraway screen, protein kinase C- ϵ , protein kinase C- η , and ErbB2 individually provided resistance to PLX4720 in the absence of ERK1/2 re-activation (Johannessen *et al.*, 2010). Studies from Nazarian *et al.* (2010) implicate the platelet-derived growth factor receptor β (PDGFR β) receptor tyrosine kinase in the resistance to PLX4032. PDGFR β is upregulated in PLX4032-resistant patient biopsies. Furthermore, resistant cell lines with upregulated PDGFR β were dependent on PDGFR β for proliferation and survival despite PLX4032 inhibition of the ERK pathway remained intact. Consistent with an ERK1/2-independent resistance mechanism, the growth of PDGFR β -overexpressing resistant cells was insensitive to MEK inhibitors. However, the mechanism underlying PDGFR β -mediated resistance to PLX4032 is currently unclear.

Other studies implicate the phosphatidylinositol 3 kinase-Akt signaling pathway in resistance to RAF inhibitors. Elevated Akt activity was detected in two out of three cell lines with acquired resistance to PLX4032 (Tap *et al.*, 2010). In addition, enhanced Akt phosphorylation is acquired and associated with increased expression of IGF-1 receptor or depletion of phosphatase and tensin homolog in relapsed tumor samples from PLX4032-treated patients (Villanueva *et al.*, 2010). Consistent with activation of the Akt pathway providing resistance to RAF inhibitors, expression of constitutively active Akt3 inhibited PLX4720-induced apoptosis in three-dimension by preventing upregulation of the pro-apoptotic BH3-only protein, Bim-EL, and Bmf, following B-RAF inhibition (Shao and Aplin, 2010). Notably, MEK inhibitors in combination with either an IGF-1 receptor inhibitor or a phosphatidylinositol-3 kinase inhibitor induces cell death in RAF inhibitor acquired resistant cell populations (Villanueva *et al.*, 2010). Importantly, these data have provided new strategies to overcome resistance to RAF inhibitors and the combination of MEK and Akt inhibitors is being pursued in clinical trials.

CHROMATIN-REGULATING EVENTS

Acquired mutations are rare events and time is required for the resistant cells to outgrow from the tumor mass. To account for the ability of cells to withstand drug toxicity in the short term, it has been proposed that reversible “drug-tolerant” states exist before the acquisition of permanent resistance (Sharma *et al.*, 2010). Tolerance is drug-induced suggesting an adaptive response to targeted therapies. In addition, tolerance is reversible and a fraction of tolerant cells revert to a sensitive state during a “drug holiday”. Thus, a second round of drug treatment will promote tumor cell death and the cycle will repeat until secondary mutations lead to permanent resistance. When challenging tumor cell lines with various anticancer agents, Settleman and colleagues detected sub-populations of tumor cell lines that show higher (>100-fold) drug tolerance than the remainder of the cells (Sharma *et al.*, 2010). This reversible drug-tolerant state is transiently acquired through elevated IGF-1 receptor signaling and chromatin remodeling mediated by enhanced expression of JARID1A, a histone demethylase (Sharma *et al.*, 2010). JARID1A associates with histone deacetylases and treatment of M14 melanoma cells with inhibitors of either IGF-1 receptor or histone deacetylases in combination with the RAF inhibitor, AZ628, prevents the emergence of AZ628-tolerant cells. Melanoma cells are known for their plasticity, and these studies indicate that an adaptive chromatin regulation response to targeted therapies may contribute ultimately to the acquisition of a resistant state.

CONCLUDING REMARKS

There is a clear need to build on the initial success of the PLX4032 trial in melanoma. As an increasing number of patient-matched pre-treatment, during treatment and post-treatment samples become available, state-of-the-art genomic and proteomic approaches need to be used to reveal the prominence of the mechanisms described above and reveal to our knowledge, previously unreported ways to promote resistance to RAF inhibitors. The findings are translational and will drive the specification of next generation RAF inhibitors and combinational strategies to provide lasting treatment to melanoma sufferers.

CONFLICT OF INTEREST

The authors state no conflict of interest.

ACKNOWLEDGMENTS

Work in the Aplin lab is supported by grants from the National Institutes of Health (GM067893, CA125103), the American Cancer Society (RSG-08-03-01-CSM), and the Pennsylvania Department of Health (AF0301).

NOTE ADDED IN PROOF: Further emphasizing re-activation of the ERK1/2 pathway as a mechanism of resistance to PLX4032, Garraway and colleagues recently identified an acquired MEK1 mutation (C121S) in one patient who relapsed on a PLX4032 trial (Wagle *et al.*, 2011). Expression of MEK1-C121S in mutant B-RAF melanoma cells conferred resistance to both RAF and allosteric MEK inhibitors, suggesting that molecular intervention downstream of MEK activation may be required in some patients with acquired resistance to RAF inhibitors.

REFERENCES

- Bhatt KV, Spofford LS, Aram G *et al.* (2005) Adhesion control of cyclin D1 and p27Kip1 levels is deregulated in melanoma cells through BRAF-MEK-ERK signaling. *Oncogene* 12:3459–71
- Bollag G, Hirth P, Tsai J *et al.* (2010) Clinical efficacy of a RAF inhibitor needs broad target blockade in BRAF-mutant melanoma. *Nature* 467:596–9
- D’Abaco GM, Hooper S, Paterson H *et al.* (2002) Loss of Rb overrides the requirement for ERK activity for cell proliferation. *J Cell Sci* 115:4607–16
- Davies H, Bignell GR, Cox C *et al.* (2002) Mutations of the BRAF gene in human cancer. *Nature* 417:949–54
- Dhomen N, Marais R (2007) New insight into BRAF mutations in cancer. *Curr Opin Genet Dev* 17:31–9
- Emery CM, Vijayendran KG, Zipsper MC *et al.* (2009) MEK1 mutations confer resistance to MEK and B-RAF inhibition. *Proc Natl Acad Sci* 106:20411–6
- Flaherty KT (2010) Narrative review: BRAF opens the door for therapeutic advances in melanoma. *Ann Intern Med* 153:587–91
- Flaherty KT, Puzanov I, Kim KB *et al.* (2010) Inhibition of mutated, activated BRAF in metastatic melanoma. *New Eng J Med* 363:809–19
- Hatzivassiliou G, Song K, Yen I *et al.* (2010) RAF inhibitors prime wild-type RAF to activate the MAPK pathway and enhance growth. *Nature* 464:431–5
- Johannessen CM, Boehm JS, Kim SY *et al.* (2010) COT drives resistance to RAF inhibition through MAP kinase pathway reactivation. *Nature* 468:968–72
- Kaplan FM, Mastrangelo MJ, Aplin AE (2010) The wrath of RAFs: rogue behavior of B-RAF kinase inhibitors. *J Invest Dermatol* 130:2669–71
- Kefford R, Arkenau H, Brown M *et al.* (2010) Phase I/II study of GSK2118436, a selective inhibitor of oncogenic mutant BRAF kinase, in patients with metastatic melanoma and other solid tumors. *J Clin Oncol* 28:15s (abstract 8503)
- Montagut C, Sharma SV, Shioda T *et al.* (2008) Elevated CRAF as a potential mechanism of acquired resistance to BRAF inhibition in melanoma. *Cancer Res* 68:4853–61
- Nazarian R, Shi H, Wang Q *et al.* (2010) Melanomas acquire resistance to B-RAF (V600E) inhibition by RTK or N-RAS upregulation. *Nature* 468:973–7
- Paraiso KHT, Fedorenko IV, Cantini LP *et al.* (2010) Recovery of phospho-ERK activity allows melanoma cells to escape from BRAF inhibitor therapy. *Br J Cancer* 102:1724–30
- Poulikakos PI, Zhang C, Bollag G *et al.* (2010) RAF inhibitors transactivate RAF dimers and ERK signalling in cells with wild-type BRAF. *Nature* 464:427–30
- Shao Y, Aplin A (2010) Akt3-mediated resistance to apoptosis in B-RAF targeted melanoma cells. *Cancer Res* 70:6670–81
- Sharma S, Lee D, Li B *et al.* (2010) A chromatin-mediated reversible drug-tolerant state in cancer cell subpopulations. *Cell* 141:69–80
- Smalley KS, Lioni M, Dalla Palma M *et al.* (2008) Increased cyclin D1 expression can mediate BRAF inhibitor resistance in BRAF V600E-mutated melanomas. *Mol Cancer Ther* 7:2876–83
- Smalley KSM, Sondak VK (2010) Melanoma - an unlikely poster child for personalized cancer therapy. *New Eng J Med* 363:876–8
- Tap W, Gong K, Dering J *et al.* (2010) Pharmacodynamic characterization of the efficacy signals due to selective BRAF inhibition with PLX4032 in malignant melanoma. *Neoplasia* 12:637–49
- Villanueva J, Vultur A, Lee JT *et al.* (2010) Acquired resistance to BRAF inhibitors mediated by a RAF kinase switch in melanoma can be overcome by cotargeting MEK and IGF-1R/PI3K. *Cancer Cell* 18:683–95
- Wagle N, Emery C, Berger MF *et al.* (2011) Dissecting therapeutic resistance to RAF inhibition in melanoma by tumor genomic profiling. *J Clin Oncol*; e-pub ahead of print 7 March 2011; doi:10.1200/JCO.2010.33.2312
- Whittaker S, Kirk R, Hayward R *et al.* (2010) Gatekeeper mutations mediate resistance to BRAF-targeted therapies. *Sci Trans Med* 2:35ra41