



previously [1,12]. Sixteen hours after peritoneal injection of sterilized 2% sodium caseinate in saline, peritoneal exudates were collected in a plastic tube and centrifuged at  $120 \times g$  for 5 min at  $4^\circ\text{C}$ . The cell pellet was treated for 30 s with 0.2% NaCl solution to lyse red cells, and then promptly mixed with an equal volume of 1.6% NaCl solution to restore the isotonicity. The cell suspension was re-centrifuged at  $120 \times g$  for 5 min at  $4^\circ\text{C}$ , and the resulting pellet of cells, more than 98% of which were PMN, was suspended in  $\text{Ca}^{2+}$ -free KRP (pH 7.4) and kept in an ice-bath until use.

#### 2.4. Assay of $\text{H}_2\text{O}_2$ generation

The rate of  $\text{H}_2\text{O}_2$  release from cells was measured by recording the rate of formation of an HRP- $\text{H}_2\text{O}_2$  complex, as reported previously [20]. A plastic cuvette of 10 mm light path containing 1.6 ml of cell suspension in the medium described below was attached to a windmill cell mixer and maintained at  $37^\circ\text{C}$  by means of a jacketed holder connected to a constant temperature water bath. The assay medium contained  $6 \mu\text{M}$  HRP,  $3 \times 10^6$  cells/ml, and 5 mM glucose in  $\text{Ca}^{2+}$ -free KRP. First  $\text{H}_2\text{O}_2$  release in the resting state was recorded and then formation of the HRP- $\text{H}_2\text{O}_2$  complex was followed at 417-403 nm in a double-beam spectrophotometer (Hitach model 556) after addition of various concentrations of CSE or related compounds as concentrated solutions in distilled water (10-15 mg/ml).

#### 2.5. Assay of $\text{O}_2^-$ -generation

The rate of  $\text{O}_2^-$  release from PMN was measured in the cuvette described above. The assay medium contained  $15 \mu\text{M}$  63%-acetylated cytochrome *c*, 5  $\mu\text{g}/\text{ml}$  catalase,  $3 \times 10^6$  PMN/ml, and 5 mM glucose in 1.6 ml of KRP. The reduction of acetylated cytochrome *c* was measured at  $37^\circ\text{C}$  at 550-540 nm after addition of various concentrations of sulfatide or related compound in a double beam spectrophotometer as described previously [19].

### 3. RESULTS

When CSE was added to the incubation medium, ferricytochrome *c* was precipitated, probably due to ionic interaction of CSE with cytochrome *c*. However, CSE did not give a precipitate with acetylated ferricytochrome *c*, and so this modified




Fig.1. Trace (left) of  $\text{O}_2^-$  formation followed as reduction of acetylated ferricytochrome *c* on addition of CSE to PMN. The reaction mixture contained  $15 \mu\text{M}$  63%-acetylated ferricytochrome *c*, 5  $\mu\text{g}/\text{ml}$  catalase,  $3 \times 10^6$  cells/ml, and 5 mM glucose in 1.6 ml of  $\text{Ca}^{2+}$ -free KRP buffer (pH 7.4). Reduction of cytochrome *c* was measured at  $37^\circ\text{C}$  at 550-540 nm. After recording the trace in the resting state, CSE was added at 0.1 mg/ml to the cell suspension. Trace (right) of  $\text{O}_2^-$  formation by phagocytosing PMN on the addition of heat-killed *E. coli* (about 100 bacteria/cell).




Fig.2. Trace (left) of  $\text{H}_2\text{O}_2$  formation by PMN by followed as formation of the HRP- $\text{H}_2\text{O}_2$  complex on addition of CSE (0.2 mg/ml). The reaction mixture contained  $6 \mu\text{M}$  HRP,  $3 \times 10^6$  cells/ml, and 5 mM glucose in 1.6 ml of  $\text{Ca}^{2+}$ -free KRP buffer. Trace (right) of  $\text{H}_2\text{O}_2$  formation by phagocytosing PMN. The reaction medium was as described above. *Escherichia coli* were added to cells in the same ratio of bacteria to cell as for fig.1. Formation of the HRP- $\text{H}_2\text{O}_2$  complex was measured at  $37^\circ\text{C}$  at 417-403 nm.

cytochrome *c* was employed for assay of  $O_2^-$  formation by PMN treated with CSE. The spectrophotometric trace in fig.1 shows the formation of  $O_2^-$  by PMN upon addition of CSE solution at a final concentration of 0.1 mg/ml. Marked stimulation of  $O_2^-$  formation was observed in the cells soon after the addition of CSE. In contrast, there was a short lag time before  $O_2^-$  began to be released when the cells were exposed to bacteria (heat-killed *E. coli*; about 100 bacteria/cell). Figure 2 shows the formation of  $H_2O_2$  from PMN on

addition of CSE at 0.2 mg/ml. No precipitate was formed after formation of the HRP- $H_2O_2$  complex until the concentration of added CSE was increased to 1.0 mg/ml.  $H_2O_2$  formation was more rapid in CSE-treated cells than in phagocytizing cells, and only the latter showed a lag time of about 20 s after started addition of *E. coli* before  $H_2O_2$  generation. Figure 3 shows a plot of the rate  $H_2O_2$  formation by PMN against the CSE concentration added to the cell suspension. The rate of  $H_2O_2$  formation increased with increase in CSE

Table 1

$H_2O_2$  generation by PMN upon addition of CSE and its related compounds

CSE and related compounds	Structure	Rate of $H_2O_2$ generation (nmol $H_2O_2 \cdot \text{min}^{-1} \cdot 10^7 \text{ cells}^{-1}$ )
CSE (galactocerebroside-3'-sulfate)		7.52
Galactocerebroside-6'-sulfate		0.56
Galactose-6-sulfate		neg
Seminolipid		neg
Cholesterol sulfate		14.1

The rate of  $H_2O_2$  formation by PMN was measured as for fig.2. CSE and related compounds were added at 0.1 mg/ml to suspension of PMN.




Fig.3. Plot of the rate of H<sub>2</sub>O<sub>2</sub> generation by PMN against the CSE concentration added to the cell suspension. The reaction mixture was the same condition as for fig.2, except that it contained various concentrations of CSE.

concentration to a plateau at 0.68 mg/ml (1.4 mg/10<sup>7</sup> cells).

Next, the stimulatory effects of CSE and related compounds on the O<sub>2</sub><sup>-</sup> and H<sub>2</sub>O<sub>2</sub> generating system of PMN were tested. Table 1 summarizes results on H<sub>2</sub>O<sub>2</sub> formation by PMN upon addition of CSE and related sulfated compounds at a concentration of 0.125 mg/ml (0.55 mg/10<sup>7</sup> cells). Galactocerebroside-6'-sulfate has a stimulatory effect on PMN, causing formation of 0.56 nmol H<sub>2</sub>O<sub>2</sub> · min<sup>-1</sup> · 10<sup>7</sup> cells<sup>-1</sup>, although this is far less than the formation of 7.52 nmol H<sub>2</sub>O<sub>2</sub> · min<sup>-1</sup> · 10<sup>7</sup> cells<sup>-1</sup> induced by CSE. This indicates the necessity of some particular sulfated carbohydrate structure for the stimulation. Galactose-6-sulfate had no ability to induce H<sub>2</sub>O<sub>2</sub> generation, indicating that the hydrophobic moiety of CSE may be necessary for stimulation of PMN, because of its attraction to the PMN cell membrane. However, another sulfated glycolipid, seminolipid, which has a hydrophobic chain of the glyceride type, had no effect on PMN when added at a similar concentration to that of galactocerebroside-6'-sulfate. Cholesterol sulfate induced marked stimulation of H<sub>2</sub>O<sub>2</sub> formation.

#### 4. DISCUSSION

It is widely accepted that the hydroxyl radical (·OH) splits carbon-carbon double bonds of lipids by a chain scission reaction because of its higher redox potential than those of other oxygen intermediates such as O<sub>2</sub><sup>-</sup> and H<sub>2</sub>O<sub>2</sub> [14,15]. ·OH is thought to be formed by a Haber-Weiss reaction of O<sub>2</sub><sup>-</sup> with H<sub>2</sub>O<sub>2</sub> [5,21]. Superoxide (O<sub>2</sub><sup>-</sup>) is dismuted to form H<sub>2</sub>O<sub>2</sub> at high velocity ( $k = 10^5 \text{ M}^{-1} \cdot \text{s}^{-1}$ ) even in the absence of superoxide dismutase. O<sub>2</sub><sup>-</sup> reacts with many substances acting as either an oxidizing or reducing agent [14]. H<sub>2</sub>O<sub>2</sub> is a reactive oxidizing agent and is converted to an even stronger oxidizing agent (HOCl) in the myeloperoxidase-H<sub>2</sub>O<sub>2</sub>-Cl system [22,23]. When phagocytosing cells, including PMN, monocytes [2] and macrophages [3,6], come in contact with infectious bacteria, their oxidative metabolism is promptly activated with generation of these toxic oxygen intermediates. This is an important host-defense mechanism. It is known that some hydrophobic agents induce marked enhancement of oxidative metabolism leading to formation of greater amounts of the oxygen intermediates than those formed on stimulation by phagocytosis [12,19,24].

As shown in the present experiments, brain sulfatide and cholesterol sulfate greatly induce oxidative metabolism of leucocytes, resulting in generation of the oxygen radicals that may attack the membrane. It is particularly interesting that galactose-6-sulfate, an analogue of the sugar moiety of sulfatide, was not stimulatory and also that seminolipid, a sulfatide with the glyceride but not the ceramide structure of CSE was not stimulatory. These findings suggest that the interaction between sulfatide and the leucocyte cell membrane necessary for induction of H<sub>2</sub>O<sub>2</sub> formation requires some particular forms of not only hydrophobic but also anionic hydrophilic structures.

Tissues of the nervous system, and particularly the myelin membrane, are rich in sulfatide. In various types of demyelinating diseases, such as multiple sclerosis, post-vaccinal encephalomyelitis, and experimental autoallergic central or peripheral demyelination, a number of phagocytes, including polymorphonuclear leucocytes, monocytes and macrophages, appear in demyelinating lesions [25,26]. These phagocytes appearing in the lesions may be in an active state, in all probability like

those reported in cases of many inflammatory diseases. These activated cells possibly release  $O_2^-$  and other membrane-toxic oxygen intermediates, which in turn may release sulfatide molecules from the membranes in a state that causes a second activation of the phagocytes leading to a further release of oxygen intermediates. Thus, a myelin-destructive chain reaction may be evoked by the interaction between sulfatide and phagocytes. Further investigations with myelin itself or modified or artificial sulfatide-containing membranes are needed to clarify this problem.

## REFERENCES

- [1] Sbarra, A.J. and Karnovsky, M.L. (1959) *J. Biol. Chem.* 234, 1355–1362.
- [2] Cline, M.J. and Lehrer, R.I. (1968) *Blood* 32, 423–435.
- [3] Rossi, F., Romeo, D. and Patriarca, P. (1972) *J. Reticuloendothel. Soc.* 12, 127–149.
- [4] Babior, B.M., Kipnes, R.S., Curnutte, J.T. (1973) *J. Clin. Invest.* 52, 741–744.
- [5] Tauber, A.I. and Babior, B.M. (1977) *J. Clin. Invest.* 60, 374–379.
- [6] Kaneda, M., Kakinuma, K., Yamaguchi, T. and Shimada, K. (1980) *J. Biochem.* 88, 1159–1165.
- [7] Kakinuma, K., Boveris, A., and Chance, B. (1977) *FEBS Lett.* 74, 295–299.
- [8] Dewald, B., Baggiolini, M., Curnutte, J.T. and Babior, B.M. (1979) *J. Clin. Invest.* 63, 21–29.
- [9] Yamaguchi, T., Sato, K., Shimada, K. and Kakinuma, K. (1982) *J. Biochem.* 91, 31–40.
- [10] Graham, R.C., Karnovsky, M.J., Shafer, A.W., Glass, E.A. and Karnovsky, M.L. (1967) *J. Cell Biol.* 32, 629–647.
- [11] Rossi, F. and Zatti, M. (1968) *Biochim. Biophys. Acta* 153, 296–299.
- [12] Kakinuma, K. (1974) *Biochim. Biophys. Acta* 348, 76–85.
- [13] Repine, J.E., White, J.G., Clauson, C.C. and Holmes, B.M. (1974) *J. Clin. Invest.* 54, 83–90.
- [14] Ciba Found. Symp. 65 'Oxygen Free Radicals and Tissue Damage' (1979) *Excerpta Medica*, pp. 1–370, Amsterdam, New York.
- [15] Rodgers, M.A.J. and Powers, E.L. eds. (1981) in: *Oxygen and Oxy-Radicals in Chemistry and Biology*, pp. 1–793, Academic Press, New York.
- [16] Nagai, Y. and Iwamori, M. (1980) in: *Structure and Function of Gangliosides* (Svennerholm, L., Dreyfus, H. and Urban, P.-F. eds) pp. 13–21, Plenum, New York and London.
- [17] Taketomi, T. and Yamakawa, T. (1964) *J. Biochem.* 55, 87–89.
- [18] Yoshizawa, T. and Nagai, Y. (1974) *Japan. J. Exp. Med.* 44, 465–471.
- [19] Kakinuma, K. and Minakami, S. (1978) *Biochim. Biophys. Acta*, 538, 50–59.
- [20] Kakinuma, K., Yamaguchi, T., Kaneda, M., Shimada, K., Tomita, Y. and Chance, B. (1979) *J. Biochem.* 86, 87–95.
- [21] Haber, F. and Weiss, J. (1934) *Proc. R. Soc. London, Ser. A* 147, 332–351.
- [22] Agner, K. (1941) *Acta Physiol. Scand. Suppl.* VIII, 2, 3–62.
- [23] Klebanoff, S.J. and Clark, R.A. (1978) *The Neutrophil: Function and Clinical Disorders*, pp. 409–443, Elsevier, Amsterdam and New York.
- [24] Kakinuma, K., Hatae, T. and Minakami, S. (1976) *J. Biochem.* 79, 795–802.
- [25] Paterson, P.Y. (1976) in: *Textbook of Immunopathology* (Miescher, P.E. and Muller-Eberhard, H.J. eds) 2nd ed., Vol. I, pp. 179–213, Grune and Stratton, New York.
- [26] Arnason, B.G.W. (1975) in: *Peripheral Neuropathy* (Dyck, D.J., Thomas, P.K. and Lambert, E.H. eds) pp. 1110–1148, W.B. Saunders Co., Philadelphia.