

Meta Analysis

# High uric acid level predicts left atrial thrombus or spontaneous echo contrast detected by transesophageal echocardiography: Meta-analysis and systematic review

En-Yuan Zhang<sup>a,b,d</sup>, Lu Kou<sup>b,d</sup>, Min Li<sup>a</sup>, Chee Yuan Ng<sup>c</sup>, Jian-Ping Zhao<sup>a</sup>,  
Guang-Ping Li<sup>a</sup>, Tong Liu<sup>a,\*</sup>

<sup>a</sup> Tianjin Key Laboratory of Ionic-Molecular Function of Cardiovascular Disease, Department of Cardiology, Tianjin Institute of Cardiology, Second Hospital of Tianjin Medical University, Tianjin 300211, China

<sup>b</sup> Tianjin Chest Hospital, Tianjin 300051, China

<sup>c</sup> Cardiac Arrhythmia Service, Massachusetts General Hospital, 55 Fruit St., Boston, MA 02114, USA

Received 23 August 2015

Available online 6 April 2016

## Abstract

**Background:** Recent observational studies have suggested that the patients with hyperuricemia have a higher risk of having left atrial thrombus (LATH) or left atrial spontaneous echo contrast (LASEC) by transesophageal echocardiography (TEE), while the ultimate predictive value of a high uric acid (UA) level on LATH/LASEC remained obscure.

**Methods:** We searched the PubMed and Cochrane clinical trials databases up to July 2015. Following screening the 369 initially identified studies, we analyzed six observational studies with 2381 patients.

**Results:** The meta-analysis of these studies showed that an elevated serum UA level was associated with a higher likelihood of LATH/LASEC ( $OR = 1.59$ ,  $95\%CI 1.13-2.23$ ,  $P = 0.008$ ), while significant differences exist among individual trials ( $P < 0.00001$  and  $I^2 = 85\%$ ). Sensitivity analysis failed to find any heterogeneity.

**Conclusion:** An elevated UA level was associated with a higher risk of detecting a left atrial abnormality represented by LATH/LASEC.

© 2016 Chinese Medical Association. Production and hosting by Elsevier B.V. on behalf of KeAi Communications Co., Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

**Keywords:** Uric acid; Left atrium; Thrombus; Transesophageal echocardiography; Atrial fibrillation

\* Corresponding author. Tel.: +86 22 88328368; fax: +86 22 28261158.

E-mail address: [liutongdoc@126.com](mailto:liutongdoc@126.com) (T. Liu).

<sup>d</sup> The two authors contribute equally to the work.

Peer review under responsibility of Chinese Medical Association.



Production and Hosting by Elsevier on behalf of KeAi

## Introduction

An atrial thrombus forming in the left atrial appendage (LAA) in the setting of atrial fibrillation results in a series of adverse outcomes including stroke, myocardial infarction and other severe embolism events. Transesophageal echocardiography (TEE)

<http://dx.doi.org/10.1016/j.cdtm.2016.02.002>

2095-882X/© 2016 Chinese Medical Association. Production and hosting by Elsevier B.V. on behalf of KeAi Communications Co., Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

is frequently used to look for an LAA thrombus and can be helpful when conducting risk stratification of patients for stroke.<sup>1</sup>

Hyperuricemia has been shown to be an independent predictor of cardiovascular events.<sup>2</sup> Other studies have investigated the relationship between high levels of serum uric acid (UA) and a left atrial thrombus/left atrial spontaneous echo contrast (LATH)/(LASEC) as detected by TEE and there are conflicting results.<sup>3–8</sup> We conducted a comprehensive meta-analysis and systematic review to investigate the potential relationship between the serum UA level and left atrial thrombi detected by TEE.

## Methods

We conducted this analysis according to the guidelines of the Meta-analysis of Observational Studies in Epidemiology Group (MOOSE).<sup>9</sup> Prospective or retrospective observational studies were enrolled with the primary objective of analyzing the association between serum levels of UA and LATH/LASEC. Titles and abstracts of all articles were evaluated and studies were rejected if the following inclusion criteria were not met: 1) human subjects with TEE and corresponding result of LATH/LASEC reported, 2) UA presented as continuous variable, 3) retrospective/prospective cohort studies, 4) baseline data available, 5) detail of LA abnormality detected by TEE mentioned, 6) sample size of more than 50 patients from individual studies.

### Search strategies

We performed a search of the on-line databases of PubMed and the Cochrane Controlled Trials Register Databases to July 2015 to identify relevant studies. We used the following keywords: “uric acid”, “hyperuricemia” and “atrial thrombus”, “spontaneous echo contrast”, and “left atrium abnormality”. Titles and abstracts as well as the reference lists of all the identified reports were independently examined by two reviewers (E.Z and T.L.) in order to include potentially relevant studies. The two reviewers agreed on the inclusion/exclusion status in 90% of the reviewed studies. Disagreements were resolved after discussion with a third reviewer (L.K.). There was no language restriction for the inclusion of studies. Additionally, we conducted a manual search using review articles, bibliographies of original papers, and abstracts of the scientific sessions of the American Heart Association, the European Society of Cardiology, the Heart Rhythm Society, and the American College of Cardiology from the past two years.

### Quality assessment

To limit heterogeneity secondary to differences between study designs, the quality of each study was evaluated according to the guidelines developed by the Evidence-Based Medicine Working Group<sup>10</sup> and the United States Preventive Task Force.<sup>11</sup> We applied the point score system that assessed the following characteristics: (1) clear description of inclusion and exclusion criteria, (2) study sample representative for the mentioned population, (3) clear description of sample selection, (4) full specification of clinical and demographic variables, (5) complete clinical data such as renal function and echocardiogram parameters, (6) no loss of follow-up, (7) cohort study, (8) clear definition of outcomes and outcome assessment, (9) temporality (assessment of UA level before TEE), and (10) adjustment of possible confounders in multivariate analysis. If one of these key points was not mentioned clearly in a study, we considered it not to have been performed properly. Therefore, the possibility of underestimation of the reported characteristics may be present.

### Data extraction

Two blind investigators (E.Z. and T.L.) independently performed data extraction with a standard data extraction form to determine eligibility for inclusion. The following collected information was tabulated: 1) publication details, first author's last name and the publication year, 2) characteristics of included studies, the study population, definition of LA abnormality, quality score, cohort design, risk estimate, and the patients' characteristics, 3) baseline data of the studied population, sample size, age, gender, paroxysmal atrial fibrillation (PAF) (%), LATH/LASEC (%), uric acid ( $\mu\text{mol/L}$ ), creatinine ( $\mu\text{mol/L}$ ), left atrium diameter, left ventricle ejection fraction, diabetes mellitus (%), hypertension (%), body mass index ( $\text{kg/m}^2$ ), and left atrial appendage flow velocity ( $\text{cm/s}$ ).

### Statistical analysis

The magnitude of the association between high serum UA level and the LATH/LASEC detected by TEE was measured by adjusted *OR*, or by an *OR* with 95% confidence intervals (*CI*s). One study<sup>7</sup> only provided a value of *OR* by univariate analysis because there was no association between the serum UA level and the LATH/LASEC following univariate analysis. We used the inverse variance method to weight studies

for the combined overall statistics. Finally, we examined the heterogeneity with the standard  $\chi^2$  test of heterogeneity. An  $I^2 > 50\%$  indicates at least moderate statistical heterogeneity.<sup>12</sup> A pooled effect was calculated with a random-effects model when the  $\chi^2$ -test for heterogeneity was found to be significant, through which we could take within-study and between-study variance into account. Sensitivity analysis was done by dropping studies and checking the consistency of the overall effect estimate. Statistical significance for the treatment effect was defined at  $P$  values  $< 0.05$ . Publication bias was evaluated using a funnel plot. All the analyses above were performed using Review Manager Version 5.3 (Revman; The Cochrane Collaboration, Oxford, UK).

## Results

We identified a total of 369 records following the primary article search. After screening the titles and abstracts, 363 studies were discarded because they were either review articles, laboratory studies, irrelevant to the current analysis, had insufficient statistics, or were non-cohort studies. After detailed evaluation, six

observational studies met all inclusion criteria (Fig. 1).<sup>3–8</sup> Overall, there were 2381 patients involved in our analysis. The average UA concentration ranged from 279.5  $\mu\text{mol/L}$ <sup>5</sup> to 368.8  $\mu\text{mol/L}$ .<sup>6</sup> The characteristics of each included study and baseline data of the patients from corresponding studies are depicted in Tables 1 and 2 respectively. Meta-analysis of the included studies demonstrated that a higher level of UA was associated with a higher incidence of LA abnormality ( $OR = 1.59$ , 95%CI 1.13–2.23,  $P = 0.008$ , Fig. 2), while there are significant differences between the individual studies ( $P < 0.00001$  and  $I^2 = 85\%$ ). Sensitivity analysis was performed to seek the origin of heterogeneity. However, removing any single study did not reduce the heterogeneity. The study from Numa et al<sup>4</sup> set the broadest definition of high risk in TEE, which included LATH, dense LASEC, low LAA flow velocity, and severe aortic atherosclerosis. However, when we exclude this study, the results remained positive, which further supported the close relationship between the high level of UA and the high incidence of LATH/LASEC ( $OR = 1.67$ , 95%CI 1.10–2.54,  $P = 0.02$ , Fig. 3). Elimination of any study did not alter the positive association. The funnel plot (Fig. 4) suggested that there was little publication bias,




Fig. 1. Flow diagram of study selection process.

Table 1  
Characteristics of studies included in meta-analysis.

| Investigator | Publication year | Patients (n) | Definition of LA abnormality  | Quality score | Cohort study design | Risk estimate | Patients characteristics                       |
|--------------|------------------|--------------|---|---------------|---------------------|---------------|--|
| Liu          | 2014             | 525          | LATH/SEC  | 7             | N/A                 | aOR           | AF   |
| Numa         | 2014             | 115          | TEE thromboembolic risk (TEE risk: low left atrial appendage flow, spontaneous echo contrast, thrombi, or aortic atherosclerosis)   | 8             | retrospective       | aOR           | CHA2DS2-VASc score = 0 or 1<br>Non-valvular AF |
| Ozturk       | 2014             | 207          | LATH (presence of a clearly defined echogenic intracavitary mass with an echo texture different from that of the underlying endocardium and not due to the pectinate muscle)                                  | 7             | N/A                 | aOR           | Mitral stenosis and sinus rhythm               |
| Tang         | 2014             | 1359         | Well-circumscribed echogenic mass with a LATH (unique echo texture contrasting with the adjacent or underlying myocardium) SEC (atrial blood pool in slow turbulent blood flow)                               | 8             | retrospective       | aOR           | Non-valvular AF                                |
| Zhao         | 2015             | 90           | LATH (a fixed or mobile echogenic mass within the left atrium) SEC (a dense persistent echogenic swirling pattern that can be identified in all TEE views regardless of ultrasound frequency or gain setting) | 8             | retrospective       | OR            | Non-valvular AF                                |
| Akpek        | 2015             | 85           | SEC (a pattern of slowly swirling intracavity echo densities imaged with gain settings which were adjusted to distinguish background noise)   | 8             | retrospective       | aOR           | Mitral stenosis                                |

AF: atrial fibrillation; aOR: adjusted odd ratio; N/A: not applicable.

LATH: left atrial thrombus; SEC: spontaneous echo contrast; TEE: transesophageal echocardiography.

Table 2  
Baseline clinical characteristics of included studies.

| Investigator | Publication year | PAF (%)        | LATH/SEC (%) | Age (years) | Male (%) | UA (μmol/L) | LAD (mm) | LVEF (%) | CR (μmol/L) | BMI (kg/m <sup>2</sup> ) | DM (%) | HTN (%) | LAAfv (cm/s) |
|--------------|------------------|----------------|--------------|-------------|----------|-------------|----------|----------|-------------|--------------------------|--------|---------|--------------|
| Liu          | 2014             | 83.0           | 10.9         | 60.8        | 64.2     | 353.9       | 37.9     | N/A      | N/A         | N/A                      | N/A    | N/A     | N/A          |
| Numa         | 2014             | 59.1           | 28.3         | 55.4        | 100      | 363.0       | 41.2     | 58.8     | 74.3        | 23.4                     | N/A    | N/A     | 49.1         |
| Ozturk       | 2014             | SR             | 10.1         | 41.9        | 20.8     | 279.5       | 44.8     | 60.8     | 79.6        | N/A                      | 24.6   | 29.0    | 35.3         |
| Tang         | 2014             | 65.1           | 4.5          | 58.2        | 70.8     | 368.8       | 39.0     | 63.4     | 83.6        | 26.2                     | 13.4   | 51.7    | N/A          |
| Zhao         | 2015             | 56.7           | 27.3         | 60.1        | 60.0     | 325.7       | 40.4     | 57.7     | 77.5        | N/A                      | 17.8   | 52.2    | N/A          |
| Akpek        | 2015             | 39 AF<br>46 SR | 51.8         | 46.0        | 34.1     | 323.2       | N/A      | 62.9     | 84.1        | 22.7                     | 28.2   | 47.1    | N/A          |

PAF: paroxysmal atrial fibrillation; SR: sinus rhythm; LATH: left atrial thrombus; SEC: spontaneous echo contrast; LAD: left atrium diameter; LVEF: left ventricle ejection fraction; BMI: body mass index; DM: diabetes mellitus; HTN: hypertension; LAAfv: left atrial appendage flow velocity; N/A: not applicable.

UA: uric acid 1 mg/dl = 59.5 μmol/L; CR: creatinine 1mg/dl = 88.41 μmol/L; CHA2DS2–VASc: Congestive heart failure, hypertension, Age >75 years, diabetes mellitus, stroke, vascular disease, Age 65–74 years, sex category.

although the small number of studies made it difficult to interpret.

## Discussion

Our results showed that higher levels of UA could predict LATH/LASEC that were detected by TEE. However, when our investigation was restricted to patients with atrial fibrillation, we found that high serum UA level did not predict LATH/LASEC.

As a final enzymatic product of purine metabolism, UA depends on the activity of xanthine oxidase,<sup>13</sup> which participates in the formation of free radical superoxide anions. A study by Dudley et al showed that the xanthine oxidase activity in LAA of the AF group was 4.4 times greater than in the control group<sup>14</sup> resulting in local cellular damage and thrombus formation. The development of LATH/LASEC in AF patients may depend on the type and duration of AF. However, this association was not defined in the enrolled studies.




Fig. 2. Forest plot of association between high uric acid levels and left atrial thrombus or spontaneous echo contrast detected by transesophageal echocardiography.




Fig. 3. Forest plot of association between high uric acid levels and left atrial thrombus or spontaneous echo contrast detected by transesophageal echocardiography excluding the study from Numa.<sup>7</sup>




Fig. 4. Funnel plot of the meta-analysis.

Our previous research<sup>15</sup> had showed that there was a close relationship between the presence of AF and elevated serum UA levels in hypertensive patients. Inflammation and oxidative stress have been clearly associated with both progression of AF<sup>16,17</sup> and left atrial abnormality.<sup>13,18</sup> However, it is unclear if there is a direct causal relationship between elevated serum UA levels and AF. We aimed to determine if abnormal serum UA concentrations participate in the development of

LATH/LASEC detected by TEE. To the best of our knowledge, there was no clear evidence supporting the association between higher serum UA levels and the detection of LA abnormalities among patients with AF despite the slight difference found in the total population when the patients with mitral stenosis and sinus rhythm were taken into consideration in the present study.

LATH/LASEC is an important risk factor for stroke in patients with non-valvular AF. A strong association between higher plasma UA levels on admission and poor coronary blood flow was also found in ST-elevation myocardial infarction patients receiving primary PCI. Hypercoagulability and thrombus formation may be reflective of abnormal UA concentrations.<sup>19</sup> Zapolski et al<sup>20</sup> also indicated that elevated serum UA could result in a prothrombotic state. Other epidemiological studies demonstrated a link between hyperuricemia and an increased risk of ischemic stroke and mortality.<sup>21,22</sup> On the other hand, Chamorro et al<sup>23</sup> reported that UA may provide an additional therapeutic effect in the basis of thrombolysis among patients with acute stroke. In addition, some researchers presumed UA elevation had a compensatory protective mechanism with antioxidant action.<sup>24</sup> The conflicting data encouraged us to weigh



the merits and demerits of higher UA in AF patients more carefully.

LATH is a common source of thromboembolism and can be seen frequently in patients with AF<sup>25</sup> and valve disease (especially mitral stenosis).<sup>5,8</sup> However, the precise mechanism of thrombus formation in LA in the setting of high serum levels of UA is unclear. The vascular damage induced by hyperuricemia may be important as indicated by the excessive oxidant generation and inflammatory response in vascular endothelium.<sup>26,27</sup> UA could promote tissue inflammation and deterioration of endothelial function, which may be responsible for AF and accompanying LA thrombus formation.<sup>28</sup> Additionally, hyperuricemia could induce proliferation of vascular smooth muscle cell and reduce the production of vascular nitric oxide leading to development of turbulent flow and thrombus formation.<sup>29</sup> Moreover, blood flow velocity in LAA may be another precursor of LATH. It was found to have an inverse correlation with serum UA levels among 130 patients with AF.<sup>30</sup>

The mechanism of LATH formation in patients with mitral stenosis involves inflammation, intra-atrial blood stasis, and abnormal platelet size and activation.<sup>31,32</sup> Exposure to high UA levels leads to a hypercoagulable condition and further accelerates the LATH formation.<sup>5</sup>

## Conclusion

Our comprehensive meta-analysis shows that increased serum UA levels could be considered an independent predictor for LATH or LASEC detected by TEE. This finding leads to the question of whether pharmacologic reduction of uric acid can reduce the risk of stroke. Large prospective studies will be needed to address this issue.

## Limitations

There were several limitations in this meta-analysis. First, the number of enrolled studies was relatively small and limited to an Asian population, despite the large sample size. Second, medications that may have influenced the serum levels of UA were not reported except for two studies,<sup>6,7</sup> and Tang et al<sup>6</sup> excluded patients on diuretics or allopurinol. Third, the disparity in age and sex ratio may have affected the UA level as there are hormonal fluctuations of UA among women at menarche and menopause.<sup>33</sup> Fourth, Numa et al<sup>4</sup> only included patients at low-intermediate clinical risk. Fifth, Ozturk<sup>5</sup> enrolled patients with mitral stenosis and sinus rhythm, which was significantly different from the other

studies. Finally, the endpoints defined by individual studies differ. One study included patients with a low LAA flow and aortic atherosclerosis as well.<sup>4</sup>

## Acknowledgments

This work was supported by a grant (No. 81270245 to T.L.) from the National Natural Science Foundation of China.

## References

1. Sasahara E, Nakagawa K, Hirai T, et al. Clinical and transesophageal echocardiographic variables for prediction of thromboembolic events in patients with nonvalvular atrial fibrillation at low-intermediate risk. *J Cardiol*. 2012;60:484–488.
2. Niskanen LK, Laaksonen DE, Nyssönen K, et al. Uric acid level as a risk factor for cardiovascular and all-cause mortality in middle-aged men: a prospective cohort study. *Arch Intern Med*. 2004;164:1546–1551.
3. Liu F, Wei W, Xue Y, Zhan X, Fang X, Liao H. Predictive value of serum uric acid level for left atrial thrombus or spontaneous echo contrast in patients with atrial fibrillation. *JACC*. 2014;64(16 Suppl C):C235.
4. Numa S, Hirai T, Nakagawa K, et al. Hyperuricemia and transesophageal echocardiographic thromboembolic risk in patients with atrial fibrillation at clinically low-intermediate risk. *Circ J*. 2014;78:1600–1605.
5. Ozturk D, Celik O, Akin F, et al. Usefulness of the uric acid and CHA2DS2-VASC score in prediction of left atrial thrombosis in patients with mitral stenosis and sinus rhythm. *Cardiol J*. 2015;22:336–342.
6. Tang RB, Dong JZ, Yan XL, et al. Serum uric acid and risk of left atrial thrombus in patients with nonvalvular atrial fibrillation. *Can J Cardiol*. 2014;30:1415–1421.
7. Zhao J, Liu T, Korantzopoulos P, et al. Red blood cell distribution width and left atrial thrombus or spontaneous echo contrast in patients with non-valvular atrial fibrillation. *Int J Cardiol*. 2015;180:63–65.
8. Akpek M, Sahin O, Sarli B, et al. Association of left atrial spontaneous Echo contrast to uric acid in patients with mitral stenosis. *Echocardiography*. 2015;32:1477–1482.
9. Stroup DF, Berlin JA, Morton SC, et al. Meta-analysis of observational studies in epidemiology: a proposal for reporting. Meta-analysis of Observational Studies in Epidemiology (MOOSE) group. *JAMA*. 2000;283:2008–2012.
10. Levine M, Walter S, Lee H, Haines T, Holbrook A, Moyer V. Users' guides to the medical literature. IV. How to use an article about harm. Evidence-Based Medicine Working Group. *JAMA*. 1994;271:1615–1619.
11. Harris RP, Helfand M, Woolf SH, et al. Methods work group, third US preventive services task force. Current methods of the US preventive services task force: a review of the process. *Am J Prev Med*. 2001;20(3 suppl 1):21–35.
12. Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ*. 2003;327:557–560.
13. Strazzullo P, Puig JG. Uric acid and oxidative stress: relative impact on cardiovascular risk? *Nutr Metab Cardiovasc Dis*. 2007;17:409–414.

14. Dudley Jr SC, Hoch NE, McCann LA, et al. Atrial fibrillation increases production of superoxide by the left atrium and left atrial appendage: role of the NADPH and xanthine oxidases. *Circulation*. 2005;112:1266–1273.
15. Liu T, Zhang X, Korantzopoulos P, Wang S, Li G. Uric acid levels and atrial fibrillation in hypertensive patients. *Intern Med*. 2011;50:799–803.
16. Patel P, Dokainish H, Tsai P, Lakkis N. Update on the association of inflammation and atrial fibrillation. *J Cardiovasc Electro-physiol*. 2010;21:1064–1070.
17. Burashnikov A, Antzelevitch C. New developments in atrial antiarrhythmic drug therapy. *Nat Rev Cardiol*. 2010;7:139–148.
18. Bergamini C, Ciccoira M, Rossi A, Vassanelli C. Oxidative stress and hyperuricaemia: pathophysiology, clinical relevance, and therapeutic implications in chronic heart failure. *Eur J Heart Fail*. 2009;11:444–452.
19. Akpek M, Kaya MG, Uyarel H. The association of serum uric acid levels on coronary flow in patients with STEMI undergoing primary PCI. *Atherosclerosis*. 2011 Nov;219:334–341.
20. Zapolski T, Waciński P, Kondracki B. Uric acid as a link between renal dysfunction and both pro-inflammatory and prothrombotic state in patients with metabolic syndrome and coronary artery disease. *Kardiol Pol*. 2011;69:319–326.
21. Chao TF, Liu CJ, Chen SJ, et al. Hyperuricemia and the risk of ischemic stroke in patients with atrial fibrillation — could it refine clinical risk stratification in AF? *Int J Cardiol*. 2014;170:344–349.
22. Li M, Hou W, Zhang X, Hu L, Tang Z. Hyperuricemia and risk of stroke: a systematic review and meta-analysis of prospective studies. *Atherosclerosis*. 2014;232:265–270.
23. Chamorro A, Amaro S, Castellanos M, et al. Safety and efficacy of uric acid in patients with acute stroke (URICO-ICTUS): a randomised, double-blind phase 2b/3 trial. *Lancet Neurol*. 2014;13:453–460.
24. Glantzounis GK, Tsimoyiannis EC, Kappas AM, Galaris DA. Uric acid and oxidative stress. *Curr Pharm Des*. 2005;11:4145–4151.
25. Rathi VK, Reddy ST, Anreddy S, et al. Contrast-enhanced CMR is equally effective as TEE in the evaluation of left atrial appendage thrombus in patients with atrial fibrillation undergoing pulmonary vein isolation procedure. *Heart Rhythm*. 2013;10:1021–1027.
26. Wang H, Jacobs Jr DR, Gaffo AL, Gross MD, Goff Jr DC, Carr JJ. Longitudinal association between serum urate and subclinical atherosclerosis: the coronary artery risk development in young adults (CARDIA) study. *J Intern Med*. 2013;274:594–609.
27. Rothenbacher D, Kleiner A, Koenig W, Primatesta P, Breitling LP, Brenner H. Relationship between inflammatory cytokines and uric acid levels with adverse cardiovascular outcomes in patients with stable coronary heart disease. *PLoS One*. 2012;7:e45907.
28. Nishida K, Chiba K, Iwasaki YK, et al. Atrial fibrillation-associated remodeling does not promote atrial thrombus formation in canine models. *Circ Arrhythm Electrophysiol*. 2012;5:1168–1175.
29. Kanellis J, Kang DH. Uric acid as a mediator of endothelial dysfunction, inflammation, and vascular disease. *Semin Nephrol*. 2005;14:39–42.
30. Murat Ç, Emre Y, Yalçın G, et al. Serum uric acid level is correlated with decreased blood flow velocity of left atrial appendage in patients with atrial fibrillation. *JACC*. 2013:113.
31. Kaya MG, Akpek M, Elcik D, et al. Relation of left atrial spontaneous echocardiographic contrast in patients with mitral stenosis to inflammatory markers. *Am J Cardiol*. 2012;109:851–855.
32. Apek M, Kaya MG, Yarlioglu M, et al. Relationship between platelet indices and spontaneous echo contrast in patients with mitral stenosis. *Eur J Echocardiogr*. 2012;2011:865–870.
33. Stockl D, Doring A, Thorand B, Heier M, Belcredi P, Meisinger C. Reproductive factors and serum uric acid levels in females from the general population: the KORA F4 study. *PLoS One*. 2012;7:e32668.

Edited by Wei-Zhu Liu