

Letters to the Editor

Am. J. Hum. Genet. 69:664–665, 2001

Female Patient Showing Hypohidrotic Ectodermal Dysplasia and Immunodeficiency (HED-ID)

To the Editor:

Amorphous mutations of the *NEMO* gene (Smahi et al. 2000) lead to incontinentia pigmenti (IP [MIM 308300]), an X-linked dominant disorder in which heterozygous female patients exhibit skin lesions along the Blaschko lines, cicatrization of the retina, and hypodontia (Landy and Donnai 1993). The majority of hemizygous male fetuses die in utero. In the December 2000 issue of the *Journal*, Zonana et al. (2000) reported that hypomorphic mutations in the *NEMO* gene lead to an apparently novel X-linked recessive entity, hypohidrotic ectodermal dysplasia and immunodeficiency (HED-ID). HED-ID is characterized by features—ectodermal dysplasia and recurrent infections accompanied by dysgammaglobulinemia—distinct from those of IP. Zonana et al. (2000) stated that HED-ID affects only males and that female heterozygotes do not exhibit clinical manifestations of immunodeficiency. Independently of Zonana et al., in the March issue of the *Journal*, Aradhya et al. (2001) reported that hypomorphic mutations in *NEMO* lead to an “atypical form of IP in males” whose clinical features, including immunodeficiency, overlap those of HED-ID. We report a female patient with features of HED-ID and a heterozygous hypomorphic *NEMO* mutation to show that female heterozygotes can be overtly affected and that the wide phenotypic spectrum of female heterozygotes should be taken into account when counseling families with HED-ID.

The proband presented with hyperpigmented macules in a streaky configuration on the trunk and extremities at age 3 wk. Histologic examination revealed irregular basal melanosis and a slight increase in the number of melanocytes in the epidermis, but there was no evidence of the pigmentary incontinence typically observed in IP. The areas of skin involvement stabilized by age 5 mo. At that time, the proband and the mother had been reported as the first familial cases of “linear and whorled nevoid hypermelanosis” (Akiyama et al. 1994). At 4 years of age, the patient began to experience recurrent infections, including respiratory-tract infections, cervical lymphadenitis, otitis media, cellulitis, and soft-tissue abscesses. The patient had a history of chronic lung infections with secondary bronchiectasis and pulmonary hypertension at 8 years of age and developed renovascular hypertension at 10 years of age. Immunoglobulin determinations at age 10 years showed abnormally high levels of IgD, 1,520 mg/dL (normal range 1.8–8.5 mg/dL), and IgE levels elevated to 17,600 IU/mL (normal range <250 IU/mL). At 11 years of age, she exhibited eczematous, dry skin with hyperpigmentation along the Blaschko lines; decreased sweating; periorbital wrinkling of the skin; thin scalp hair and sparse eyebrows; conical incisors; and hypodontia with four permanent teeth missing. The patient died shortly after cardiac catheterization at 11 years of age. The proband’s mother manifested hyperpigmented macules in a streaky configuration on the trunk and the extremities at several weeks of age. The mother, 44 years of age, never developed clinical manifestations of immunodeficiency. The younger sister of the proband developed hyperpigmentation along the Blaschko lines in infancy but is otherwise healthy at 7 years of age.

After obtaining informed consent, a genomic DNA sample from the proband’s peripheral blood leukocytes was examined. Direct sequencing of the entire coding region of the *NEMO* gene (Smahi et al. 2000) revealed insertion of a cytosine (dupC1161) within a wild-type run of seven cytosines. This insertion resulted in a novel amino acid at codon 390–393 and a premature stop at codon 394. This mutation truncates the protein and deleted a putative zinc-finger domain. X-inactivation analysis of the proband’s peripheral blood leukocytes based on the methylation status of the androgen-receptor gene (Parrish et al. 1996) revealed that the normal and the mutated chromosomes were randomly inactivated. We were unable to perform molecular analyses of the mother and the younger sister.

The dupC1161 mutation has been documented in three other pedigrees (Zonana et al. 2000; Aradhya et al. 2001). The phenotypic spectrum of the obligate carrier females in these families, including the one in this report, is very broad: HED-ID in the proband of the family described here, linear and whorled skin hyperpigmentation in the mother and the younger sister, “typical IP skin-pigmentation and retinal abnormalities” in families XL267 and XL344 (Aradhya et al. 2001), and “normal” (HED-ID-4, II-2; Zonana et al. 2000).

Variable phenotype in females with the same mutation could be explained by variations in expression due to differences in X inactivation among families, as suggested by Aradhya (2000). If so, an X-inactivation study could be helpful in predicting phenotypic outcome in a female heterozygote with the mutation. However, the proband in the family described here (HED-ID phenotype) and the female heterozygote in family XL344 (IP-like phenotype; Aradhya et al. 2001) yielded the same random X-inactivation pattern. We therefore suspect that an X-inactivation study would have limited predictive value.

To summarize, hypomorphic mutations in the *NEMO* gene can lead to the devastating HED-ID phenotype in female heterozygotes as well as males. The very broad phenotypic spectrum of female heterozygotes, ranging from normal to overt immunodeficiency, poses a challenge during counseling of families with HED-ID.

Acknowledgments

This work was supported, in part, by a grant from Pharmacia Fund for Growth & Development Research and the Ministry of Education, Culture, Sports, Science & Technology of Japan. We would like to thank Drs. Akira Sakaguchi and Masashi Akiyama for referring the families.

KENJIRO KOSAKI,^{1,2} NORIKO SHIMASAKI,³

HIROYUKI FUKUSHIMA,² MITSUHIRO HARA,³

TSUTOMU OGATA,¹ AND NOBUTAKE MATSUO¹

¹Department of Pediatrics and ²Pharmacia Fund for Growth & Development Research, Keio University School of Medicine, Tokyo; and ³Shimizu City Hospital, Shimizu, Japan

Electronic-Database Information

Accession numbers and the URL for data in this article are as follows:

Online Mendelian Inheritance in Man (OMIM), <http://www.ncbi.nlm.nih.gov/Omim/> (for IP [MIM 308300] and HED-ID [MIM 300291])

References

- Akiyama M, Aranami A, Sakaki Y, Ebihara T, Sugiura M (1994) Familial linear and whorled nevoid hypermelanosis. *J Am Acad Dermatol* 30:831–833
- Aradhya S, Courtois G, Rajkovic A, Lewis RA, Levy M, Israel A, Nelson F (2001) Atypical forms of incontinentia pigmenti in male individuals result from mutations of a cytosine tract in exon 10 of *NEMO* (*IKK-γ*). *Am J Hum Genet* 68: 765–771
- Landy SJ, Donnai D (1993) Incontinentia pigmenti (Bloch-Sulzberger syndrome). *J Med Genet* 30:53–59
- Parrish JE, Scheuerle AE, Lewis RA, Levy ML, Nelson DL (1996) Selection against mutant alleles in blood leukocytes

is a consistent feature in incontinentia pigmenti type 2. *Hum Mol Genet* 5:1777–1783

Smahi A, Courtois G, Vabres P, Yamaoka S, Heuertz S, Munnich A, Israel A, et al (2000) Genomic rearrangement in *NEMO* impairs NF-κB activation and is a cause of incontinentia pigmenti: the International Incontinentia Pigmenti (IP) Consortium. *Nature* 405:466–472

Zonana J, Elder ME, Schneider LC, Orlow SJ, Moss C, Golabi M, Shapira SK, Farndon PA, Wara DW, Emmal SA, Ferguson BM (2000) A novel X-linked disorder of immune deficiency and hypohidrotic ectodermal dysplasia is allelic to incontinentia pigmenti and due to mutations in *IKK-gamma* (*NEMO*). *Am J Hum Genet* 67:1555–1562

Address for correspondence and reprints: Dr. Kenjiro Kosaki, Division of Medical Genetics, Department of Pediatrics, Keio University School of Medicine, 35 Shinanomachi, Shinjuku-ku, Tokyo 160-8582, Japan. E-mail: kkosaki@sc.itc.keio.ac.jp

© 2001 by The American Society of Human Genetics. All rights reserved. 0002-9297/2001/6903-0021\$02.00

Am. J. Hum. Genet. 69:665–666, 2001

Reply to Kosaki et al.

To the Editor:

The letter from Kosaki et al. (2001 [in this issue]) reports the first instance of immunodeficiency in a female heterozygous for a hypomorphic mutation of the X-linked *NEMO* gene. *NEMO* hypomorphic mutations result, in males, in the disorder hypohidrotic ectodermal dysplasia with immunodeficiency (HED-ID) (Zonana et al. 2000; Aradhya et al. 2001; Doffinger et al. 2001; Jain et al. 2001). In the 17 families previously reported, only males have had clinical immunodeficiency. As expected, owing to Lyonization, carrier females either were normal or had defects of their teeth, skin, or hair. A single female carrier, heterozygous for the same mutation reported in the family studied by Kosaki et al., had an elevated immunoglobulin A level (Zonana et al. 2000) but did not have immunodeficiency.

Null mutation of the *NEMO* gene generally results in prenatal lethality in hemizygous males and in incontinentia pigmenti in heterozygous females. Affected females are immunologically normal, presumably owing to preferential survival and proliferation of cells expressing the normal allele (skewed X inactivation). Female carriers with hypomorphic mutations have shown both random and skewed X inactivation, indicating that at least some of the mutations do not affect T- and B-cell survival or proliferation (Aradhya et al. 2001; Doffinger et al. 2001; Jain et al. 2001). Rarely, females can fully manifest X-linked recessive disorders, including other X-linked immunodeficiencies, such as Wiskott-Al-