

single transmembrane helix, long unstructured linker followed by two cytoplasmic domains, called C2A and C2B, which are separated by a short flexible linker. The reports of the intramolecular interactions between C2A and C2B domains in solution are inconsistent. In order to determine relative rearrangements of C2A and C2B domains in the presence of synaptotagmin ligands, lipids, SNARE proteins and SNARE complex, single molecule fluorescence resonance energy transfer (sm FRET) was used. C2A and C2B domains were each labeled with a single fluorophore, tethered to the surface and conformational dynamics monitored using smFRET. Results suggest that the relative rearrangement of Synaptotagmin III C2A and C2B domains in the presence and absence of calcium is dynamic and influenced by interactions with liposomes, SNARE proteins, and SNARE complex.

1842-Pos Board B686

Fusion Activity of HIV GP41 Fusion Domain is Related to its Secondary Structure in a Cholesterol-Dependent Fashion

Alex L. Lai, Yinling Li, Lukas K. Tamm.

University of Virginia, Charlottesville, VA, USA.

The HIV gp41 fusion domain plays a critical role in membrane fusion during the viral entry process. A thorough understanding of the relationship between the structure and activity of the fusion domain in different environments helps to formulate mechanistic models on how it might function in mediating membrane fusion. We examined the secondary structure of the fusion domain in small unilamellar vesicles composed of different lipid compositions by circular dichroism spectroscopy. The results show that the secondary structure switches from alpha helix below approx. 30% cholesterol to beta structure above this threshold. The lipid and content mixing activities of the fusion domain were examined by standard FRET fusion assays. Although the fusion activities increase with higher percentages of cholesterol, the helical conformation supports fusion in the absence of cholesterol and the beta-sheet conformation supports fusion in the presence of >30% cholesterol. Our results also show that the fusion domain has a higher activity in more negatively charged membranes, which can be explained by its higher binding affinity to such membranes.

1843-Pos Board B687

Membrane Fusion in Autophagy

David Taresté¹, Ai Yamamoto², Thomas Melia¹.

¹Yale School of Medicine, New Haven, CT, USA, ²Columbia University, New York City, NY, USA.

Macroautophagy is a pathway for the sequestration and degradation of cytosolic material when the cell is faced with stress conditions like starvation or infection. This cytosolic material is captured within a double-membraned vesicle (the autophagosome) which forms de novo and ultimately traffics to the lysosome for degradation. How membranes come together to form the autophagosome is unknown, but recent studies have suggested a possible role for the autophagosome-associated protein ATG8 in membrane tethering and fusion. Here we develop novel membrane platforms and protein-lipid coupling strategies to test fundamental aspects of ATG8 function, including the involvement of associated factors (ATG3 and ATG7) and the role of different lipid populations such as phosphatidylethanolamine. We compare the functionality of ATG8 to its mammalian homologue, MAP1LC3 and to a variety of mutants to establish which regions of the protein contribute to each activity. We utilize a range of different membrane dynamics assays including lipid-mixing, real-time membrane tethering, vesicle aggregation and cryo-electron microscopy. Our results suggest ATG8 may act as a "molecular glue", bringing membranes and possibly protein cargo together, but cast doubt on the physiologic relevance of ATG8 as a membrane fusion protein.

1844-Pos Board B688

Hemifusion of Giant Lipid Vesicles by a Small Transient Osmotic Depletion Pressure

Chang-chun Lee, Shuo Qian, Yen Sun, Huey W. Huang.

Rice University, Houston, TX, USA.

We observed hemifusion of giant unilamellar vesicles (GUVs) by injecting a small volume of 5% polyethylene glycol solution in the vicinity for a short time. The induced transient osmotic depletion attraction between two bilayers was smaller than the corresponding van der Waals force, yet hemifusion readily occurred for some lipid compositions. This might imply that the energy barrier for hemifusion is very low if the lipid compositions were primed for the reaction. The barrier for hemifusion is then the task of bringing two membranes into contact. We found that hemifusion is a stochastic event very much like pore formation by tension or by pore-forming peptides. This method provides an operational definition for hemifusogenicity which so far has been defined only qualitatively. Surprisingly the configuration of hemifusion depends on the lipid composition, ranging from a stalk-like hemifusion to a large hemifusion dia-

phragm. The formation of a large hemifusion diaphragm requires trans-bilayer lipid mixing in the participating GUVs. We also found that hemifusogenicity correlates with the water activity for the stalk phase in the lipid water mixture. The effect of cholesterol on hemifusogenicity was clearly demonstrated.

1845-Pos Board B689

Direct Visualization of Large and Protein-free Hemifusion Diaphragms

Joerg Nikolaus¹, Martin T. Stoeckl¹, Dieter Langosch², Rudolf Volkmer³, Andreas Herrmann¹.

¹Humboldt University Berlin, Berlin, Germany, ²Technische Universität München, München, Germany, ³Charité, Berlin, Germany.

Fusion of cellular membranes is an ubiquitous biological process requiring remodeling of two phospholipid bilayers. Very likely, merging of membranes proceeds via similar sequential intermediates. Contacting membranes form a stalk between the proximal leaflets which expands radially into a hemifusion diaphragm (HD) and subsequently open to a fusion pore. While considered to be a key intermediate in fusion, direct experimental verification of this structure is difficult due to its transient nature. Using confocal fluorescence microscopy we have investigated the fusion of giant unilamellar vesicles (GUVs) containing fluorescent membrane protein anchors and fluorescent lipid analogues in the presence of divalent cations. Time resolved imaging revealed that fusion was preceded by displacement of peptides and lipid analogues from the GUV-GUV contact region being of several μm in size. A detailed analysis of contact regions which could be stabilized at lower cation concentration showed that peptides were completely sequestered from this site as expected for an HD. Lateral distribution of lipid analogues in attached GUVs in three different experimental setups as well as fluorescence recovery after photobleaching (FRAP) measurements were consistent with the formation of an HD but not with the presence of two closely approached bilayers.

1846-Pos Board B690

Covalent Inhibition of HIV Membrane Fusion

Amy Jacobs.

State University of New York at Buffalo, Buffalo, NY, USA.

Peptide inhibitors consisting of sequences from the six helix bundle structure of the fusogenic portion (gp41) of the HIV envelope glycoprotein have been successfully implemented in preventing HIV entry. These peptides block entry by binding to regions of HIV gp41 that are transiently exposed during the fusion stage of viral entry. We have successfully designed and tested peptide analogs composed of chemical spacers and reactive moieties positioned strategically to facilitate the covalent attachment of the peptide to gp41. We have utilized the covalent peptide to show evidence for the trapping of a pre-six helix bundle fusion intermediate by a covalent reaction with a specific anti-HIV-1 peptide. This is the first demonstration in live cells of the trapping of an intermediate conformation of a viral envelope glycoprotein during the fusion process. The permanent specific attachment of the covalent inhibitor is projected to improve the long-term sustainability of peptide entry inhibitor therapy and to serve as an important tool in probing the conformational changes that occur to the envelope protein complex during viral entry.

1847-Pos Board B691

Leaky Fusion Between Intraluminal Vesicles Of Late Endosomes That Have An Unusual Lipid Composition As A Mechanism Of Endosomal Escape By Cell-penetrating Peptides

Sung-Tae Yang.

NICHHD, Bethesda, MD, USA.

It has recently been recognized that endocytosis/pinocytosis plays a major role in the delivery of various bioactive molecules such as proteins and nucleic acids by HIV Tat-derived peptide (TAT). In this work we explored the mechanism of TAT escape from endosomes using protein-free liposomes. We found that TAT induces both membrane leakage and fusion of liposomes mimicking late endosomal lipid composition and the extent increased with an increase in the content of bis(monoacylglycerol)phosphate (BMP), which is a characteristic lipid of late endosomal membrane. The TAT-induced membrane fusion of BMP-containing liposomes was promoted by acidic pH and elevated Ca^{2+} concentration. In contrast, liposomes in which BMP was replaced by its structural isomer phosphatidylglycerol (PG) were much less affected by TAT. While there was no significant difference between BMP and PG in the binding affinity of TAT, effects of BMP and PG on the L_a to inverted H_{II} phase transition of egg PE suggested that BMP is more fusogenic than PG. In addition, modifications of liposome composition that inhibited TAT-induced lipid mixing (incorporation of either PEG-lipid or LPC) also inhibited TAT-induced leakage. Based on these results, we propose that TAT induced leaky fusion between BMP-containing bilayers of late endosomal membranes first deliver TAT into the intraluminal vesicles and then, upon vesicle fusion with the limiting membranes, release the peptide into cytosol.