

Available online at [www.sciencedirect.com](http://www.sciencedirect.com)**ScienceDirect**

Radiology of Infectious Diseases 3 (2016) 74–78

[www.elsevier.com/locate/jrid](http://www.elsevier.com/locate/jrid)

Case report

# Supratentorial extradural epidermoid cyst associated with brain abscesses

Ruisheng Li <sup>a</sup>, Xiaoliang Liu <sup>a</sup>, Kang Yuan <sup>a</sup>, Meng Law <sup>b</sup>, Bo Gao <sup>a,\*</sup><sup>a</sup> Department of Radiology, Yantai Yuhuangding Hospital, Yantai, Shandong 264000, China<sup>b</sup> Department of Radiology, Keck School of Medicine, University of Southern California, Los Angeles, CA 90033, USA

Received 16 March 2016; revised 21 March 2016; accepted 23 March 2016

Available online 28 March 2016

## Abstract

We describe the unique CT and MRI findings of a young woman presenting with cerebral abscesses resulting from left temporal epidermoid cyst secondary infection confirmed at operation. Early neurosurgical intervention is advocated to prevent the ongoing risk of significant complications. The significances of pre-operative imaging evaluation and procedure planning are also discussed.

© 2016 Beijing You'an Hospital affiliated to Capital Medical University. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

**Keywords:** Epidermoid cyst; Cerebral abscess; Magnetic resonance imaging; Diffusion-weighted imaging

## 1. Introduction

Epidermoid cysts are slow-growing congenital lesions of ectodermal origin, representing approximately 1% of all primary intracranial tumors. Intracranial epidermoid cysts are considered to be congenital in etiology, though trauma or surgery is also known to cause epidermoid cysts in brain, spine or elsewhere [1]. Patients with epidermoid cyst occasionally experience complications such as aseptic meningitis and delayed postoperative hemorrhage [2]. In this article, we report a unique case of a patient presenting with headache and fever, found to have an epidermoid cyst in middle fossa associated with an ipsilateral temporal lobe abscess.

## 2. Materials and methods

A 29-year-old female initially caught a cold (the highest temperature reached as high as 39 °C) and presented to our hospital with duration of over twenty days of fever and headache, her symptom of headache associated with nausea

and vomiting aggravated in recent three days. She had been affected with hearing loss from childhood without any proper treatment. Physical examinations: temperature 36.6 °C, pulse 78 per minute, heart rate 20 per minute, blood pressure 129/78 mmHg. General examinations were within normal limits except of her bilateral conductive hearing loss. No focal neurological deficit was elicited. On admission, her white blood cell count was 9720 cells/mcL, neutrophils 64.7%, erythrocyte sedimentation rate (ESR) 19 mm/h, electrolytes and coagulation test were normal. Cerebrospinal fluid (CSF) routine analyses: white blood cells  $65 \times 10^6/L$ , mononuclear cell 90%, glucose 4.3 mmol/L, protein 487 mg/L, chloride 119.4 mmol/L, pressure 265 mm H<sub>2</sub>O, negative of bacterial cultures. So ceftriaxone sodium was prescribed to counter infect, mannitol to reduce her increased intracranial pressure and oxiracetam injection to protect neurons.

Cranial CT examination was immediately performed showing a well-defined bulging inhomogeneous hypo-attenuated round mass located in the left middle cranial fossa and intruding into intracranial parenchyma with ring-like hyper-attenuated thin wall (Fig. 1A). An ill-defined hypo-attenuated round lesion with slightly peripheral edema was also noticed in left temporal lobe on the higher level plane. Remarkable bony defect of left temporal bone with sclerotic margin was also demonstrated on bony window (Fig. 1B, C).

\* Corresponding author.

E-mail address: [gygb2004@163.com](mailto:gygb2004@163.com) (B. Gao).

Peer review under responsibility of Beijing You'an Hospital affiliated to Capital Medical University.




Fig. 1. Axial brain CT image showing a round bulging hypoattenuated mass in the left middle cranial fossa and intruding into intracranial parenchyma with ring-like hyper-attenuated thin wall (A). Axial bone window and oblique-coronal reformatted CT images demonstrating remarkable bony defect of left temporal bone with surrounding erosion and sclerosis and communicating with sphenoid sinus and left posterior ethmoid sinus with mucosal thickening (B, C).

In order to further characterize the lesion and associated intracranial extension, Cranial MRI examination was also performed. Moderate signal intensity on T1WI and T2WI images in the anterior part with hyperintense foci in the central components and hypointensities on fat-suppressed T1WI, behind which another oval hypointense cystic mass communicating each other via a defect between them two were demonstrated evidently (Fig. 2A, B). Axial T2WI revealed heterogeneous hyperintense components in the lesion and fluid–fluid level of hyperintensity in the posterior part, coronal T2WI showing its extradural origin relative to left skull base and parasellar location, hyperintense edema in proximal white matter of left temporal lobe was also noted (Fig. 2C, D). Grossly hyperintensity in the anterior part and mild hyperintensity in the posterior part are shown on diffusion-weighted imaging (DWI), in keeping with pus (Fig. 2E). Post-contrast T1WI after administering gadolinium showed thin-wall rim enhancement in both of the two parts (Fig. 2F).

Based on the complicated anatomies around the lesion and deteriorated clinical status, the patient was undertaken excision of skull base epidermoid cyst and drainage of intracranial abscess through the approach of posterior wall maxillary sinus with the aid of imaging navigation system. The operation was navigated with ENT dedicated imaging navigation system (Stealth Station Landmax 4; Medtronic Navigation, Louisville, Colorado) and digital high-resolution nasal endoscopes (Karl Storz Endoscopy, El Segundo, California, USA). Under general anesthesia, the patient was underwent a posterior-lateral approach of left maxillary sinus wall osteotomy and was exposed the anterior-inferior portion of the mass after entering into pterygopalatine fossa (PPF). A reddish soft cystic mass filling with white bean curd-like internal components was discovered, which was in accordance with the diagnosis of cholesteatoma (Fig. 3). After expanding the opening we performed intracystic incision and eliminated surrounding cystic wall with the aid of endonasal endoscopy guidance. During the procedure, we found the cholesteatoma enclosing left internal carotid artery and compressing left cavernous sinus, which was also communicating with left sphenoid sinus. The tumor

destroyed sphenoid greater wing and involved pterygoid process, however PFP was spared. When gently compressing internal carotid artery upward, yellow fluid flowed out from the back, which was considered the concomitant intracranial abscess. Pus was drained and aspirated, then was washed with normal solution. With the guidance of endonasal endoscopy, grossly thoroughly excision of the lesion and open the ostiums of maxillary, sphenoid sinuses and adequately drainage were performed. Histopathological examination demonstrated keratinized squamous epithelium-lined with a large amount of inflammatory cells infiltration, which was conformed to epidermoid cyst and associated with infection (Fig. 4).

Antibiotic treatment was entailed one week after surgery, the patient has resolved from her symptoms of headache and fever. The patient's post-operative course and subsequent clinical follow-up one month later were uneventful, and MRI follow-up showed normal drainage of left maxillary sinus, closure of intracranial abscess and without prominent residual and recurrence (Fig. 5A, B).

### 3. Discussion

Intracranial epidermoid tumors are congenital inclusion cysts with an outer capsule and inner cystic fluids. Intracranial epidermoid cysts are usually located in the cerebellopontine angle and parasellar region, as well as petrous apex, chiasmal region, brain stem or intraventricular cavity. Intracranial epidermoid cysts usually have a benign clinical process and biological nature. Microscopically, the capsule of the epidermoid cyst consists of a layer of avascular stratified squamous epithelium tissue, the contents are mainly composed of keratin in concentric layers and cholesterol in a solid crystalline state. Most of epidermoid cysts are subdural origin or occasionally located in epidural space, the latter of which usually forming into round, oval or massive shapes with well-demarcated boundary [3], as of in this case.

The imaging features of epidermoid cyst are variable and largely depend on their content, usually multiple layers of keratin debris, dense debris in cystic centers and intraluminal




Fig. 2. T1-weighted axial MRI demonstrating mixed signal intensities mass with hyperintense foci in the central components and hypointensities on fat suppressed T1-weighted imaging, behind which another oval hypointense cystic structure communicating each other via a defect between them two (A, B); T2-weighted axial MRI revealing the lesion with heterogeneous hyperintense components and fluid–fluid level of hyperintensity in the posterior part, T2-weighted coronal MRI showing its extradural origin relative to left skull base and parasellar location, hyperintense edema in white matter of left temporal lobe is also noted (C, D); Remarkable hyperintensity in the anterior part and mild hyperintensity in the posterior part are shown on diffusion-weighted imaging (E); Contrast enhancement T1-weighted axial MRI after administered gadolinium showing thin-wall rim enhancement in both of two parts (F).




Fig. 3. Intraoperative photographs, the lesion is characteristic of yellowish cyst fluid, cyst wall and significant erosion of the left temporal bone are observed, after resection of the lesion greasy appearance content is also observed.




Fig. 4. Histopathological examination showing epidermoid cyst-keratinized squamous epithelium-lined with a large amount of inflammatory cells infiltration (H&E,  $\times 100$ ).

calcifications. On CT, epidermoid cysts usually appear as hypodense heterogeneous mass, without contrast enhancement and mostly hyperdense compared to cerebrospinal fluid. The associated bony defect or tract may be visible on plain films on occasion, however which is best delineated on CT




Fig. 5. MRI follow-up one month later showing the closure of intracranial abscess and unresolved vasogenic edema in adjoining white matter of the left temporal lobe, without residual or recurrence of tumor is disclosed (A, B).

scan. Multi-planar reformations of CT scan reveal the exact location of the lesion, its relationship with adjacent structures and any related cranial vault thinning or focal defect. The lesion does not enhance with contrast unless infected. As in this case, the superinfection of epidermoid cyst and associated intracranial abscess should attribute to its communication with paranasal sinuses and the inflammatory reaction intrude into the mass resulting in secondary infection, which was better demonstrated on CT than MRI does. In the critically ill patient requiring urgent intervention, multi-detector CT with its inherent capabilities of multi-planar reformat and bone structure demonstration, rapidly provide comparative imaging information, making it a useful alternative to MRI. MRI has been advocated as the imaging modality of best choice in assessing epidermoid cysts, yet the position, their inhomogeneous content and rare occurrence would lead to difficulties in establishing the right diagnosis. The typical presentations can vary according to the content of the cyst and comprise a wide range of T1 and T2 hyper- and hypointensities without contrast enhancement. DWI can contribute to further characterization and differentiation, showing diffusion restricted on diffusion-weighted images and ADC maps due to its inherent viscosity of internal fluid which usually includes debris, keratin, water, and cholesterol [4]. However, if infected, the desmoplastic thickening of the adjacent dura would generally show contrast enhancement of linear patterns. As in this case, rim-like enhancement of the associated middle fossa abscesses and the infected epidermoid cyst is clearly seen on post-contrast T1WI. On the other side, epidermoid cyst and abscess are usually hypointense on T1WI but rarely can be hyperintense on T1WI and can mimic each other in different pulse sequences including DWI [5]. Preoperative cross sectional imaging combined with advanced techniques would be essential to assess its location, optimize the surgical approach, and to evaluate any related complications [6].

Prompt neurosurgical intervention is often necessary to reverse presenting neurologic deficits, and treat systemic complications from severe intracranial infection secondary to abscess formation or bacterial meningitis. Definitive treatment involves complete removal of the related tumor and appropriate aspiration, drainage or excision of any accompanying abscesses. The preferred surgical treatment for epidermoid tumors in the skull base is the endoscopic endonasal approach (EEA) [7]. This innovative, minimally invasive technique uses the nose and nasal cavities as natural corridors to access hard-to-reach or previously inoperable tumors. Giant tumors that extend into multiple compartments are formidable, and the goal of total removal of the capsule would be quite challenging [8].

#### 4. Conclusion

We have presented a quite rare case with extradural epidermoid cyst and associated intracranial abscess. Early neurosurgical intervention is advocated to prevent the ongoing risk of serious complications. Pre-operative imaging workup plays a fatal role in procedure planning and prognosis assessment. Clarifying the pathogenesis would lead to effective preventive measures.

#### Fundings

This work was supported by National Natural Science Foundation of China (81471645), Natural Science Foundation of Shandong Province (ZR2014HL084), Shandong Provincial Key Research & Development Project (2015GSF118185), and Yantai Science & Technology Development Plan Program (2014WS005).

#### Acknowledgment

We appreciate the great help from our ENT department, Dr. Hua Zhang for providing the intraoperative photographs and

from our pathology department, Dr. Zhiqiang Lang for reading the pathological images. And we also give sincere thanks to Dr. Wende Gibbs, from neuroradiology department, Keck School of Medicine of USC, for revising the manuscript.

## References

- [1] Kaido T, Okazaki A, Kurokawa S, Tsukamoto M. Pathogenesis of intraparenchymal epidermoid cyst in the brain: a case report and review of the literature. *Surg Neurol* 2003;59(3):211–6.
- [2] Kalfas F, Ramanathan D, Mai J, Schwartz S, Sekhar LN. Petrous bone epidermoid cyst caused by penetrating injury to the external ear: case report and review of literature. *Asian J Neurosurg* 2012;7(2):93–7.
- [3] Shellenberger DL, Roehm PC, Gantz B. Large extradural epidermoid tumor of the temporal bone and posterior fossa cranium. *Otol Neurotol* 2006;27(7):1043–4.
- [4] Hu XY, Hu CH, Fang XM, Cui L, Zhang QH. Intraparenchymal epidermoid cysts in the brain: diagnostic value of MR diffusion-weighted imaging. *Clin Radiol* 2008;63(7):813–8.
- [5] Chatterjee S, Deepak KS, Banuprakash S, Saini J, Kesavadas C, Kapilamoorthy TR, et al. T1 hyperintense cystic lesion-an epidermoid or an abscess? *Clin Neuroradiol* 2010;20(1):66–9.
- [6] von Koch CS, Young G, Chin CT, Lawton MT. Magnetic resonance imaging/spectroscopy of an intraaxial epidermoid: similarity to an abscess. Case illustration. *J Neurosurg* 2002;97(2):492.
- [7] Best J, Schneider JS, Turner JH. Endoscopic endonasal resection of a giant middle fossa epidermoid cyst. *Allergy Rhinol (Providence)* 2015;6(3):195–7.
- [8] Aboud E, Abolfotoh M, Pravdenkova S, Gokoglu A, Gokden M, Al-Mefty O. Giant intracranial epidermoids: is total removal feasible? *J Neurosurg* 2015;122(4):743–56.