

HOSTED BY

Contents lists available at [ScienceDirect](http://www.elsevier.com/locate/jtcm)

Journal of Traditional and Complementary Medicine

journal homepage: <http://www.elsevier.com/locate/jtcm>

Review article

Rosa damascena as holy ancient herb with novel applications



Mohaddese Mahboubi*

Department of Microbiology, Research Center of Barij Essence Pharmaceutical Company, Kashan, Iran

ARTICLE INFO

Article history:

Received 13 June 2015

Received in revised form

31 August 2015

Accepted 15 September 2015

Available online 30 October 2015

Keywords:

Rosa damascena

Traditional

Chemical composition

Pharmaceutical application

Gol-E-Mohammadi

ABSTRACT

Rosa damascena as an ornamental plant is commonly known as “Gole Mohammadi” in Iran. Iranian people have been called this plant, the flower of Prophet “Mohammad”. *R. damascena* is traditionally used for treatment of abdominal and chest pains, strengthening the heart, menstrual bleeding, digestive problems and constipation.

This paper reviews the ethnopharmacology, phytochemistry and pharmaceutical investigations on *R. damascena*.

All relevant databases and local books on ethnopharmacology of *R. damascena* were probed without limitation up to 31st March 2015 and the results of these studies were collected and reviewed.

R. damascena has an important position in Iranian traditional medicine. It is economically a valuable plant with therapeutic applications in modern medicine. The antimicrobial, antioxidant, analgesic, anti-inflammatory, anti-diabetic and anti-depressant properties of *R. damascena* have been confirmed. Citronellol and geraniol as the main components of *R. damascena* essential oil are responsible for pharmacological activities.

Overall, *R. damascena* as holy ancient plant with modern pharmacological investigations should be more investigated as traditional uses in large preclinical and clinical studies.

Copyright © 2016, Center for Food and Biomolecules, National Taiwan University. Production and hosting by Elsevier Taiwan LLC. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Rosa damascena Mill is the hybrid between *R. gallica* and *R. Phoenicia* and is the member of Rosaceae family with more than 200 species and 18,000 cultivars around the world. *R. damascena* as the king of flowers has been the symbol of love, purity, faith and beauty since the ancient times. It was originated from Iran and essential oil extracting from its flowers has been started since 7th century A.D. It was brought to Europe and has been cultivated in European countries. Nowadays, Bulgaria and Turkey are the main producers of *R. damascena* essential oil in the world and the Bulgarian *R. damascena* oil is the known best ones. The Iranian people called it, Gol-E-Mohammadi (the flower of Prophet “Mohammad”) (Fig. 1). *R. damascena* flowers should handpick daily at early morning and are used immediately after harvesting for industrial purposes.

2. Different commercial derivations from *R. damascena*

Nowadays, different commercial products are produced from rose flowers. They are including:

2.1. Rose essential oil

Rose essential oil or liquid gold is expensive yellow pale essential oil that is extracted by different commercial methods. Bulgaria, Turkey and Morocco are the main producers of rose essential oil in the world.¹ The essential oil yield from Bulgarian rose varied from 0.032–0.049% (w/w).²

2.2. Rose water

Rose water or colorless liquid water with common name Golab (in Iran) due to its calming and relaxing properties is used in religious ceremonies like washing the God House in Mecca (Saudi Arabia) and also for flavoring the foods in Iran.

* Tel.: +98 8644465112; fax: +98 8644465187.

E-mail addresses: mahboubi1357@yahoo.com, Mahboubi@barjessence.com.

Peer review under responsibility of The Center for Food and Biomolecules, National Taiwan University.



Fig. 1. *Rosa damascena* flowers, buds and leaves.

2.3. Dried flowers

The dried buds and petals of rose are sold in groceries as flavor and laxative agents. In Iran, dried flowers are used as laxative agent and flavoring in foods.

2.4. Rose hips

Rose hips are the berry like fruits under the petals of flower. They are rich of vitamins, minerals and fatty acids, polyphenols, carotenoids and tannins^{3,4}. The vitamin C content of rose hips is higher than citrus fruits.

2.5. Rose concrete

Rose concrete, a red orange Vaseline mass is extracted by non polar solvent extraction and is the main material for production of rose absolute.

2.6. Rose absolute

Rose absolute or the ethanol extract of rose concrete has an orange red liquid with rose aroma.⁵

3. Chemical composition of *R. damascena*

Citronellol, geraniol, nerol, phenyl ethyl alcohol, nonadecane, nonadecene, eicosane, heneicosane, tricosane, α -guaiene, geranyl acetate and eugenol have been reported from different parts of

world (Table 1). Phenyl ethyl alcohol (71%) as the main component of rose essential oil was reported from Pakistan.⁶ Citronellol, nonadecane and geraniol were the main components of rose essential oil from central of Iran^{7,8} while the chemical composition of rose essential oil has been changed to triacosane (24.6%), 1-nonadecene (18.6%), n-tricosane (16.7%) and geraniol (15.5%) from north of Iran.⁹ Citronellol, geraniol, nonadecane and nerol were reported by Bulgarian authors as the main components.¹⁰ There was low variability in chemical composition of Bulgarian rose oil.¹¹

The chemical composition of Turkian rose essential oil^{12,13} was similar to chemical compositions of rose essential oil from central of Iran.^{7,8}

Different chemotypes of rose essential oil were reported from India,^{14,15} furthermore one study has been reported citronellol, geraniol, nonadecane¹⁵ as the main components of Indian rose essential oil.

The chemical compositions of rose essential oil change under different conditions; one of the most factors is the used plant's organ. For example, citronellol is obtained with a better yield in petals. Other factor is the freshness or dryness of petals that can affect on chemical compositions of rose essential oil.^{12,15} The use of acidic solutions like sulfuric acid solutions in extraction method removes some compounds such as geraniol, linalool, geraniol acetate, cis farnesol, nerol and 2-actamine and decreases the percentage of citronellol.⁷ Therefore, writing a documented procedure for harvesting of rose fresh petals, time of harvesting, condition of preservation and extracting the essential oil can help to produce the rose essential oil with high quality.

In total, the percentage of citronellol (20–34%), nerol (5–12%), geraniol (5–22%), paraffins c17 (1–2.5%), paraffins C19 (8–15%) and paraffins C21 (3.0–5.5%) are the best criteria for producing a high quality rose essential oil. In other word, the citronellol/geraniol ratio should be between 1.25–1.3. 2-octamine in essential oil decreases the quality of rose essential oil while citronellol is responsible for quality.⁷

There are some studies on chemical compositions of rose water and rose absolute.^{13,15,16} Phenyl ethyl alcohol (78.4%), citronellol (9.9%), nonadecane (4.4%) and geraniol (3.7%) have been reported as the main compounds of rose absolute.¹³ Geraniol (30.7%), citronellol (29.4%), phenyl ethyl alcohol (23.7%) and nerol (16.1%) have been reported as the main components of rose water¹³ while heneicosane (19.7%), nonadecane (13%), tricosane (11.3%), citronellol (7.1%), geraniol (2%) and citronellal (2.2%) also have been reported as the main component of rose water.¹⁵ In study that is published in 2015,¹⁶ the analysis of 10 rose water samples from Shiraz (Iran) revealed the presence of phenylethyl alcohol, geraniol and β -citronellol as the main components of rose water samples.

Table 1
Chemical composition of rose essential oil from different part of the world.

Main components	Origin	References
Phenyl ethyl alcohol (70.9%), citronellol (3.7%), rhodinol (2.7%), citranellyl acetate (2.5%), eugenol (1.6%), geraniol (1.5%)	Pakistan	[6]
Citronellol (23%), nonadecane (16%), geraniol (16%), heneicosane (5%)	Iran-Kashan	[8]
Citronellol (14.5–47.5%), nonadecane (10.5–40.5%), geraniol (5.5–18%), heneicosane (7–14%)	Iran-Kashan	[7]
Citronellol (48.2%), geraniol (17%), β -phenyl ethyl benzoate (5.4%) and phenyl ethyl alcohol (5.1%)	Iran-Kashan	[25]
Citronellol (35.2%), geraniol (22.2%), nonadecane (13.8%), nerol (10.3%)	Turkey	[13]
Citronellol (15.9–35.3%), geraniol (8.3–32.3%), nerol (4–9.6%), nonadecane (4.5–16%), heneicosane (2.6–7.9%)	India	[15]
Linalool (3.4%), nerol (3.1%), geraniol (15.5%), 1-nonadecene (18.6%), n-tricosane (16.7%), n-pentacosane (5.1%), n-hexa triacosane (24.6%)	Iran-Guilan	[9]
Citronellol (38.7%), geraniol (17.2%), nerol (8.3%), nonadecane (7.2%)	Turkey	[24]
Citronellol (24.5–42.9%), nonadecane (6.4–18.9%), geraniol (2.1–18.1%), ethanol (0–13.4%), heneicosane (2.3–8.9%), nerol (0.75–7.6%) and 1-nonadecene (1.8–5.4%)	Turkey	[12]
Phenyl ethyl alcohol (27.2%), octadecane (10.5%), hexadecane (7.8%)	India	[14]
Citronellol (23.4%), geraniol (19.0%), nonadecane (11.9%), nerol (7.5%)	Bulgaristan	[10]
Citronellol (23–28%), geraniol (14–20%), nonadecane (11–16%), nerol (6–11%), linalool (8%) and heneicosane (7%)	Saudi Arabia	[93]

4. Traditional uses of rose products

In Iranian traditional Medicines, the decoction of flowers was used for treatment of chest and abdominal pains, menstrual bleeding and digestive ailments (gentle laxative for constipation). It has been famous as cardiogenic agent for strengthening the heart.¹⁷ Rose essential oil was extracted by AveSina (Famous Iranian Scientist) in 10th century and was used for treatment of different ailments. Rose water traditionally was used as antiseptic agent for eye washing¹⁸ and mouth disinfecting¹⁹ and as antispasmodic agent for alleviating the abdominal pains, and bronchial and chest congestions.

The decoctions of dried rose water was used as diuretic and was recommended for relieving the fever, breast pain and menstrual problems.²⁰ In Iranian traditional medicine, rose petals were cooked with sugar or honey and used for cooling the mind and body. Rose hips were prescribed as blood purifier and also are used by Iranian people with bread.

5. Modern research on *R. damascena*

5.1. Antimicrobial activity

5.1.1. In vitro

Methanol and aqueous extracts of rose petals showed antiviral activity against HIV infection by targeting different stages of HIV replicative cycle. Kaempferol and its derivatives affect on viral protease and gp120/CD4.²¹ The antiviral activity of main components of rose essential oil, citronellol, geraniol were confirmed against HSV-1, *Haemophilus parainflunzae* type 3.²²

Furthermore, the acceptable antibacterial activity of rose essential oil were confirmed against *Xanthomonas axonopodis* spp. *Vesicatoria*²³, *Chromobacterium violaceum* and *Erwinia carotovora* strains,¹³ *Staphylococcus aureus*,²⁴ *Bacillus cereus*, *Staphylococcus epidermidis*, *Pseudomonas fluorescens*,¹⁸ *Pseudomonas aeruginosa*^{10,24,25}, *Escherichia coli*, *Proteus vulgaris*, *Klebsiella pneumoniae*, *Candida albicans*, *Enterococcus faecalis*,²⁴ *Enterococcus faecium* and *Salmonella typhimurium*^{24,25} were less sensitive to rose essential oil. Bulgarian rose oil has been shown no antimicrobial activity.²

Alcoholic and aqueous extracts of rose petals showed higher antibacterial activity than that of petroleum ether extract. *E. coli* was resistant to rose petal ethanol extract while its aqueous extract showed more sensitivity.²⁶ Ethanol extract also showed antimicrobial activity against methicillin resistant *S. aureus*, *S. typhimurium*, *B. cereus*, *C. albicans*.²⁷ The antibacterial activity of rose petal acetone extract was confirmed against *E. coli* and *B. subtilis*. This activity was higher than its aqueous extract.²³ *A. niger* exhibited less sensitivity to rose petal ethanol extract.²⁷

The antibacterial activity of rose water and rose absolute was confirmed against *E. coli*, *P. aeruginosa*, *B. subtilis*, *S. aureus*, *Chromobacterium violaceum* and *Erwinia carotovora* strains.^{13,23,28} The antibacterial activity of rose absolute, rose essential oil has been higher than rose water and rose extracts.¹³

Therefore, *R. damascena* like other holy medicinal plants²⁹ had antimicrobial activity.

5.1.2. In clinics

The effectiveness of herbal mouthwash containing rose extract in treatment of recurrent aphthous stomatitis were confirmed in two weeks randomized double blind, placebo-controlled clinical trial on fifty patients. The clinical results of mouthwash on pain, size and the number of ulcers was significantly different with placebo group. In other word, rose mouthwash was more effective than that of placebo group.³⁰

The anti-infective and anti-inflammatory effects of ophthacare[®], a product containing different herbs in combination with *R. damascena* were confirmed in patients with conjunctivitis, dry eye, acute dacryocystitis, pterygium or pinguecula disorders.^{31,32}

In poultry industry, dried rose dreg (by-product) decreased the occurrence of pathogen microorganisms including mesophilic aerobic bacteria, *Enterococci*, *Enterobacteriaceae* and *S. aureus* without any effect on broiler performance and feed conversion ratio during the production period.³³

Indeed, the antimicrobial activity of rose extracts especially rose essential oil and absolute is related to chemical components especially geraniol, citronellol and nerol or synergistic effects between these components. The antibacterial^{34–38} and antifungal^{39–43} activities of geraniol were confirmed against a large number of microorganisms. Also, the synergistic effect between citronellol, geraniol and nerol were demonstrated against Gram positive, Gram negative bacteria.²⁸ In other hand, the antimicrobial activity of rose extracts is related to chemical components of extracts and their synergistic or antagonistic effects.

5.2. Anticancer activity of *R. damascena*

The anti-tumor, anti-carcinogenic⁴⁴ and cytotoxic effects of *R. damascena*^{45,46} against cancer cells were confirmed. The geraniol as the main compounds of *R. damascena* acts via different mechanisms. It induces the apoptosis in cancer cells and increases the expression of apoptotic protein Bak,⁴⁷ arrests the G₀/G₁ phase of cell cycle and reduces cdk2 activity,⁴⁸ inhibits the 3-hydroxy-3-methylglutaryl-CoA (HMG-CoA) reductase⁴⁹ and ornithine decarboxylase activity⁵⁰ that finally causes the death in cancer cells.

5.3. Relaxant and anti-depressant activity of *R. damascena*

R. damascena via stimulating the β -adrenergic receptors,⁵¹ inhibiting the histamine H₁ receptors and blocking the calcium channels of tracheal chain,⁵² inhibiting the KCl-related contraction and electrical field stimulation⁸ shows the relaxant activity.

R. damascena aqueous and ethanol extracts affect on respiratory system of guinea pigs by inhibition of tachykinin and decrease the number of citric acid induced coughs,⁵² therefore, it shows bronchodilatory and antitussive effects.

Other sub-fractions of *R. damascena* such as its ethyl acetate fraction shows inhibitory effects on muscarinic receptors and induces relaxant effects on tracheal smooth muscles.⁵³ Therefore, *R. damascena* can be used as antitussive agent in clinics. Verifying this effect needs more clinical studies.

The anti-depressant effects of *R. damascena* aqueous extract were confirmed in animal models.⁵⁴ Rose absolute exhibits the anti-depressant activity by decreasing the lipid peroxidation and increasing the antioxidants in cerebral cortex.⁵⁵ Furthermore, depression is one reason for libido. As mentioned before, it is believed that rose essential oil and rose water bring happiness, self-confidence and are known as sensual and aphrodisiac agents. It has been confirmed that rose essential oil helps to infertility and libido via increasing the diameters of seminiferous tubules, sperm count and motility and enhancing the testosterone production.⁵⁶ Furthermore, administration of rose essential oil improves sexual dysfunction and symptoms of depression in male patients suffering from major depressive disorders (consuming the selective serotonin-reuptake inhibitors) higher than placebo group after week 2 to week 8.⁵⁷ In other world, patients with depression may benefits from rose extracts by their antidepressant activity and anti-libido effects.

5.4. Antioxidant activity of *R. damascena*

Decoctions,⁵⁸ aqueous extract,⁵⁹ essential oil,^{60–62} absolute,¹³ methanol,⁶³ ethanol extracts^{17,62} of rose petals have been shown antioxidant activity in different systems. The antioxidant activity of rose absolute with higher amount of carotene, α , β , γ -tocopherols was higher than that of rose essential oil and rose water.¹³

The antioxidant activity of *R. damascena* is not related to anthocyanin level,⁶⁴ but is correlated to total phenolic, flavonol contents of *R. damascena*.⁶³ Leaf methanol extract of rose with high amount of (+)-catechin and (–)-epicatechin as phenolic compounds has been shown antioxidant activity higher than that of BHT, trolox and BHT.⁶³

The benefit effects of rose essential oil against formaldehyde inhalation on reproductive system are related to the antioxidant activity of rose essential oil. Pretreatment with rose essential oil has been decreased the abnormal sperm and increased the sperm counts in rats.⁶⁰ Rose hips as herbal teas are consumed as strong antioxidant beverages.⁶⁵

Therefore, the benefit effects of *R. damascena* in scavenging of free radicals introduce it as good beverage for helping the health condition. In Iranian cultures, rose water was added to cold beverages as refreshing agent.

5.5. Analgesic, anti-inflammatory activities of *R. damascena*

The analgesic, anti-inflammatory effects of *R. damascena* ethanol,⁶⁶ chloroform extracts⁶⁷ have been shown in animal models, while rose essential oil has been exhibited no analgesic and anti-inflammatory effects.⁶⁷ Indeed, the component(s) that have analgesic effects in ethanol extract were not found in rose essential oil.

There are confusing results about the beneficial effects of rose hips in treatment of patients suffering from knee or hip osteoarthritis.^{22,68–70} It has been shown rose hip powder (10 g) for 1 month showed no anti-inflammatory or antioxidant effects in rheumatoid arthritis patients,⁶⁸ while others have been identified rose hips as anti-inflammatory agent.^{22,69,70} Unsaturated fatty acids, triterpenic acids or unidentified compounds and their synergistic effects exhibit anti-inflammatory effect⁷¹ via inhibiting cyclooxygenase 1 and 2.^{3,4}

5.6. Other pharmacological activities of *R. damascena*

The hypnotic effects of *R. damascena* different extracts (ethanol, aqueous extracts, and ethyl acetate, aqueous and n-butanol fractions) were comparable to diazepam.^{72–74} The best hypnotic effects of extracts were reported for ethyl acetate fraction. The hypnotic effects of ethyl acetate fraction may be related to the affinity of flavonoids in extracts to benzodiazepine receptors.⁷⁴

R. damascena is benefit to Alzheimer and dementia patients due to inhibitory effects on amyloid β formation, induction of neurite outgrowth^{75,76} and anti-cholinesterase activity.⁷⁷

The protective effects of *R. damascena* ethanol extract on memory performance of scopolamine-induced memory deficits rats were confirmed. The antioxidant effects of rose extract was the cause of memory enhancing.⁷⁸

Rose essential oil,⁷⁹ ethanol and aqueous extracts have been shown anti-seizure effects because of their anticonvulsant effects⁸⁰ and reducing of epileptic seizures. The affinity of flavonoids on GABAergic system in brain has been proposed as one of probable mechanisms because flavonoids enhance the effect of benzodiazepines on GABA receptors.⁸¹

As its traditional uses, the usefulness of *R. damascena* ethanol extract for treatment of digestive disorders has been shown.

R. damascena decreases ileum movements dose-dependently, probably through stimulating the β -adrenergic and opioid receptors and voltage-dependent calcium channels.⁸²

Rose essential oil has been used traditionally for treatment of cardiac diseases via massage on the skin. Nowadays, it has been shown *R. damascena* aqueous extract increases the heart rate and contractility in guinea pig via stimulatory effect on β -adrenoceptor⁸³ and suppressing the activity of ACE (angiotensin- I- converting) enzyme.⁸⁴

The efficacy of *R. damascena* extract on primary dysmenorrheal syndrome (PMS) was confirmed in double blind cross over clinical trial on 92 single girls. *R. damascena* extract with no side effects decreased the average of pain density in PMS such as mefenamic acid.⁸⁵

R. damascena methanol extract like other ethnobotanical holy plants such as *Trichilia emetica*, *Opilia amentacea* showed anti-diabetic activity.⁸⁶ *R. damascena* methanol extract inhibited α -glucosidase enzyme and suppressed carbohydrate absorption from small intestine. So, it reduces the postprandial glucose level, therefore it had anti-diabetic effects.⁸⁷

R. damascena methanol extract had moderate effects on reduction of total cholesterol, triglyceride and low density lipoprotein and plaques formation.⁸⁷ It has no effect on high density lipoproteins levels. Therefore, the anti hyperlipidemic effects of *R. damascena* methanol extract is caused through inhibiting the activities of pancreatic lipase and HMG COA reductase.^{88,89}

Waste rose petals as by product of rose essential oil's industries have been explored as a source of immune-modulating peptic polysaccharids.⁹⁰ Further studies are required to confirm the applications of waste rose petal as immunomodulator agent.

6. Toxicity

Oral LD₅₀ of *R. damascena* and rose absolute was >5 g/Kg in rats and dermal LD₅₀ of *R. damascena* was >2.5 g/kg in rabbits.⁹¹ Rose essential oil may cause sensitization in sensitive persons.

The potential toxic effects of *R. damascena* infusion in dogs at doses 90–1440 mg/kg/day (0.5–8 times of human uses) for 10 successive days revealed a minimal nephrotoxic or hepatotoxic effects. Therefore, it may have hepatotoxic effects at extraordinary high doses.⁹²

7. Conclusion

R. damascena Mill is one of the most important plants from *Rosaceae* family is a holy ancient plant with long historical uses in Iranian traditional medicine. Although, people in folklore, use rose as rose water or dried rose petals, but the majority application of *R. damascena* is producing of rose water and rose essential oil for application in religious ceremonies, cooking of some foods and in high grade perfumes. Rose essential oil is the important industrial product for different applications. The rose essential oil yield is very low (0.3–0.4 ml/kg). Ecological, geographical and environmental conditions, soil composition, harvesting and storage condition, distillation methods may affect on chemical composition of rose essential oil. The best rose essential oil is the oil with high amounts of monoterpenes. Therefore, finding the methods for increasing the oil yield and decreasing the expense of oil production are important issues for future.

The modern investigations on *R. damascena* have been confirmed the antiviral, antibacterial, anticancer, antidepressant, antioxidant, analgesic, anti-inflammatory, anticonvulsant activities and its relaxant and hypnotic effects.

Although, many studies demonstrate the medicinal applications of rose *in vitro* and in animal models, but other large preclinical and



Fig. 2. Some Iranian products from *Rosa damascena*.

clinical studies are needed for evaluating its potencies on different patients. In other word, there is low attention to therapeutic applications of rose products. In Iran, there are three famous products from BarijEssence Pharmaceutical Company (Fig. 2). GOL-E-Ghand Majoon has been produced according to traditional perscription and is prescribed for constipation as laxative agent. GOL-E-SORKH oral drop is used as anti-depressant and aphrodisiac agents. Gol-E-Mohammadi oil (the petal of rose in vegetable oil) is prescribed for ear pain and hemmorroid ailments.

There is a good situation for evaluating the potencies of *R. damascena* as traditional applications and planning the new products as the results of these researches.

Conflict of interest statement

We declare that we have no conflict of interest.

References

- Tosun I, Gonullu MT, Gunay A. Anaerobic digestion of residues from rose oil production. Paper Presented at: ISWA 2002 World Environmental Congress and Exhibition, 8–12 July 2002; Istanbul.
- Kovatcheva N, Zheljzakov VD, Astatkie T. Productivity, oil content, composition, and bioactivity of oil-bearing rose accessions. *Hort Sci.* 2011;46:710–714.
- Jäger AK, Eldeen IS, van Staden J. COX-1 and -2 activity of rose hip. *Phytother Res.* 2007;21:1251–1252.
- Wenzig EM, Widowitz U, Kunert O, et al. Phytochemical composition and in vitro pharmacological activity of two rose hip (*Rosa canina* L.) preparations. *Phytomedicine.* 2008;15:826–835.
- Aydinli M, Tutuş M. Production of rose absolute from rose concrete. *Flav Fragr J.* 2003;18:26–31.
- Khan MA, Rehman S. Extraction and analysis of essential oil of *Rosa* species. *Int J Agric Biol.* 2005;7:973–974.
- Sadraei H, Asghari G, Emami S. Inhibitory effect of *Rosa damascena* Mill flower essential oil, geraniol and citronellol on rat ileum contraction. *Res Pharm Sci.* 2013;8:1–7.
- Sadraei H, Asghari G, Emami S. Inhibitory effect of *Rosa damascena* Mill flower essential oil, geraniol and citronellol on rat ileum contraction. *Res Pharm Sci.* 2013;8:1–7.
- Yassa N, Masoomi F, Rankouhi SE, Hadjiakhoondi A. Chemical composition and antioxidant activity of the extract and essential oil of *Rosa damascena* from Iran, population of Guilan. *Daru.* 2009;17:175–180.
- Gochev V, Jirovets L, Wlcek K, et al. Chemical composition and antimicrobial activity of historical rose oil from Bulgaria. *J Essent Oil Bear Plants.* 2009;12:1–6.
- Rusanov K, Kovacheva N, Rusanova M, Atanassov I. Low variability of flower volatiles of *Rosa damascena* Mill. plants from rose plantations along the Rose Valley, Bulgaria. *Indust Crops Prod.* 2012;37:6–10.
- Bayrak A, Akgül A. Volatile oil composition of Turkish rose (*Rosa damascena*). *J Sci Food Agric.* 1994;64:441–448.
- Ulusoy S, Boşgelmez-Tınaz G, Seçilmiş-Canbay H. Tocopherol, carotene, phenolic contents and antibacterial properties of rose essential oil, hydrosol and absolute. *Curr Microbiol.* 2009;59:554–558.
- Perumal K, Sambanda Moorthy TA, Savitha JS. Characterization of essential oil from offered temple flower *Rosa damascena* Mill. *Asian J Exp Biol Sci.* 2012;3:330–334.
- Verma SR, Padalia CR, Chauhan A. Chemical investigation of the volatile components of shade-dried petals of damask rose (*Rosa damascena* Mill.). *Arch Biol Sci.* 2011;63:1111–1115.

16. Moein M, Zarshenas MM, Delnavaz S. Chemical composition analysis of rose water samples from Iran. *Pharm Biol.* 2014;52:1358–1361.
17. Shahriari S, Yasa N, Mohammadirad A, Khorasani R, Abdollahi M. In vivo antioxidant potentials of *Rosa damascena* petal extract from Guilan, Iran, comparable to α -tocopherol. *Int J Pharmacol.* 2006;3:187–190.
18. Gochev V, Wlcek K, Buchbauer G, et al. Comparative evaluation of antimicrobial activity and composition of rose oils from various geographic origins, in particular Bulgarian rose oil. *Nat Prod Commun.* 2008;3:1063–1068.
19. Akhmadieva A, Zaichkina SI, Ruzieva RK, Ganassi EE. The protective action of a natural preparation of anthocyan (pelargonidin-3, 5-diglucoside). *Radio-biologiya.* 1992;33:433–435.
20. Foster S, Duke JA. *Rosa rugosa* Thunb. In *Medicinal Plants*. 1990. New York, NY.
21. Mahmood N, Piacente S, Pizzi C, Burke A, Khan AI, Hay AJ. The anti-HIV activity and mechanisms of action of pure compounds isolated from *Rosa damascena*. *Biochem Biophys Res Commun.* 1996;229:73–79.
22. Orhan IE, Ozcelik B, Kartal M, Kan Y. Antimicrobial and antiviral effects of essential oils from selected Umbelliferae and Labiatae plants and individual essential oil components. *Turk J Biol.* 2012;36:239–246.
23. Basim E, Basim H. Antibacterial activity of *Rosa damascena* essential oil. *Fito-terapia.* 2003;74:394–396.
24. Jirovetz L, Eller G, Buchbauer G, et al. Chemical composition, antimicrobial activities and odor descriptions of some essential oils with characteristic floral-rosy scent and of their principal aroma compounds. *Recent Res Dev Agron Hort.* 2006;2:1–12.
25. Mahboubi M, Kazempour N, Khamechian T, Fallah MH, Kermani MM. Chemical composition and antimicrobial activity of *Rosa damascena* Mill essential oil. *J Biol Act Prod Nat.* 2011;1:19–26.
26. Hirulkar NB. Antimicrobial activity of rose petals extract against some pathogenic bacteria. *Int J Pharm Biol Arch.* 2010;1:478–484.
27. Talib WH, Mahasneh AM. Antimicrobial, cytotoxicity and phytochemical screening of Jordanian plants used in traditional medicine. *Molecules.* 2010;15:1811–1824.
28. Andogan BC, Baydar H, Kaya S, Demirci M, Özbaşar D, Mumcu E. Antimicrobial activity and chemical composition of some essential oils. *Arch Pharm Res.* 2002;25:860–864.
29. Konaté K, Yomalan K, Sytar O, Brestic M. Antidiarrheal and antimicrobial profiles extracts of the leaves from *Trichilia emetica* Vahl. (Meliaceae). *Asian Pac J Trop Biomed.* 2015;5:242–248.
30. Hoseinpour H, Peel SA, Rakhshandeh H, et al. Evaluation of *Rosa damascena* mouthwash in the treatment of recurrent aphthous stomatitis: a randomized, double-blinded, placebo-controlled clinical trial. *Quintessence Int.* 2011;42:483–491.
31. Biswas N, Gupta S, Das G, et al. Evaluation of Ophthacare® eye drops—a herbal formulation in the management of various ophthalmic disorders. *Phytother Res.* 2001;15:618–620.
32. Mitra S, Sundaram R, Venkatarangana M, et al. Anti-inflammatory, antioxidant and antimicrobial activity of ophthacare brand, an herbal eye drops. *Phytomedicine.* 2000;7:123–127.
33. Aktan S, Sagdic O. Dried rose (*Rosa damascena* Mill.) dreg: an alternative litter material in broiler production. *South Afr J Anim Sci.* 2004;34:75–79.
34. Friedman M, Henika PR, Levin CE, Mandrell RE. Antibacterial activities of plant essential oils and their components against *Escherichia coli* O157: H7 and *Salmonella enterica* in apple juice. *J Agric Food Chem.* 2004;52:6042–6048.
35. Inouye S, Takizawa T, Yamaguchi H. Antibacterial activity of essential oils and their major constituents against respiratory tract pathogens by gaseous contact. *J Antimicrob Chem.* 2001;47:565–573.
36. Scortichini M, Rossi MP. In vitro susceptibility of *Erwinia amylovora* (Burrill) Winslow et al. to geraniol and citronellol. *J Appl Bacteriol.* 1991;71:113–118.
37. Si W, Gong J, Tsao R, et al. Antimicrobial activity of essential oils and structurally related synthetic food additives towards selected pathogenic and beneficial gut bacteria. *J Appl Microbiol.* 2006;100:296–305.
38. Togashi N, Inoue Y, Hamashima H, Takano A. Effects of two terpene alcohols on the antibacterial activity and the mode of action of farnesol against *Staphylococcus aureus*. *Molecules.* 2008;13:3069–3076.
39. Maruyama N, Takizawa T, Ishibashi H, et al. Protective activity of geranium oil and its component, geraniol, in combination with vaginal washing against vaginal candidiasis in mice. *Biol Pharm Bull.* 2008;31:1501–1506.
40. Suppakul P, Miltz J, Sonneveld K, Bigger SW. Antimicrobial properties of basil and its possible application in food packaging. *J Agric Food Chem.* 2003;51:3197–3207.
41. Van Zyl RL, Seatholo ST, Van Vuuren SF, Viljoen AM. The biological activities of 20 nature identical essential oil constituents. *J Essent Oil Res.* 2006;18:129–133.
42. Viollon C, Chaumont J-P. Antifungal properties of essential oils and their main components upon *Cryptococcus neoformans*. *Mycopathologia.* 1994;128:151–153.
43. Zhang Z-Z, Li Y-B, Qi L, Wan X-C. Antifungal activities of major tea leaf volatile constituents toward *Colletorichum camelliae* Masea. *J Agric Food Chem.* 2006;54:3936–3940.
44. Zu Y, Yu H, Liang L, et al. Activities of ten essential oils towards *Propionibacterium acnes* and PC-3, A-549 and MCF-7 cancer cells. *Molecules.* 2010;15:3200–3210.
45. Zamiri-Akhlagh A, Rakhshandeh H, Tayarani-Najaran Z, Mousavi SH. Study of cytotoxic properties of *Rosa damascena* extract in human cervix carcinoma cell line. *Avicenna J Phytomed.* 2011;1:74–77.
46. Venkatesan B, Subramanian V, Tumala A, Vellaichamy E. Rapid synthesis of biocompatible silver nanoparticles using aqueous extract of *Rosa damascena* petals and evaluation of their anticancer activity. *Asian Pac J Trop Med.* 2014;7(suppl 1(0)):S294–S300.
47. Burke YD, Stark MJ, Roach SL, Sen SE, Crowell PL. Inhibition of pancreatic cancer growth by the dietary isoprenoids farnesol and geraniol. *Lipids.* 1997;32:151–156.
48. Wiseman DA, Werner SR, Crowell PL. Cell cycle arrest by the isoprenoids perillyl alcohol, geraniol, and farnesol is mediated by p21Cip1 and p27Kip1 in human pancreatic adenocarcinoma cells. *J Pharmacol Exp Ther.* 2007;320:1163–1170.
49. Elson CE. Suppression of mevalonate pathway activities by dietary isoprenoids: protective roles in cancer and cardiovascular disease. *J Nutr.* 1995;125(6 suppl):1666S–1672S.
50. Carnesecci S, Schneider Y, Ceraline J, et al. Geraniol, a component of plant essential oils, inhibits growth and polyamine biosynthesis in human colon cancer cells. *J Pharmacol Exp Ther.* 2001;298:197–200.
51. Boskabady M, Kiani S, Rakhshandeh H. Relaxant effects of *Rosa damascena* on guinea pig tracheal chains and its possible mechanism (s). *J Ethnopharmacol.* 2006;106:377–382.
52. Shafei MN, Rakhshandeh H, Boskabady MH. Antitussive effect of *Rosa damascena* in guinea pigs. *Iran J Pharm Res.* 2010;2:231–234.
53. Rakhshandeh H, Boskabady MH, Mossavi Z, Gholami M, Saberi Z. The differences in the relaxant effects of different fractions of *Rosa damascena* on guinea pig tracheal smooth muscle. *Iran J Basic Med Sci.* 2010;13:126–132.
54. Dolati K, Rakhshandeh H, Shafei MN. Antidepressant-like effect of aqueous extract from *Rosa damascena* in mice. *Avicenna J Phytomed.* 2011;1:91–97.
55. Naziroğlu M, Kozlu S, Yorgancigil E, Uğuz AC, Karakuş K. Rose oil (from *Rosa damascena* Mill.) vapor attenuates depression-induced oxidative toxicity in rat brain. *J Nat Med.* 2013;67:152–158.
56. Momeni T, Shahrokhi N. *Essential Oils and Their Therapeutic Actions*. Tehran, Iran: Tehran Univ. Press; 1991. in Persian.
57. Farnia V, Shirzadifar M, Shakeri J, et al. *Rosa damascena* oil improves SSRI-induced sexual dysfunction in male patients suffering from major depressive disorders: results from a double-blind, randomized, and placebo-controlled clinical trial. *Neuropsychiatr Dis Treat.* 2015;11:625–635.
58. Vinokur Y, Rodov V, Reznick N, et al. Rose petal Tea as an antioxidant rich beverage: cultivar effects. *J Food Sci.* 2006;71:S42–S47.
59. Cho E, Yokozawa T, Rhyu D, Kim S, Shibahara N, Park J. Study on the inhibitory effects of Korean medicinal plants and their main compounds on the 1, 1-diphenyl-2-picrylhydrazyl radical. *Phytomedicine.* 2003;10:544–551.
60. Köse E, Sarıalmaz M, Taş U, et al. Rose oil inhalation protects against formaldehyde induced testicular damage in rats. *Andrologia.* 2012;44(s1):342–348.
61. Wei A, Shibamoto T. Antioxidant activities and volatile constituents of various essential oils. *J Agric Food Chem.* 2007;55:1737–1742.
62. Yassa N, Masoomi F, Rankouhi SR, Hadjiakhoondi A. Chemical composition and antioxidant activity of the extract and essential oil of *Rosa damascena* from Iran, population of Guilan. *Daru.* 2009;17:175–180.
63. Baydar NG, Baydar H. Phenolic compounds, antiradical activity and antioxidant capacity of oil-bearing rose (*Rosa damascena* Mill.) extracts. *Ind Crops Prod.* 2013;41:375–380.
64. Vanderjagt T, Ghattas R, Vanderjagt D, Crossey M, Glew R. Comparison of the total antioxidant content of 30 widely used medicinal plants of New Mexico. *Life Sci.* 2002;70:1035–1040.
65. Halvorsen BL, Holte K, Myhrstad MC, et al. A systematic screening of total antioxidants in dietary plants. *J Nutr.* 2002;132:461–471.
66. Hajshamsi V, Ghannadi A, Hajiloo M. Analgesic and anti-inflammatory effects of *Rosa damascena* hydroalcoholic extract and its essential oil in animal models. *Iran J Pharm Res.* 2010;9:163.
67. Rakhshandeh H, Vahdati-Mashhadian N, Dolati K, Hosseini M. Antinociceptive effect of *Rosa damascena* in Mice. *J Biol Sci.* 2008;8:176–180.
68. Kirkeskov B, Christensen R, Bügel S, et al. The effects of rose hip (*Rosa canina*) on plasma antioxidative activity and C-reactive protein in patients with rheumatoid arthritis and normal controls: a prospective cohort study. *Phytomedicine.* 2011;18:953–958.
69. Willich S, Rossnagel K, Roll S, et al. Rose hip herbal remedy in patients with rheumatoid arthritis- a randomised controlled trial. *Phytomedicine.* 2010;17:87–93.
70. Winther K, Apel K, Thamsborg G. A powder made from seeds and shells of a rose-hip subspecies (*Rosa canina*) reduces symptoms of knee and hip osteoarthritis: a randomized, double-blind, placebo-controlled clinical trial. *Scand J Rheumatol.* 2005;34:302–308.
71. Larsen E, Kharazmi A, Christensen LP, Christensen SB. An antiinflammatory galactolipid from Rose Hip (*Rosa canina*) that inhibits chemotaxis of human peripheral blood neutrophils in vitro. *J Nat Prod.* 2003;66:994–995.
72. Rakhshandeh H, Hosseini M. Potentiation of pentobarbital hypnosis by *Rosa damascena* in mice. *Indian J Exp Biol.* 2006;44:910.
73. Rakhshandeh H, Hosseini M, Dolati K. Hypnotic effect of *Rosa damascena* in mice. *Iran J Pharm Res.* 2010;181–185.
74. Rakhshandeh H, Shakeri MT, Ghasemzadeh MR. Comparative hypnotic effect of *Rosa damascena* fractions and diazepam in Mice. *Iran J Pharm Res.* 2010:193–197.
75. Awale S, Tohda C, Tezuka Y, Miyazaki M, Kadota S. Protective effects of *Rosa damascena* and its active constituent on A β (25–35)-induced neuritic atrophy. *Evid-Based Complement Altern Med.* 2011;2011:1–8.

76. Efsandiary E, Karimpour M, Mardani M, et al. Novel effects of *Rosa damascena* extract on memory and neurogenesis in a rat model of Alzheimer's disease. *J Neurosci Res*. 2014;92:517–530.
77. Jazayeri SB, Amanlou A, Ghanadian N, Pasalar P, Amanlou M. A preliminary investigation of anticholinesterase activity of some Iranian medicinal plants commonly used in traditional medicine. *Daru*. 2014;22:17.
78. Mohammadpour T, Hosseini M, Naderi A, et al. Protection against brain tissues oxidative damage as a possible mechanism for the beneficial effects of *Rosa damascena* hydroalcoholic extract on scopolamine induced memory impairment in rats. *Nutr Neurosci*. 2014;18:329–336.
79. Ashrafzadeh F, Rakhshandeh H, Mahmodi E. *Rosa damascena* oil: an adjunctive therapy for pediatric refractory seizures. *Iran J Child Neurol*. 2007;1:13–17.
80. Hosseini M, Ghasemzadeh RM, Sadeghnia H, Rakhshandeh H. Effects of different extracts of *Rosa damascena* on pentylenetetrazol-induced seizures in mice. *J Chin Integr Med*. 2011;9:1118–1124.
81. Kheirabadi M, Moghimi A, Rakhshandeh H, Rassouli MB. Evaluation of the anticonvulsant activities of *Rosa damascena* on the PTZ induced seizures in wistar rats. *J Biol Sci*. 2008;8:426–430.
82. Sedighi M, Rafeian-Kopaei M, Noori-Ahmadabadi M, Godarzi I, Baradaran A. In vitro impact of hydro-alcoholic extract of *Rosa damascena* Mill. On rat ileum contractions and the mechanisms involved. *Int J Prev Med*. 2014;5:767–775.
83. Boskabady MH, Shafei MN, Saberi Z, Amini S. Pharmacological effects of *Rosa damascena*. *Iran J Basic Med Sci*. 2011;14:295.
84. Kwon E-K, Lee D-Y, Lee H, et al. Flavonoids from the buds of *Rosa damascena* inhibit the activity of 3-hydroxy-3-methylglutaryl-coenzyme a reductase and angiotensin I-converting enzyme. *J Agric Food Chem*. 2009;58:882–886.
85. Bani S, Hasanpour S, Mousavi Z, Mostafa Garehbaghi P, Gojazadeh M. The effect of *Rosa damascena* extract on primary dysmenorrhea: a double-blind cross-over clinical trial. *Iran Red Crescent Med J*. 2014;16:e14643.
86. Konaté K, Yomalan K, Sytar O, et al. Free radicals scavenging capacity, antidiabetic and antihypertensive activities of flavonoid-rich fractions from leaves of *Trichilia emetica* and *Opilia amentacea* in an animal model of type 2 diabetes mellitus. *J Evid Based Complement Altern Med*. 2014;2014:13.
87. Gholamhoseinian A, Shahouzehi B, Joukar S, Iranpoor M. Effect of *Quercus infectoria* and *Rosa damascena* on lipid profile and atherosclerotic plaque formation in rabbit model of hyperlipidemia. *Pak J Biol Sci*. 2012;15:27–33.
88. Gholamhoseinian A, Shahouzehi B, Shariffar F. Inhibitory effect of some plant extracts on pancreatic lipase. *Int J Pharmacol*. 2010;6:18–24.
89. Gholamhoseinian A, Sharifi-Far F, Shahouzehi B. Inhibitory activity of some plant methanol extracts on 3-Hydroxy-3-Methylglutaryl coenzyme a reductase. *Int J Pharmacol*. 2010;6:705–711.
90. Slavov A, Kiyohara H, Yamada H. Immunomodulating pectic polysaccharides from waste rose petals of *Rosa damascena* Mill. *Int J Biol Macromol*. 2013;59:192–200.
91. Lis-Balchin M. *Aromatherapy Science: A Guide for Healthcare Professionals*. Pharmaceutical Press; 2006.
92. Akbari M, Kazerani HR, Kamrani A, Mohri M. A preliminary study on some potential toxic effects of *Rosa damascena* Mill. *Iran J Veter Res*. 2013;14:232–236.
93. Kürkçüoğlu M, Abdel-Megeed A, Başer KHC. The composition of Taif rose oil. *J Essent Oil Res*. 2013;25:364–367.