

JOURNAL OF APPROXIMATION THEORY 2, 384–388 (1969)

Simultaneous Best Approximations with Two Polynomials

DAVID A. SPRECHER

*Department of Mathematics, University of California,
Santa Barbara, California 93106*

1. THE PROBLEM

The following problem was posed by T. J. Rivlin [1]: Let \mathcal{C} stand for the space of continuous real valued functions on the interval $[0, 1]$; let $E_j(f)$ represent the Chebyshev degree of approximation to f by algebraic polynomials of degree j . Characterize the n -tuples $\{p_0, p_1, \dots, p_{n-1}\}$ of algebraic polynomials p_j , where $\deg p_j = j$, which have the property that there is a function $f \in \mathcal{C}$ such that

$$E_j(f) = \|f - p_j\| \quad (1)$$

for $j = 0, 1, 2, \dots, n - 1$, the norm on the right side being the uniform norm.

In this note we prove the following related

THEOREM. *Given polynomials p_m and p_n , $0 \leq m < n$, there is a function $f \in \mathcal{C}$ which satisfies (1) for the integers $j = m$ and $j = n$ if and only if the polynomial $p_n - p_m$ changes sign at least $m + 1$ times in $[0, 1]$.*

The linear case $m = 0, n = 1$ was proved by Deutsch, Morris and Singer in [2]. Our only excuse for reproving this case is that our proof is very short. The necessity of the condition in the theorem was stated and proved by Rivlin [1].

2. PROOF OF THE CASE $m = 0, n = 1$

To verify the sufficiency of the condition let us consider polynomials $p_0(x) = c$ and $p_1(x) = ax + b$ ($a \neq 0$) such that $at + b = c$ for some $t, 0 < t < 1$. Then the equality

$$\frac{1}{2}[p_1(t - \delta) + p_1(t + \delta)] = c$$

holds for each number δ , and we fix a δ such that $0 < \delta < \min \{t, 1 - t\}$. Let us suppose that $a > 0$ and put

$$\alpha = \max \{|a + b - c|, |b - c|\}.$$

If $f \in \mathcal{C}$ is the piecewise linear function with vertices

$$(0, c - \alpha) \quad (t - \delta, c - \alpha) \quad (t + \delta, c + \alpha) \quad (1, p_1(t + \delta)),$$

(see Fig. 1), then (1) is easily seen to hold for $j = 0, 1$. The desired f in case $a < 0$ is obtained from the above construction by the substitution $x = 1 - y$.

The necessity of the asserted sign change is demonstrated as follows: let a function $f \in \mathcal{C}$ and polynomials p_0, p_1 satisfy (1). If we put

$$g_j = f - p_j \quad (j = 0, 1),$$

then it is an elementary observation that

$$\max g_j + \min g_j = 0 \quad (j = 0, 1), \tag{2}$$




FIG. 1.

where the maximum and minimum are taken over $[0, 1]$. If, say, $g_1 > g_2$ on $(0, 1)$, then $\max g_1 > \max g_2$ and $\min g_1 > \min g_2$, so that

$$\max g_1 + \min g_1 > \max g_2 + \min g_2.$$

But this implies that (2) fails to be true for g_1 or g_2 . Clearly, one arrives at the same conclusion if $g_2 > g_1$ on $(0, 1)$ and the necessity of the sign change is hereby proved.

3. PROOF OF THE CASE $m = 0, n > 1$

Let $p_n(x_0) = c$ for some point $0 < x_0 < 1$. Then there are points $0 < x_1 < x_0 < x_2 < 1$ such that

$$\frac{1}{2}[p_n(x_1) + p_n(x_2)] = c;$$

we may assume without loss of generality that $p_n(x_2) > c$. We introduce the polynomials

$$\begin{aligned} p_n^-(x) &= p_n(x) - [p_n(x_1) + \|p_n\| - c], \\ p_n^+(x) &= p_n(x) + [p_n(x_1) + \|p_n\| - c], \end{aligned} \tag{3}$$

satisfying

$$\frac{1}{2}[p_n^-(x) + p_n^+(x)] = p_n(x).$$

Let

$$k = \begin{cases} n+3 & \text{if } n \text{ is odd,} \\ n+4 & \text{if } n \text{ is even,} \end{cases}$$

and fix points

$$x_1 = u_1 < u_2 < u_2 < \dots < u_{k-1} < u_k = x_2$$

(see Fig. 2). The function

$$g(x) = \begin{cases} \frac{u_{2j} - x}{u_{2j} - u_{2j-1}} p_n^-(x) + \frac{x - u_{2j-1}}{u_{2j} - u_{2j-1}} p_n^+(x) \\ \quad \left(u_{2j-1} \leq x \leq u_{2j}, j = 1, 2, \dots, \frac{k}{2} \right) \\ \frac{u_{2j+1} - x}{u_{2j+1} - u_{2j}} p_n^+(x) + \frac{x - u_{2j}}{u_{2j+1} - u_{2j}} p_n^-(x) \\ \quad \left(u_{2j} \leq x \leq u_{2j+1}, j = 1, 2, \dots, \frac{k}{2} - 1 \right) \end{cases} \quad (4)$$

which is defined and continuous on the interval $[x_1, x_2]$, is best approximated in the class of all n -degree algebraic polynomials by $p_n(x)$. This follows from the facts that

$$p_n(u_{2j-1}) - g(u_{2j-1}) = p_n(u_{2j-1}) - p_n^-(u_{2j-1}) = p_n(x_1) + \|p_n\| - c \\ \left(j = 1, 2, \dots, \frac{k}{2} \right),$$

$$p_n(u_{2j}) - g(u_{2j}) = p_n(u_{2j}) - p_n^+(u_{2j}) = -[p_n(x_1) + \|p_n\| - c] \\ \left(j = 1, 2, \dots, \frac{k}{2} - 1 \right),$$

and

$$\|g - p_n\| = p_n(x_1) + \|p_n\| - c,$$

as shown by a simple calculation. We let

$$f(x) = \begin{cases} \frac{x_1 - x}{x_1} p_n(x) + \frac{x}{x_1} p_n^-(x) & (0 \leq x \leq x_1) \\ g(x) & (x_1 \leq x \leq x_2) \\ \frac{1 - x}{1 - x_2} p_n^+(x) + \frac{x - x_2}{1 - x_2} p_n(x) & (x_2 \leq x \leq 1) \end{cases}$$

Then

$$\|f - p_n\| = \|g - p_n\|$$

so that $E_n(f) = \|f - p_n\|$. To see that $E_0(t) = \|f - c\|$ we observe from (3) that

$$f(x_1) = p_n^-(x_1) = c - \|p_n\|,$$

$$f(x_2) = p_n^+(x_2) = c + \|p_n\|,$$

and a routine calculation shows that

$$-\|p_n\| \leq f(x) - c \leq \|p_n\| \quad (0 \leq x \leq 1).$$

Necessity is proved just as in the previous case.




FIG. 2.

4. PROOF OF THE CASE $m > 0, n > 1$

Consider arbitrary polynomials p_m and p_n , with $1 \leq m < n$, such that $p_n - p_m$ changes sign at least $m + 1$ times in $[0, 1]$. Then there are points

$$0 < t_1 < t_2 < \dots < t_{m+1} < 1$$

such that $p_m(t_j) = p_n(t_j)$ for $j = 1, 2, \dots, m + 1$. Letting $t_0 = 0$ and $t_{m+2} = 1$, we put

$$\alpha_j = \max_{t_j \leq x \leq t_{j+1}} |p_m(x) - p_n(x)|,$$

$$\alpha = \frac{1}{2} \min \{\alpha_0, \alpha_1, \dots, \alpha_{m+1}\},$$

$$\beta = \max \{\|p_m\|, \|p_n\|\}.$$

Suppose $p_n(x) \geq p_m(x)$ for $t_1 \leq x \leq t_2$ and consider the polynomials

$$p_n^-(x) = p_n(x) - \beta,$$

$$p_n^+(x) = p_n(x) + \beta,$$

$$p_m^-(x) = p_m(x) - \alpha - \beta,$$

$$p_m^+(x) = p_m(x) + \alpha + \beta.$$

Then each of the polynomials $p_n^- - p_m^-$ and $p_n^+ - p_m^+$ also changes sign at least $m + 1$ times in $[0, 1]$. Thus, there are $m + 2$ points x_j ,

$$x_j < t_j < x_{j+1}, \quad j = 1, 2, \dots, m + 1,$$

such that

$$p_m^-(x_{2k-1}) = p_n^-(x_{2k-1}) \quad (5)$$

and

$$p_m^+(x_{2k}) = p_n^+(x_{2k}) \quad (6)$$

for all admitted values of k . The point x_2 can be so chosen that

$$p_m^-(x) \leq p_n^-(x) < p_n^+(x) \leq p_m^+(x), \quad (x_1 \leq x \leq x_2).$$

On the interval $[x_1, x_2]$ the situation is now similar to that in Case 3. Thus, let g be defined as in (4). Then p_n is its best approximation in the class of algebraic polynomials of degree n .

Put for each $x \in [0, 1]$,

$$q^-(x) = \max\{p_m^-(x), p_n^-(x)\},$$

$$q^+(x) = \min\{p_m^+(x), p_n^+(x)\}.$$

Then

$$q^-(x_{2k-1}) = p_m^-(x_{2k-1}) = p_n^-(x_{2k-1}),$$

$$q^+(x_{2k}) = p_m^+(x_{2k}) = p_n^+(x_{2k})$$

(see (5) and (6)). In a manner similar to that in Case 3 we construct a function $f \in \mathcal{C}$ such that

$$q^-(x) \leq f(x) \leq q^+(x), \quad (7)$$

$$f(x) = g(x) \quad \text{for } x_1 \leq x \leq x_2, \quad (8)$$

$$f(x_{2k-1}) = q^-(x_{2k-1}),$$

$$f(x_{2k}) = q^+(x_{2k}). \quad (9)$$

The conditions (7) and (8) guarantee that p_n is also the best approximation of f by algebraic polynomials of degree n . Conditions (7) and (9) show that the best approximation to f from among the algebraic polynomials of degree m is p_m : the calculations justifying this conclusion are routine and therefore omitted here.

For the necessity of the asserted sign changes we refer this time to Rivlin's proof in [1].

REFERENCES

1. T. J. RIVLIN in Proc. Coll. on Abstract Spaces and Approximation (Oberwolfach, July 1968), Birkhäuser Verlag (to appear.)
2. F. DEUTSCH, P. D. MORRIS AND I. SINGER, On a problem of T. J. Rivlin in approximation theory. *J. Approx. Theory* (to appear).