



Journal of Traditional and Complementary Medicine

Journal homepage <http://www.jtcm.org>

Adlay (薏苡 yì yǐ; "soft-shelled job's tears"; the seeds of *Coix lachryma-jobi* L. var. *ma-yuen* Stapf) is a Potential Cancer Chemopreventive Agent toward Multistage Carcinogenesis Processes

Ching-Chuan Kuo^{1,2,*}, Huang-Hui Chen¹, Wenchang Chiang^{3,*}

¹ National Institute of Cancer Research, National Health Research Institutes, Tainan;

² Institute of Clinical Pharmacy and Pharmaceutical Sciences, National Cheng Kung University Medical College, Tainan;

³ Graduate Institute of Food Science and Technology, National Taiwan University, Taipei, Taiwan, R.O.C.

ABSTRACT

Adlay (薏苡 yì yǐ; "soft-shelled job's tears", the seeds of *Coix lachryma-jobi* L. var. *ma-yuen* Stapf) is a grass crop that has long been used in traditional Chinese medicine (TCM) and as a nourishing food in China for the treatment of warts, chapped skin, rheumatism, neuralgia, inflammatory, and neoplastic diseases. In addition, adlay also has been said to have stomachic, diuretic, antiphlogistic, anodynic, and antispasmodic effects. Carcinogenesis is a multistage process that begins with exposure of viruses or chemicals that are found in the environment. Chemoprevention refers to the use of natural or synthetic, non-toxic chemical substances to reverse, repress, or prevent carcinogenesis. In this review, we summarize recent research attempting to study the chemopreventive blocking and suppressing potential of adlay and its active components in scavenging electrophiles and reactive oxygen species, antimutagenicity, enhancing Nrf2-mediated detoxification and antioxidant effect, altering carcinogen metabolism, suppressing proliferation, decreasing inflammation, and enhancing antitumor immunity. In addition, several active components with diverse chemopreventive properties have been also mentioned in this review article.

Keywords: Adlay, Traditional Chinese medicine, Cancer chemoprevention, Blocking agent, Suppressing agent

Adlay (*Coix lachryma-jobi* L. var. *ma-yuen* Stapf.)

Adlay (薏苡 yì yǐ; the seeds of *Coix lachryma-jobi* L. var. *ma-yuen* Stapf.), also named coix seeds, Chinese pearl barley, pearl barley, semen coicis, yokuinin, 薏苡仁 (yì yǐ rén), and 薏米 (yì mǐ), belongs to the family Gramineae. It is an annual or perennial herb, 100 ~ 180 cm high, flowering from July to September and fruiting

from September to October. The adlay seed consists of four parts from outside to inside including the hull, testa, bran, and endosperm (polished adlay) (Figure 1). Adlay is widely planted in Taiwan, China, and Japan, where it is considered a healthy food supplement.

According to the first pharmaceutical monograph in ancient China, *The Divine Husbandman's Herbal Foundation Canon* (神農本草經 shén nóng běn cǎo jīng), adlay is considered as a top grade (上品 shàng

*Correspondence to:

Ching-Chuan Kuo, Ph.D., National Institute of Cancer Research, National Health Research Institutes, No. 367, Shengli Road, Tainan, Taiwan, R.O.C.
Phone: +886-6-7000123 ext. 65115; Fax: +886-6-2083427; Email: cckuo@nhri.org.tw

Wenchang Chiang, Ph.D., Graduate Institute of Food Science and Technology, National Taiwan University, No. 1, Sec. 4, Roosevelt Road, Taipei, Taiwan, R.O.C. Phone: +886-2-33664115; Fax: +886-2-23638673; Email: chiang@ntu.edu.tw



Figure 1. The photograph of adlay plant, adlay seed and its fractions, including adlay hull, adlay testa, dehulled adlay, adlay bran, and polished adlay.

pǐn) of Traditional Chinese Medicine (TCM). People received adlay for a long period can nourish life (養命 yǎng mìng), boost qi (益氣 yì qì), rejuvenate the body (輕身 qīng shēn), and extend life (增年 zēng nán). Since adlay can make body healthy and prolong life with non-toxic properties, it has been thought to be the longevity of monarch herb (不老長生之君藥), and generally form the basis of dietary functional foods. Adlay traditionally has long been used in China for the treatment of warts, chapped skin, rheumatism, neuralgia, and inflammatory diseases. In addition, adlay also has been said to have stomachic, diuretic, antiphlogistic, anodynic, antispasmodic, and antitumor effects.

Although medicinal functionality of adlay has been around for thousands of years, recently, it is interesting to note that integration exists between traditional medicine and modern medicine because a number of biological activities of adlay have been elucidated by scientific investigation, including antioxidant/free radical scavenging (Kuo *et al.*, 2001; Kuo *et al.*, 2002; Yao *et al.*, 2011; Yu *et al.*, 2011; Chen *et al.*, 2012a; Wang *et al.*, 2012), anti-inflammatory (Otsuka *et al.*, 1988; Huang *et al.*, 2009a; Huang *et al.*, 2009b), anti-mutagenic (Huang and Chiang, 1999; Chen *et al.*, 2011), anti-tumor (Tanimura, 1961; Numata *et al.*, 1994; Kuo *et al.*, 2001; Chang *et al.*, 2003; Hung and Chang, 2003; Shih *et al.*, 2004; Lee *et al.*, 2008; Chung *et al.*, 2010; Chung *et al.*, 2011b; Li *et al.*, 2011), anti-allergic (Hsu

et al., 2003; Chen *et al.*, 2010; Chen *et al.*, 2012b; Chen *et al.*, 2012c), hypolipidemic (Kim *et al.*, 2004; Yu *et al.*, 2004; Huang *et al.*, 2005; Yeh *et al.*, 2006; Son *et al.*, 2008; Yu *et al.*, 2011), hypocholesterolemic (Wang *et al.*, 2012), hypoglycemic (Takahashi *et al.*, 1986; Huang *et al.*, 2005; Yeh *et al.*, 2006; Lin *et al.*, 2010), anti-obesity (Kim *et al.*, 2004; Huang *et al.*, 2005; Kim *et al.*, 2007), anti-ulcer (Chung *et al.*, 2011a), prebiotic activity (Chiang *et al.*, 2000), abortifacient (Tzeng *et al.*, 2005), hormonal modulating (Kondo *et al.*, 1988; Chang *et al.*, 2006; Hsia *et al.*, 2006; Hsia *et al.*, 2007; Hsia *et al.*, 2008; Hsia *et al.*, 2009), osteoporosis preventing (Yang *et al.*, 2008), and antimicrobial (Ishiguro *et al.*, 1993) effect. Since cancer mortality and incidence worldwide has increased rapidly in recent decades, thus, this review article focused on chemopreventive role of adlay in blocking carcinogenesis progress, preventing cancer formation and progression.

Multidrug Carcinogenesis

Carcinogenesis is a multistage process that begins with exposure of viruses or chemicals that are found in the environment. This process involves various molecular and cellular events that transform a normal cell to a malignant neoplastic cell. It can be conceptually identified as tumor initiation, tumor promotion, and tumor progression (Figure 2) (Panayiotidis,




Figure 2. Multistage carcinogenesis and strategies for cancer chemoprevention.

2008). Tumor initiation is a step where the normal cell undergoes irreversible genetic and epigenetic alterations to generate an initiated cell (Lengauer *et al.*, 1998). Oxidative stress plays a major role in the initiation of carcinogenesis (Khansari *et al.*, 2009). Oxidative stress can be induced through a number of endogenous sources, such as the production of free radical by mitochondrial oxidative phosphorylation, and exogenous sources, including ionizing radiations, ultraviolet exposure and environmental carcinogens. In fact, chemical carcinogens cannot damage DNA until they are metabolized by cytochrome P-450 in cells and converted to reactive electrophiles. Reactive oxygen species (ROS) can react with DNA as well as chromatin proteins, resulting in several types of DNA damage, including depurination, depyrimidination, single- and double-strand breaks, base modifications, and DNA-protein cross linkages (Barzilai and Yamamoto, 2004; Fruehauf and Meyskens, 2007). In addition, carcinogen-DNA adduct formation is an important molecular mechanism that contributes to chemical carcinogenesis (Windmill *et al.*, 1997). ROS-mediated DNA damage contributes to the initiation of carcinogenesis through increasing a cell's mutation rate, influencing the activation of protooncogene and tumor-suppressor gene,

and promoting the oncogenic phenotype (Campos *et al.*, 2007). Tumor promotion involves the clonal expansion of initiated cells to form an actively premalignant, proliferating lesion. Direct DNA interaction and damage is not involved in this process. Tumor promoters generally alter gene expression resulting in decrease of apoptosis cell death and increase of cell proliferation (Klaunig and Kamendulis, 2004). Tumor progression involved in the expression of the malignant, aggressive, invasive, and metastatic phenotype. During this process, further genetic and epigenetic alterations may occur resulting in the formation of neoplasms.

Cancer Chemoprevention

The prominent and effective strategy to inhibit cancer formation is elimination, reverse or delay of genetic and epigenetic alterations induced by chemical carcinogens (Hail *et al.*, 2008). Chemoprevention refers to the use of natural or synthetic, non-toxic chemical substances to reverse, repress, or prevent carcinogenesis. These chemopreventive agents can be broadly categorized into two types: blocking agents and suppressing agents (Figure 2) (Wattenberg, 1985; Manson *et al.*, 2000). Blocking agents generally inhibit the initiation of carcinogenesis. Blocking agents

alter carcinogen metabolism and enhance carcinogen detoxification, resulting in elimination of carcinogens reaching their target site to induce DNA damage (Surh, 2003). In addition, blocking agents effectively scavenge electrophiles and ROS to prevent DNA alterations. Furthermore, blocking agents induce DNA repair machinery to reduce DNA damage as well as genetic mutations. Suppressing agents inhibit pre-malignant or malignant conversion of initiated cells during the stages of tumor promotion and progression. They efficiently modulate gene expression, including oncogenes and tumor-suppressor genes, resulting in suppressing cell proliferation, encouraging apoptosis, and inhibiting angiogenesis (Hayes and McMahon, 2001).

Adlay Acts as a Blocking Agent for Cancer Chemoprevention

Scavenge Electrophiles and Reactive Oxygen Species

Reactive oxygen species (ROS) are induced through a variety of endogenous and exogenous sources. Overwhelming of antioxidant and DNA repair mechanisms in a cell by ROS may result in oxidative stress and oxidative damage to a cell. Oxidative damage resulting from ROS generation can participate in all stages of the carcinogenesis process (Klaunig *et al.*, 2011). Therefore, elimination of ROS or electrophiles by chemopreventive blocking agents is an important strategy to prevent the malignant development. Kuo *et al.* have demonstrated that the methanol extract of adlay hull (AHM) is a highly effective ROS scavenger that inhibits free radical-generating enzymes, blocks tumor promoter-generated oxidative processes in neutrophil-like leukocytes, and exhibits a cytoprotective effect on cultured cells exposed to tert-butyl hydroperoxide (Kuo *et al.*, 2001). Furthermore, active component that represent strong antioxidant activity were isolated from the methanol extract of adlay hull to be coniferyl alcohol, syringic acid, ferulic acid, syringaresinol, 4-ketopinonesinol, and mayuenolide (Kuo *et al.*, 2002).

Antimutagenicity

It is well-documented that cancer begins after a mutational episode in a single cell and then it progressively transforms to malignancy in multiple stages through sequential acquisition of additional mutations (Khan and Pelengaris, 2006). The Ames Salmonella typhimurium mutagenicity assay (Ames test) is a short-term bacterial reverse mutation assay specifically designed to detect a wide range of

chemical substances that can produce genetic damage leading to gene mutations (Mortelmans and Zeiger, 2000). Several lines of evidence showed that the compound prevents mutagenesis caused by known carcinogens in the Ames test indicates a possible role in chemoprevention (Weisburger, 2001). Notably, Huang and Chiang previously investigated the antimutagenic activity of different portions of adlay, including adlay hull, adlay testa, adlay bran, dehulled adlay, and polished adlay. Among them, the acetone extract of adlay hull and adlay bran possesses most potent antimutagenic activity against Benzo(a)pyrene (B[a]P), 2-amino-3-methylimidazo[4,5-f]quinoline (IQ), and 4-nitroquinoline -N-oxide (NQNO) induced mutagenesis in Salmonella typhimurium TA98 (Huang and Chiang, 1999). Recently, Chen *et al.* further identified the antimutagenic constituents from adlay hull by used Ames antimutagenic activity-guide isolation procedures, and found that p-hydroxybenzaldehyde, vanillin, syringaldehyde, trans-coniferylaldehyde, sinapaldehyde, and coixol exert great antimutagenic activity in suppressing IQ-induced mutagenicity in Salmonella typhimurium TA98 (Chen *et al.*, 2011). In addition to antimutagenic activity, trans-coniferylaldehyde and sinapaldehyde also exhibit potent effect in scavenging of 2,2'-diphenyl-1-picrylhydrazyl (DPPH) radicals and inhibiting 12-O-tetradecanoylphorbol-13-acetate (TPA) stimulated superoxide anion generation in neutrophil-like leukocytes (Chen *et al.*, 2011).

Enhance Nrf2-mediated Detoxification and Antioxidant Effect

Recently, a potential chemopreventive strategy involving the induction of a battery of cytoprotective genes has represented a recent focus of attention in cancer research. NF-E2-related factor 2 (Nrf2), a member of the Cap'n'Collar (CNC) family of transcription factors that share a highly conserved basic region-leucine zipper (bZIP) structure mainly regulates transcriptional activation through antioxidant responsive element (ARE) (Huang *et al.*, 2002). Nrf2-mediated cytoprotective genes are thought to protect cells against carcinogenesis and attenuate cancer development via neutralization of ROS or carcinogens (Klaunig *et al.*, 2010; Kwak and Kensler, 2010). The gene families regulated by the Nrf2/ARE pathway include phase II metabolizing/detoxifying enzymes [e.g. aldo keto reductase (AKR), glutathione S-transferases (GST), UDP-glucuronoyltransferase (UGT), NADP(H) quinone oxidoreductase 1 (NQO1), heme oxygenase-

1(HO-1)], and antioxidants [e.g. superoxide dismutase (SOD), catalase, ferritin, s-glutamyl cysteine synthetase modifier subunit (GCLM), γ-glutamyl cysteine synthetase catalytic subunit (GCLC), glutathione peroxidase (GPx), glutathione reductase (GR), thioredoxin, thioredoxin reductase (TR), peroxiredoxins, metallothionein, *etc.*] (Zhang and Gordon, 2004; Giudice and Montella, 2006). Therefore, one of the most prominent strategies of cancer chemoprevention is to protect cells or tissues against various carcinogens and carcinogenic metabolites, both exogenous and endogenous, through the induction of metabolizing/detoxifying and antioxidant enzymes. We recently found that 4-ketopinoresinol, trans-coniferylaldehyde and sinapaldehyde, the active component isolated by free radical scavenge or antimutagenicity, significantly induced Nrf2/ARE-driven luciferase activity (Chen *et al.*, 2011; Chen *et al.*, 2012a). Further study demonstrated that 4-ketopinoresinol strongly induces Nrf2-mediated detoxifying/antioxidant proteins, including HO-1, AKR1C1, AKR1C2, AKR1C3, GR, TR, GCLC, GCLM, *etc.*, via the PI3K/AKT pathway, and at least in part, 4-ketopinoresinol -induced HO-1 expression contributes to a cellular defense mechanism against hydrogen peroxide-induced cell injury (Chen *et al.*, 2012a). These results suggested that the cytoprotective effects induced by 4-ketopinoresinol may come from direct scavenging ROS directly (Kuo *et al.*, 2002) and activating of PI3K/AKT/Nrf2/HO-1 axis (Chen *et al.*, 2012a). Future studies evaluating the efficacy of 4-ketopinoresinol as a potential chemopreventive agent, using different carcinogenesis animal models, are warranted.

Alter Carcinogen Metabolism

Mammalian cytochrome P-450 (CYP) enzymes are phase I metabolizing enzymes, generally catalyze the oxidative metabolism of various xenobiotics, including drug, chemical carcinogens, *etc.* (Rendic, 2002). Despite the phase I oxidative metabolic reactions of chemicals result in the formation of more-water-soluble and less-toxic metabolites, however, some CYP enzymes, such as CYP1A1, CYP3A, and CYP2E1, have been indicated to involve in the metabolic activation of carcinogens, such as benzo[a]pyrene, N-nitrosodimethylamine, or aflatoxin B1 (Gonzalez and Gelboin, 1994). In addition, CYP1A2 have been addressed to activate a tobacco pre-carcinogen to a carcinogen [*i.e.* 4-methylnitrosamino-1-(3-pyridyl)-1-butanone; NNK] (Smith *et al.*, 1996). Yao *et al.* demonstrated that the ethanolic extract of

adlay bran (ABE) significantly suppresses CYP1A1 and CYP1A2 activities in the liver and CYP1A1 activity in the lungs of rats after 4 weeks of feeding (Yao *et al.*, 2011). This change may have resulted in a decrease in the metabolism of chemical carcinogens. Indeed, a recent study reported that rats fed the ethyl acetate fraction of ABE showed reduced 1,2-dimethylhydrazine (DMH)-induced colon carcinogenesis (Chung *et al.*, 2010). Because DMH is a pre-carcinogen that requires activation by liver CYP to form DNA-reactive metabolites, therefore, ABE may have chemopreventive effect against colon carcinogenesis in the initiation stage.

Adlay Acts as a Suppressing Agents for Cancer Chemoprevention

Suppress Proliferation

Dysregulated proliferation is a well-established hallmark of carcinogenesis (Hanahan and Weinberg, 2000). The earliest antitumor study of adlay was demonstrated by Tanimura's group at 1961. They reported that the growth of Ehrlich ascites sarcoma was inhibited by adlay (Tanimura, 1961; Ukita and Tanimura, 1961). Adlay is also applied as an adjuvant to treat lung, stomach, breast, colon, and cervical cancers (Chang *et al.*, 2003; Hung and Chang, 2003; Woo *et al.*, 2007; Hu *et al.*, 2009). In addition to tumor transplantation experiment, Chang *et al.* used NNK-induced lung tumorigenesis model to investigate the chemopreventive potential of adlay seed. As the result, mice received the powder of adlay seed significantly reduced the number of surface lung tumors (Chang *et al.*, 2003).

The active compound with anticancer property was first identified by Tanimura's group to be a oil component, coixenolide (Tanimura, 1961; Ukita and Tanimura, 1961). Therefore, it is important to know the distribution of this compound in adlay seed. Huang and Chiang demonstrated that among different part of adlay seed, adlay bran contained the highest amount of coixenolide (473 ppm) (Huang and Chiang, 1999). In addition to coixenolide, other oil components, including palmitic acid, stearic acid, oleic acid, and linoleic acid, also showed potent growth inhibitory activity on a transplantable mouse tumor (Numata *et al.*, 1994). These results suggested that oil components of adlay play a critical role of anticancer efficacy. Recently, Kanglaite injection (KLT), an aqueous microemulsion of an oil extracted from adlay seed, have been developed using

the latest and most complex modern technologies (Lu *et al.*, 2008). KLT is a new biphasic extended-spectrum anticancer medicine, the food and drug administration (FDA) of United States also approved a phase II trial of KLT to test its efficacy in treating patients with stage IV of non-small-cell lung cancer or combination of KLT with gemcitabine to investigate the therapeutic efficacy in pancreatic cancer patients (<http://www.clinicaltrials.gov>). The mechanisms contributed to antitumor effect of neutral lipid components, at least in part, through cell cycle arrest and apoptosis induction involving upregulation of cyclin-dependent kinase inhibitor p21 and inhibition of antiapoptotic protein Bcl-2 expression (Bao *et al.*, 2005).

In addition to oil extract, Chiang's group recently identified several active components from adlay bran with anticancer growth inhibitory ability, including: (i) caffeic acid and chlorogenic acid toward gastric cancer cell growth (Chung *et al.*, 2011a); (ii) coixspirolactam D, coixspirolactam E, coixspiroenone, coixspirolactam A, coixspirolactam C, and coixlactam toward breast cancer cells growth (Chung *et al.*, 2011b); and (iii) coixspirolactam A, coixspirolactam B, coixspirolactam C, coixlactam, and methyl dioxindole-3-acetate toward non-small cell lung and colorectal cancer cell growth (Lee *et al.*, 2008). The mechanism underlying anticancer effect of these component merit for further investigation.

Decrease Inflammation

A causal association between inflammation and cancer has long been suspected. Multiple lines of compelling evidence from clinical, epidemiologic and laboratory studies support that inflammation plays a critical role in the promotion and progression stages of carcinogenesis (Fitzpatrick, 2001).

1. The Nuclear Factor κ B (NF- κ B)

NF- κ B comprises a family of transcription factors involved in the regulation of a wide variety of biological responses. NF- κ B plays a well-known function in the regulation of immune responses and inflammation, but growing evidences support a major role in oncogenesis. NF- κ B regulates the expression of genes involved in many processes that play a key role in the development and progression of cancer such as proliferation, migration and apoptosis (Wang and Cho, 2010). Thus, clinical trials with drugs that block NF- κ B are currently in progress with promising result (<http://www.clinicaltrials.gov>). Woo *et al.* showed that anti-neoplastic activity of an adlay extract in breast MDA-MB-231 xenografts (Woo *et al.*, 2007). Using gene array

technology, adlay extract significantly inhibits NF- κ B-dependent gene transcription, including downregulation of COX-2 and matrix metalloproteinases (MMP), in these cells. In addition, adlay extract also inhibits activity of protein kinase C, a major mediator of signal transduction and activator of NF- κ B (Woo *et al.*, 2007). Likewise, treatment with KLT decreased the level of NF- κ B in the nucleus in a dose-dependent manner, and KLT markedly decreased the expression of I κ B α , IKK and EGFR in the cytoplasm of lung tumor cells (Pan *et al.*, 2012).

2. Cyclooxygenase-2 (COX-2)

Aberrant upregulation of COX-2, a key player in inflammatory signaling, is frequently observed in various precancerous and malignant tissues. Therefore, the normalization of inappropriately overamplified signaling cascades implicated in chronic inflammation-associated carcinogenesis by use of COX-2 specific inhibitors has been recognized as a rational and pragmatic strategy in molecular target-based cancer prevention (Surh and Kundu, 2007). Huang *et al.* demonstrated that the ethanolic extract of adlay hull (AHE) and adlay testa (ATE) possessed potent anti-inflammatory activity through inhibiting lipopolysaccharide (LPS)-induced COX-2 expression in RAW264.7 macrophage (Huang *et al.*, 2009a; Huang *et al.*, 2009b). The active COX-2 inhibiting components were identified to be ceramide and naringenin in AHE (Huang *et al.*, 2009a), and gallic acid, caffeic acid, and ferulic acid in ATE (Huang *et al.*, 2009b), respectively. In the in vivo investigation, rats received dehulled adlay significantly reduced the expression level of COX-2 and the number of preneoplastic aberrant crypt foci (ACF) in azoxymethane (AOM)-induced colon carcinogenesis animal model (Shih *et al.*, 2004). Li *et al.* further showed that adlay bran and its ethanolic extract and residue play an important role in suppressing ACF formation in an early stage of colon carcinogenesis (Li *et al.*, 2011). In addition, Chung *et al.* demonstrated that the ethyl acetate fraction of ABE significantly suppresses DMH-induced preneoplastic lesion of the colon in F344 rats through an anti-inflammatory pathway, involved COX-2 inhibition (Chung *et al.*, 2010). Moreover, Hung and Chang also suggested that the methanolic extract of adlay seed also inhibits cancer growth and prevents lung tumorigenesis through COX-2 suppression (Hung and Chang, 2003).

Enhance Antitumor Immunity

Cytotoxic T lymphocytes and natural killer cells are

essential effectors of anti-tumor immune responses *in vivo* (Seino *et al.*, 2006). Miyai's group have investigated the changes in lymphocyte subsets in seven volunteers before, during (four weeks) and after taking six adlay seed tablets. As the results, the level of CD3+CD56+ (MHC-non restricted cytotoxic T cells) markedly increased at four weeks. The level of CD16+CD57- (the mature, most active natural killer cells) also increased at three weeks (Hidaka *et al.*, 1992; Kaneda *et al.*, 1992). These results indicate that adlay seeds increase peripheral cytotoxic lymphocytes and may be effective not only to viral infection through the enhancement of cytotoxic activity but also to boost antitumor immunity. In addition, it is interesting to note that KLT boosted anticancer immunity in Lewis lung carcinoma-bearing C57BL/6 mice through stimulating T cell proliferation and IL-2 secretion (Pan *et al.*, 2012).

CONCLUSION REMARKS

In summary, adlay is a potential chemopreventive TCM to block multistage carcinogenesis with anti-initiation, -promotion and -progression properties. The chemopreventive functionality of adlay involving in scavenging electrophiles and reactive oxygen species, antimutagenicity, enhancing Nrf2-mediated detoxification and antioxidant effect, altering carcinogen metabolism, suppressing proliferation, decreasing inflammation, and enhancing antitumor immunity. Oil extract, especially KLT, have been proved to test its therapeutic efficacy in clinical setting.

Otherwise, several active components with diverse chemopreventive properties have been identified, including coniferyl alcohol, syringic acid, ferulic acid, syringaresinol, 4-ketopinoresinol, and mayuenolide (scavenge electrophiles and reactive oxygen species); p-hydroxybenzaldehyde, vanillin, syringaldehyde, trans-coniferylaldehyde, sinapaldehyde, and coixol (antimutagenicity); 4-ketopinoresinol, trans-coniferylaldehyde and sinapaldehyde (enhance Nrf2-mediated detoxification and antioxidant effect); coixenolide, Kanglaite injection (KLT; neutral lipid components), caffeic acid, chlorogenic acid, coixspirolactam D, coixspirolactam E, coixspiroenone, coixspirolactam A, coixspirolactam C, coixlactam, coixspirolactam A, coixspirolactam B, coixspirolactam C, coixlactam, and methyl dioxindole-3-acetate (suppressing proliferation); ceramide and naringenin, gallic acid, caffeic acid, and ferulic acid (decrease inflammation); and KLT (enhance antitumor immunity). The underlying mechanism of these active components

is merit for further investigation.

ACKNOWLEDGEMENTS

The study was supported by grants from the National Health Research Institutes (CA-101-PP-05), the National Science Council (NSC98-2320-B-400-003-MY3), and the Department of Health (DOH101-TD-C-111-004), Taiwan, R.O.C.

ABBREVIATIONS

ABE	the ethanolic extract of adlay bran
ACF	aberrant crypt foci
AHE	the ethanolic extract of adlay hull
AHM	the methanol extract of adlay hull
AKR	aldo keto reductase
AOM	azoxymethane
ATE	the ethanolic extract of adlay testa
Ames test	the Ames Salmonella typhimurium mutagenicity assay
ARE	antioxidant responsive element
B[a]P	Benzo(a)pyrene
COX-2	Cyclooxygenase-2
CYP	cytochrome P-450
DMH	1,2-dimethylhydrazine
DPPH radicals	2,2'-diphenyl-1-picrylhydrazyl radicals
FDA	the food and drug administration
GCLC	g-glutamyl cysteine synthetase catalytic subunit
GCLM	g-glutamyl cysteine synthetase modifier subunit
GPx	glutathione peroxidase
GR	glutathione reductase
GST	glutathione S-transferases
HO-1	heme oxygenase-1
IQ	2-amino-3-methylimidazo[4,5-f]quinoline
KLT	Kanglaite injection
LPS	lipopolysaccharide
NNK	4-methylnitrosamino-1-(3-pyridyl)-1-butanone
NQNO	4-nitroquinoline -N-oxide
NQO1	NADP(H) quinone oxidoreductase 1
Nrf2	NF-E2-related factor 2
ROS	reactive oxygen species
SOD	superoxide dismutase
TCM	Vraditional Chinese Medicine
TPA	12-O-tetradecanoylphorbol-13-acetate
TR	thioredoxin reductase
UGT	UDP-glucuronoyltransferase

REFERENCES

- Bao, Y., Yuan, Y., Xia, L., Jiang, H., Wu, W., Zhang, X., 2005. Neutral lipid isolated from endosperm of Job's tears inhibits the growth of pancreatic cancer cells via apoptosis, G2/M arrest, and regulation of gene expression. *J Gastroenterol Hepatol* 20, 1046-1053.
- Barzilai, A., Yamamoto, K., 2004. DNA damage responses to oxidative stress. *DNA repair* 3, 1109-1115.

- Campos, A.C., Molognoni, F., Melo, F.H., Galdieri, L.C., Carneiro, C.R., D'Almeida, V., Correa, M., Jasiulionis, M.G., 2007. Oxidative stress modulates DNA methylation during melanocyte anchorage blockade associated with malignant transformation. *Neoplasia* 9, 1111-1121.
- Chang, H.C., Huang, Y.C., Hung, W.C., 2003. Antiproliferative and chemopreventive effects of adlay seed on lung cancer in vitro and in vivo. *J Agric Food Chem* 51, 3656-3660.
- Chang, L.L., Wun, A.W., Hung, C.T., Hsia, S.M., Chiang, W., Wang, P.S., 2006. Effects of crude adlay hull acetone extract on corticosterone release from rat zona fasciculata-reticularis cells. *Naunyn Schmiedebergs Arch Pharmacol* 374, 141-152.
- Chen, H.H., Chen, Y.T., Huang, Y.W., Tsai, H.J., Kuo, C.C., 2012a. 4-Ketopinosin, a novel naturally occurring ARE activator, induces the Nrf2/HO-1 axis and protects against oxidative stress-induced cell injury via activation of PI3K/AKT signaling. *Free Radic Biol Med* 52, 1054-1066.
- Chen, H.H., Chiang, W., Chang, J.Y., Chien, Y.L., Lee, C.K., Liu, K.J., Cheng, Y.T., Chen, T.F., Kuo, Y.H., Kuo, C.C., 2011. Antimutagenic constituents of adlay (*Coix lachryma-jobi* L. var. *ma-yuen* Stapf) with potential cancer chemopreventive activity. *J Agric Food Chem* 59, 6444-6452.
- Chen, H.J., Hsu, H.Y., Chiang, W., 2012b. Allergic immune-regulatory effects of Adlay Bran on an OVA-immunized mice allergic model. *Food Chem Toxicol*.
- Chen, H.J., Lo, Y.C., Chiang, W., 2012c. Inhibitory effects of adlay bran (*Coix lachryma-jobi* L. var. *ma-yuen* Stapf) on chemical mediator release and cytokine production in rat basophilic leukemia cells. *J Ethnopharmacol* 141, 119-127.
- Chen, H.J., Shih, C.K., Hsu, H.Y., Chiang, W., 2010. Mast cell-dependent allergic responses are inhibited by ethanolic extract of adlay (*Coix lachryma-jobi* L. var. *ma-yuen* Stapf) testa. *J Agric Food Chem* 58, 2596-2601.
- Chiang, W., Cheng, C., Chiang, M., Chung, K.T., 2000. Effects of dehulled adlay on the culture count of some microbiota and their metabolism in the gastrointestinal tract of rats. *J Agric Food Chem* 48, 829-832.
- Chung, C.P., Hsia, S.M., Lee, M.Y., Chen, H.J., Cheng, F., Chan, L.C., Kuo, Y.H., Lin, Y.L., Chiang, W., 2011a. Gastroprotective activities of adlay (*Coix lachryma-jobi* L. var. *ma-yuen* Stapf) on the growth of the stomach cancer AGS cell line and indomethacin-induced gastric ulcers. *J Agric Food Chem* 59, 6025-6033.
- Chung, C.P., Hsu, C.Y., Lin, J.H., Kuo, Y.H., Chiang, W., Lin, Y.L., 2011b. Antiproliferative lactams and spiroenone from adlay bran in human breast cancer cell lines. *J Agric Food Chem* 59, 1185-1194.
- Chung, C.P., Hsu, H.Y., Huang, D.W., Hsu, H.H., Lin, J.T., Shih, C.K., Chiang, W., 2010. Ethyl acetate fraction of adlay bran ethanolic extract inhibits oncogene expression and suppresses DMH-induced preneoplastic lesions of the colon in F344 rats through an anti-inflammatory pathway. *J Agric Food Chem* 58, 7616-7623.
- Fitzpatrick, F.A., 2001. Inflammation, carcinogenesis and cancer. *Int Immunopharmacol* 1, 1651-1667.
- Fruehauf, J.P., Meyskens, F.L., Jr., 2007. Reactive oxygen species: a breath of life or death? *Clinical cancer research : an official journal of the American Association for Cancer Research* 13, 789-794.
- Giudice, A., Montella, M., 2006. Activation of the Nrf2-ARE signaling pathway: a promising strategy in cancer prevention. *Bioessays* 28, 169-181.
- Gonzalez, F.J., Gelboin, H.V., 1994. Role of human cytochromes P450 in the metabolic activation of chemical carcinogens and toxins. *Drug Metab Rev* 26, 165-183.
- Hail, N., Jr., Cortes, M., Drake, E.N., Spallholz, J.E., 2008. Cancer chemoprevention: a radical perspective. *Free Radic Biol Med* 45, 97-110.
- Hanahan, D., Weinberg, R.A., 2000. The hallmarks of cancer. *Cell* 100, 57-70.
- Hayes, J.D., McMahon, M., 2001. Molecular basis for the contribution of the antioxidant responsive element to cancer chemoprevention. *Cancer letters* 174, 103-113.
- Hidaka, Y., Kaneda, T., Amino, N., Miyai, K., 1992. Chinese medicine, Coix seeds increase peripheral cytotoxic T and NK cells. *Biotherapy* 5, 201-203.
- Hsia, S.M., Chiang, W., Kuo, Y.H., Wang, P.S., 2006. Downregulation of progesterone biosynthesis in rat granulosa cells by adlay (*Coix lachryma-jobi* L. var. *ma-yuen* Stapf.) bran extracts. *Int J Impot Res* 18, 264-274.
- Hsia, S.M., Kuo, Y.H., Chiang, W., Wang, P.S., 2008. Effects of adlay hull extracts on uterine contraction and Ca²⁺ mobilization in the rat. *Am J Physiol Endocrinol Metab* 295, E719-726.
- Hsia, S.M., Tseng, Y.W., Wang, S.W., Kuo, Y.H., Huang, D.W., Wang, P.S., Chiang, W., 2009. Effect of adlay (*Coix lachryma-jobi* L. var. *ma-yuen* Stapf.) hull extracts on testosterone release from rat Leydig cells. *Phytother Res* 23, 687-695.
- Hsia, S.M., Yeh, C.L., Kuo, Y.H., Wang, P.S., Chiang, W., 2007. Effects of adlay (*Coix lachryma-jobi* L. var. *ma-yuen* Stapf.) hull extracts on the secretion of progesterone and estradiol in vivo and in vitro. *Exp Biol Med (Maywood)* 232, 1181-1194.
- Hsu, H.Y., Lin, B.F., Lin, J.Y., Kuo, C.C., Chiang, W., 2003. Suppression of allergic reactions by dehulled adlay in association with the balance of TH1/TH2 cell responses. *J Agric Food Chem* 51, 3763-3769.
- Hu, S.H., Xiao, X.N., Yi, X., Zhang, J.P., 2009. Study progress of coix seed. *Shizhen Guoyi Guoyao* 20, 1059-1060.
- Huang, B.W., Chiang, M.T., Yao, H.T., Chiang, W., 2005. The effect of adlay oil on plasma lipids, insulin and leptin in rat. *Phytomedicine* 12, 433-439.
- Huang, D.W., Chung, C.P., Kuo, Y.H., Lin, Y.L., Chiang, W., 2009a. Identification of compounds in adlay (*Coix lachryma-jobi* L. var. *ma-yuen* Stapf) seed hull extracts that inhibit lipopolysaccharide-induced inflammation in RAW 264.7 macrophages. *J Agric Food Chem* 57, 10651-10657.
- Huang, D.W., Kuo, Y.H., Lin, F.Y., Lin, Y.L., Chiang, W., 2009b. Effect of Adlay (*Coix lachryma-jobi* L. var. *ma-yuen* Stapf) Testa and its phenolic components on Cu²⁺-treated low-density lipoprotein (LDL) oxidation and lipopolysaccharide (LPS)-induced inflammation in RAW 264.7 macrophages. *J Agric Food Chem* 57, 2259-2266.
- Huang, H.C., Nguyen, T., Pickett, C.B., 2002. Phosphorylation of Nrf2 at Ser-40 by protein kinase C regulates antioxidant response element-mediated transcription. *J Biol Chem* 277, 42769-42774.
- Huang, S.-L., Chiang, W., 1999. Composition of the different fractions of adlay seed and the demutagenic effect of their acetone extract. *Food Sci. (Chinese)* 26, 121-130.
- Hung, W.C., Chang, H.C., 2003. Methanolic extract of adlay seed suppresses COX-2 expression of human lung cancer cells via inhibition of gene transcription. *J Agric Food Chem* 51, 7333-7337.
- Ishiguro, Y., Okamoto, K., Sakamoto, H., Sonoda, Y., 1993. Antimicrobial substances coixindens A and B in etiolated seedlings of adlay. *Nippon Nogei Kagaku Kaishi* 67, 1405-1410.
- Kaneda, T., Hidaka, Y., Kashiwai, T., Tada, H., Takano, T., Nishiyama, S., Amino, N., Miyai, K., 1992. [Effect of coix seed on the changes in peripheral lymphocyte subsets]. *Rinsho Byori* 40, 179-181.
- Khan, M., Pelengaris, S., 2006. The molecular biology of cancer. Black-well Publishing. Massachusetts.
- Khansari, N., Shakiba, Y., Mahmoudi, M., 2009. Chronic inflammation and oxidative stress as a major cause of age-related diseases and cancer. *Recent patents on inflammation & allergy drug discovery* 3, 73-80.
- Kim, S.O., Yun, S.J., Jung, B., Lee, E.H., Hahm, D.H., Shim, I., Lee, H.J., 2004. Hypolipidemic effects of crude extract of adlay seed (*Coix lachrymajobi* var. *mayuen*) in obesity rat fed high fat diet: relations of TNF-alpha and leptin mRNA expressions and serum lipid levels. *Life Sci* 75, 1391-1404.
- Kim, S.O., Yun, S.J., Lee, E.H., 2007. The water extract of adlay seed (*Coix lachrymajobi* var. *mayuen*) exhibits anti-obesity effects through neuroendocrine modulation. *Am J Chin Med* 35, 297-308.
- Klaunig, J.E., Kamendulis, L.M., 2004. The role of oxidative stress in carcinogenesis. *Annual review of pharmacology and toxicology* 44, 239-267.
- Klaunig, J.E., Kamendulis, L.M., Honcevar, B.A., 2010. Oxidative

- stress and oxidative damage in carcinogenesis. *Toxicol Pathol* 38, 96-109.
- Klaunig, J.E., Wang, Z., Pu, X., Zhou, S., 2011. Oxidative stress and oxidative damage in chemical carcinogenesis. *Toxicol Appl Pharmacol* 254, 86-99.
- Kondo, Y., Nakajima, K., Nozoe, S., Suzuki, S., 1988. Isolation of ovulatory-active substances from crops of Job's tears (*Coix lacryma-jobi* L. var. *ma-yuen* STAPF.). *Chem Pharm Bull (Tokyo)* 36, 3147-3152.
- Kuo, C.C., Chiang, W., Liu, G.P., Chien, Y.L., Chang, J.Y., Lee, C.K., Lo, J.M., Huang, S.L., Shih, M.C., Kuo, Y.H., 2002. 2,2'-Diphenyl-1-picrylhydrazyl radical-scavenging active components from adlay (*Coix lacryma-jobi* L. var. *ma-yuen* Stapf) hulls. *J Agric Food Chem* 50, 5850-5855.
- Kuo, C.C., Shih, M.C., Kuo, Y.H., Chiang, W., 2001. Antagonism of free-radical-induced damage of adlay seed and its antiproliferative effect in human histolytic lymphoma U937 monocytic cells. *J Agric Food Chem* 49, 1564-1570.
- Kwak, M.K., Kensler, T.W., 2010. Targeting NRF2 signaling for cancer chemoprevention. *Toxicol Appl Pharmacol* 244, 66-76.
- Lee, M.Y., Lin, H.Y., Cheng, F., Chiang, W., Kuo, Y.H., 2008. Isolation and characterization of new lactam compounds that inhibit lung and colon cancer cells from adlay (*Coix lacryma-jobi* L. var. *ma-yuen* Stapf) bran. *Food Chem Toxicol* 46, 1933-1939.
- Lengauer, C., Kinzler, K.W., Vogelstein, B., 1998. Genetic instabilities in human cancers. *Nature* 396, 643-649.
- Li, S.C., Chen, C.M., Lin, S.H., Chiang, W., Shih, C.K., 2011. Effects of adlay bran and its ethanolic extract and residue on preneoplastic lesions of the colon in rats. *J Sci Food Agric* 91, 547-552.
- Lin, M.H., Wu, M.C., Lu, S., Lin, J., 2010. Glycemic index, glycemic load and insulinemic index of Chinese starchy foods. *World J Gastroenterol* 16, 4973-4979.
- Lu, Y., Li, C.S., Dong, Q., 2008. Chinese herb related molecules of cancer-cell-apoptosis: a minireview of progress between Kanglaite injection and related genes. *J Exp Clin Cancer Res* 27, 31.
- Manson, M.M., Gescher, A., Hudson, E.A., Plummer, S.M., Squires, M.S., Prigent, S.A., 2000. Blocking and suppressing mechanisms of chemoprevention by dietary constituents. *Toxicology letters* 112-113, 499-505.
- Mortelmans, K., Zeiger, E., 2000. The Ames Salmonella/microsome mutagenicity assay. *Mutat Res* 455, 29-60.
- Numata, M., Yamamoto, A., Moribayashi, A., Yamada, H., 1994. Antitumor components isolated from the Chinese herbal medicine *Coix lacryma-jobi*. *Planta Med* 60, 356-359.
- Otsuka, H., Hirai, Y., Nagao, T., Yamasaki, K., 1988. Anti-inflammatory activity of benzoxazinoids from roots of *Coix lacryma-jobi* var. *ma-yuen*. *J Nat Prod* 51, 74-79.
- Pan, P., Wu, Y., Guo, Z.Y., Wang, R., Wang, Y.J., Yuan, Y.F., 2012. Antitumor activity and immunomodulatory effects of the intraperitoneal administration of Kanglaite in vivo in Lewis lung carcinoma. *J Ethnopharmacol* 143, 680-685.
- Panayiotidis, M., 2008. Reactive oxygen species (ROS) in multistage carcinogenesis. *Cancer letters* 266, 3-5.
- Rendic, S., 2002. Summary of information on human CYP enzymes: human P450 metabolism data. *Drug Metab Rev* 34, 83-448.
- Seino, K., Motohashi, S., Fujisawa, T., Nakayama, T., Taniguchi, M., 2006. Natural killer T cell-mediated antitumor immune responses and their clinical applications. *Cancer Sci* 97, 807-812.
- Shih, C.K., Chiang, W., Kuo, M.L., 2004. Effects of adlay on azoxymethane-induced colon carcinogenesis in rats. *Food Chem Toxicol* 42, 1339-1347.
- Smith, T.J., Guo, Z., Guengerich, F.P., Yang, C.S., 1996. Metabolism of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK) by human cytochrome P450 1A2 and its inhibition by phenethyl isothiocyanate. *Carcinogenesis* 17, 809-813.
- Son, B.K., Kim, J.Y., Lee, S.S., 2008. Effect of adlay, buckwheat and barley on lipid metabolism and aorta histopathology in rats fed an obesogenic diet. *Ann Nutr Metab* 52, 181-187.
- Surh, Y.J., 2003. Cancer chemoprevention with dietary phytochemicals. *Nature reviews. Cancer* 3, 768-780.
- Surh, Y.J., Kundu, J.K., 2007. Cancer preventive phytochemicals as speed breakers in inflammatory signaling involved in aberrant COX-2 expression. *Curr Cancer Drug Targets* 7, 447-458.
- Takahashi, M., Konno, C., Hikino, H., 1986. Isolation and hypoglycemic activity of coixans A, B and C, glycans of *Coix lacryma-jobi* var. *ma-yuen* seeds. *Planta Med*, 64-65.
- Tanimura, A., 1961. Studies on the antitumor components in the seeds of *Coix lacryma-jobi* L. var. *ma-yuen* (Roman.) Stapf. II. The structure of coixenolide. *Chem Pharm Bull (Tokyo)* 9, 47-53.
- Tzeng, H.P., Chiang, W., Ueng, T.H., Liu, S.H., 2005. The abortifacient effects from the seeds of *Coix lacryma-jobi* L. var. *ma-yuen* Stapf. *J Toxicol Environ Health A* 68, 1557-1565.
- Ukita, T., Tanimura, A., 1961. Studies on the antitumor components in the seeds of *Coix lacryma-jobi* L. var. *ma-yuen* (Roman.) Stapf. I. Isolation and antitumor activity of coixenolide. *Chem Pharm Bull (Tokyo)* 9, 43-46.
- Wang, H., Cho, C.H., 2010. Effect of NF-kappaB signaling on apoptosis in chronic inflammation-associated carcinogenesis. *Curr Cancer Drug Targets* 10, 593-599.
- Wang, L., Sun, J., Yi, Q., Wang, X., Ju, X., 2012. Protective Effect of Polyphenols Extract of Adlay (*Coix lacryma-jobi* L. var. *ma-yuen* Stapf) on Hypercholesterolemia-Induced Oxidative Stress in Rats. *Molecules* 17, 8886-8897.
- Wattenberg, L.W., 1985. Chemoprevention of cancer. *Cancer research* 45, 1-8.
- Weisburger, J.H., 2001. Antimutagenesis and anticarcinogenesis, from the past to the future. *Mutat Res* 480-481, 23-35.
- Windmill, K.F., McKinnon, R.A., Zhu, X., Gaedigk, A., Grant, D.M., McManus, M.E., 1997. The role of xenobiotic metabolizing enzymes in arylamine toxicity and carcinogenesis: functional and localization studies. *Mutat Res* 376, 153-160.
- Woo, J.H., Li, D., Wilsbach, K., Orita, H., Coulter, J., Tully, E., Kwon, T.K., Xu, S., Gabrielson, E., 2007. Coix seed extract, a commonly used treatment for cancer in China, inhibits NFkappaB and protein kinase C signaling. *Cancer Biol Ther* 6, 2005-2011.
- Yang, R.S., Chiang, W., Lu, Y.H., Liu, S.H., 2008. Evaluation of osteoporosis prevention by adlay using a tissue culture model. *Asia Pac J Clin Nutr* 17 Suppl 1, 143-146.
- Yao, H.T., Lin, J.H., Chiang, M.T., Chiang, W., Luo, M.N., Lii, C.K., 2011. Suppressive effect of the ethanolic extract of adlay bran on cytochrome P-450 enzymes in rat liver and lungs. *J Agric Food Chem* 59, 4306-4314.
- Yeh, P.H., Chiang, W., Chiang, M.T., 2006. Effects of dehulled adlay on plasma glucose and lipid concentrations in streptozotocin-induced diabetic rats fed a diet enriched in cholesterol. *Int J Vitam Nutr Res* 76, 299-305.
- Yu, F., Gao, J., Zeng, Y., Liu, C.X., 2011. Effects of adlay seed oil on blood lipids and antioxidant capacity in hyperlipidemic rats. *J Sci Food Agric* 91, 1843-1848.
- Yu, Y.M., Chang, W.C., Liu, C.S., Tsai, C.M., 2004. Effect of young barley leaf extract and adlay on plasma lipids and LDL oxidation in hyperlipidemic smokers. *Biol Pharm Bull* 27, 802-805.
- Zhang, Y., Gordon, G.B., 2004. A strategy for cancer prevention: stimulation of the Nrf2-ARE signaling pathway. *Mol Cancer Ther* 3, 885-893.