




Cairo University
Bulletin of Faculty of Pharmacy, Cairo University

www.elsevier.com/locate/bfopcu
www.sciencedirect.com



REVIEW ARTICLE

Naturally occurring didemnaketals: Structural elucidation, features, and pharmacological activities



Gamal A. Mohamed ^a, Sabrin R. M. Ibrahim ^{b,*}, Diaa T. A. Youssef ^{c,d}

^a Department of Pharmacognosy, Faculty of Pharmacy, Al-Azhar University, Assiut Branch, Assiut 71524, Egypt

^b Department of Pharmacognosy, Faculty of Pharmacy, Assiut University, Assiut 71526, Egypt

^c Department of Natural Products and Alternative Medicine, Faculty of Pharmacy, King Abdulaziz University, Jeddah 21589, Saudi Arabia

^d Department of Pharmacognosy, Faculty of Pharmacy, Suez Canal University, Ismailia 41522, Egypt

Received 13 December 2014; accepted 11 February 2015

Available online 3 March 2015

KEYWORDS

Didemnaketals;
Didemnum;
 Ascidian;
 Spectral data;
 Bioactivities

Abstract Didemnaketals are rare terpenoid class reported only in the marine ascidian *Didemnum* species. They possess important biological activities like HIV-1 protease inhibitor, protein kinase inhibition, antimicrobial, antiproliferative, and cytotoxic. A compilation of new naturally occurring didemnaketals reported during 1991–2014 is provided with available physical and spectral data: mp, $[\alpha]_D$, UV, IR, ^1H and ^{13}C NMR as well as biological activities and references.

© 2015 Production and hosting by Elsevier B.V. on behalf of Faculty of Pharmacy, Cairo University.

Contents

1. Introduction	70
2. The structural features of didemnaketals	71
3. The absolute stereochemistry of didemnaketals	72
4. Biological activities of didemnaketals	72
4.1. Potent HIV-1 protease inhibitor activity.	72
4.2. Protein kinase inhibitory activity.	75
4.3. Antimicrobial activity.	75
4.4. Antiproliferative and cytotoxic activities	75
5. Spectral data.	75
6. Conflict of interest.	75
References	76

* Corresponding author. Tel.: +20 88 2141330; fax: +20 88 2332776.

E-mail address: sabrinshaur@gmail.com (S.R.M. Ibrahim).

Peer review under responsibility of Faculty of Pharmacy, Cairo University.

<http://dx.doi.org/10.1016/j.bfopcu.2015.02.003>

1110-0931 © 2015 Production and hosting by Elsevier B.V. on behalf of Faculty of Pharmacy, Cairo University.




Figure 1

1. Introduction

The number of natural products isolated from marine organisms increases rapidly and many new compounds are discovered every year.¹ A large proportion of natural compounds have been isolated from marine invertebrates such as sponges, ascidians, bryozoans, and molluscs, and some of them are currently in clinical trials.² Ascidians or sea-squirts are cosmopolitan, exclusively marine invertebrates, which represent the most highly evolved group of animals commonly investigated by marine natural product chemists.³ Attention has focused on ascidians because of their biologically active metabolites. The chemistry of ascidians has become one of the most active fields of marine natural products. It has been demonstrated that these sea creatures are prolific producers of unusual structures with significant bioactivities.⁴ The compounds isolated from genus *Didemnum* are diverse and possess a wide range of biological activities such as antiplasmodial,⁵ antiviral,⁶ cytotoxic,⁷

and protein kinase inhibitors.⁸ Didemnaketals are very rare linear heptaprenoids, isolated from the marine *Didemnum* species. To date, seven spiroketals, didemnaketals A–G (1–7) (Fig. 1) were isolated from the marine *Didemnum* species collected from different geographical locations including Aulup-tagel Island Palau^{9,10} and Nabq/Sharm El-Sheikh on the Egyptian Red Sea coast.^{8,11} Two of them, didemnaketals A (1) and B (2) were reported as a result of oxidation and methanolysis of polar isoethonic ester, didemnaketal C (3) during long storage of the ascidian sample in MeOH.^{9,10,12} While, didemnaketals C–G (4–7) were isolated from fresh samples of marine ascidian *Didemnum* species.^{8,9,11} Here, we listed the didemnaketals that have appeared in the literature over the past few decades, with their biological activities, physical constants, spectral data, and references. For each compound these data were listed in the following order: name; structure; melting point (°C); optical rotation (concentration, solvent); UV (solvent, λ_{\max} nm, log ϵ); IR (medium, absorption band





Figure 1 (continued)

in cm^{-1}); ^1H NMR (spectrometer frequency, solvent, chemical shift values in ppm, starting with H-3 and listed in order); ^{13}C NMR (spectrometer frequency, solvent, chemical shift in ppm values starting with C-1 and listed in order); molecular formula; calculated molecular weight; and reference(s) (Tables 1–3). The main aim of this review is for rapid identification of isolated didemnaketals by comparison of spectral data as well as to draw attention of the natural product chemists for evaluation of biological and pharmacological activities of the isolated didemnaketals. The high-light of pharmacological activity may possibly draw the attention of the synthetic chemists for design of new drugs using the known didemnaketals.

2. The structural features of didemnaketals

Didemnaketals are a rare class of polyisoprenoids, characterized by the presence of bicyclic ketal region from C-12 to C-20 and *penta*-ester side chain.^{13,14} The linear heptaprenoid structure contains a number of intriguing structural features

such as a spiroketal moiety and two side chains with eight chiral centers.^{13,15} Didemnaketals **D** (4) and **E** (5) isolated from *Didemnum* sp. collected from Sharm El-sheikh on the Red Sea coast,⁸ differ from didemnaketal **A** isolated from the magenta ascidian *Didemnum* sp. collected at Auluptagel island, Palauin,¹⁰ in lacking the methyl at C-6 and the hydroxy at C-21, as well as possessing an additional substituent at C-20. They showed new moieties at C-6 (methyl ester in **D** (4) and ethyl ester in **E** (5) and an oxygen functionality at C-20 (acetoxyl in **D** (4) and hydroxy in **E** (5)). The new moieties at C-20 represent an unprecedented and complicated spiroketal/ketal (in 4) and spiroketal/hemiketal (in 5) moieties, which are not observed in the previously described spiroketal systems.⁸ Moreover, didemnaketals **F** (6) and **G** (7) are characterized by the absence of the methyl functionality at C-2 and terminal methyl ester moiety of the spiroketal skeleton. Instead, they possess a terminal methyl ketone moiety at the left part of the molecule. Additionally, **6** possesses a *di*-substituted double bond at C-2/C-3, while **7** has a secondary alcohol at C-3. Also, they have spiroketal/hemiketal functionality at the right part of the molecule.^{8,11}




Figure 1 (continued)

3. The absolute stereochemistry of didemnaketals

The gross structures of didemnaketals **A–G** (**1–7**) were determined by extensive 1D and 2D NMR spectroscopic analyses.^{8–11}

The absolute stereochemistry at the stereogenic centers of didemnaketals **A–C** (**1–3**) was determined to be *5S,6S,7R,8R,10S,11S,12S,14S,16S,18R,20S,21S* (didemnaketal **A**) and *5S,6S,7R,8R,10S,11S,12S,14S,16S,18R,20S,21S,26S* (didemnaketals **B** and **C**),¹⁴ using a combination of degradation and derivatization experiments, chiral shift methods, as well as synthesis of the C9–C28 subunit containing the spiroketal moiety through a highly convergent strategy in didemnaketal **B** (**2**).^{12,14} The configuration of didemnaketals **D** and **E** was assigned to be *5S,6S,7R,8R,10S,11S,12R,14S,16S,18R* on the basis of NOESY correlations and comparison of the NMR spectral data with those of previously reported didemnaketals.⁸ Fuwa and coworkers revised Faulkner's stereochemical assignment of didemnaketals **B** through its total synthesis.¹⁶ The stereochemical relationship of the C5/C6, C6/C7, and C7/C8 stereogenic centers was correlated by NMR spectro-

scopic analyses on suitable acetonide derivatives. X-ray crystallographic analysis is used for assignment of the relative configuration of the C10–C20 spiroacetal domain. The absolute configuration of the C5, C8, C11, and C21 stereogenic centers was determined by application of the modified Mosher analysis, while that of the C20 and C26 stereogenic centers was assigned on the basis of the phenyl glycine methyl ester (PGME) method.¹⁶ The geometry of the C2–C3 double bond was confirmed to be *E* by NOE experiment as well as by a large ³*J*_{H,H} value (³*J*_{H₂,H₃} = 16.5 Hz).^{11,16} The absolute configuration of didemnaketals **F** (**6**) and **G** (**7**) was determined to be *5S,6S,7R,8R,10S,11S,12S,14R,16S,18R*,¹¹ based on Fuwa's stereochemical revision of didemnaketals **B** (**2**) as well as extensive spectroscopic studies.¹⁶

4. Biological activities of didemnaketals

4.1. Potent HIV-1 protease inhibitor activity

Replication of human immunodeficiency virus (HIV) entails expression of several viral polyproteins which require the pres-

Table 1 NMR spectroscopic data of didemnaketals **A (1)**, **D (4)**, and **E (5)**.

No.	Didemnaketals A (1)		Didemnaketals D (4)		Didemnaketals E (5)	
	δ_{H} (<i>J</i> in Hz) ^a	δ_{C} ^a	δ_{H} (<i>J</i> in Hz) ^b	δ_{C} ^b	δ_{H} (<i>J</i> in Hz) ^a	δ_{C} ^a
1	-	168.1	-	-	-	176.7
2	-	130.2	-	175.3	-	131.7
3	6.71 ddd (5, 7, 2)	136.1	-	133.1	5.15 t (6.2)	127.2
4	2.49 ddd (16, 5, 5) 2.53 ddd (16, 7, 7)	30.8	5.23 t (6.0)	128.2	2.03 m	33.3
5	4.83 m	72.2	2.24 m, 2.17 m	34.6	3.45 dt (11.6, 7.1)	72.9
6	2.09 m	36.7	3.51 dt (11.5, 7.3)	75.1	2.29 dt (11.6, 6.9)	41.8
7	5.16 dd (9, 2)	72.9	3.00 m	42.2	4.85 t (9.1)	67.9
8	5.22 br t (10)	70.6	4.95 m	69.0	3.95 m	69.3
9	1.24 m, 1.61 m	32.2	4.12 m	70.9	1.64 m, 1.25 m	36.4
10	1.84 m	29.4	1.68 m, 1.23 m	37.9	1.94 m	33.7
11	4.78 dd (6, 6)	78.2	2.26 m	35.5	5.00 m	74.7
12	3.69 m	69.2	5.08 dd (6.5, 6.3)	76.8	3.67 m	73.9
13	0.81 ddd (12, 12, 12) 1.45 br d (13)	34.5	4.08 m	75.5	1.05 m, 0.80 m	36.6
14	1.93 m	24.9	1.27 m, 0.77 m	37.7	1.69 m	25.4
15	0.98 br m 1.67 dd (14, 3)	43.9	1.78 m	27.6	1.62 m, 1.08 m	45.0
16	-	98.4	1.08 m, 1.67 m	46.2	-	98.8
17	1.45 br d (13) 1.59 br d (13)	40	-	100.3	2.10 m, 1.95 m	46.7
18	1.99 m	24.3	2.22 m, 2.07 m	47.6	1.68 m	24.6
19	1.34 br dd (14, 3) 1.71 ddd (14, 11, 5)	31.8	1.75 m	25.8	2.20 m	33.9
20	3.89 ddd (11, 5, 3)	67.5	1.62 m, 1.42 m	37.7	-	99.7
21	4.14 dd (5, 5)	79.1	-	101.5	2.99 d (13.5) 2.80 d (13.5)	41.1
22	-	209.3	3.07 d (14.1) 3.05 d (14.1)	42.3	-	202.6
23	2.32 s	28.1	-	204.4	2.16 s	20.6
24	1.82 s	12.5	2.15 s	21.1	1.63 s	17.0
25	0.98 d	9.8	1.71 s	17.6	-	173.2
26	0.98 d	16.1	-	178.2	0.88 d (6.8)	14.8
27	0.86 d (7)	22.0	0.88 d (6.5)	15.1	0.92 d (6.9)	22.2
28	1.08 m	20.6	0.95 d (7.2)	22.7	0.94 d (6.5)	19.6
29	3.72 s	51.8	0.92 d (6.5)	22.5	3.65 s	51.7
30	-	173.8	3.68 s	55.3	-	170.3
31	2.28 br m	27.5	-	170.4	2.15 m	26.6
32	1.08 m	8.3	2.24 m	26.4	1.05 t (7.5)	7.7
33	-	170.2	1.02 t (6.3)	8.4	-	169.9
34	1.98 s	20.8	-	171.4	2.05 s	21.3
35	-	172.4	2.06 s	21.2	-	173.2
36	2.13 m	43.1	-	178.2	2.15 m	41.8
37	2.04 m	25.2	2.32 m	40.9	2.85 m	33.3
38	0.93 d	22.3	2.90 m	35.4	1.30 d (6.6)	14.3
39	0.93 d	22.4	1.22 d (7.0)	17.5	1.30 d (6.6)	14.3
40	-	172.7	1.22 d (7.0)	17.5	-	168.6
41	2.24 m	43.6	-	172.2	1.98 m	34.2
42	2.08 m	25.5	2.31 m	38.1	1.92 m	30.1
43	0.98 d	22.5	2.03 m	31.1	1.19 d (6.5)	17.2
44	0.98 d	22.5	0.98 (d, 7.0)	20.0	1.19 d (6.5)	17.2
45	-	-	0.98 (d, 7.0)	20.0	4.15 q (6.9)	60.1
46	-	-	3.67 s	51.9	1.25 t (6.9)	14.3
47	-	-	-	172.2	-	-

^a Recorded in CDCl₃ at 400 and 100 MHz.^b Recorded in CD₃OD at 500 and 125 MHz.

ence of a virus-specific protease for their maturation. Inhibition of this enzyme results in immature viral particles and inhibition of viral replication *in vitro*. Inhibition of HIV-1 protease is one

of the most effective ways to treat AIDS, but the emergence of drug resistant viral strains mandates new approaches.¹³ Didemnaketals **A (1)** and **B (2)** were found to inhibit HIV-1 protease

Table 2 NMR data of didemnaketals **B (2)** and **C (3)** (CDCl₃, 500 and 125 MHz).

No.	Didemnaketal B (2)		Didemnaketal C (3)	
	δ_{H} (<i>J</i> in Hz)	δ_{C}	δ_{H} (<i>J</i> in Hz)	δ_{C}
1	-	168.0	-	168.0
2	-	130.3	-	130.2
3	6.69 t (7)	135.5	6.66 t (6)	135.9
4	2.47 dt (16, 7) 2.51 dt (16, 5)	30.9	2.41 m, 2.50 m	30.9
5	4.83 m	72.3	4.79 m	72.2
6	1.96 m	37.0	1.93 m	36.9
7	5.12 dd (9, 3)	72.7	5.10 t (9)	72.7
8	5.23 dd (9, 8)	70.6	5.18 m	70.5
9	1.22 m, 1.66 m	32.9	1.14, m, 1.60 m	33.0
10	1.90 m	29.5	1.90 m	29.6
11	4.77 dd (6, 5)	78.6	4.71 t (6)	78.5
12	3.85 m	69.1	3.84 m	69.0
13	0.80 q (14) 1.44 m	35.1	0.78 m, 1.40 m	35.0
14	2.06 m	24.8	2.04 m	24.4
15	1.00 d (7) 1.67 m	44.3	0.96 m	44.3
16	-	98.3	-	98.4
17	1.54 br d (14) 1.64 m	40.7	1.39 m, 1.54 m	40.6
18	1.91 m	24.5	2.05 m	24.7
19	1.12 m 1.58 br d (14)	29.98	1.07 m, 1.56 m	29.6
20	3.87 m	66.6	3.84 m	66.3
21	4.05 d (3)	78.0	4.03 br s	77.6
22	-	133.0	-	133.0
23	5.48 t (7)	126.3	5.41 t (7)	125.9
24	2.08 m, 2.08 m	24.9	1.91 m, 2.04 m	25.1
25	1.25 m, 1.38 m	36.5	1.20 m, 1.39 m	36.2
26	2.00 m,	30.04	1.90 m	29.4
27	2.13 m 2.29 dd (7, 5)	41.5	1.90 m, 2.24 m	41.7
28	-	173.5	-	173.8
29	1.84 s	12.6	1.80 m	12.6
30	0.97 d (7)	9.9	0.94 d (7)	9.9
31	0.98 d (7)	16.2	0.97 d (7)	16.1
32	0.86 d (7)	22.1	0.82 d (6)	22.1
33	1.07 d (6)	20.84	1.04 d (7)	20.8
34	1.62 s	13.3	1.57 br s	13.6
35	0.96 d (7)	19.6	0.91 d (7)	19.5
36	3.73 s	51.8	3.70 s	51.9
37	-	173.7	-	173.3
38	2.35 m, 2.35 m	27.6	2.22 m, 2.28 m	27.6
39	1.09 t (7)	8.9	1.06 t (7)	8.9
40	-	170.1	-	170.3
41	1.98 s	20.77	1.95 s	20.8
42	-	172.3	-	172.4
43	2.13 d (7)	43.2	2.12 m	43.1
44	2.06 m	25.3	2.05 m	25.2
45	0.94 d (6)	22.5	0.90 d (7)	22.5
46	0.94 d (6)	22.5	0.95 d (7)	22.5
47	-	172.7	-	172.8
48	2.25 d (7)	43.7	2.22 m	43.6
49	2.12 m	25.6	2.00 m	25.6
50	0.99 d (6)	22.6	0.90 d (7)	22.5
51	0.99 d (6)	22.6	0.95 d (7)	22.5
52	3.66 s	51.3	4.46 t (7)	59.5
53	-	-	3.17 t (7)	50.0

Table 3 NMR data of didemnaketals **F (6)** and **G (7)** (CDCl₃, 600 and 150 MHz).

No.	Didemnaketals F (6)		Didemnaketals G (7)	
	δ_{H} (J in Hz)	δ_{C}	δ_{H} (J in Hz)	δ_{C}
1	-	198.5	-	207.2
2	6.11 d (16.5)	133.0	2.61 dd (14.5, 7.3) 2.45 dd (14.5, 7.5)	50.1
3	6.75 dt (16.5, 6.8)	145.8	4.28 quin (7.3)	67.5
4	1.97 m	29.9	1.65 m, 0.87 m	36.1
5	3.45 dt (11.6, 7.3)	72.3	3.45 dt (11.5, 6.9)	72.3
6	2.30 dt (11.6, 6.9)	41.6	2.95 dt (11.6, 6.5)	41.7
7	4.83 t (9.6)	67.9	4.83 t (9.8)	67.9
8	4.30 m	67.5	3.90 m	67.5
9	1.70 m, 1.25 m	36.8	1.75 m, 1.29 m	36.9
10	1.92 m	34.1	1.95 m	33.9
11	5.02 dd (6.8, 6.3)	76.5	5.01 dd (6.5, 6.3)	76.8
12	3.86 m	75.3	3.86 m	75.3
13	1.48 m, 0.88 m	36.1	1.47 m, 0.85 m	36.1
14	1.72 m	25.5	1.70 m	26.6
15	1.62 m, 1.08 m	44.8	1.63 m, 1.08 m	44.7
16	-	98.7	-	98.7
17	1.72 m, 1.56 m	42.0	1.74 m, 1.55 m	41.1
18	1.75 m	24.6	1.75 m	24.6
19	2.15 m, 1.74 m	33.1	2.15 m	32.0
20	-	99.8	-	99.8
21	2.95 m, 2.86 m	41.2	2.85 m, 2.72 m	41.1
22	-	202.5	-	202.5
23	2.19 s	21.3	2.06 s	21.0
24	2.24 s	29.7	2.16 s	31.2
25	-	172.6	-	175.8
26	0.90 d (6.5)	14.8	0.88 d (6.5)	14.8
27	0.91 d (6.8)	22.2	0.92 d (6.8)	22.0
28	0.97 d (6.5)	19.8	0.90 d (6.5)	20.5
29	-	170.5	-	170.2
30	2.16 q (7.3)	26.6	2.18 q (7.3)	26.8
31	1.03 t (7.3)	7.5	1.03 t (6.9)	7.5
32	-	170.2	-	170.5
33	2.05 s	21.0	2.01 s	21.3
34	-	176.6	-	176.6
35	2.13 m	41.8	2.33 m	39.8
36	2.87 m	33.9	2.85 m	33.2
37	1.20 d (6.8)	17.2	1.22 d (6.8)	17.2
38	1.20 d (6.8)	17.2	1.22 d (6.8)	17.2
39	-	168.6	-	168.6
40	2.03 m	33.2	2.05 m	33.1
41	1.97 m	31.2	1.97 m	29.7
42	0.95 d (6.3)	22.0	0.93 d (6.3)	22.2
43	0.95 d (6.3)	22.0	0.93 d (6.3)	22.2
44	4.11 q (6.8)	60.4	3.64 s	51.8
45	1.27 t (6.8)	14.3	-	-

with an IC₅₀ of 2 and 10 μM , respectively.¹⁰ So, several synthetic studies of the didemnaketals especially **A (1)** have been conducted to determine the structural components necessary for activity; as well as stable analogs that may provide a unique and potent mode of HIV-1 protease inhibition by an unusual mechanism.¹³ Unfortunately, didemnaketals **C (3)** did not inhibit HIV-1 in a peptidolysis assay.^{14,17}

4.2. Protein kinase inhibitory activity

Didemnaketals **D (4)** and **E (5)** displayed moderate protein kinase inhibitory activity against CDK5, CK1, DyrK1A and, GSK3 at concentration 10 $\mu\text{g}/\text{mL}$. The IC₅₀ of both

compounds were higher than 10 $\mu\text{g}/\text{mL}$. Such activity is not sufficient to pursue with these compounds into the *in vivo* studies.⁸

4.3. Antimicrobial activity

Didemnaketals **D (4)** and **E (5)** were evaluated for their antimicrobial potential against Gram-positive bacteria, Gram-negative bacteria, and yeast using the disk diffusion method. Didemnaketals **D (4)** showed moderate activity (11 mm, inhibition zone) toward *Staphylococcus aureus* ATCC 6538, while didemnaketals **E (5)** showed moderate activity (11 mm, inhibition zone) against *Bacillus subtilis* ATCC CC33.⁸ While, didemnaketals **F (6)** showed strong antimicrobial activity against *Escherichia coli* ATCC 25922 and *Candida albicans* ATCC 14053 with inhibition zones of 20 and 24 mm at a concentration of 100 $\mu\text{g}/\text{disk}$. Didemnaketals **G (7)** showed moderate activity against *C. albicans* with an inhibition zone of 17 mm at the same concentration.¹¹

4.4. Antiproliferative and cytotoxic activities

Didemnaketals **F (6)** and **G (7)** displayed moderate antiproliferative activity against HeLa cells with IC_{50s} of 49.9 and 14.0 μM , respectively using sulforhodamine B (SRB) assay. In contrast, cytotoxic effects were not observed with didemnaketals **F (6)** at concentrations up to 50 μM compared with paclitaxel as a positive control (IC₅₀ of 0.0017 μM).¹¹

5. Spectral data

Didemnaketals **A (1)**: Clear oil; $[\alpha]_{\text{D}} -11.0^{\circ}$ (*c* 0.8, CHCl₃); IR (CHCl₃) ν_{max} 3490, 1735, 1712 cm^{-1} ; HRFABMS *m/z* 825.5030 (calcd for C₄₄H₇₃O₁₄, [M + H]⁺, 825.5000).¹⁰

Didemnaketals **B (2)**: Clear oil; IR (CHCl₃) ν_{max} 3500, 1735, 1710 cm^{-1} ; HRFABMS *m/z* 933.5951 (calcd for C₅₂H₈₅O₁₄, [M – OH]⁺, 933.59393).^{10,16}

Didemnaketals **C (3)**: Colorless oil; $[\alpha]_{\text{D}} + 48$ (*c* 0.067, CH₂Cl₂); UV (CH₃CN) λ_{max} 213 nm; IR (film) ν_{max} 3440, 2955, 1740, 1715 cm^{-1} ; HRFABMS *m/z*: 1067.5522 (calcd for C₅₃H₈₈O₁₈SNa, 1067.5589).⁹

Didemnaketals **D (4)**: Colorless oil, $[\alpha]_{\text{D}} -51.9^{\circ}$ (*c* 1.5, CHCl₃); UV (MeOH) λ_{max} (log ϵ) 246 (4.08) nm; IR (KBr) ν_{max} 1732, 1715, 1296, 1050 cm^{-1} ; HRESIMS *m/z* 911.4923 (calcd for C₄₇H₇₅O₁₇, [M + H]⁺, 911.4926).⁸

Didemnaketals **E (5)**: Colorless oil, $[\alpha]_{\text{D}} -46.3^{\circ}$ (*c* 1.5, CHCl₃); UV (MeOH) λ_{max} (log ϵ) 234 (4.18) nm; IR (KBr) ν_{max} 3495, 1735, 1710, 1291, 1056 cm^{-1} ; HRESIMS *m/z* 883.5029 (calcd for C₄₆H₇₅O₁₆, [M + H]⁺, 883.5027).⁸

Didemnaketals **F (6)**: Colorless oil; $[\alpha]_{\text{D}} - 72.8^{\circ}$ (*c* 1.5, CHCl₃); UV (MeOH) λ_{max} (log ϵ) 252 (3.93) nm; IR (film) ν_{max} 3460, 1735, 1720, 1675, 1650, 1070 cm^{-1} ; HRESIMS *m/z* 853.4950 (calcd for C₄₅H₇₃O₁₅ [M + H]⁺, 853.4949).¹¹

Didemnaketals **G (7)**: Colorless oil; $[\alpha]_{\text{D}} -85.7^{\circ}$ (*c* 1.5, CHCl₃); UV (MeOH) λ_{max} (log ϵ) 258 (3.86) nm; IR (film) ν_{max} 3465, 1731, 1718, 1065 cm^{-1} ; HRESIMS *m/z* 857.4899 (calcd for C₄₄H₇₃O₁₆, [M + H]⁺, 857.4898).¹¹

6. Conflict of interest

None.

References

1. Faulkner DJ. Marine natural products. *Nat Prod Rep* 2002;**19**:1–48.
2. Proksch P, Edrada RA, Ebel R. Drugs from the sea current status and microbiological implications. *Appl Microbiol Biotechnol* 2002;**59**:125–34.
3. Blunt JW, Copp BR, Hu WP, Munro MHG, Northcote PT, Prinsep MR. Marine natural products. *Nat Prod Rep* 2008;**25**:35–94.
4. Blunt JW, Copp BR, Munro MHG, Northcote PT, Prinsep MR. Marine natural products. *Nat Prod Rep* 2010;**27**:165–237.
5. Wright AD, Goclik E, König GM, Kaminsky R. Antiplasmodial and antitrypanosomal decahydroquinoline derivatives from the tropical marine tunicate *Didemnum* sp. *J Med Chem* 2002;**45**:3067–72.
6. Mitchell SS, Rhodes D, Bushman FD, Faulkner DJ. Cyclodidemniserinol trisulfate, a sulfated serinolipid from the Palauan ascidian *Didemnum guttatum* that inhibits HIV-1 integrase. *Org Lett* 2000;**2**:1605–7.
7. Segraves NL, Lopez S, Johnson TA, Said SA, Fu X, Schmitz FJ, Pietraszkiewicz H, Crews P. Structures and cytotoxicities of faspaplysin and related alkaloids from two marine phyla-Faspaplysinopsis sponges and *Didemnum* tunicates. *Tetrahedron Lett* 2003;**44**:3471–5.
8. Mohamed GA, Ibrahim SRM, Badr JM, Youssef DTA. Didemnaketals D and E, bioactive terpenoids from a Red Sea ascidian *Didemnum* species. *Tetrahedron* 2014;**70**:35–40.
9. Pika J, Faulkner DJ. A reinvestigation of the didemnaketals from the Palauan ascidian *Didemnum* sp. *Nat Prod Lett* 1995;**7**:291–6.
10. Potts BCM, Faulkner DJ. Didemnaketals A and B, HIV-1 protease inhibitors from the ascidian *Didemnum* sp. *J Am Chem Soc* 1991;**113**:6321–2.
11. Shaala LA, Youssef DTA, Ibrahim SRM, Mohamed GA, Badr AL, Risinger AL, Mooberry SL. Didemnaketals F and G, new bioactive spiroketals from a Red Sea ascidian *Didemnum* species. *Mar Drugs* 2014;**12**:5021–34.
12. Fuwa H, Noji S, Sasaki M. Convergent assembly of the spiroacetal subunit of didemnaketal B. *Org Lett* 2010;**12**:5354–7.
13. Fan X, Flentke GR, Rich DH. Inhibition of HIV-1 protease by a subunit of didemnaketal A. *J Am Chem Soc* 1998;**120**:8893–4.
14. Salomon CE, Williams DH, Lobkovsky E, Clardy JC, Faulkner DJ. Relative and absolute stereochemistry of the didemnaketals, metabolites of a Palauan ascidian *Didemnum* sp. *Org Lett* 2002;**4**:1699–702.
15. Ito H, Inoue T, Iguchi K. Toward the synthesis of didemnaketal B: A convergent synthesis of the C9–C28 subunit. *Org Lett* 2008;**10**:3873–6.
16. Fuwa H, Muto T, Sekine K, Sasaki M. Total Synthesis and structure revision of didemnaketal B. *Chem Eur J* 2014;**20**:1848–60.
17. Hyland LJ, Dayton BD, Moore ML, Shu AYL, Heys JR, Meek TD. A radiometric assay for HIV-1 protease. *Anal Biochem* 1990;**188**:408–15.