



Contents lists available at ScienceDirect

Journal of Pediatric Surgery CASE REPORTS

journal homepage: www.jpascasereports.com

A case of prenatally detected hepatic cyst communicating with the hepatic duct[☆]



Shinsuke Katsuno^{a,*}, Minoru Horisawa^b, Hiroyuki Fujisaki^c, Takashi Arikawa^c,
Katsuhiko Kotake^c, Tsuyoshi Kurokawa^c, Toshiaki Nonami^c

^a Department of Pediatric Surgery, Saishukan Hospital, 111 Nishimuramae, Shikata, Kita Nagoya-shi, Aichi 481-0004, Japan

^b Department of Pediatric Surgery, Anjo Kosei Hospital, 28 Higashihokute, Anjocho, Anjo-shi, Aichi 446-8602, Japan

^c Department of Gastroenterological Surgery, Aichi Medical University School of Medicine, 1-1 Yazakokarimata, Nagakute-shi, Aichi 480-1195, Japan

ARTICLE INFO

Article history:

Received 9 December 2013

Received in revised form

4 September 2014

Accepted 4 September 2014

Key words:

Hepatic cyst

Communication with bile duct

Intrahepatic biliary cyst

ABSTRACT

Here, we describe the case of a 9-year-old girl. During the patient's prenatal period, her mother had suffered domestic violence perpetrated by the patient's father. On maternal ultrasonography and magnetic resonance imaging, an intraabdominal cystic lesion was detected in the fetus at around the 30th prenatal week. The patient was delivered normally, and there were no evident anomalies on her body. Computed tomography with drip infusion cholangiography and percutaneous retrograde transhepatic cholangiography demonstrated an intrahepatic cyst of approximately 3 cm in diameter, which was located at S₅–S₈ and communicated with the confluence of the bilateral hepatic ducts. The cyst is clinically conjectured to be a solitary intrahepatic biliary cyst. However, it remains possible that the cyst is a ciliated hepatic foregut cyst or indicates hepatic injury that may have occurred as a result of domestic violence to the mother. Careful, long-term observation of the patient will be continued to ensure that any malignant transformation is not missed.

© 2014 The Authors. Published by Elsevier Inc. All rights reserved.

The use of ultrasonography or magnetic resonance imaging (MRI) can prenatally visualize various congenital diseases as intraabdominal cystic lesions. These congenital diseases include congenital biliary dilatation, hydronephrosis, renal cyst, ovarian cyst, mesenteric cyst, intestinal atresia, and meconium peritonitis. However, the incidence of congenital hepatic cysts is considered to be very low (0.01%) [1]. In this report, we describe an extremely rare case of a prenatally detected hepatic cyst communicating with the hepatic duct.

1. Case report

The patient is a girl who is currently 9 years old. During the patient's prenatal period, her mother had suffered domestic violence perpetrated by the patient's father. In the 23rd gestational week, an epigastric cystic lesion was detected in the fetal abdomen using maternal ultrasonography. In the 30th gestational week, MRI

revealed an intrahepatic or extrahepatic cystic lesion in the fetus (Fig. 1). On the sixth day of the 38th gestational week, the patient was born by scheduled cesarean section, which was performed because the mother had previously undergone cesarean delivery.

The baby girl weighed 2675 g at birth and had an Apgar score of 9 points at both 1 and 5 min after birth. There were no apparent anomalies on her body. Her respiratory and circulation systems functioned normally. Her abdomen was soft and flat, without any palpable masses, and her extremities were normal, in both appearance and function.

All blood data were consistent with a normal neonate at birth, and there was no evidence of jaundice, liver dysfunction, or inflammation (Table 1).

After birth, an abdominal ultrasound of the patient revealed an irregular cystic lesion that measured approximately 3 cm in diameter. The lesion was located in segments 5–8 (S₅–S₈) of the liver, had a clear border, showed posterior echo enhancement, and appeared to contain biliary sludge. We observed mild dilatation of the intrahepatic bile ducts near the porta hepatis. The cyst was adjacent to both the gallbladder and the middle hepatic vein, which was mildly displaced to the left on Doppler ultrasonography. No blood flow was detectable in the cyst, and no dilation of the extrahepatic bile duct was evident (Fig. 2).

[☆] This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/3.0/>).

All authors had access to the data and a role in writing the manuscript.

* Corresponding author. Tel.: +81 568 21 0811; fax: +81 568 22 7494.

E-mail address: skatsuno_pedsurg@world.odn.ne.jp (S. Katsuno).




Fig. 1. Maternal ultrasonography and magnetic resonance imaging before the birth of the patient. An intrahepatic or extrahepatic cystic lesion is apparent in the fetus (arrows). Hydramnios is not seen.

We first suspected a congenital biliary dilatation. However, this possibility was soon rejected for the following reasons: (1) the cyst was intrahepatic and adjacent to the middle hepatic vein, (2) no dilatation was observed in the extrahepatic bile duct, (3) the patient did not have jaundice or liver dysfunction, and (4) the patient did not have hyperamylasemia or inflammations. Accordingly, we made preparations for further examination: computed tomography (CT), upper gastrointestinal series, hepatobiliary scintigraphy, magnetic resonance cholangiopancreatography (MRCP), and percutaneous antegrade transhepatic cholangiography (PTC).

On drip infusion cholangiography CT (DIC-CT), influxes of contrast medium were observed in the cyst and the gallbladder. No other lesions were detected in the liver, and no dilation of the extrahepatic bile duct was observed. A three-dimensional DIC-CT image reconstruction displayed the enhanced cyst, located just between the bilateral hepatic ducts. The cyst had an enhanced density that was almost identical to all major components of the bile duct system, including the bilateral hepatic ducts, cystic duct, and common bile duct, except the gallbladder (Fig. 3).

Upper gastrointestinal series showed good passage through the upper gastrointestinal tract and no enlargement of the duodenal C-loop.

Hepatobiliary scintigraphy with technetium-99m Sn-N-pyr-idoxyl-5-methyltryptophan (99mTc-PMT) demonstrated normal accumulation of radioisotope in the porta hepatis, a cold area nearby and normal excretion of radioisotope to the duodenum (Fig. 4).

Table 1
Laboratory data at birth.

WBC	12,200/mm ³
RBC	390 × 10 ⁴ /mm ³
Hb	14.7 g/dl
Ht	43.7%
Plt	36.5 × 10 ⁴ /mm ³
CRP	0.01 mg/dl
TP	5.6 g/dl
Alb	3.7 g/dl
T.Bil	2.44 mg/dl
D.Bil	0.56 mg/dl
AST	26 IU/L
ALT	6 IU/L
ALP	544 IU/L
LDH	258 IU/L
AMY	10 IU/L
Lipase	7 mg/dl

MRCP demonstrated an irregular cystic mass, which had an intensity that was almost identical to the intensities of the gallbladder and the common bile duct. The connections between the cyst and bile ducts were not obvious (Fig. 5).

PTC was performed under general endotracheal anesthesia 2 months after the patient's birth. A greenish bilious fluid (7 mL) was aspirated from the cyst; it was not mucinous. Although the contrast medium sequentially revealed the cyst, the bilateral main hepatic ducts, gallbladder, common bile duct, and duodenum, some branches of the main bile ducts could not be identified. The presence of a very narrow communication between the cyst and the confluence of the bilateral hepatic ducts was confirmed by PTC (Fig. 6).

Biochemical investigation of the fluid aspirated during PTC did not indicate reflux of pancreatic juice. Instead, results indicated the presence of bile in the cyst (Table 2). The day after PTC was performed, ultrasonography revealed that the cyst had entirely recovered its original size and configuration.

Because the patient did not manifest any symptoms and blood panels did not suggest any disorders, no additional testing was performed. At present, the patient is healthy and displays normal growth and development. She has no clinical symptoms or abnormal characteristics (for example, jaundice, abdominal pain, or liver dysfunction). The cyst has not changed in size, although the patient has grown considerably, and the intrahepatic duct dilatation has already disappeared from ultrasound results. Currently, the patient is thought to have a variety of congenital hepatic cyst communicating with the bile ducts. She has been carefully followed up as an outpatient.

2. Discussion

Although congenital intraabdominal cystic lesions occasionally arise in fetuses or neonates, most of these lesions are extrahepatic and, indeed, lesions in the liver are extremely rare [1]. We suggest three possible etiologies for our case.

First, this case appears to be consistent with a solitary intrahepatic biliary cyst (SIBC). Nonparasitic liver cysts are classified into solitary or multiple cysts [2], and the solitary liver cysts (SLCs) are further categorized as simple solitary liver cysts or SIBCs [3] (Fig. 7). A few etiologies have been suggested for SLCs, namely, they may result from the obstruction of an aberrant bile duct caused by infection, vascular disruption of the fetus, or overdevelopment of the intralobular bile ducts [2–8]. In addition, Soyer et al. [2] hypothesized that insufficient drainage of the hypoplastic bile ducts accompanied by stenosis in some regions may cause dilatation of the bile pool, which results in an SIBC.

SLCs are usually asymptomatic—their symptoms do not manifest until patients are in their forties or fifties [2] and neonatal cases are extremely rare [4]. The antero-inferior segment of the right hepatic lobe is the most frequent site of SLC [2,4], which is consistent with the findings in our case. Furthermore, it is possible that SLCs develop connections to the biliary tracts.

Soyer et al. [2] reported a case of a newborn with SIBC that required an urgent laparotomy because of its progressive enlargement. They were able to demonstrate the presence of a connection between the cyst and the biliary tree on retrograde cholangiography via the common bile duct or antegrade cavigraphy; however, we found that antegrade cavigraphy could accurately determine that the cyst was near the bilateral hepatic ducts and, indeed, had a short connection with their confluence. In our case, DIC-CT could also indicate the connection. These findings suggest that SIBC may originate from the biliary system.

Alternatively, the patient's lesion may be a ciliated hepatic foregut cyst (CHFC), which is an extremely rare congenital lesion of




Fig. 2. Abdominal ultrasonography at birth. (A), (B) An irregular cystic lesion of approximately 3 cm in diameter, extending from the S₅ to S₈ of the liver with a clear border, posterior echo enhancement and sludge. (C) The cyst is located adjacent to the middle hepatic vein (MHV), which is mildly displaced to the left on Doppler ultrasonography.




Fig. 3. DIC-CT. (A), (B) Influxes of contrast medium into the cyst and the gallbladder are apparent. (C) A three-dimensional image reconstruction demonstrates the enhanced cyst just between the bilateral hepatic ducts. The cyst has an enhanced density, almost identical to all major components of the bile duct system, including the bilateral hepatic ducts, the cystic duct, and the common bile duct, except the gallbladder. GB: gallbladder; CBD: common bile duct; CD: cystic duct; RHD: right hepatic duct; LHD: left hepatic duct.




Fig. 4. Hepatobiliary scintigraphy with technetium-99m Sn-N-5-methyltryptphan demonstrates normal accumulation of radioisotope in the porta hepatis (B), a cold area nearby (A) and normal excretion of radioisotope to the duodenum (C).




Fig. 5. MRCP shows an irregular cystic mass, which has an intensity that was almost identical to the intensities of the gallbladder and the common bile duct. The connections between the cyst and the bile ducts are not evident. GB: gallbladder; CBD: common bile duct.




Fig. 6. PTC sequentially visualized the cyst, the bilateral hepatic ducts, the gallbladder, the common bile duct, and the duodenum. The presence of a very narrow communication between the cyst and the confluence of the bilateral hepatic ducts was confirmed by the PTC (arrow head). GB: gallbladder; CBD: common bile duct; Duo: duodenum.

the liver. Indeed, the findings in our case resemble CHFC. CHFC was first described by Friedreich [9] in 1857 and was given its present name by Wheeler and Edmondson [10] in 1987, based on the histopathological features of viscous or mucinous fluid contained in the cyst, pseudostratified ciliated columnar epithelium, bands of smooth muscle fibers, subepithelial connective tissue, and a fibrous external capsule.

CHFCs are believed to arise from a remnant of the embryonic foregut. At the end of the third gestational week, the ventral cranial foregut develops into the esophagus, the trachea, the bronchial and bronchiolar tree, and the lung buds. During the fourth gestational week, the ventral caudal foregut is altered into the liver

diverticulum. An abnormal bronchiolar budding could migrate into the liver through the pleuroperitoneal canal, forming a CHFC [10–13]. Therefore, the histopathological findings of CHFC are similar to those of an esophageal or a bronchogenic cyst.

Yoshida et al. [14] analyzed 80 cases of CHFC and found that the average age of presentation was 51.3 years (range, 5–82 years), the

Table 2
Biochemical data of the aspirated fluid.

T.Bil	9.4 mg/dl
D.Bil	7.21 mg/dl
AST	24 IU/L
ALT	1 IU/L
ALP	1159 IU/L
γ-GTP	1373 IU/L
AMY	1 IU/L
Lipase	13 mg/dl




Fig. 7. Classification of nonparasitic liver cyst.

male to female ratio was 1:1, and that the majority of CHFCs were localized in the left lobe (65.2%) of the liver, especially in S₄ (54.5%)—an observation that differs from the present case.

One case of surgically treated CHFC communicating with the gallbladder has been reported [15]. In our case, the lesion communicates with the hepatic duct and does not contain viscous or mucinous fluid. Although CHFC is believed to be a benign entity, three cases have been described in which a squamous cell carcinoma ultimately developed [16–18].

As an alternative to SIBC and CHFC, domestic violence may have been responsible for the lesion in our patient. Of all pregnant women, 22–35% suffer from domestic violence [19], and 17% experience blunt traumas of some kind, including motor vehicle accidents, falls, and domestic violence [20,21]. However, Baskett et al. [22] reported that the incidence of significant fetal trauma during pregnancy was rare (2.0%) and is more likely to result from labor or assisted vaginal delivery than from injuries to the mother.

Although the fetus is protected from direct force by maternal structures and amniotic fluid [23], a blunt trauma to the abdomen during pregnancy can result in intracranial hemorrhage, hemothorax, hemopericardium, low body weight, distress, or the death of fetus if the impact greatly increases the pressure in the uterus [24–26].

There have been no reports of congenital anomaly of the liver resulting from fetal injuries sustained during domestic violence. The patient's mother had suffered domestic violence perpetrated by her husband. The mother had received repetitive blows to her abdomen, even during the patient's gestational period. It is possible that these assaults induced a rupture of the hepatic duct, resulting in a biloma in the hepatic parenchyma or an intrahepatic hematoma in the fetus, which could rupture the bile duct as part of the degenerative process.

To confirm the diagnosis of our case, histopathological evaluation based on fine-needle or surgical biopsy will likely to be necessary in the future. However, except for the ultrasound findings, no abnormal test results or clinical symptoms have manifested; therefore, more invasive examinations have not been performed. Although the majority of intraabdominal cystic diseases can be surgically treated after birth, the close proximity of the cyst and the hepatic duct presents an obstacle to treatment in our case, and we are carefully considering the treatment options. If the cyst increases in size, a cysto-enterostomy with Roux-en Y anastomosis may be considered. When our patient grows up and her hepatic ducts thicken, a partial hepatectomy accompanied by suturing the fistula may be an appropriated choice of treatment. Until then, we will ensure careful long-term observation so that serious complications, such as a malignant transformation, are missed.

3. Conclusions

We presented an extremely rare case of a congenital hepatic cyst communicating with the hepatic duct, which was especially suspected of SIBC. To confirm the diagnosis, histopathological evaluation will be essential in the future. We must keep on careful long-term observation on her not to miss malignant transformation.

Conflicts of interest statement

There are no potential conflicts of interest to disclose.

Statement of informed consent

Written informed consent was obtained from the patient's parent for publication of this case report and accompanying images.

A copy of the written consent is available for review by the Editor-in-Chief of this journal on request.

Acknowledgments

We would like to show my greatest appreciation to Prof. Hisami Ando at Department of Pediatric Surgery, Nagoya University Graduate School of Medicine, whose enormous advice and insightful comments were invaluable during the course of examining the patient.

References

- [1] Foley PT, Sithasanan N, McEwing R, Lipsett J, Ford WD, Furness M. Enteric duplications presenting as antenatally detected abdominal cysts. Is delayed resection appropriate? *J Pediatr Surg* 2003;38:1810–3.
- [2] Soyer T, Karnak I, Senocak ME. Congenital solitary intrahepatic biliary cyst in a newborn: report of a case. *Surg Today* 2007;37:521–4.
- [3] Brown DK, Kimura K, Sato Y, Pringle KC, Abu-Yousef MM, Soper RT. Solitary intrahepatic biliary cyst: diagnostic and therapeutic strategy. *J Pediatr Surg* 1990;25:1248–9.
- [4] Berg C, Baschat AA, Geipel A, Krapp M, Germer U, Smrcek JM, et al. First-trimester diagnosis of fetal hepatic cyst. *Ultrasound Obstet Gynecol* 2002;19:287–9.
- [5] Macken MB, Wright JR, Lau H, Cooper MC, Grantmyre EB, Thompson DL, et al. Prenatal sonographic detection of congenital hepatic cyst in third trimester after normal second-trimester sonographic examination. *J Clin Ultrasound* 2000;28:307–10.
- [6] Howard ER. Cysts. In: Howard ER, Stringer MD, Colombani PM, editors. *Surgery of liver, bile ducts and pancreas in children*. 2nd ed. London: Arnold; 2002. p. 239.
- [7] Narasimharao KL, Chatterjee H. Congenital solitary cyst of neonatal liver. *Indian Pediatr* 1984;21:345–6.
- [8] Pitt HA. Liver cysts: non-parasitic. In: Carter D, Russell RCG, Pitt HA, editors. *Hepatobiliary and pancreatic surgery*. 5th ed. London: Chapman and Hall Medical; 1996. p. 93–7.
- [9] Friedreich N. Cyste mit timmerepithelial in der leber. *Virchows* 1857;11:466–9.
- [10] Wheeler DA, Edmondson HA. Ciliated hepatic foregut cyst. *Am J Surg Pathol* 1984;8:467–70.
- [11] Terada T, Nakamura Y, Kono N, Ueda K, Kadoya M, Matsui O. Ciliated hepatic foregut cyst. A mucus histochemical and ultrastructural study in three cases in comparison with normal bronchi and intrahepatic bile ducts. *Am J Surg Pathol* 1990;14:356–63.
- [12] Gray SW, Skandalakis JE. *Embryology for surgeons. The embryologic basis for the treatment of congenital defects*. Philadelphia, PA: W.B. Saunders Company; 1972. p. 217–383.
- [13] Chatelain D, Chailley-Heu B, Teris B, Molas G, Le Cae A, Vilgrain V, et al. The ciliated hepatic foregut cyst, an unusual bronchiolar foregut malformation: a histological, histochemical, and immunohistochemical study of 7 cases. *Hum Pathol* 2000;31(2):241–6.
- [14] Yoshida Y, Yoshida K, Daa T, Kashima K, Kondo Y, Urabe S, et al. Two cases of ciliated hepatic foregut cyst. *Shindanbyouri* 2005;22(1):36–9 [Article in Japanese].
- [15] Kolestas T, Tzioufa V, Michalopoulos A, Apostolidis S, Papadopoulos B, Hytioglou P. Ciliated hepatic foregut cyst communicating with the gallbladder. *Virchows Arch* 2005;446(2):200–1.
- [16] Fulanetto A, Dei Tos AP. Squamous cell carcinoma arising in a ciliated hepatic foregut cyst. *Virchows Arch* 2002;441:296–8.
- [17] Vick DJ, Goodman ZD, Ishak KG. Squamous cell carcinoma arising in a ciliated hepatic foregut cyst. *Arch Pathol Lab Med* 1999;123:1115–7.
- [18] De Lajarte-Thirouard AS, Rioux-Leclercq N, Boudjema K, Gandon Y, Ramée MP, Turlin B. Squamous cell carcinoma arising in a hepatic foregut cyst. *Pathol Res Pract* 2002;198:697–700.
- [19] McLee SV, Anwar R. A study of battered women presenting in an emergency department. *Am J Public Health* 1989;79:65–6.
- [20] Rudloff U. Trauma in pregnancy. *Arch Gynecol Obstet* 2007;276:101–17.
- [21] Nancy BG. Blunt trauma in pregnancy. *Am Fam Physician* 2004;70:1303–10.
- [22] Baskett TF, Allen VM, O'Connell CM, Allenn AC. Fetal trauma in term pregnancy. *Am J Obstet Gynecol* 2007;197:499e1–7.
- [23] Baker DP. Trauma in the pregnant patient. *Surg Clin North Am* 1982;62:275–89.
- [24] Lavery JP, Staten-McCormick M. Management of moderate to severe trauma in pregnancy. *Obstet Gynecol Clin North Am* 1995;22:69–90.
- [25] Hill DA, Lense JJ. Abdominal trauma in pregnant patient. *Am Fam Physician* 1996;53:1269–74.
- [26] Goldman SM, Wagner LK. Radiologic management of abdominal trauma in pregnancy. *Am J Radiol* 1996;166:763–7.