



Available online at www.sciencedirect.com



Journal of Computational and Applied Mathematics 180 (2005) 413–423

JOURNAL OF
COMPUTATIONAL AND
APPLIED MATHEMATICS

www.elsevier.com/locate/cam

A note on two identities arising from enumeration of convex polyominoes

Victor J.W. Guo^{a,*}, Jiang Zeng^{a,b}

^aCenter for Combinatorics, LPMC, Nankai University, Tianjin 300071, People's Republic of China

^bInstitut Girard Desargues, Université Claude Bernard (Lyon I), F-69622, Villeurbanne Cedex, France

Received 18 April 2004; received in revised form 4 November 2004

Abstract

Motivated by some binomial coefficients identities encountered in our approach to the enumeration of convex polyominoes, we prove some more general identities of the same type, one of which turns out to be related to a strange evaluation of ${}_3F_2$ of Gessel and Stanton.

© 2004 Elsevier B.V. All rights reserved.

MSC: 05A15; 05A19

Keywords: Binomial coefficients identities; Generating function

1. Introduction

In our elementary approach to the enumeration of convex polyominoes with an $(m+1) \times (n+1)$ minimal bounding rectangle [5], we encountered the following two interesting identities:

$$\sum_{a=1}^m \sum_{b=1}^n \binom{m+n-a+b-1}{m-a} \binom{m+n+a-b-1}{n-b} = \frac{mn}{2(m+n)} \binom{m+n}{m}^2, \quad (1)$$

* Corresponding author. Tel.: +86 222 3502 180; fax: +86 222 3509 272.

E-mail addresses: jwguo@eyou.com (V.J.W. Guo), zeng@desargues.univ-lyon1.fr (J. Zeng).

$$\begin{aligned} & \sum_{a=1}^{m-2} \sum_{b=1}^{n-2} \binom{m+n+a-b-1}{m+a+1} \binom{m+n-a+b-1}{n+b+1} \\ &= \binom{m+n}{m}^2 + \binom{m+n}{m-1} \binom{m+n}{n-1} + \frac{mn}{2(m+n)} \binom{m+n}{m}^2 - \binom{2m+2n}{2n}. \end{aligned} \quad (2)$$

Although the single-sum case of binomial coefficients identities is well-studied, the symbolic manipulation of binomial multiple-sum identities depends on the performance of computers (see, for example, [2]). Therefore, formulas of binomial double-sums are still a challenge both for human and computer.

In this paper, we will give some generalizations of the above two formulas, our main results are the following two theorems.

Theorem 1. For $m, n \in \mathbb{N}$ and any number $\alpha \neq 0$, we have

$$\begin{aligned} & \sum_{a=1}^m \sum_{b=1}^n \binom{(1+\alpha)m-a+b-1}{m-a} \binom{(1+\alpha^{-1})n+a-b-1}{n-b} \\ &= \frac{mn}{(1+\alpha)(m+\alpha^{-1}n)} \binom{(1+\alpha)m}{m} \binom{(1+\alpha^{-1})n}{n}. \end{aligned} \quad (3)$$

Theorem 2. For $m, n, r \in \mathbb{N}$ and any number $\alpha \neq 0$, we have

$$\begin{aligned} & \sum_{a=0}^{m-r-2} \sum_{b=0}^{n-r-2} \binom{(1+\alpha)m-a+b-1}{m-r-2-a} \binom{(1+\alpha^{-1})n+a-b-1}{n-r-2-b} \\ &+ \sum_{a=0}^{m+r} \sum_{b=0}^{n+r} \binom{(1+\alpha)m-a+b-1}{m+r-a} \binom{(1+\alpha^{-1})n+a-b-1}{n+r-b} \\ &= \frac{2mn}{(1+\alpha)(m+\alpha^{-1}n)} \binom{(1+\alpha)m}{m} \binom{(1+\alpha^{-1})n}{n} \\ &+ \sum_{k=-r}^r (r-|k|+1) \binom{(1+\alpha)m}{m-k} \binom{(1+\alpha^{-1})n}{n-k}. \end{aligned} \quad (4)$$

We use essentially the generating function techniques, that is, to prove $A = B$ we show that their generating functions are equal. Two proofs of Theorem 1 will be given in Section 2, and Theorem 2 will be proved in Section 3. In Section 4, we derive some interesting special cases of Theorems 1 and 2.

2. Proofs of Theorem 1

2.1. First proof of Theorem 1

Multiplying the left-hand side of (3) by $x^m y^n$ and summing over $m \geq 0$ and $n \geq 0$ we obtain the generating function $F(x, y)$, which can be written after exchanging the order of summation as:

$$F(x, y) := \sum_{a,b=1}^{\infty} x^a y^b \sum_{m,n=0}^{\infty} \binom{(1+\alpha)m+a\alpha+b-1}{m} \binom{(1+\alpha^{-1})n+a+b\alpha^{-1}-1}{n} x^m y^n.$$

Summing the two inner sums by the following classical formula (see [6, p. 146]) and [4, (9)]:

$$\sum_{n=0}^{\infty} \binom{\alpha + \beta n}{n} w^n = \frac{z^{\alpha+1}}{(1-\beta)z + \beta}, \quad \text{where } w = \frac{z-1}{z^\beta}$$

and then summing the two resulted geometric series over a and b we obtain

$$F(x, y) = \frac{uv(u-1)(v-1)}{(1+\alpha-\alpha u)(1+\alpha^{-1}-\alpha^{-1}v)(uv-u-v)^2},$$

where

$$x = \frac{u-1}{u^{1+\alpha}}, \quad y = \frac{v-1}{v^{1+\alpha^{-1}}}. \quad (5)$$

Now, using the fact that $\sum_{k \geq 0} kx^k = x/(1-x)^2$ we have

$$\begin{aligned} F(x, y) &= \frac{uv}{(1+\alpha-\alpha u)(1+\alpha^{-1}-\alpha^{-1}v)} \sum_{k=0}^{\infty} k(u-1)^k(v-1)^k \\ &= \sum_{k=0}^{\infty} \sum_{m,n=k}^{\infty} k \binom{(1+\alpha)m}{m-k} \binom{(1+\alpha^{-1})n}{n-k} x^m y^n. \end{aligned}$$

Comparing with (3), it remains to check the following identity:

$$\sum_{k=0}^{\min\{m,n\}} k \binom{(1+\alpha)m}{m-k} \binom{(1+\alpha^{-1})n}{n-k} = \frac{mn}{(1+\alpha)(m+\alpha^{-1}n)} \binom{(1+\alpha)m}{m} \binom{(1+\alpha^{-1})n}{n}. \quad (6)$$

Writing

$$\begin{aligned} k \binom{(1+\alpha)m}{m-k} \binom{(1+\alpha^{-1})n}{n-k} &= \frac{(m+\alpha^{-1}k)(n+\alpha k)}{(1+\alpha)(m+\alpha^{-1}n)} \binom{(1+\alpha)m}{m-k} \binom{(1+\alpha^{-1})n}{n-k} \\ &\quad - \frac{(m+\alpha^{-1}(k+1))(n+\alpha(k+1))}{(1+\alpha)(m+\alpha^{-1}n)} \binom{(1+\alpha)m}{m-k-1} \\ &\quad \times \binom{(1+\alpha^{-1})n}{n-k-1}, \end{aligned}$$

Eq. (6) follows by summing over k from 0 to $\min\{m, n\}$. \square

Remark. Notice that

$$\sum_{m=0}^{\infty} \binom{(1+\alpha)m}{m} x^m = \frac{u}{1+\alpha-\alpha u}, \quad \sum_{n=0}^{\infty} \binom{(1+\alpha^{-1})n}{n} x^n = \frac{v}{1+\alpha^{-1}-\alpha^{-1}v}.$$

Multiplying the two sides of (3) by $(1+\alpha)(m+\alpha^{-1}n)$, we see that (3) is equivalent to

$$\left[(1+\alpha)x \frac{\partial}{\partial x} + (1+\alpha^{-1})y \frac{\partial}{\partial y} \right] F(x, y) = xy \frac{d}{dx} \left(\frac{u}{1+\alpha-\alpha u} \right) \frac{d}{dy} \left(\frac{v}{1+\alpha^{-1}-\alpha^{-1}v} \right). \quad (7)$$

It is then possible to give another proof of (3) by checking (7), which is left to the interested reader.

2.2. Second proof of Theorem 1

Replacing b by $k + 1$ and writing the k -sum in standard hypergeometric notation we can write the left-hand side as

$$\begin{aligned} L &:= \sum_{a=1}^m \sum_{k \geq 0} \binom{(1+\alpha)m-a+k}{m-a} \binom{(1+\alpha^{-1})n+a-2-k}{\alpha^{-1}n+a-1} \\ &= \sum_{a=1}^m \binom{(1+\alpha)m-a}{m-a} \binom{(1+\alpha^{-1})n+a-2}{n-1} \\ &\quad \times {}_3F_2 \left[\begin{matrix} 1-n, 1, (1+\alpha)m+1-a \\ \alpha m+1, -(1+\alpha^{-1})n-a+2 \end{matrix}; 1 \right]. \end{aligned}$$

Applying the transformation [1, p. 142]:

$${}_3F_2 \left[\begin{matrix} -N, a, b \\ d, e \end{matrix}; 1 \right] = \frac{(e-b)_N}{(e)_N} {}_3F_2 \left[\begin{matrix} -N, b, d-a \\ d, 1+b-e-N \end{matrix}; 1 \right]$$

to the above ${}_3F_2$ we get

$$\begin{aligned} L &= \sum_{a=1}^m \binom{(1+\alpha)m-a}{m-a} \binom{(1+\alpha^{-1})n+a-2}{n-1} \frac{(-(1+\alpha)(m+\alpha^{-1}n)+1)_{n-1}}{(-(1+\alpha^{-1})n-a+2)_{n-1}} \\ &\quad \times {}_3F_2 \left[\begin{matrix} 1-n, (1+\alpha)m+1-a, \alpha m \\ \alpha m+1, (1+\alpha)m+\alpha^{-1}n+1 \end{matrix}; 1 \right]. \end{aligned}$$

Expanding the ${}_3F_2$ as a k -sum and exchanging the order with a -sum yields

$$\begin{aligned} L &= \binom{(1+\alpha)m}{m} \binom{(1+\alpha^{-1})n-2}{n-1} \frac{(-(1+\alpha)(m+\alpha^{-1}n)+1)_{n-1}}{(2-(1+\alpha^{-1})n)_{n-1}} \\ &\quad \times \sum_{k \geq 0} \frac{(1-n)_k (\alpha m)_k ((1+\alpha)m+1)_k}{(\alpha m+1)_k ((1+\alpha)m+\alpha^{-1}n+1)_k k!} \frac{m}{(1+\alpha)m+k} \times {}_2F_1 \left[\begin{matrix} 1-m, 1 \\ 1-(1+\alpha)m-k \end{matrix}; 1 \right] \end{aligned} \tag{8}$$

$$\begin{aligned} &= \binom{(1+\alpha)m}{m} \binom{(1+\alpha^{-1})n-2}{n-1} \frac{(-(1+\alpha)(m+\alpha^{-1}n)+1)_{n-1}}{(2-(1+\alpha^{-1})n)_{n-1}} \\ &\quad \times \frac{m}{\alpha m+1} {}_3F_2 \left[\begin{matrix} 1-n, \alpha m, (1+\alpha)m+1 \\ \alpha m+2, (1+\alpha)m+\alpha^{-1}n+1 \end{matrix}; 1 \right]. \end{aligned} \tag{9}$$

The theorem then follows by applying Gessel and Stanton's formula [3, (1.9)]:

$${}_3F_2 \left[\begin{matrix} -sb+s+1, b-1, -N \\ b+1, s(-N-b)-N \end{matrix}; 1 \right] = \frac{(1+s+sN)_N b(N+1)}{(1+s(b+N))_N (b+N)}$$

with $N = n - 1$, $b = \alpha m + 1$ and $s = -1 - \alpha^{-1}$. \square

Remark. If $\alpha = 1$, we can also evaluate the ${}_3F_2$ in (9) by applying Dixon's formula [1, p. 143]:

$$\begin{aligned} {}_3F_2 & \left[\begin{matrix} a, b, c \\ 1+a-b, 1+a-c \end{matrix}; 1 \right] \\ &= \frac{\Gamma(1+a-b)\Gamma(1+a-c)\Gamma(1+\frac{a}{2})\Gamma(1+\frac{a}{2}-b-c)}{\Gamma(1+a)\Gamma(1+\frac{a}{2}-b)\Gamma(1+\frac{a}{2}-c)\Gamma(1+a-b-c)}, \end{aligned}$$

and if $m = n$, we can apply Whipple's formula [1, p. 149]

$$\begin{aligned} {}_3F_2 & \left[\begin{matrix} a, 1-a, c \\ d, 1+2c-d \end{matrix}; 1 \right] \\ &= \frac{2^{1-2c}\pi\Gamma(d)\Gamma(1-2c+d)}{\Gamma(\frac{1}{2}+\frac{a}{2}+c-\frac{d}{2})\Gamma(\frac{a}{2}+\frac{d}{2})\Gamma(1-\frac{a}{2}+c-\frac{d}{2})\Gamma(\frac{1}{2}-\frac{a}{2}+\frac{d}{2})}. \end{aligned}$$

3. Proof of Theorem 2

Consider the generating function

$$G_r(x, y) := \sum_{m,n=-r}^{\infty} \sum_{a=0}^{m+r} \sum_{b=0}^{n+r} \binom{(1+\alpha)m - a + b - 1}{m+r-a} \binom{(1+\alpha^{-1})n + a - b - 1}{n+r-b} x^m y^n.$$

Using (5), as the first proof of Theorem 1, we have

$$G_r(x, y) = \frac{uv(u-1)^{-r}(v-1)^{-r}}{(1+\alpha-\alpha u)(1+\alpha^{-1}-\alpha^{-1} v)(uv-u-v)^2}. \quad (10)$$

Replacing r by $-r-2$ in (10), we obtain

$$\begin{aligned} G_{-r-2}(x, y) &:= \sum_{m,n=r+2}^{\infty} \sum_{a=0}^{m-r-2} \sum_{b=0}^{n-r-2} \binom{(1+\alpha)m - a + b - 1}{m-r-2-a} \\ &\quad \times \binom{(1+\alpha^{-1})n + a - b - 1}{n-r-2-b} x^m y^n \\ &= \frac{uv(u-1)^{r+2}(v-1)^{r+2}}{(1+\alpha-\alpha u)(1+\alpha^{-1}-\alpha^{-1} v)(uv-u-v)^2}. \end{aligned}$$

On the other hand, for $-r \leq k \leq r$, we have

$$\sum_{m,n=-r}^{\infty} \binom{(1+\alpha)m}{m-k} \binom{(1+\alpha^{-1})n}{n-k} x^m y^n = \frac{uv(u-1)^k(v-1)^k}{(1+\alpha-\alpha u)(1+\alpha^{-1}-\alpha^{-1}v)}.$$

It is routine to verify the following identity:

$$\begin{aligned} G_r(x, y) + G_{-r-2}(x, y) &= \frac{2uv(u-1)(v-1)}{(1+\alpha-\alpha u)(1+\alpha^{-1}-\alpha^{-1}v)(uv-u-v)^2} \\ &\quad + \sum_{k=-r}^r (r-|k|+1) \frac{uv(u-1)^k(v-1)^k}{(1+\alpha-\alpha u)(1+\alpha^{-1}-\alpha^{-1}v)}. \end{aligned}$$

The result then follows from Theorem 1. \square

4. Some consequences

4.1. Consequences of Theorem 1

Replacing α , m , and n by q/p , pm , and qn , respectively, in Theorem 1, we obtain

Corollary 3. *For positive integers m , n , p , and q , there holds*

$$\begin{aligned} &\sum_{a=1}^{pm} \sum_{b=1}^{qn} \binom{pm+qm-a+b-1}{pm-a} \binom{pn+qn+a-b-1}{qn-b} \\ &= \frac{pqmn}{(p+q)(m+n)} \binom{pm+qm}{pm} \binom{pn+qn}{pn}. \end{aligned}$$

Exchanging p and m , and q and n , respectively, Corollary 3 may be written as follows:

$$\begin{aligned} &\sum_{a=1}^{pm} \sum_{b=1}^{qn} \binom{pm+qm-a+b-1}{b-1} \binom{pn+qn+a-b-1}{a-1} \\ &= \frac{pqmn}{(p+q)(m+n)} \binom{pm+pn}{pm} \binom{qm+qn}{qm}. \end{aligned}$$

By the Chu–Vandermonde formula, we have

$$\begin{aligned} &\sum_{a=1-pn}^{pm} \sum_{b=1}^{qn} \binom{pm+qm-a+b-1}{pm-a} \binom{pn+qn+a-b-1}{qn-b} \\ &= \frac{pqn}{p+q} \binom{pm+qm+pn+qn}{pm+pn}. \end{aligned}$$

Therefore, by Corollary 3, we have

$$\begin{aligned} & \sum_{a=1-pn}^0 \sum_{b=1}^{qn} \binom{pm + qm - a + b - 1}{pm - a} \binom{pn + qn + a - b - 1}{qn - b} \\ &= \frac{pqn}{p+q} \binom{pm + qm + pn + qn}{pm + pn} - \frac{pqmn}{(p+q)(m+n)} \binom{pm + qm}{pm} \binom{pn + qn}{pn}. \end{aligned}$$

Replacing a by $1 - a$, we obtain

$$\begin{aligned} & \sum_{a=1}^{pn} \sum_{b=1}^{qn} \binom{pm + qm + a + b - 2}{pm + a - 1} \binom{pn + qn - a - b}{qn - b} \\ &= \frac{pqn}{p+q} \binom{pm + qm + pn + qn}{pm + pn} - \frac{pqmn}{(p+q)(m+n)} \binom{pm + qm}{pm} \binom{pn + qn}{pn}. \end{aligned}$$

Dividing both sides by $\binom{pm + qm}{pm}$, we get

$$\begin{aligned} & \sum_{a=1}^{pn} \sum_{b=1}^{qn} \frac{(pm + qm + 1)_{a+b-2}}{(pm + 1)_{a-1}(qm + 1)_{b-1}} \binom{pn + qn - a - b}{qn - b} \\ &= \frac{pqn(pm + qm + 1)_{pn+qn}}{(p+q)(pm + 1)_{pn}(qm + 1)_{qn}} - \frac{pqmn}{(p+q)(m+n)} \binom{pn + qn}{pn}. \end{aligned}$$

Replacing p, q, m, n , by $m, n, x, 1$, respectively, we have

Corollary 4. For $m, n \in \mathbb{N}$, there holds

$$\begin{aligned} & \sum_{a=1}^m \sum_{b=1}^n \frac{(mx + nx + 1)_{a+b-2}}{(mx + 1)_{a-1}(nx + 1)_{b-1}} \binom{m + n - a - b}{m - a} \\ &= \frac{mn(mx + nx + 1)_{m+n}}{(m+n)(mx + 1)_m(nx + 1)_n} - \frac{mnx}{(m+n)(1+x)} \binom{m + n}{m}. \end{aligned}$$

Letting $x \rightarrow \infty$, we obtain

Corollary 5. For $m, n \in \mathbb{N}$, there holds

$$\sum_{a=1}^m \sum_{b=1}^n \binom{m + n - a - b}{m - a} \frac{(m + n)^{a+b-2}}{m^{a-1}n^{b-1}} = \frac{(m + n)^{m+n-1}}{m^{m-1}n^{n-1}} - \frac{mn}{m+n} \binom{m + n}{m}.$$

4.2. Consequences of Theorem 2

Replacing α , m , and n by q/p , pm , and qn , respectively, in (4), we obtain

$$\begin{aligned}
& \sum_{a=1}^{pm-r-1} \sum_{b=1}^{qn-r-1} \binom{pm+qm-a+b-1}{pm-r-1-a} \binom{pn+qn+a-b-1}{qn-r-1-b} \\
& + \sum_{a=0}^{pm+r} \sum_{b=0}^{qn+r} \binom{pm+qm-a+b-1}{pm+r-a} \binom{pn+qn+a-b-1}{qn+r-b} \\
& = \frac{2pqmn}{(p+q)(m+n)} \binom{pm+qm}{pm} \binom{pn+qn}{pn} \\
& + \sum_{k=-r}^r (r-|k|+1) \binom{pm+qm}{pm-k} \binom{pn+qn}{qn-k}. \tag{11}
\end{aligned}$$

Namely,

$$\begin{aligned}
& \sum_{a=1}^{pm-r-1} \sum_{b=1}^{qn-r-1} \binom{pm+qm-a+b-1}{pm-r-1-a} \binom{pn+qn+a-b-1}{qn-r-1-b} \\
& + \sum_{a=-pm-r}^0 \sum_{b=-qn-r}^0 \binom{pm+qm+a-b-1}{pm+r+a} \binom{pn+qn-a+b-1}{qn+r+b} \\
& = \frac{2pqmn}{(p+q)(m+n)} \binom{pm+qm}{pm} \binom{pn+qn}{pn} \\
& + \sum_{k=-r}^r (r-|k|+1) \binom{pm+qm}{pm-k} \binom{pn+qn}{qn-k}. \tag{12}
\end{aligned}$$

By the Chu–Vandermonde formula, we have

$$\begin{aligned}
& \sum_{a=-pm-r}^0 \sum_{b=1}^{qm-r-1} \binom{pm+qm+a-b-1}{pm+r+a} \binom{pn+qn-a+b-1}{qn+r+b} \\
& + \sum_{a=-pm-r}^0 \sum_{b=-qn-r}^0 \binom{pm+qm+a-b-1}{pm+r+a} \binom{pn+qn-a+b-1}{qn+r+b} \\
& = \frac{(pm+r+1)q}{p+q} \binom{(p+q)(m+n)}{pm+pn} \tag{13}
\end{aligned}$$

and

$$\begin{aligned}
& \sum_{a=-pm-r}^0 \sum_{b=1}^{qm-r-1} \binom{pm+qm+a-b-1}{pm+r+a} \binom{pn+qn-a+b-1}{qn+r+b} \\
& \times \sum_{a=1}^{pn-r-1} \sum_{b=1}^{qm-r-1} \binom{pm+qm+a-b-1}{pm+r+a} \binom{pn+qn-a+b-1}{qn+r+b} \\
& = \frac{(qm-r-1)p}{p+q} \binom{(p+q)(m+n)}{pm+pn}. \tag{14}
\end{aligned}$$

Summarizing (12)–(14) and replacing $r+1$ by r , we get

Corollary 6. *For positive integers m, n, p, q , and r , there holds*

$$\begin{aligned}
& \sum_{a=1}^{pm-r} \sum_{b=1}^{qn-r} \binom{pm+qm-a+b-1}{pm-r-a} \binom{pn+qn+a-b-1}{qn-r-b} \\
& + \sum_{a=1}^{pn-r} \sum_{b=1}^{qm-r} \binom{pn+qn-a+b-1}{pn-r-a} \binom{pm+qm+a-b-1}{qm-r-b} \\
& = \frac{2pqmn}{(p+q)(m+n)} \binom{pm+qm}{pm} \binom{pn+qn}{pn} - r \binom{(p+q)(m+n)}{pm+pn} \\
& + \sum_{k=1-r}^{r-1} (r-|k|) \binom{pm+qm}{pm-k} \binom{pn+qn}{qn-k}. \tag{15}
\end{aligned}$$

And for the case r is negative, a similar formula can be deduced from (11).

For the $p=q=1$ and $m=n=1$ cases, we obtain the following two corollaries:

Corollary 7. *For positive integers m, n , and r , we have*

$$\begin{aligned}
& \sum_{a=1}^{m-r} \sum_{b=1}^{n-r} \binom{2m-a+b-1}{m-r-a} \binom{2n+a-b-1}{n-r-b} \\
& = \frac{mn}{2(m+n)} \binom{2m}{m} \binom{2n}{n} - \frac{r}{2} \binom{2m+2n}{m+n} + \frac{r}{2} \binom{2m}{m} \binom{2n}{n} \\
& + \sum_{k=1}^{r-1} (r-k) \binom{2m}{m-k} \binom{2n}{n-k}.
\end{aligned}$$

Corollary 8. For positive integers m, n , and r , we have

$$\begin{aligned} & \sum_{a=1}^{m-r} \sum_{b=1}^{n-r} \binom{m+n-a+b-1}{m-r-a} \binom{m+n+a-b-1}{n-r-b} \\ &= \frac{mn}{2(m+n)} \binom{m+n}{m}^2 - \frac{r}{2} \binom{2m+2n}{2m} + \frac{r}{2} \binom{m+n}{m}^2 \\ &+ \sum_{k=1}^{r-1} (r-k) \binom{m+n}{m-k} \binom{m+n}{n-k}. \end{aligned}$$

Furthermore, when $r = 1$ and $r = 2$ we obtain the following:

$$\begin{aligned} & \sum_{a=1}^{m-2} \sum_{b=1}^{n-2} \binom{2m-a+b-1}{m-a-2} \binom{2n+a-b-1}{n-b-2} \\ &= \binom{2m}{m} \binom{2n}{n} + \binom{2m}{m-1} \binom{2n}{n-1} + \frac{mn}{2(m+n)} \binom{2m}{m} \binom{2n}{n} - \binom{2m+2n}{m+n}, \\ & \sum_{a=1}^{m-1} \sum_{b=1}^{n-1} \binom{2m-a+b-1}{m-a-1} \binom{2n+a-b-1}{n-b-1} \\ &= \frac{1}{2} \binom{2m}{m} \binom{2n}{n} + \frac{mn}{2(m+n)} \binom{2m}{m} \binom{2n}{n} - \frac{1}{2} \binom{2m+2n}{m+n}, \\ & \sum_{a=1}^{m-1} \sum_{b=1}^{n-1} \binom{m+n-a+b-1}{m-a-1} \binom{m+n+a-b-1}{n-b-1} \\ &= \frac{1}{2} \binom{m+n}{m}^2 + \frac{mn}{2(m+n)} \binom{m+n}{m}^2 - \frac{1}{2} \binom{2m+2n}{2m}, \end{aligned}$$

and Eq. (2).

We end this paper with one more identity of the same type

Theorem 9. There holds

$$\sum_{a=1}^m \sum_{b=1}^n \binom{x+m-a+b-1}{n+b-1} \binom{x+a-b-1}{n-b} = \frac{mn}{2x+m} \binom{2x+m}{2n}.$$

Proof. Replacing x by $-x-m+n$, one sees that the theorem is equivalent to

$$2 \sum_{a=1}^m \sum_{b=1}^n \binom{x+a-1}{n+b-1} \binom{x+m-a}{n-b} = m \binom{2x+m-1}{2n-1}. \quad (16)$$

Now, changing a to $m + 1 - a$ and b to $1 - b$, respectively, we obtain

$$\sum_{a=1}^m \sum_{b=1}^n \binom{x+a-1}{n+b-1} \binom{x+m-a}{n-b} = \sum_{a=1}^m \sum_{b=1-n}^0 \binom{x+m-a}{n-b} \binom{x+a-1}{n+b-1}.$$

So we can rewrite the left-hand side of (16) as follows:

$$\begin{aligned} & \sum_{a=1}^m \sum_{b=1}^n \binom{x+a-1}{n+b-1} \binom{x+m-a}{n-b} + \sum_{a=1}^m \sum_{b=1-n}^0 \binom{x+m-a}{n-b} \binom{x+a-1}{n+b-1} \\ &= \sum_{a=1}^m \sum_{b=1-n}^n \binom{x+a-1}{n+b-1} \binom{x+m-a}{n-b} \\ &= m \binom{2x+m-1}{2n}, \end{aligned}$$

where the last step follows from Chu–Vandermonde’s formula. \square

Acknowledgements

We thank Christian Krattenthaler for his helpful comments, which, in particular, led to the second proof of Theorem 1.

References

- [1] G. Andrews, R. Askey, R. Roy, Special Functions, Encyclopedia of Mathematics and its Applications, vol. 71, Cambridge University Press, Cambridge, UK, 1999.
- [2] G. Andrews, P. Paule, Some questions concerning computer-generated proofs of a binomial double-sum identity, *J. Symbolic Comput.* 16 (1993) 147–153.
- [3] I. Gessel, D. Stanton, Strange evaluations of hypergeometric series, *SIAM. J. Math. Anal.* 13 (2) (1982) 295–308.
- [4] H.W. Gould, Some generalizations of Vandermonde’s convolution, *Amer. Math. Monthly* 63 (1956) 84–91.
- [5] V.J.W. Guo, J. Zeng, The number of convex polyominoes and the generating function of Jacobi polynomials, available at arXiv: math.CO/0403262, 2004.
- [6] G. Pólya, G. Szegö, Problems and Theorems in Analysis I, Classical in Mathematics, Springer, Berlin, 1978.