



Dicer inactivation in osteoprogenitor cells compromises fetal survival and bone formation, while excision in differentiated osteoblasts increases bone mass in the adult mouse

Tripti Gaur^a, Sadiq Hussain^a, Rajini Mudhasani^a, Isha Parulkar^a, Jennifer L. Colby^a, Dana Frederick^a, Barbara E. Kream^b, Andre J. van Wijnen^a, Janet L. Stein^a, Gary S. Stein^a, Stephen N. Jones^a, Jane B. Lian^{a,*}

^a Department of Cell Biology and Cancer Center, University of Massachusetts Medical School, Worcester, MA 01655, USA

^b Department of Medicine, University of Connecticut Health Center, Farmington, CT 06030-1850, USA

ARTICLE INFO

Article history:

Received for publication 24 June 2009

Revised 23 December 2009

Accepted 5 January 2010

Available online 15 January 2010

Keywords:

Dicer ablation

MicroRNAs

Bone formation

Col1a-Cre

Osteocalcin-Cre

High bone mass

Osteoblast differentiation

Bone development

ABSTRACT

MicroRNA attenuation of protein translation has emerged as an important regulator of mesenchymal cell differentiation into the osteoblast lineage. A compelling question is the extent to which miR biogenesis is obligatory for bone formation. Here we show conditional deletion of the *Dicer* enzyme in osteoprogenitors by *Col1a1*-Cre compromised fetal survival after E14.5. A mechanism was associated with the post-commitment stage of osteoblastogenesis, demonstrated by impaired ECM mineralization and reduced expression of mature osteoblast markers during differentiation of mesenchymal cells of ex vivo deleted *Dicer*^{c/c}. In contrast, in vivo excision of *Dicer* by *Osteocalcin*-Cre in mature osteoblasts generated a viable mouse with a perinatal phenotype of delayed bone mineralization which was resolved by 1 month. However, a second phenotype of significantly increased bone mass developed by 2 months, which continued up to 8 months in long bones and vertebrae, but not calvaria. Cortical bone width and trabecular thickness in *Dicer*^{Δoc/Δoc} was twice that of *Dicer*^{c/c} controls. Normal cell and tissue organization was observed. Expression of osteoblast and osteoclast markers demonstrated increased coupled activity of both cell types. We propose that Dicer generated miRs are essential for two periods of bone formation, to promote osteoblast differentiation before birth, and control bone accrual in the adult.

© 2010 Elsevier Inc. All rights reserved.

Introduction

For normal development of a mineralized skeleton and renewal of bone throughout life, membranous and endochondral bone formation is tightly regulated by osteogenic signaling pathways (Wnt, TGFβ/BMP, Notch) and transcription factors directing cell specification (Lian et al., 2006; Soltanoff et al., 2009). Following commitment of mesenchymal stem cells to osteoprogenitors, further differentiation requires the temporal activation of genes coding proteins responsible for forming the bone ECM, promoting mineralization and supporting the metabolic activities of osteoblasts. Gene regulatory mechanisms essential for bone formation have been elaborated in relation to epigenetic and chromatin alteration of the gene, post-transcriptional control of mRNA from splice variants and post-translational biochemical modifications of protein activity. MicroRNA (miR), are small non-coding RNA molecules that regulate gene expression through post-transcriptional degradation or translational inhibition by binding to

their target mRNAs. The regulatory function of miRs in controlling levels and/or translation of target mRNAs during in vivo bone formation has not been examined.

Only a small number of miRs that function during skeletal development and in relation to disease have been identified (Kobayashi et al., 2008; Li et al., 2009; Luzi et al., 2008; Mizuno et al., 2008a; Nakasa et al., 2008). MicroRNAs have been characterized in bone and cartilage and in association with rheumatoid arthritis synovial tissue (Kobayashi et al., 2008; Li et al., 2008b; Luzi et al., 2008; Mizuno et al., 2008b; Nakasa et al., 2008; Tuddenham et al., 2006). Significantly, two studies identified miRs which were both modulated at the onset of induced osteoblastogenesis by BMP2 in human mesenchymal stromal cells and C2C12 myogenic cells (Li et al., 2008b; Oskowitz et al., 2008). Profiling of BMP2 during bone formation in vivo (Kobayashi et al., 2008) and during stages of osteoblast differentiation in vitro (Li et al., 2009) has also implicated mature miRs in the regulation of transcripts that contribute to skeletal activities. In addition to these studies in bone forming osteoblasts, miRs regulating the bone resorbing osteoclasts are now known (Sugatani and Hruska, 2009). These findings raise a compelling question as to the requirement for miRs contributing to activation of bone formation and regulating suppression of the osteoblast phenotype in non-osseous cells. Here we addressed whether

* Corresponding author. Department of Cell Biology, University of Massachusetts Medical School, 55 Lake Avenue North, Worcester, MA 01655, USA. Fax: +1 508 856 6800.

E-mail address: jane.lian@umassmed.edu (J.B. Lian).

miRNAs are required for activation of bone formation and maintenance of osteoblast function *in vivo*.

One approach for understanding tissue specific requirements for miRNAs is the conditional deletion of the enzymes Drosha and Dicer (Bernstein et al., 2003). Drosha processes the primary miRNAs encoded in genes to 60–70 nucleotide long precursor miRNA (pre-miRNA) in the cytoplasm which are recognized and cleaved by an RNase III endonuclease protein, Dicer, to generate 21–23 nucleotide mature miRNAs that assemble into a ribonucleoprotein silencing complex (RISC) to bind to 3' untranslated region (UTR) of the target mRNAs (Jaskiewicz and Filipowicz, 2008). Dicer plays a critical role in a vast range of physiologic processes, including embryonic development, cell growth and phenotype differentiation through regulation of microRNA maturation (Blakaj and Lin, 2008). Dicer functions are essential for embryo development and survival as a Dicer-null mouse (Bernstein et al., 2003) and a hypomorphic Dicer mutant (Yang et al., 2005) results in lethality in the first week of embryogenesis. Thus, conditional ablation of Dicer activity has been studied in various tissues, demonstrating a pivotal role of Dicer-dependent processing of microRNAs for cell fate specification, differentiation and tissue morphogenesis in all the tissues examined, among which include skin follicles, heart, brain, immune cells, lung, spermatogenesis and female germline cells (Andl et al., 2006; Chen et al., 2008; Cobb et al., 2005; da Costa Martins et al., 2008; Damiani et al., 2008; Davis et al., 2008; Harris et al., 2006; Harvey et al., 2008; Hayashi et al., 2008; Kanellopoulou et al., 2005; Korolov et al., 2008; Muljo et al., 2005; Murchison et al., 2007; Nagaraja et al., 2008; O'Rourke et al., 2007; Schaefer et al., 2007; Yi et al., 2006).

Limited information is available for Dicer and microRNA requirements during skeletal development. Limb mesoderm-specific deletion of *Dicer* using *prx1*-Cre in mice resulted in massive cell death and reduction in size, but did not affect skeletal patterning (Harfe et al., 2005). However, the requirement for Dicer processed mature miRNAs in bone tissue has remained unexplored. In this study, excision of *Dicer* in mice was performed at two stages of bone formation using *Col1a1*-Cre and *Osteocalcin* (OC)-Cre which resulted in distinct phenotypes identifying different requirements for miR biogenesis. Ablation of Dicer in osteoprogenitors (using *Col1a1*-Cre for conditional deletion) prevents their differentiation and compromises fetal survival at E15.5. OC-Cre excision of *Dicer* delays perinatal bone formation which is resolved during post-natal growth. However, in *Dicer*^{Δoc/Δoc} adult mice a striking increase in both trabecular and cortical bone mass was found. This deregulated increase in bone tissue formation was accompanied by a coupled increase in bone resorption to support vascularization of the thickened bone. A contributing mechanism to this anabolic phenotype is the increased synthesis of bone matrix proteins including collagen type I, a target of several microRNAs in osteoblasts. Our studies establish that miRNAs are required to initiate osteoblast maturation during development and to regulate bone mass in adult mice.

Materials and methods

Conditional excision of *Dicer*

For conditional deletion of *Dicer*, *Dicer*^{c/c} mice (Mudhasani et al., 2008) were crossed with *Col1a1*-Cre (2.3 kb promoter of *collagen type I*) (Liu et al., 2004) and OC-Cre (*Osteocalcin*) (Chiang et al., 2009; Yuan et al., 2008) mouse line obtained from Dr. Thomas Clemens (University of Alabama at Birmingham, AL). *Dicer*^{c/c} mice were crossed with *Ink4a/Arf*^{-/-} mice for deletion of the *Ink4a/Arf* locus to obtain senescence-resistance (Serrano et al., 1996). The mice were maintained at the University of Massachusetts by IACUC approved procedures. Genotyping was carried out as previously described (Chiang et al., 2009; Liu et al., 2004; Mudhasani et al., 2008).

MicroCT (μ CT) analysis

MicroCT analysis was performed by the University of Massachusetts Medical School Musculoskeletal Center for Imaging Core facility. Bones were fixed in periodate-lysine-paraformaldehyde (PLP) fixative (Miao and Scutt, 2002) from *Dicer*^{c/c} and *Dicer*^{Δoc/Δoc} mice. After dehydration to 70% alcohol, femurs were scanned at 10 μ m voxel resolution (μ CT 40; Scanco Medical AG, Brüttisellen, Wangen-Brüttisellen, Switzerland). Image reconstruction was performed by Scanco software version 5.0. For trabecular bone 100 contiguous slices below the growth plate were selected for contouring inside the endosteal edge for analyses of various bone parameters. Cortical parameters were analyzed from 50 cross-sectional slices at the mid-diaphysis region. Parameters were obtained using threshold range 220–1000.

Skeletal staining and histology

Whole skeletal staining was performed by standard procedures using Alcian blue and Alizarin red stains for cartilage and bone tissue (Lufkin et al., 1992). Separate embryos, day 2 pups of femurs from 2 mo old mice were fixed in PLP, paraffin-embedded, and sequential staining of sections was performed by von Kossa stain for mineral and Toluidine blue for cellular detail. Cytochemical detection of bone-specific alkaline phosphatase (Alk Phos) and osteoclast-specific tartrate resistant acid phosphatase (TRAP) was performed (Lengner et al., 2004). Images were captured using a Zeiss Axioskop 40 (Mikron, San Marcos, CA, USA) microscope with a CCD camera.

Ex vivo osteoblast differentiation

Bone marrow stromal cells (BMSC) were isolated from *Dicer*^{c/c} mice in α MEM supplemented with 20% fetal bovine serum (Hyclone, Logan, UT, USA). Cells were transduced at 60% confluency with Ad-Cre or Ad-GFP (control virus) for 3 h for excision of *Dicer*, then cultured in α MEM with 50 μ g/ml ascorbic acid and 10 mM β -glycerol phosphate (SIGMA-Aldrich, St. Louis, MO, USA) to induce the osteoblast differentiation. Calvarial osteoblasts were isolated from newborn *Dicer*^{Δoc/Δoc} by collagenase P digestion, genotyped and cultured for osteoblast differentiation (Pratap et al., 2003). Cells were harvested at indicated time points for Alk Phos activity, mineral staining, and gene expression analyses. Staining for senescence-associated β -Galactosidase (SA- β Gal) was performed by a standard procedure (Dimri et al., 1995). Excision of *Dicer* was detected by PCR with forward (5'-CCGACCAGCCTTGT-TACCTG-3') and reverse primers (5'-CGGTGTTTCCTTGAATACTT-3') using GAPDH as internal control (Applied Biosystems). Experiments were repeated at least twice with similar results.

RNA isolation and quantitative real time PCR

Cells were harvested in 300 μ l TRIzol reagent (Invitrogen, Carlsbad, California, USA). Total RNA isolated as per the manufacturer's instructions (Invitrogen, Carlsbad, California, USA) and treated with RNase-free DNase. The reverse transcription reaction was performed on 1 μ g of RNA using the First Strand Synthesis Kit (Invitrogen, Carlsbad, California, USA). Relative transcript levels were measured by quantitative PCR in 25 μ l reaction volume using ABI PRISM 7300 sequence detection system (Applied Biosystems, Foster City, CA, USA), following the recommended protocol for SYBR-Green, and normalized with GAPDH levels (Applied Biosystems, Foster City, CA, USA). The primers used for amplification are described in Table 1.

For detection of let-7a and miR-29b, mirVANA qRT-PCR miRNA detection kit along with primer sets for each microRNA (Applied Biosystems/Ambion, Foster City, CA, USA) were used, following the manufacturer's procedure. The miR levels were normalized using U6 primers.

Table 1

List of primers used for quantitative real time PCR.

Gene	Primer sequence
<i>Runx2</i>	For 5' CGCCCTCCCTGAACCT 3' Rev 5' TGCTGCCTGGGATCTGTA 3'
<i>Alkaline phosphatase</i>	For 5' TGTGCGAGAGAAAGGAGA 3' Rev 5' GTTTCAGGGCAATTTTCAAGGT 3'
<i>Osteocalcin</i>	For 5' CTGACAAAGCCTTCATGTCCAA 3' Rev 5' GCGCCGGAGTCTGTCTACTA 3'
<i>Collagen1a1</i>	For 5' CCAAGGAAAAGAGCACGTC 3' Rev 5' AGGTCAGCTGGATAGCGACATC 3'
<i>Collagen1a2</i>	For 5' GTCTAGTCGATGGCTGCTC 3' Rev 5' CAATGTCAGAGGTGCAATG 3'
<i>Collagen3a1</i>	For 5' AGGCCAGTGCCAATGTAAAG 3' Rev 5' CTCCATTCCCAGTGTGTT 3'
<i>Collagn5a3</i>	For 5' AGGGACCAACTGGGAAGAGT 3' Rev 5' AAAAGTCAGAGGCAGCCACAT 3'
<i>Bone sialoprotein</i>	For 5' GCACTCCAAGTCCCAAGA 3' Rev 5' TTTTGGAGCCCTGCTTCTG 3'
<i>Osteopontin</i>	For 5' ACTCCAATGTCCTACAGTCG 3' Rev 5' TGAGGTCCTCATCTGTGGCAT 3'
<i>TRAP</i>	ABI mm00475698_m1
<i>RANKL</i>	ABI mm00437135_m1
<i>Dicer</i>	For 5' GGTGGTTCGTTTIGATTGGC 3' Rev 5' GGCAGTGTGATTGTGACTC 3'
<i>p21</i>	For 5' TTGCACTCTGGTGTCTGAGC 3' Rev 5' TCTGCGCTGGAGTGATAGA 3'

Results

In vivo deletion of *Dicer* in hypertrophic chondrocytes and osteoblast lineage cells induces embryonic lethality

To establish the role of *Dicer*-dependent miRNAs in bone growth and differentiation, we used a *Dicer* conditional mouse model (*Dicer^{c/c}*) in which Cre-mediated excision leads to deletion of the *Dicer* PAZ domain (Mudhasani et al., 2008). The *collagen type 1* (2.3 kb) promoter driving Cre recombinase (*Col1a1-Cre*) which is robustly expressed in committed osteoprogenitors, osteoblasts and hypertrophic chondrocytes (Liu et al., 2004) was used to address *Dicer* functions during skeletal development. Because no live mice with homozygous deletion of *Dicer* were recovered at birth, embryos were examined for skeletal deformities. At E14.5 Cre-positive *Dicer^{Δcol1/Δcol1}* fetal pups represented 24% of the total population of 49 embryos from 8 litters and were smaller but viable (Fig. 1A, upper panels). Skeletal staining, however, revealed a deformed cartilage skeleton without bone tissue in *Dicer^{Δcol1/Δcol1}*, whereas *Dicer^{c/c}* embryos showed early bone formation (Fig. 1A, lower panels). E15.5 *Dicer^{Δcol1/Δcol1}* pups represented 23% of the population but were partly resorbed (Fig. 1B). Thus, after E14.5 fetal survival is compromised by *Dicer* excision.

We analyzed cellular details of *Dicer^{c/c}* and *Dicer^{Δcol1/Δcol1}* fetal pups to identify specific functions of *Dicer* during embryonic bone development. In control mice at E14.5 (Fig. 1C) intramembranous and endochondral bone formation was detected by mineral deposition and alkaline phosphatase (Alk Phos) positive osteoblasts and hypertrophic chondrocytes in the mandible, clavicles and ribs (Fig. 1D, 1–4). In *Dicer^{Δcol1/Δcol1}* embryos, bone tissue was either completely absent or significantly reduced as reflected by fewer Alk Phos positive cells.

Mineralized tissue formation appears to be compromised as no von Kossa staining occurred in cartilaginous or osseous tissues (Fig. 1D, 5–8). Thus, *Dicer* ablation by *Col1a1-Cre* inhibits further differentiation of




Fig. 1. *In vivo* excision of *Dicer* by *Col1a1-Cre* impairs bone formation at E14.5 and induces embryonic lethality. (A) Skeletal deformities observed at E14.5. Upper panel shows smaller but viable *Dicer^{Δcol1/Δcol1}* embryos at E14.5. Alcian blue/Alizarin red staining (lower panel) reveals lack of mineralizing clavicles (see arrow in *Dicer^{c/c}* and *Dicer^{wt/Δcol1}*) and a deformed cartilaginous skeleton in *Dicer^{Δcol1/Δcol1}*. (B) Images of E15.5 fetal pups show partial resorption of *Dicer^{Δcol1/Δcol1}* pup. (C) Embryo sections are compared at E14.5 and (D) detailed (boxed areas) at higher magnification for *Dicer^{c/c}* (1–4) and *Dicer^{Δcol1/Δcol1}* (5–8), showing severely impaired mineralization of mandible mesenchyme that surrounds Meckel's cartilage (1, 5), clavicle (2, 6) and reduced Alk Phos staining in the lower jaw (3, 7) and ribs (4, 8) of *Dicer^{Δcol1/Δcol1}* embryos.

committed osteoblasts between E14.5 and E15.5, when there is a rapid induction of mineralizing elements in control mice.

Fetal lethality at the onset of bone formation that results from lack of *Dicer* in *Col1a1* expressing cells may be due in part because a marrow cavity, which supports hematopoiesis, is not formed. Alternatively, bone formation may only be delayed in the absence of *Dicer*; however, lethality at E15.5 does not allow this assessment. We conclude that loss of miR processing at E14.5 prevents maturation to hypertrophic chondrocytes and osteoblasts that results in failure to produce mineralized matrices.

Ex vivo deletion of Dicer abrogates miR processing and results in defective osteoblast differentiation

To identify the stage of osteoblastogenesis blocked in *Dicer*^{Δcol1/Δcol1} mice, bone marrow stromal cells (BMSCs) from *Dicer*^{c/c} mice at 8 weeks were differentiated into the osteoblast lineage. Following transduction of BMSCs by adenovirus Cre (Ad-Cre), the *Dicer* locus showed >90% excision (Fig. 2A). To confirm loss of miRs, we examined miR-29b and let-7a expression (Fig. 2B), which are known to influence osteoblast differentiation and tissue formation (Harfe et al., 2005; Li et al., 2009). We find that in differentiating control Ad-GFP cells, miR-29b and let-7a are upregulated at the matrix maturation stage (day 19) and down-regulated during mineralization (day 27). In *Dicer*-ablated cells, let-7a levels were suppressed throughout differentiation (Fig. 2B), indicating loss of miRs in the absence of *Dicer*. No difference in cellularity was observed between Ad-GFP and Ad-Cre treated cells from initial plating to confluency, as shown by Toluidine blue staining at day 5 (Fig. 2C) and normal expression of histone H4 RNA (Supplementary Fig. S1A). Also, pro-apoptotic Bax and anti-apoptotic Bcl2 were expressed at comparable levels in both Ad-GFP control and Ad-Cre treated samples (Supplementary Fig. S1B) suggesting normal growth and survival properties in absence of *Dicer* in BMSCs. While control cells exhibited the normal differentiation profile of increased Alk Phos and ECM mineralization (Fig. 2C, upper panels), BMSCs lacking *Dicer* activity (Ad-Cre) showed markedly reduced Alk Phos activity and mineral deposition as early as day 12, reflecting failed maturation (Fig. 2C, lower panels).

The mechanism of inhibited differentiation was interrogated by examining expression of genes that contribute to osteoblast maturation. *Runx2*, the early marker of a committed osteoblast, and *Col1a1* are equivalently expressed between Ad-GFP and Ad-Cre cultures at day 12 and 19 (for *Col1a1*). However, neither *Runx2* nor *Col1a1* is further upregulated in Ad-Cre cells as differentiation progresses (Fig. 2D). Markers related to mineral deposition, *Alk Phos*, *Osteopontin* (*OP*), *Osteocalcin* and *Bone sialoprotein* (*BSP*) were not induced in *Dicer*-ablated cells (Fig. 2D). Our results show that osteoblast defects occur after cells are committed to the bone phenotype by *Runx2*; thus, evidence is provided that commitment and proliferative expansion of osteoprogenitors are not affected by *Dicer* excision in BMSCs. These findings suggest that a group of miRs processed by *Dicer* are required for maturation to the fully differentiated osteoblast phenotype.

Absence of Dicer induces differentiation defects independent of cell senescence

Recent reports indicate an increase in senescence of mouse embryonic fibroblasts upon *Dicer* deletion (Mudhasani et al., 2008). To explore the possibility of a similar secondary effect in BMSCs, we performed senescence-associated β-Gal staining on growing cells after *Dicer* excision by Ad-Cre. Increased senescence was observed in *Dicer*-ablated cells compared to control cells by histochemical staining (Fig. 3A, left panel) and confirmed by 2.4 fold higher expression of p21, a marker of senescence (Fig. 3A, right panel). To exclude a role for cell senescence in reduced osteogenesis of *Dicer*-ablated cells, BMSCs were obtained from *Dicer*^{c/c}; *Ink4a/Arf*^{-/-} mice for osteoblast differentiation following treatment with Ad-GFP (control) and Ad-Cre.

Analyses were performed on day 7 post infection. An 80% reduction of *Dicer* mRNA was found in BMSCs infected with Ad-Cre (Fig. 3B). Also, p21 expression by qRT-PCR confirmed that osteoblast cell senescence is not affected by *Dicer* excision in BMSCs from senescence-resistant mice (Fig. 3B). *Dicer*-ablated BMSCs could not undergo matrix maturation and mineralization based on inhibited Alk Phos, mineral deposition, and *OC* mRNA levels (Fig. 3B). Thus, removal of senescence did not rescue *Dicer*-induced differentiation defects. This strengthens the conclusion that defects in osteoblast differentiation are not secondary to premature cell death, but are directly related to absence of miRs.

Deletion of Dicer in osteoblasts and osteocytes delays perinatal bone formation

Osteocalcin is highly expressed in mature surface osteoblasts and osteocytes. To further explore in vivo mechanisms for *Dicer* control of bone formation during mineralization, we inactivated *Dicer* by *OC*-Cre which is robustly expressed in mature osteoblasts in a matrix undergoing active deposition of mineral (Chiang et al., 2009; Yuan et al., 2008). *Dicer*^{Δoc/Δoc} mice were viable, with no apparent skeletal defects at birth by skeletal staining and radiography (data not shown). Femur histology (day 2 mice) showed normal cellular organization of the growth plate and bone formation (data not shown). However, we observed a delay in endochondral bone formation in the tail vertebrae, which are the last bones to develop. *Dicer*^{Δoc/Δoc} vertebrae had fewer hypertrophic chondrocytes and weaker Alk Phos activity in the bone collar compared to *Dicer*^{c/c} (Fig. 4A, 1, 2). Therefore, we examined the fetal skeleton at E17.5 and found that the craniofacial region and ribs displayed strikingly reduced Alk Phos activity in *Dicer*^{Δoc/Δoc} (Fig. 4A, 3–6). These results suggest delayed bone formation and an inherent defect in osteoblastogenesis.

Cellular mechanisms for *Dicer*^{Δoc/Δoc} induced bone defects were examined by *ex vivo* differentiation of calvarial osteoblasts from newborn mice. A severe reduction in Alk Phos activity was observed after day 14, while control *Dicer*^{c/c} calvarial cells reached peak levels in activity and mRNA (Figs. 4B, C). Significantly, *Dicer*^{Δoc/Δoc} cells showed no induction of *Runx2* and *Osterix*, two transcription factors essential for further differentiation. In contrast, expression of *Col1a1*, the major bone protein was upregulated in *Dicer*-ablated cells (Fig. 4C), consistent with the decrease in let-7a (Fig. 4C) which targets collagens (Li et al., 2009). A block in osteoblast differentiation occurs at the mineralization stage as reflected by lack of *OC* expression after day 14 (Fig. 4C). All the markers except *Col1a1* were at normal levels in proliferating cells, when *OC* promoter activity driving Cre expression is at low basal levels. These findings are analogous to *ex vivo* excision of *Dicer*^{c/c} BMSCs (Fig. 2). Thus, mechanisms controlled by *Dicer* in mature osteoblasts support post-transcriptional regulation of target genes through miRs that are essential for mineralization of the bone ECM. Such targets would include well-characterized enzymes and matrix proteins (Li et al., 2009).

Dicer excision in mature osteoblasts/osteocytes increases bone mass in adult mice

Mice after birth were continuously monitored up to 8 months and exhibited similar growth rates (data not shown). Unlike the severe defects observed at late gestation in *Dicer*^{Δoc/Δoc} mice, radiography revealed that bone formation after birth (post-natal day 2 and 4 weeks age) was normal (data not shown). However, at 2 months of age increased radio-opacity was observed in femurs, spine and tail vertebrae in *Dicer*^{Δoc/Δoc} compared to the *Dicer*^{c/c} mice (Supplementary Fig. S2). Cortical bone width increased and trabecular bone of the vertebrae became very dense. Interestingly, no differences could be observed in radiographs of calvarial bones at 2 months of age between




Fig. 2. Contributing mechanisms for defective osteoblast differentiation in the absence of Dicer. Dicer excision in BMSCs from *Dicer^{+/c}* mice by Ad-Cre, followed by differentiation into osteoblasts. Ad-GFP was used as control for transduction of cells. (A) Genomic PCR of Ad-GFP (control) and Ad-Cre treated cells identified >90% excision of *Dicer* locus on day 5 (*GAPDH*, internal control). (B) Lack of miR-29b (red lines) and let-7a (blue lines) expression in the absence of Dicer. Solid lines (—) show control Ad-GFP treated cells; dashed lines (---), Ad-Cre Dicer-ablated cells. (C) Cellularity (Tol blue) was not affected by *Dicer* excision, but differentiation (Alk Phos activity) and mineral (von Kossa stain) were severely impaired. (D) Expression patterns of early markers of committed osteoprogenitors, *Runx2* and *Col1a1*, and components for the onset and regulation of mineralization by the non-collagenous proteins, OC, BSP and OP, reflect impaired maturation of osteoblasts in *Dicer*-excised (Ad-Cre) cells.

Dicer^{Δoc/Δoc} and *Dicer^{c/c}* mice (Supplementary Fig. S2). Hence, calvariae were not characterized further by μ CT for detailed analyses.

At 1, 2, 4 and 8 months of age, cortical and trabecular bone structure of femurs was evaluated by μ CT analysis (Fig. 5). Single sagittal sections show increasing trabecular thickness (Fig. 5A) and widened cortex (Fig. 5C) from 2 to 8 months in *Dicer^{Δoc/Δoc}* compared to *Dicer^{c/c}*.

The trabeculae bone volume (BV/TV) indicates at 1 month age, an 11.4% significant decrease compared to controls (Fig. 5B), but at 2 months BV/TV was not significantly different. At 2 months, trabecular parameters were beginning to change and during the next 2 months of growth, a 5-fold increase in BV/TV was observed. Quantitative parameters specific for trabecular bone showed significant increases in




Fig. 3. *Dicer* excision-induced defects in osteoblast differentiation occur in the cellular senescence-resistant *Dicer*^{c/c}; *Ink4a/Arf*^{-/-} mice. (A) SA-βGal staining of control (Ad-GFP) or *Dicer*-ablated (Ad-Cre) BMSCs at day 5 shows increased senescence in the absence of *Dicer*. qRT-PCR analysis shows the cellular senescence marker p21. (B) BMSCs from *Dicer*^{c/c}; *Ink4a/Arf*^{-/-} mice were transduced with Ad-Cre, and cultured for osteogenesis. qRT-PCR shows reduced *Dicer* mRNA after Ad-Cre-mediated excision, and p21 expression shows rescue of premature cell senescence in cells lacking *Dicer*. Alk Phos/von Kossa staining identifies reduced osteoblast differentiation from day 14 to 28 in absence of *Dicer*. qRT-PCR analysis of mRNAs at days 7–21. Solid lines (—) show control Ad-GFP treated cells; dashed lines (---), Ad-Cre *Dicer*-ablated cells. The data are presented as mean ± SD, n = 3.

the thickness (53%) and number (25%) in *Dicer*^{Δoc/Δoc} at 4 months and consistent with other parameters (Fig. 5B and Supplementary Fig. S3A).


Evidence of increased cortical bone mass was found earlier at 1 month in *Dicer*^{Δoc/Δoc} (29% higher cortical bone area) with decreased cortical porosity through 4 months, although not significantly changed from *Dicer*^{c/c} at 8 months age (Figs. 5C, D). Other cortical parameters reflected maintenance of high bone mass in the cortex (Supplementary Fig. S4). Notably the periosteal area was more active than the endosteal surface of the cortex in *Dicer*^{Δoc/Δoc} (Supplementary Fig. 4S) which accounted for normal marrow space. However, at 8 months the endocortical surface had a greater response in contributing to bone formation than at earlier ages (Supplementary Fig. S4). Overall, at 8 months when the control *Dicer*^{c/c} mice have decreased trabecular and cortical bone mass, *Dicer*^{Δoc/Δoc} continued to accumulate cortical bone from 4 to 8 months (Fig. 5C).

In summary, the delayed bone formation phenotype observed in *Dicer*^{Δoc/Δoc} neonates was resolved during post-natal growth and reversed at puberty age (8 weeks). This reversal of the phenotype

documented in long bone, lumbar and tail vertebrae is well underway from 1 to 2 months, stabilized by 4 months and continues to increase bone mass up to 8 months in *Dicer*^{Δoc/Δoc}. Although a significantly higher accumulation of bone tissue formed by the endochondral process of bone formation and the periosteal surface of long bone, membranous calvarial bone was spared from excessive bone formation.

Increased bone mass is represented by normal cellular and tissue organization

To identify the possible changes in cellular organization of bone tissue caused by high bone mass phenotype in absence of *Dicer* in mature osteoblasts, we performed histological and molecular studies of the femurs at 2 month age *Dicer*^{c/c} and *Dicer*^{Δoc/Δoc} mice when the phenotype is reversed. The Toluidine blue stained sections of cortical bone (femur mid-shaft region shown in Fig. 6A, left panels) reveals individual osteocytes in their lacunae with blood vessel channels throughout in both groups, but thicker bone in *Dicer*^{Δoc/Δoc} mice



which appears to have 3 zones of accumulated bone. The most mature bone (deposited from birth), which takes up less Toluidine blue stain, is on the endosteal surface, while the middle zone represents bone accumulated during rapid post-natal growth and is seen as vascularized. The third layer of recently laid down bone formed from the periosteal surface has rows of osteocytes in individual lacunae with fewer blood vessel channels. This organization of new growth more from the periosteum is supported by μ CT parameters (Supplementary Fig. S4). Histomorphometric quantitation of osteocytes in the cortical bone ($n=4$ fields of bone along the cortex in $n=3$ mice) indicates that while the total number of osteocytes is 2-fold (data not shown) higher because of increased bone volume, the osteocyte density per unit area is similar in $Dicer^{c/c}$ and $Dicer^{\Delta oc/\Delta oc}$ (Fig. 6A, right panel).

We had previously identified (Li et al., 2009) that a significant number of microRNAs are upregulated from days 21 to 28, the mineralization stage when the ECM is mineralized and cells are transitioned from osteoblast to osteocytes in MC3T3 cells (Sudo et al., 1983). At least 6 upregulated microRNAs, based on bioinformatic studies, target collagen type I that constitutes greater than 90% of adult bone ECM. Therefore we tested the hypothesis that a contributing mechanism to the observed phenotype is a stimulated production of type I collagen that accelerated the transition to osteocytes. Total cellular RNA was prepared from long bone flushed of marrow at 4 months when the phenotype of $Dicer^{\Delta oc/\Delta oc}$ was fully developed. qRT-PCR analyses of multiple collagen transcripts at 4 months of age revealed that the expression of bone-specific collagens (Col1a1, Col1a2 and Col5) were upregulated; whereas Col3a1, known to be expressed in immature bone, is at low and equivalent levels in $Dicer^{\Delta oc/\Delta oc}$ and $Dicer^{c/c}$ mice (Fig. 6B), indicating an involvement of collagen proteins as part of the mechanism for high bone mass in $Dicer^{\Delta oc/\Delta oc}$ mice. Although the Runx2 transcription was not significantly different between the two groups (data not shown), reflecting that commitment to the osteoblast lineage is not enhanced, the Osterix transcriptional regulator of later differentiation and osteocalcin reflecting a mineralized matrix are increased by >30% in $Dicer^{\Delta oc/\Delta oc}$ mice. We also examined expression of *Sclerostin* mRNA levels, a specific product of osteocytes that is an inhibitor of Wnt signaling positive effects on bone formation (ten Dijke et al., 2008). However, we find no statistical difference between $Dicer^{c/c}$ and $Dicer^{\Delta oc/\Delta oc}$ mice (data not shown), further indicating that osteocyte function is normal.

Several high bone mass (HBM) mouse models, such as the LRP5 activating mutation and *Sclerostin* null mice display an uncoupling of bone turnover which leads to increased bone tissue deposition (Babji et al., 2003; Li et al., 2008a). To examine this possibility in $Dicer^{\Delta oc/\Delta oc}$ mice, osteoclast-specific tartrate resistant acid phosphatase (TRAP) staining was performed on bone sections from 2 month-old mice (Fig. 6C). Overall, there is an increase in the number of osteoblasts in the metaphyses because there is an increase in bone trabecular surfaces. This finding suggests that bone remodeling is coupled to ensure resorption of calcified cartilage in the primary spongiosa and conversion to lamellar from newly secreted woven bone. Furthermore, increased osteoclasts are observed in cutting cones in cortical bone of $Dicer^{\Delta oc/\Delta oc}$ mice to vascularize the thickened tissue (Fig. 6D). Consistent with these observations, the mRNA levels of TRAP (originating from osteoclast) and RANKL, the osteoblast coupling factor for osteoclast differentiation, were both upregulated in bone tissues from $Dicer^{\Delta oc/\Delta oc}$ mice (Fig. 6E). These findings indicate that the phenotype of $Dicer^{\Delta oc/\Delta oc}$ mice is not because of a reduction in bone resorption. Rather, a coupling of increased bone remodeling to increased bone formation is revealed.

Discussion

Control of mesenchymal differentiation is an intricate process requiring the interplay of cell signaling, transcription, and post-transcriptional mechanisms. Our studies now show that miR-mediated control of gene expression is critical for osteogenesis and bone homeostasis. This conclusion is based on several key findings that establish functions of Dicer in bone tissue. First, we have identified the Dicer enzyme as essential for membranous and endochondral bone formation in mid-stage embryos (E14.5). Second, deletion of *Dicer* by *OC-Cre* in mature osteoblasts delayed bone formation before birth. However, this phenotype was reversed post-natally and adult mice showed a dramatic increase in bone mass compared to controls. Third, in the $Dicer^{\Delta oc/\Delta oc}$ post-natal skeleton a continual accrual of trabecular and cortical bone occurs up to age 8 months with normal bone remodeling such that the thickened tissue is vascularized. Fourth, *ex vivo* studies indicate that the cellular mechanisms which are impaired by Dicer deletion in osteoprogenitor cells are related to post-proliferative differentiation of committed osteoblasts. We conclude from these observations that Dicer function is rate-limiting for the processing of specific miRs at distinct developmental stages of bone tissue formation in long bones and vertebral bodies. In summary, our findings suggest that Dicer processes a distinct set of miRs in the fetal skeleton necessary for osteogenesis, whereas in the adult skeleton miRs present in mature osteoblasts are potent negative regulators of excessive bone formation. Thus, Dicer produced mature miRs in osteoblast lineage cells have potent activities in regulating cell determination, differentiation and tissue homeostasis and exhibit a specificity for specific anatomical bone tissues.

Mechanisms contributing to the phenotype of delayed osteoblast differentiation before birth were revealed by *ex vivo* characterization of defective cell differentiation along the osteoblast lineage. We find that the critical stage for which miRs are required in mesenchymal cells occurs early in the osteoblast maturation sequence, shortly after commitment to the osteoblast lineage. Neonatal calvarial cells from $Dicer^{\Delta oc/\Delta oc}$ exhibited a block in osteoblast maturation, consistent with *in vitro* Dicer-ablated bone marrow derived osteoprogenitors from $Dicer^{c/c}$ mice. In both cell models, Runx2 was found at equivalent levels prior to differentiation; and, proliferation and multi-layering of cells are not compromised, as Alk Phos positive bone nodules are present throughout the culture. However, the nodules are defective in their ability to mineralize, in part compromised by further induction of Alk Phos and absence of non-collagenous proteins (bone sialoprotein and osteocalcin) that are normally induced in post-proliferative osteoblasts to support mineralization of the bone ECM (Alford and Hankenson, 2006; Malaval et al., 2008; Robey and Boskey, 2008). These findings suggest that a critical set of miRs must be expressed in post-proliferative osteoblasts to indirectly activate bone-related proteins required to produce a mineralizing ECM.

While inhibition of osteoblast differentiation is similar in *ex vivo* cultured cells from our two mouse models, the striking differences in the *in vivo* phenotypes can be explained by the distinct expression patterns of the promoters used in our studies. The *Col1a1* promoter is active in pre-osteoblasts, the inner cell layer of the periosteum that will become a bone collar (Liu et al., 2004). Thus, Dicer ablation by *Col1a1-Cre* between E14.5 and E15.5, when there is a rapid induction of mineralizing elements in control mice, compromises differentiation of osteoprogenitor cells exhibiting Dicer loss-of-function. *Osteocalcin-Cre* mice are well documented for robust activity in the post-natal mouse (Chiang et al., 2009; Yuan et al., 2008). There are distinct

Fig. 4. *In vivo* excision of *Dicer* by *OC-Cre* causes perinatal skeletal deformities that differ from abnormalities in post-natal mice. (A) Delayed osteoblast differentiation comparing $Dicer^{c/c}$ (1, 3, 5) and $Dicer^{\Delta oc/\Delta oc}$ (2, 4, 6). Alk Phos/Methyl Green staining in newborn caudal vertebrae (1, 2); E17.5 craniofacial regions (3, 4) and ribs (5, 6). (B) Reduced *ex vivo* differentiation of Dicer-ablated osteoblasts (right panel) revealed by Alk Phos (left panel). (C) Mechanism of defective $Dicer^{\Delta oc/\Delta oc}$ osteoblast maturation. Reduced osteoblast markers *Runx2*, *Osterix*, *Alk Phos* and *OC* after the proliferation period (day 7). Reduced *let-7a* expression and consequent upregulation of *Col1a1* confirm the loss of Dicer processing.




Fig. 5. High bone mass phenotype in adult mouse caused by *Dicer* excision by OC-Cre. (A) Femur mid-bone sagittal μ CT sections show mineralized trabecular and cortical bone at the indicated ages. (B) Quantitation of trabecular bone parameters. Reduced trabecular bone in femurs of 1 month *Dicer*^{Δoc/Δoc} (%BV/TV) is significantly normalized by 2 months and significantly increased at 4 and 8 months compared to control *Dicer*^{c/c} bone. (C) Transverse scan of femurs at mid-diaphysis. (D) Quantitation of cortical bone parameters shows significant increase in cortical bone area (a volume parameter) with decreased porosity, especially at 2, 4 and 8 months in *Dicer*^{Δoc/Δoc} mice. Values are mean \pm SD ($n = 3$ mice), *** p -value < 0.001 , ** p -value < 0.005 , * p -value < 0.05 .




Fig. 6. Normal bone content in *Dicer^{Δoc/Δoc}* adult mice. (A) Left panels—Toluidine blue stained sections of mid-shaft cortical bone from control and *Dicer^{Δoc/Δoc}* femurs at age 2 months. Right panel—Normal density of osteocytes by osteocyte counts at 2 months of age. (B) Total cellular RNA prepared from 4 month age tibia bone ($n = 4$ mice per group) for gene expression of bone-specific collagen (*Col1a1*, *Col1a2* and *Col5a3*) and the immature collagen (*Col3a1*). (C) Increased osteoclast activity coupled to increased bone formation shown by TRAP staining of whole femur at 2 month and (D) magnified trabeculae showing TRAP positive cells. (E) Expression of *TRAP* and *RANKL* mRNA by qRT-PCR. Values are mean \pm SD ($n = 3$ mice), * p -value < 0.05 .

differences between the pre- and post-natal bone phenotype in the *Dicer^{Δoc/Δoc}* mice. Before birth, bone formation is delayed, and ex vivo osteoprogenitor cells from these mice exhibit a similar phenotype as ex vivo *Dicer* excision of bone marrow mesenchymal cells, both defective in differentiation to mature osteoblasts. The histology of bone from E17.5 shows few mature osteoblasts, and suggests that the immature cells expressing low basal *osteocalcin* may have different functioning miRs than those expressed in mature osteoblasts in a

mineralized matrix after birth. In contrast, adult mice showed increased bone accumulation in the appendicular skeleton and spine, and with a normal cellular organization.

A high bone mass phenotype in *Dicer^{Δoc/Δoc}* was evident after weaning initially in tail vertebrae and cortical bone. At 4 weeks age, cortical bone in *Dicer^{Δoc/Δoc}* increases prior to trabecular which becomes normal relative to *Dicer^{c/c}*. This phenotype of decreased trabecular bone, but increased cortical bone has been observed in

other mouse models where cortical bone compensates for weak structure of trabecular bone (Morko et al., 2005; Turner et al., 2000). However, beyond 2 months, a high bone mass phenotype was clearly established which included cortical and trabecular bone of limbs and vertebrae. Notably, the calvarium is not affected. It is possible that excessive growth of calvarial bone may be regulated by mechanisms other than through miRNAs which limit the size of other bones in the body.

A striking feature of the increased bone mass of the limbs was stimulation of newly formed bone from 1 month to after 4 months of age occurring largely from the periosteal surfaced and more on the lateral than the medial side of the limb. This observation suggests two contributing mechanistic pathways that are likely to be influenced by miRNA regulation. First, the bone extracellular matrix harbors numerous growth factors, which contribute to both proliferative expansion of pre-osteoblasts and their differentiation e.g. IGF-1, FGFs, BMPs and TGF β , which have been well studied (reviewed in Allori et al. (2008); Giustina et al. (2008); Ornitz (2005)). Any increase in their levels could potentially stimulate bone formation. Secondly, the finding of a more profound effect on long bone indicates that mechanical forces on the bone may in part be contributing to the stimulated bone formation as the periosteum is well documented for its responses to mechanical stimulation and tissue repair (Henriksen et al., 2009). Furthermore, osteocytes are the responsive “mechanotransducer cells” of bones (reviewed in Bonewald and Johnson (2008)), which can secrete factors to stimulate the activity of the surface osteoblast (Bonewald and Johnson, 2008; Heino et al., 2004; Hirao et al., 2007). The osteocytes are organized in individual lacunae at a density equivalent to control. Further, bone remodeling and osteoclast activity is not impaired and the thickened bone becomes vascularized to maintain viability of the osteocytes.

Insight into the mechanisms for the Dicer^{Δoc/Δoc} post-natal phenotype of increased bone mass is obtained from our recent profiling studies of miRs during osteoblast differentiation. We identified a large representation of miRs during the mineralization period which target collagens and proved miR-29b directly downregulated four different collagens (Li et al., 2009). After birth, resolution of the initial delay in bone formation followed by continuous increase in bone mass in Dicer^{Δoc/Δoc} adult mice may be due to loss of miRs that target bone matrix extracellular proteins. For example, we confirmed in the present studies loss of let-7a and miR-29b and following Dicer excision in osteoblasts and with concomitant increases in collagen levels in bone of the Dicer^{Δoc/Δoc} mice, including the major protein of the bone ECM (90% type I collagen). This finding of the significance of miR-29 in the regulation of collagen is consistent with reports in other tissues where miR-29 decreases in myocardial infarction which leads to fibrosis (van Rooij et al., 2008). Establishment of a collagen scaffold in bone promotes osteoblast differentiation and mineralization (Franceschi et al., 1994; Lynch et al., 1995). In the absence of Dicer, miRs that inhibit biosynthesis of matrix proteins are not processed, and consequently bone formation can progress unimpeded to produce a greater than 2-fold increase in bone volume by 4 months that is retained up to 8 months. The process driving osteoblasts to osteocyte maturation may arise from stimulated secretion of ECM that accelerates the enveloping of surface osteoblasts to further differentiate (Bonewald and Johnson, 2008). Thus, while the Dicer enzyme is essential in early osteoprogenitors for bone formation, excision of Dicer has an anabolic effect on the adult skeleton due to loss of miRs that are required to regulate bone mass.

In conclusion, our findings have now identified functions for Dicer during development of the osteoblast phenotype for tissue morphogenesis and for post-natal regulation of bone accrual. These findings provide a potential approach for stimulating bone formation with antagonists upon identification of the specific microRNAs contributing to the anabolic phenotype in adult bone.

Acknowledgments

This work was supported by NIH grants R01AR039588 (GSS), P01AR048818 (GSS), R37DE012528 (JBL), DK73324 (SNJ), CA77735 (SNJ), DK032520 (for Core facilities), and S10RR023540 from the National Center for Research Resources (JBL). RM was supported in part by an American Heart Award (0625823T) and by NIH-T32CA130807. The contents of this manuscript are solely the responsibility of the authors and do not necessarily represent the official views of the National Center for Research Resources or the National Institutes of Health.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at doi:10.1016/j.ydbio.2010.01.008.

References

- Alford, A.I., Hankenson, K.D., 2006. Matricellular proteins: extracellular modulators of bone development, remodeling, and regeneration. *Bone* 38, 749–757.
- Allori, A.C., Sillon, A.M., Warren, S.M., 2008. Biological basis of bone formation, remodeling, and repair—part I: biochemical signaling molecules. *Tissue Eng. Part B Rev.* 14, 259–273.
- Andl, T., Murchison, E.P., Liu, F., Zhang, Y., Yunta-Gonzalez, M., Tobias, J.W., Andl, C.D., Seykora, J.T., Hannon, G.J., Millar, S.E., 2006. The miRNA-processing enzyme dicer is essential for the morphogenesis and maintenance of hair follicles. *Curr. Biol.* 16, 1041–1049.
- Babji, P., Zhao, W., Small, C., Kharode, Y., Yaworsky, P.J., Bouxsein, M.L., Reddy, P.S., Bodine, P.V., Robinson, J.A., Bhat, B., Marzolf, J., Moran, R.A., Bex, F., 2003. High bone mass in mice expressing a mutant LRP5 gene. *J. Bone Miner. Res.* 18, 960–974.
- Bernstein, E., Kim, S.Y., Carmell, M.A., Murchison, E.P., Alcorn, H., Li, M.Z., Mills, A.A., Elledge, S.J., Anderson, K.V., Hannon, G.J., 2003. Dicer is essential for mouse development. *Nat. Genet.* 35, 215–217.
- Blakaj, A., Lin, H., 2008. Piecing together the mosaic of early mammalian development through microRNAs. *J. Biol. Chem.* 283, 9505–9508.
- Bonewald, L.F., Johnson, M.L., 2008. Osteocytes, mechanosensing and Wnt signaling. *Bone* 42, 606–615.
- Chen, J.F., Murchison, E.P., Tang, R., Callis, T.E., Tatsuguchi, M., Deng, Z., Rojas, M., Hammond, S.M., Schneider, M.D., Selzman, C.H., Meissner, G., Patterson, C., Hannon, G.J., Wang, D.Z., 2008. Targeted deletion of Dicer in the heart leads to dilated cardiomyopathy and heart failure. *Proc. Natl. Acad. Sci. U. S. A.* 105, 2111–2116.
- Chiang, C., Chiu, W.M., Moore, A.J., Anderson, P.H., Ghasem-Zadeh, A., McManus, J.F., Ma, C., Seaman, E., Clemens, T.L., Morris, H.A., Zajac, J.D., Davey, R.A., 2009. Mineralization and bone resorption are regulated by the androgen receptor in male mice. *J. Bone Miner. Res.* 24, 621–631.
- Cobb, B.S., Nesterova, T.B., Thompson, E., Hertweck, A., O'Connor, E., Godwin, J., Wilson, C.B., Brockdorff, N., Fisher, A.G., Smale, S.T., Merckenschlager, M., 2005. T cell lineage choice and differentiation in the absence of the RNase III enzyme Dicer. *J. Exp. Med.* 201, 1367–1373.
- da Costa Martins, P.A., Bourajjaj, M., Gladka, M., Kortland, M., van Oort, R.J., Pinto, Y.M., Molkenin, J.D., De Windt, L.J., 2008. Conditional dicer gene deletion in the postnatal myocardium provokes spontaneous cardiac remodeling. *Circulation* 118, 1567–1576.
- Damiani, D., Alexander, J.J., O'Rourke, J.R., McManus, M., Jadhav, A.P., Cepko, C.L., Hauswirth, W.W., Harfe, B.D., Strettoi, E., 2008. Dicer inactivation leads to progressive functional and structural degeneration of the mouse retina. *J. Neurosci.* 28, 4878–4887.
- Davis, T.H., Cuellar, T.L., Koch, S.M., Barker, A.J., Harfe, B.D., McManus, M.T., Ullian, E.M., 2008. Conditional loss of Dicer disrupts cellular and tissue morphogenesis in the cortex and hippocampus. *J. Neurosci.* 28, 4322–4330.
- Dimri, G.P., Lee, X., Basile, G., Acosta, M., Scott, G., Roskelley, C., Medrano, E.E., Linskens, M., Rubelj, I., Pereira-Smith, O., 1995. A biomarker that identifies senescent human cells in culture and in aging skin in vivo. *Proc. Natl. Acad. Sci. U. S. A.* 92, 9363–9367.
- Franceschi, R.T., Iyer, B.S., Cui, Y., 1994. Effects of ascorbic acid on collagen matrix formation and osteoblast differentiation in murine MC3T3-E1 cells. *J. Bone Miner. Res.* 9, 843–854.
- Giustina, A., Mazziotti, G., Canalis, E., 2008. Growth hormone, insulin-like growth factors, and the skeleton. *Endocr. Rev.* 29, 535–559.
- Harfe, B.D., McManus, M.T., Mansfield, J.H., Hornstein, E., Tabin, C.J., 2005. The RNaseIII enzyme Dicer is required for morphogenesis but not patterning of the vertebrate limb. *Proc. Natl. Acad. Sci. U. S. A.* 102, 10898–10903.
- Harris, K.S., Zhang, Z., McManus, M.T., Harfe, B.D., Sun, X., 2006. Dicer function is essential for lung epithelium morphogenesis. *Proc. Natl. Acad. Sci. U. S. A.* 103, 2208–2213.
- Harvey, S.J., Jarad, G., Cunningham, J., Goldberg, S., Schermer, B., Harfe, B.D., McManus, M.T., Benzing, T., Miner, J.H., 2008. Podocyte-specific deletion of dicer alters cytoskeletal dynamics and causes glomerular disease. *J. Am. Soc. Nephrol.* 19, 2150–2158.
- Hayashi, K., de Sousa, Chuva, Lopes, S.M., Kaneda, M., Tang, F., Hajkova, P., Lao, K., O'Carroll, D., Das, P.P., Tarakhovskiy, A., Miska, E.A., Surani, M.A., 2008. MicroRNA

- biogenesis is required for mouse primordial germ cell development and spermatogenesis. *PLoS ONE* 3, e1738.
- Heino, T.J., Hentunen, T.A., Vaananen, H.K., 2004. Conditioned medium from osteocytes stimulates the proliferation of bone marrow mesenchymal stem cells and their differentiation into osteoblasts. *Exp. Cell Res.* 294, 458–468.
- Henriksen, K., Neutzsky-Wulff, A.V., Bonewald, L.F., Karsdal, M.A., 2009. Local communication on and within bone controls bone remodeling. *Bone* 44, 1026–1033.
- Hirao, M., Hashimoto, J., Yamasaki, N., Ando, W., Tsuboi, H., Myoui, A., Yoshikawa, H., 2007. Oxygen tension is an important mediator of the transformation of osteoblasts to osteocytes. *J. Bone Miner. Metab.* 25, 266–276.
- Jaskiewicz, L., Filipowicz, W., 2008. Role of Dicer in posttranscriptional RNA silencing. *Curr. Top. Microbiol. Immunol.* 320, 77–97.
- Kanellopoulou, C., Muljo, S.A., Kung, A.L., Ganesan, S., Drapkin, R., Jenuwein, T., Livingston, D.M., Rajewsky, K., 2005. Dicer-deficient mouse embryonic stem cells are defective in differentiation and centromeric silencing. *Genes Dev.* 19, 489–501.
- Kobayashi, T., Lu, J., Cobb, B.S., Rodda, S.J., McMahon, A.P., Schipani, E., Merckenschlager, M., Kronenberg, H.M., 2008. Dicer-dependent pathways regulate chondrocyte proliferation and differentiation. *Proc. Natl. Acad. Sci. U. S. A.* 105, 1949–1954.
- Koralov, S.B., Muljo, S.A., Galler, G.R., Krek, A., Chakraborty, T., Kanellopoulou, C., Jensen, K., Cobb, B.S., Merckenschlager, M., Rajewsky, N., Rajewsky, K., 2008. Dicer ablation affects antibody diversity and cell survival in the B lymphocyte lineage. *Cell* 132, 860–874.
- Lengner, C.J., Lepper, C., van Wijnen, A.J., Stein, J.L., Stein, G.S., Lian, J.B., 2004. Primary mouse embryonic fibroblasts: a model of mesenchymal cartilage formation. *J. Cell Physiol.* 200, 327–333.
- Li, X., Ominsky, M.S., Niu, Q.T., Sun, N., Daugherty, B., D'Agostin, D., Kurahara, C., Gao, Y., Cao, J., Gong, J., Asuncion, F., Barrero, M., Warmington, K., Dwyer, D., Stolina, M., Morony, S., Sarosi, I., Kostenuik, P.J., Lacey, D.L., Simonet, W.S., Ke, H.Z., Paszty, C., 2008a. Targeted deletion of the sclerostin gene in mice results in increased bone formation and bone strength. *J. Bone Miner. Res.* 23, 860–869.
- Li, Z., Hassan, M.Q., Jafferji, M., Garzon, R., Croce, C.M., van Wijnen, A.J., Stein, J.L., Stein, G.S., Lian, J.B., 2009. Biological functions of miR-29b contribute to positive regulation of osteoblast differentiation. *J. Biol. Chem.* 284, 15676–15684.
- Li, Z., Hassan, M.Q., Volinia, S., van Wijnen, A.J., Stein, J.L., Croce, C.M., Lian, J.B., Stein, G.S., 2008b. A microRNA signature for a BMP2-induced osteoblast lineage commitment program. *Proc. Natl. Acad. Sci. U. S. A.* 105, 13906–13911.
- Lian, J.B., Stein, G.S., Javed, A., van Wijnen, A.J., Stein, J.L., Montecino, M., Hassan, M.Q., Gaur, T., Lengner, C.J., Young, D.W., 2006. Networks and hubs for the transcriptional control of osteoblastogenesis. *Rev. Endocr. Metab. Disord.* 7, 1–16.
- Liu, F., Woitte, H.W., Braut, A., Kronenberg, M.S., Lichtler, A.C., Mina, M., Kream, B.E., 2004. Expression and activity of osteoblast-targeted Cre recombinase transgenes in murine skeletal tissues. *Int. J. Dev. Biol.* 48, 645–653.
- Lufkin, T., Mark, M., Hart, C.P., Dolle, P., LeMeur, M., Chambon, P., 1992. Homeotic transformation of the occipital bones of the skull by ectopic expression of a homeobox gene. *Nature* 359, 835–841.
- Luzi, E., Marini, F., Sala, S.C., Tognarini, I., Galli, G., Brandi, M.L., 2008. Osteogenic differentiation of human adipose tissue-derived stem cells is modulated by the miR-26a targeting of the SMAD1 transcription factor. *J. Bone Miner. Res.* 23, 287–295.
- Lynch, M.P., Stein, J.L., Stein, G.S., Lian, J.B., 1995. The influence of type I collagen on the development and maintenance of the osteoblast phenotype in primary and passaged rat calvarial osteoblasts: modification of expression of genes supporting cell growth, adhesion, and extracellular matrix mineralization. *Exp. Cell Res.* 216, 35–45.
- Malaval, L., Wade-Gueye, N.M., Boudiffa, M., Fei, J., Zirngibl, R., Chen, F., Laroche, N., Roux, J.P., Burt-Pichat, B., Duboef, F., Boivin, G., Jurdic, P., Lafage-Proust, M.H., Amedee, J., Vico, L., Rossant, J., Aubin, J.E., 2008. Bone sialoprotein plays a functional role in bone formation and osteoclastogenesis. *J. Exp. Med.* 205, 1145–1153.
- Miao, D., Scutt, A., 2002. Histochemical localization of alkaline phosphatase activity in decalcified bone and cartilage. *J. Histochem. Cytochem.* 50, 333–340.
- Mizuno, Y., Yagi, K., Tokuzawa, Y., Kanesaki-Yatsuka, Y., Suda, T., Katagiri, T., Fukuda, T., Maruyama, M., Okuda, A., Amemiya, T., Kondoh, Y., Tashiro, H., Okazaki, Y., 2008a. miR-125b inhibits osteoblastic differentiation by down-regulation of cell proliferation. *Biochem. Biophys. Res. Commun.* 368, 267–272.
- Mizuno, Y., Yagi, K., Tokuzawa, Y., Kanesaki-Yatsuka, Y., Suda, T., Katagiri, T., Fukuda, T., Maruyama, M., Okuda, A., Amemiya, T., Kondoh, Y., Tashiro, H., Okazaki, Y., 2008b. miR-125b inhibits osteoblastic differentiation by down-regulation of cell proliferation. *Biochem. Biophys. Res. Commun.* 368, 267–272.
- Morko, J., Kiviranta, R., Hurme, S., Rantakokko, J., Vuorio, E., 2005. Differential turnover of cortical and trabecular bone in transgenic mice overexpressing cathepsin K. *Bone* 36, 854–865.
- Mudhasani, R., Zhu, Z., Hutvagner, G., Eischen, C.M., Lyle, S., Hall, L.L., Lawrence, J.B., Imbalzano, A.N., Jones, S.N., 2008. Loss of miRNA biogenesis induces p19Arf-p53 signaling and senescence in primary cells. *J. Cell Biol.* 181, 1055–1063.
- Muljo, S.A., Ansel, K.M., Kanellopoulou, C., Livingston, D.M., Rao, A., Rajewsky, K., 2005. Aberrant T cell differentiation in the absence of Dicer. *J. Exp. Med.* 202, 261–269.
- Murchison, E.P., Stein, P., Xuan, Z., Pan, H., Zhang, M.Q., Schultz, R.M., Hannon, G.J., 2007. Critical roles for Dicer in the female germline. *Genes Dev.* 21, 682–693.
- Nagaraja, A.K., Andreu-Vieyra, C., Franco, H.L., Ma, L., Chen, R., Han, D.Y., Zhu, H., Agno, J.E., Gunaratne, P.H., DeMayo, F.J., Matzuk, M.M., 2008. Deletion of Dicer in somatic cells of the female reproductive tract causes sterility. *Mol. Endocrinol.* 22, 2336–2352.
- Nakasa, T., Miyaki, S., Okubo, A., Hashimoto, M., Nishida, K., Ochi, M., Asahara, H., 2008. Expression of microRNA-146 in rheumatoid arthritis synovial tissue. *Arthritis Rheum.* 58, 1284–1292.
- O'Rourke, J.R., Georges, S.A., Seay, H.R., Tapscott, S.J., McManus, M.T., Goldhamer, D.J., Swanson, M.S., Harfe, B.D., 2007. Essential role for Dicer during skeletal muscle development. *Dev. Biol.* 311, 359–368.
- Ornitz, D.M., 2005. FGF signaling in the developing endochondral skeleton. *Cytokine Growth Factor Rev.* 16, 205–213.
- Oskowitz, A.Z., Lu, J., Penfornis, P., Ylostalo, J., McBride, J., Flemington, E.K., Prockop, D.J., Pochampally, R., 2008. Human multipotent stromal cells from bone marrow and microRNA: regulation of differentiation and leukemia inhibitory factor expression. *Proc. Natl. Acad. Sci. U. S. A.* 105, 18372–18377.
- Pratap, J., Galindo, M., Zaidi, S.K., Vradii, D., Bhat, B.M., Robinson, J.A., Choi, J.-Y., Komori, T., Stein, J.L., Lian, J.B., Stein, G.S., van Wijnen, A.J., 2003. Cell growth regulatory role of Runx2 during proliferative expansion of pre-osteoblasts. *Cancer Res.* 63, 5357–5362.
- Robey, P.G., Boskey, A.L., 2008. The composition of bone. In: Rosen, C.J. (Ed.), *Primer on the Metabolic Bone Diseases and Disorders of Mineral Metabolism*. American Society for Bone and Mineral Research, Washington, DC, pp. 32–38.
- Schafer, A., O'Carroll, D., Tan, C.L., Hillman, D., Sugimori, M., Llinas, R., Greengard, P., 2007. Cerebellar neurodegeneration in the absence of microRNAs. *J. Exp. Med.* 204, 1553–1558.
- Serrano, M., Lee, H., Chin, L., Cordon-Cardo, C., Beach, D., DePinho, R.A., 1996. Role of the INK4a locus in tumor suppression and cell mortality. *Cell* 85, 27–37.
- Soltanoff, C.S., Yang, S., Chen, W., Li, Y.P., 2009. Signaling networks that control the lineage commitment and differentiation of bone cells. *Crit. Rev. Eukaryot. Gene Expr.* 19, 1–46.
- Sudo, H., Kodama, H.-A., Amagai, Y., Yamamoto, S., Kasai, S., 1983. In vitro differentiation and calcification in a new clonal osteogenic cell line derived from newborn mouse calvaria. *J. Cell Biol.* 96, 191–198.
- Sugatani, T., Hruska, K.A., 2009. Impaired micro-RNA pathways diminish osteoclast differentiation and function. *J. Biol. Chem.* 284, 4667–4678.
- ten Dijke, P., Krause, C., de Gorter, D.J., Lowik, C.W., Van Bezooijen, R.L., 2008. Osteocyte-derived sclerostin inhibits bone formation: its role in bone morphogenetic protein and Wnt signaling. *J. Bone Joint Surg. Am.* 90 (Suppl 1), 31–35.
- Tuddenham, L., Wheeler, G., Ntounia-Fousara, S., Waters, J., Hajihosseini, M.K., Clark, I., Dalmay, T., 2006. The cartilage specific microRNA-140 targets histone deacetylase 4 in mouse cells. *FEBS Lett.* 580, 4214–4217.
- Turner, C.H., Hsieh, Y.F., Muller, R., Bouxsein, M.L., Baylink, D.J., Rosen, C.J., Grynblas, M.D., Donahue, L.R., Beamer, W.G., 2000. Genetic regulation of cortical and trabecular bone strength and microstructure in inbred strains of mice. *J. Bone Miner. Res.* 15, 1126–1131.
- van Rooij, E., Sutherland, L.B., Thatcher, J.E., DiMaio, J.M., Naseem, R.H., Marshall, W.S., Hill, J.A., Olson, E.N., 2008. Dysregulation of microRNAs after myocardial infarction reveals a role of miR-29 in cardiac fibrosis. *Proc. Natl. Acad. Sci. U. S. A.* 105, 13027–13032.
- Yang, W.J., Yang, D.D., Na, S., Sandusky, G.E., Zhang, Q., Zhao, G., 2005. Dicer is required for embryonic angiogenesis during mouse development. *J. Biol. Chem.* 280, 9330–9335.
- Yi, R., O'Carroll, D., Pasolli, H.A., Zhang, Z., Dietrich, F.S., Tarakhovskiy, A., Fuchs, E., 2006. Morphogenesis in skin is governed by discrete sets of differentially expressed microRNAs. *Nat. Genet.* 38, 356–362.
- Yuan, B., Takaiwa, M., Clemens, T.L., Feng, J.Q., Kumar, R., Rowe, P.S., Xie, Y., Drezner, M.K., 2008. Aberrant PheX function in osteoblasts and osteocytes alone underlies murine X-linked hypophosphatemia. *J. Clin. Invest.* 118, 722–734.