

ACADEMIC
PRESSAvailable at
WWW.MATHEMATICSWEB.ORG
POWERED BY SCIENCE @ DIRECT®

Journal of Approximation Theory 123 (2003) 125–146

<http://www.elsevier.com/locate/jat>JOURNAL OF
**Approximation
Theory**

q -Taylor theorems, polynomial expansions, and interpolation of entire functions[☆]

Mourad E.H. Ismail^{a,*} and Dennis Stanton^b^a *Department of Mathematics, University of South Florida, 4202 East Fowler Avenue, PHY 114, Tampa, FL 33620-5700, USA*^b *School of Mathematics, University of Minnesota, Minneapolis, MN 55455, USA*

Received 13 September 2002; accepted in revised form 10 April 2003

Communicated by Arno B.J. Kuijlaars

Abstract

We establish q -analogues of Taylor series expansions in special polynomial bases for functions analytic in bounded domains and for entire functions whose maximum modulus $M(r; f)$ satisfies $|\ln M(r; f)| \leq A \ln^2 r$. This solves the problem of constructing such entire functions from their values at $[aq^n + q^{-n}/a]/2$, for $0 < q < 1$. Our technique is constructive and gives an explicit representation of the sought entire function. Applications to q -series identities are given.

© 2003 Elsevier Science (USA). All rights reserved.

Keywords: q -Taylor series; Polynomial bases; Askey–Wilson operators; Integral representations; q -exponential function; Carlson's theorem

1. Introduction

Two important problems in complex function theory are the problems of expanding a function in a series of polynomials and the interpolation problem of finding an entire function from its values on a given sequence $\{x_n\}$, $x_n \rightarrow \infty$ as $n \rightarrow \infty$. The polynomial expansion problem has a long history. Whittaker [22,23] introduced the concept of basic sets of polynomials where the polynomials are

[☆] Research partially supported by NSF grants DMS 99-70865, DMS 02-03282.

*Corresponding author. Fax: +1-913-974-2700.

E-mail addresses: ismail@math.usf.edu (M.E.H. Ismail), stanton@math.umn.edu (D. Stanton).

ordered but not necessarily according to their degrees and all degrees are present. A more recent treatment of Whittaker’s approach is in the interesting monograph by Makar [16]. Boas and Buck [4] restricted the polynomials to having a generating function of a special type which guarantees that $p_n(x)$ has precise degree n , $n = 0, 1, \dots$. As a result of the specialization imposed by Boas and Buck, they have been able to obtain more refined results than those which hold for general basic sets of polynomials. The bases treated here are not of Boas and Buck type.

In this paper we solve the interpolation problem for the sequence $\{x_{2n}\}$,

$$x_n = [aq^{n/2} + q^{-n/2}/a]/2, \quad 0 < q < 1, \quad 0 < a < 1, \tag{1.1}$$

for entire functions f satisfying

$$\limsup_{r \rightarrow +\infty} \frac{\ln M(r;f)}{\ln^2 r} = c, \tag{1.2}$$

for a particular c which depends upon q . Here $M(r;f)$ is [3]

$$M(r;f) = \sup\{|f(z)| : |z| \leq r\}. \tag{1.3}$$

In the process of solving this problem we also solve the expansion problem of entire functions in two specific bases of polynomials, namely $\{\phi_n(x;a)\}$ and $\{\rho_n(x)\}$ defined in (1.9)–(1.10) and the coefficients in the expansion on $\{\phi_n(x;a)\}$ involve function evaluations at $\{x_{2n}\}$. In the case of $\{\rho_n(x)\}$, the interpolation points are

$$u_n = i(q^{n/2} - q^{-n/2})/2, \quad n = \dots, -1, 0, 1, \dots \tag{1.4}$$

Carlson’s theorem [3] states that an entire function f of order one and type less than π is uniquely determined by the sequence $\{f(n) : n = 0, 1, \dots\}$. Moreover, if $f(x)$ is entire of order 1 and type $< \ln 2$ then

$$f(x) = \sum_{n=0}^{\infty} \binom{x}{n} (\Delta^n f)(0), \tag{1.5}$$

and the series converges uniformly on compact subsets of the complex x -plane, [3, Theorem 9.10.7]. In the above $(\Delta f)(x) = f(x + 1) - f(x)$. Another representation was obtained by Ramanujan in his first notebooks, where he wrote

$$\int_0^{\infty} x^{s-1} \sum_{k=0}^{\infty} f(k)(-x)^k dx = \frac{\pi}{\sin \pi s} f(-s). \tag{1.6}$$

Hardy [9, (11.2A), p. 186] proved (1.6) by contour integration and pointed out that it holds under the assumptions in Carlson’s theorem. Therefore, Ramanujan’s formula (1.6) provides a constructive proof of Carlson’s theorem by showing how to construct the function f from $\{f(n)\}$. Therefore, in some sense, our formulas are closer in spirit to (1.5). An interesting question is to find the analogue of Ramanujan’s formula (1.6).

Ramis [19] defined an entire function f to have a q -exponential growth of order k and a finite type if there exist real numbers $K, \alpha, K > 0$, such that

$$|f(x)| < K|x|^\alpha \exp\left(\frac{k \ln^2 |x|}{2 \ln^2 q}\right).$$

Thus functions satisfying (1.2) are of q -exponential growth of order $2c \ln^2 q$.

Two of our main results are Theorems 3.1 and 3.3 which are stated and proved in Section 3. Before we can state our results we need to explain the notation used, which is mainly from [1,7]. The q -shifted factorials are

$$(a; q)_0 := 1, \quad (a; q)_n := \prod_{k=1}^n (1 - aq^{k-1}), \quad n = 1, 2, \dots, \quad \text{or } \infty, \tag{1.7}$$

while the multiple q -shifted factorials are defined by

$$(a_1, a_2, \dots, a_k; q)_n := \prod_{j=1}^k (a_j; q)_n. \tag{1.8}$$

The bases of polynomials we are interested in are defined by

$$\phi_n(\cos \theta; a) = (ae^{i\theta}, ae^{-i\theta}; q)_n = \prod_{k=0}^{n-1} [1 - 2axq^k + a^2q^{2k}], \quad a > 0 \tag{1.9}$$

$$\rho_n(\cos \theta) = (1 + e^{2i\theta})(-q^{2-n}e^{2i\theta}; q^2)_{n-1} e^{-in\theta}. \tag{1.10}$$

The motivation for considering these special bases is our desire to establish Taylor-like series where the Askey–Wilson operator plays the role of $\frac{d}{dx}$ and these polynomials play the role of monomials. The basis $\{\phi_n(x; a)\}$ was introduced in the Askey–Wilson memoir [2] but the basis $\{\rho_n(x)\}$ does not seem to have been considered before we introduced them in [12].

We now define the Askey–Wilson operator \mathcal{D}_q introduced in [2]. Given a function f we set $\check{f}(e^{i\theta}) := f(x), x = \cos \theta$, that is

$$\check{f}(z) = f((z + 1/z)/2), \quad z = e^{i\theta}. \tag{1.11}$$

In other words we think of $f(\cos \theta)$ as a function of $e^{i\theta}$. In this notation the Askey–Wilson divided difference operator \mathcal{D}_q is defined by

$$(\mathcal{D}_q f)(x) = \frac{\check{f}(q^{1/2}e^{i\theta}) - \check{f}(q^{-1/2}e^{i\theta})}{(q^{1/2} - q^{-1/2})i \sin \theta}, \quad x = \cos \theta. \tag{1.12}$$

For example with $f(x) = 4x^3 - 3x, \check{f}(z) = [z^3 + z^{-3}]/2$, and

$$\mathcal{D}_q f(x) = \frac{q^{3/2} - q^{-3/2}}{q^{1/2} - q^{-1/2}} (4x^2 - 1).$$

It is a fact that \mathcal{D}_q reduces the degree of a polynomial by one and

$$\lim_{q \rightarrow 1} \mathcal{D}_q f(x) = \frac{d}{dx} f(x), \tag{1.13}$$

at the points where f is differentiable. Furthermore, in the calculus of the Askey–Wilson operator the basis $\{\phi_n(x; a) : n \geq 0\}$ plays the role played by the monomials $\{(1 - 2ax + a^2)^n : n \geq 0\}$ in the differential and integral calculus.

Note that although we use $x = \cos \theta$, θ is not necessarily real but $e^{\pm i\theta}$ are defined as

$$e^{\pm i\theta} = x \pm \sqrt{x^2 - 1},$$

and the branch of the square root is taken such that $\sqrt{x^2 - 1} \approx x$ as $x \rightarrow \infty$. Thus $|e^{-i\theta}| = |e^{i\theta}|$ if and only if $x \in [-1, 1]$.

The action of \mathcal{D}_q on the bases in (1.9)–(1.10) is given by

$$\mathcal{D}_q \phi_n(x; a) = -\frac{2a(1 - q^n)}{1 - q} \phi_{n-1}(x; aq^{1/2}), \tag{1.14}$$

$$\mathcal{D}_q \rho_n(x) = 2q^{(1-n)/2} \frac{1 - q^n}{1 - q} \rho_{n-1}(x). \tag{1.15}$$

As already mentioned the values of an entire function f on the nonnegative integers determine f , [3], when f is of order one and type less than π . On the other hand $f(z) = \sin \pi z$ is order 1 and type π and vanishes at all the integers, so type π is a cut off point. A similar situation occurs for the interpolation points $\{x_{2k} : k = 0, 1, \dots\}$. The function $\phi_\infty(x; a)$ vanishes at all the points x_{2k} , so if $\{f(x_{2k}) : k \geq 0\}$ determine an entire function f uniquely then f is expected to grow slower than $\phi_\infty(x; a)$. It turns out that when c in (1.2) is $< 1/(2 \ln q^{-1})$ then f can be interpolated from $\{f(x_{2k})\}$ and f has a polynomial expansion in $\{\phi_n(x; a)\}$. For $f(x) = \phi_\infty(x; a)$, $c = 1/(2 \ln q^{-1})$, so the barrier, which corresponds to type π , is $1/(2 \ln q^{-1})$. This will be proved in Section 3. The uniqueness of an entire function taking prescribed values at $\{x_{2k}\}$ or $\{u_k\}$ follows from general theory of entire functions, [3], and divided difference operators [8]. Our contribution is two-fold. Firstly, we provide an explicit representation of the entire function with growth restriction. Secondly, we give alternate representation of f when f is only assumed to be analytic in a bounded domain.

In Theorem 3.1 we extend the following theorem from polynomials to entire functions. Theorem 1.1 combines results from [11,12].

Theorem 1.1. *Let $f(x)$ be a polynomial and assume that x_n is defined by (1.1). Then,*

$$f(x) = \sum_{k=0}^{\infty} f_{k,\phi} \phi_k(x; a)$$

with

$$f_{k,\phi} = \frac{(q - 1)^k}{(2a)^k (q; q)_k} q^{-k(k-1)/4} (\mathcal{D}_q^k f)(x_k).$$

In addition we have

$$f(x) = \sum_{k=0}^{\infty} f_{k,\rho} \rho_k(x),$$

where

$$f_{k,\rho} = \frac{q^{(k^2-k)/4} (1-q)^k}{2^k (q; q)_k} (\mathcal{D}_q^k f)(0).$$

In Section 4 we rewrite Theorem 3.1 in the form of a Mittag–Leffler expansion, see (4.2). This expansion turns out to be very useful in studying summation theorems for basic hypergeometric series. Some examples are given in Section 5. Section 6 contains concluding remarks and the evaluation of c in (1.2) for the q -exponential function $\mathcal{E}_q(z; \alpha)$. We have only included a few examples of the implications of the material derived here and we have avoided including technical special functions results, which will appear in a more specialized publication.

One interesting byproduct of our results is the following version of a formula of Cooper [5]

$$\begin{aligned} & \mathcal{D}_q^n f(x) \\ &= \frac{2^n q^{n(1-n)/4}}{(q^{1/2} - q^{-1/2})^n} \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q \frac{q^{k(n-k)} z^{2k-n} \check{f}(q^{(n-2k)/2} z)}{(q^{1+n-2k} z^2; q)_k (q^{2k-n+1} z^{-2}; q)_{n-k}}, \end{aligned} \tag{1.16}$$

where $z = e^{i\theta}$, $x = \cos \theta$, and

$$\begin{bmatrix} n \\ k \end{bmatrix}_q = \frac{(q; q)_n}{(q; q)_k (q; q)_{n-k}}, \tag{1.17}$$

is the q -analogue of the binomial coefficient.

The key tool used to establish the expansion of the Cauchy kernel is the theory of basic hypergeometric functions. An interesting open problem is to find a purely complex variable proof of this expansion or of the expansions of entire functions. For convenience we include the definition of a basic hypergeometric series

$$\begin{aligned} & {}_r\phi_s \left(\begin{matrix} a_1, \dots, a_r \\ b_1, \dots, b_s \end{matrix} \middle| q, z \right) \\ &= {}_r\phi_s(a_1, \dots, a_r; b_1, \dots, b_s; q, z) \\ &= \sum_{n=0}^{\infty} \frac{(a_1, \dots, a_r; q)_n}{(q, b_1, \dots, b_s; q)_n} z^n (-q^{(n-1)/2})^{n(s+1-r)}. \end{aligned} \tag{1.18}$$

Finally we use the Bailey notation

$$\begin{aligned}
 &W(a^2; a_1, \dots, a_r; q, z) \\
 &:= {}_{r+3}\phi_{r+2} \left(\begin{matrix} a^2, qa, -qa, a_1, \dots, a_r \\ a, -a, qa^2/a_1, \dots, qa^2/a_r \end{matrix} \middle| q, z \right).
 \end{aligned}
 \tag{1.19}$$

The ϕ function in (1.19) is called very well-poised.

2. Expansions of the Cauchy kernel

In this section we expand the Cauchy kernel $1/(y - x)$ in terms of $\{\phi_n(x; a)\}$ and $\{\rho_n(x)\}$. This is done in Theorem 2.1. The Cauchy kernel expansion is then used in Theorems 2.2 and 2.3 to expand entire functions in the same bases with coefficients represented by contour integrals. These integral representations are analogues of the Cauchy formulas.

It is easy to prove that

$$\mathcal{D}_q(1/\phi_n(x; a)) = \frac{2a(1 - q^n)}{1 - q} \frac{1}{\phi_{n+1}(x; aq^{-1/2})}.
 \tag{2.1}$$

It is clear that

$$\frac{1}{\cos \phi - \cos \theta} = \frac{2e^{i\phi}}{\phi_1(\cos \theta; e^{i\phi})},
 \tag{2.2}$$

and therefore if $y = \cos \phi$,

$$\mathcal{D}_q^k(y - x)^{-1} \Big|_{x=x_k} = \frac{2(-1)^k q^{k(1-k)/2} e^{i(k+1)\phi}}{a^k (ae^{i\phi}, q^{-k} e^{i\phi}/a; q)_{k+1}}.
 \tag{2.3}$$

Theorem 2.1. *The Cauchy kernel has the expansion*

$$\frac{1}{y - x} = \frac{1}{y - x} \frac{\phi_\infty(x; a)}{\phi_\infty(y; a)} - 2a \sum_{n=0}^\infty \frac{\phi_n(x; a)}{\phi_{n+1}(y; a)} q^n,$$

for all y such that $y \neq x$, and $\phi_\infty(y; a) \neq 0$. The above expansion also holds if $y = y_0$, $\phi_\infty(y_0; a) = 0$, but $x \neq y_0$ in the sense that the left-hand side at $y = y_0$ equals the limit of the right-hand side as $y \rightarrow y_0$. Moreover, the expansion of the Cauchy kernel in $\{\rho_n\}$ is

$$\begin{aligned}
 \frac{1}{y - x} &= \frac{x}{y^2 - x^2} \frac{(-qe^{2i\theta}, -qe^{-2i\theta}; q^2)_\infty}{(-qe^{2i\phi}, -qe^{-2i\phi}; q^2)_\infty} + \frac{y}{y^2 - x^2} \frac{(-e^{2i\theta}, -e^{-2i\theta}; q^2)_\infty}{(-e^{2i\phi}, -e^{-2i\phi}; q^2)_\infty} \\
 &+ 4 \sum_{n=0}^\infty \frac{y\rho_n(x)}{[(1 - q^n)^2 + 4y^2q^n]\rho_n(y)} q^n,
 \end{aligned}$$

provided that $y \neq x$ and $\rho_n(y) \neq 0$ for all $n, n = 0, 1, \dots$.

First note that $\rho_n(x)/\rho_n(y)$ is uniformly bounded if y is not a zero of $(-e^{2i\phi}, -e^{-2i\phi}; q)_\infty$, since we have

$$\lim_{N \rightarrow \infty} \frac{\rho_{2N}(x)}{\rho_{2N}(y)} = \frac{(-e^{2i\theta}, -e^{-2i\theta}; q^2)_\infty}{(-e^{2i\phi}, -e^{-2i\phi}; q^2)_\infty},$$

$$\lim_{N \rightarrow \infty} \frac{\rho_{2N+1}(x)}{\rho_{2N+1}(y)} = \frac{x(-qe^{2i\theta}, -qe^{-2i\theta}; q^2)_\infty}{y(-qe^{2i\phi}, -qe^{-2i\phi}; q^2)_\infty}.$$

Hence the second series in Theorem 2.1 converges absolutely and uniformly for x and y in compact sets.

Proof. The idea of the proof is to formally expand the Cauchy kernel using (2.3) and Theorem 1.1. The formal expansions of the Cauchy kernel in $\{\phi_n(x; a)\}$ and $\{\rho_n(x)\}$ converge, but not to the Cauchy kernel. So we evaluate the difference explicitly using the theory of basic hypergeometric functions [1,7]. These are the two terms in Theorem 2.1.

With $x = \cos \theta, y = \cos \phi$ we get

$$\frac{(q-1)^k q^{k(1-k)/4}}{(2a)^k (q; q)_k} \mathcal{D}_q^k (y-x)^{-1} \Big|_{x=x_k} = \frac{-2aq^k}{\phi_{k+1}(\cos \phi; a)},$$

using (2.3). Thus the formal expansion of the Cauchy kernel is given by

$$-2a \sum_{k=0}^{\infty} \frac{(ae^{i\theta}, ae^{-i\theta}; q)_k}{(ae^{i\phi}, ae^{-i\phi}; q)_{k+1}} q^k$$

$$= \frac{-2a}{1-2ay+a^2} {}_3\phi_2 \left(\begin{matrix} q, ae^{i\theta}, ae^{-i\theta} \\ qae^{i\phi}, qae^{-i\phi} \end{matrix} \middle| q, q \right), \tag{2.4}$$

which is the sum on the right-hand side of the first formula in Theorem 2.1. Applying the transformation [7, (III.9)] and assuming $\text{Im } \phi > 0$, we see that the last expression is given by

$$-\frac{2a}{(1-q)(1-ae^{i\phi})} {}_3\phi_2 \left(\begin{matrix} q, qe^{i(\phi-\theta)}, qe^{i(\phi+\theta)} \\ qae^{i\phi}, q^2 \end{matrix} \middle| q, ae^{-i\phi} \right)$$

$$= \frac{2e^{i\phi}}{(1-e^{i(\phi-\theta)})(1-e^{i(\phi+\theta)})}$$

$$\times \left[1 - {}_2\phi_1 \left(\begin{matrix} e^{i(\phi-\theta)}, e^{i(\phi+\theta)} \\ ae^{i\phi} \end{matrix} \middle| q, ae^{-i\phi} \right) \right].$$

The ${}_2\phi_1$ is summable by the q -analogue of Gauss’ theorem [7, (II.8)] and its sum is $(ae^{i\theta}, ae^{-i\theta}; q)_\infty / (ae^{i\phi}, ae^{-i\phi}; q)_\infty$. The result is the first expansion in Theorem 2.1. We remove the assumption $\text{Im } \phi > 0$ by analytic continuation.

To prove the second expansion, we evaluate the sum on the right-hand side of the second formula in Theorem 2.1. After the application of (1.10) the sum with which

we are concerned is found to be

$$2 \sum_{n=0}^{\infty} \frac{e^{i(n+1)\phi} (1 + e^{2i\theta}) (-q^{2-n} e^{2i\theta}; q^2)_{n-1}}{e^{in\theta} (iq^{-n/2} e^{i\phi}, -iq^{-n/2} e^{i\phi}; q)_{n+1}}. \tag{2.5}$$

We first assume that $|e^{-i\phi}| < 1$. We next sum over n even and over n odd. The even sum is

$$\begin{aligned} & \frac{2e^{i\phi}}{1 + e^{2i\phi}} \sum_{n=0}^{\infty} \frac{(-e^{2i\theta}, -e^{-2i\theta}; q^2)_n q^{2n}}{(iqe^{i\phi}, iqe^{-i\phi}, -iqe^{i\phi}, -iqe^{-i\phi}; q)_n} \\ &= \frac{2e^{i\phi}}{1 + e^{2i\phi}} {}_3\phi_2 \left(\begin{matrix} q^2, -e^{2i\theta}, -e^{-2i\theta} \\ -q^2 e^{2i\phi}, -q^2 e^{-2i\phi} \end{matrix} \middle| q^2, q^2 \right) \\ &= \frac{2e^{3i\phi} (1 + e^{-2i\phi})}{(1 - e^{2i(\phi+\theta)})(1 - e^{2i(\phi-\theta)})} \\ & \quad \times \left[1 - {}_2\phi_1 \left(\begin{matrix} e^{2i(\phi-\theta)}, e^{2i(\phi+\theta)} \\ -e^{2i\phi} \end{matrix} \middle| q^2, -e^{-2i\phi} \right) \right], \end{aligned}$$

where the transformation [7, (III.9)] was applied in the last step. Again Gauss’ theorem [7, (II.8)] sums the ${}_2\phi_1$ and we see that the even sum is

$$\frac{y}{y^2 - x^2} \left[1 - \frac{(-e^{2i\theta}, -e^{-2i\theta}; q^2)_{\infty}}{(-e^{2i\phi}, -e^{-2i\phi}; q^2)_{\infty}} \right]. \tag{2.6}$$

The odd sum can be similarly handled and can be simplified to

$$\frac{x}{y^2 - x^2} \left[1 - \frac{(-qe^{2i\theta}, -qe^{-2i\theta}; q^2)_{\infty}}{(-qe^{2i\phi}, -qe^{-2i\phi}; q^2)_{\infty}} \right]. \tag{2.7}$$

Now we remove the assumption $|e^{-i\phi}| < 1$ by analytic continuation. Equating (2.5) to the sum of (2.6) and (2.7) gives the second part of the theorem and the proof is complete. \square

Theorem 2.2. *Let f be analytic in a bounded domain D and let C be a contour within D and x belong to the interior of C . If the distance between C and the set of zeros of $\phi_{\infty}(x; a)$ is positive then*

$$\begin{aligned} f(x) &= \frac{\phi_{\infty}(x; a)}{2\pi i} \oint_C \frac{f(y)}{y - x} \frac{dy}{\phi_{\infty}(y; a)} \\ & \quad - \frac{a}{\pi i} \sum_{n=0}^{\infty} q^n \phi_n(x; a) \oint_C \frac{f(y) dy}{\phi_{n+1}(y; a)}. \end{aligned}$$

Proof. It is clear that

$$\phi_n(x; a) / \phi_{n+1}(y; a) \rightarrow \phi_{\infty}(x; a) / \phi_{\infty}(y; a)$$

uniformly in y on compact subsets not intersecting the set of zeros of $\phi_{\infty}(y; a)$. Thus we can multiply the first expansion in Theorem 2.1 by $f(y)$ and integrate with respect

to y and interchange integration and summation. The result then follows from Cauchy’s theorem. \square

Theorem 2.2 gives the q -analogue of expanding $f(x)$ around $x = (a + 1/a)/2$. From the theory of functions we know that if $f(a) = 0$ and $f^{(j)}(a) = 0$ for $1 \leq j \leq m - 1$, then the Taylor series starts with the term $f^{(m)}(a)(x - a)^m/m!$. This feature continues to hold but we have to define a q -analogue of a multiple zero.

Definition. Let $x = (a + 1/a)/2$ be a zero of $f(x)$. We say that it has q -multiplicity m if

$$f(z_k) = 0, \quad 1 \leq k \leq m - 1, \quad \text{and} \quad f(z_m) \neq 0, \quad z_k := \frac{1}{2}(aq^k + q^{-k}/a). \quad (2.8)$$

Similarly a pole x of f has q -multiplicity m if x is a zero of $1/f$ with q -multiplicity m .

It must be emphasized that the above definition is completely analogous to the definition of a multiple zero in difference equations in Hartman [10]. With this definition one can see that if $(a + 1/a)/2$ is a zero of f of q -multiplicity m then the terms corresponding to $n = 0, 1, \dots, m - 1$ in the sum in Lemma 2.2 vanish and the sum starts from $n = m$.

Theorem 2.3. Let f be analytic in a bounded domain D and let C be a contour within D and x is interior to C . If the contour C is at a positive distance from the set $\{\pm i(q^{n/2} - q^{-n/2})/2; n = 0, 1, \dots\}$, then

$$f(x) = \frac{2x}{\pi i} \oint_C \frac{yf(y)}{y-x} \frac{(-qe^{i(\theta+\phi)}, -qe^{i(\theta-\phi)}, -qe^{i(\phi-\theta)}, -qe^{-i(\theta+\phi)}; q)_\infty}{(-q, -q; q)_\infty (-e^{2i\phi}, -e^{-2i\phi}; q)_\infty} dy + \frac{2}{\pi i} \sum_{n=0}^\infty \rho_n(x) q^n \oint_C \frac{yf(y) dy}{[(1 - q^n)^2 + 4y^2 q^n] \rho_n(y)},$$

with $x = \cos \theta$ and $y = \cos \phi$.

Proof. The proof is very similar to the proof of Theorem 2.2. The only step requiring justification is the identity

$$\frac{x}{y^2 - x^2} \frac{(-qe^{2i\theta}, -qe^{-2i\theta}; q^2)_\infty}{(-qe^{2i\phi}, -qe^{-2i\phi}; q^2)_\infty} + \frac{y}{y^2 - x^2} \frac{(-e^{2i\theta}, -e^{-2i\theta}; q^2)_\infty}{(-e^{2i\phi}, -e^{-2i\phi}; q^2)_\infty} = \frac{4xy}{y-x} \frac{(-qe^{i(\theta+\phi)}, -qe^{i(\theta-\phi)}, -qe^{i(\phi-\theta)}, -qe^{-i(\theta+\phi)}; q)_\infty}{(-q, -q; q)_\infty (-e^{2i\phi}, -e^{-2i\phi}; q)_\infty}. \quad (2.9)$$

The proof of (2.9) uses the relationships [21, Chapter 21]

$$\vartheta_2(z) = 2Gq^{1/4} \cos z (-q^2 e^{2iz}, -q^2 e^{-2iz}; q^2)_\infty$$

$$= \frac{Gq^{1/4}}{2 \cos z} (-e^{2iz}, -e^{-2iz}; q^2)_\infty,$$

$$\vartheta_3(z) = G(-qe^{2iz}, -qe^{-2iz}; q^2)_\infty,$$

and the product formulas in Exercise 3, page 488 in Whittaker and Watson [21]. The notations $\mathfrak{y}_j := \mathfrak{y}_j(0)$, $G := (q^2; q^2)_\infty$ [21] were used. We omit the details. One referee kindly pointed out that (2.9) also follows from (2.16) and (5.21) of [7]. \square

We record the following equivalent form of the representation of f in Theorem 2.3

$$\begin{aligned}
 f(x) &= \frac{x}{2\pi i} \oint_C \frac{f(y)}{y^2 - x^2} \frac{(-qe^{2i\theta}, -qe^{-2i\theta}; q^2)_\infty}{(-qe^{2i\phi}, -qe^{-2i\phi}; q^2)_\infty} dy \\
 &+ \frac{1}{2\pi i} \oint_C \frac{yf(y)}{y^2 - x^2} \frac{(-e^{2i\theta}, -e^{-2i\theta}; q^2)_\infty}{(-e^{2i\phi}, -e^{-2i\phi}; q^2)_\infty} dy \\
 &+ \frac{2}{\pi i} \sum_{n=0}^\infty \rho_n(x) q^n \oint_C \frac{yf(y) dy}{[(1 - q^n)^2 + 4y^2 q^n] \rho_n(y)}.
 \end{aligned}$$

3. Expansions of entire functions

In this section we establish expansion theorems for entire functions of q -exponential growth. The expansions are in terms of the bases $\{\phi_n(x; a)\}$ and $\{\rho_n(x)\}$.

Observe that $M(r; \phi_\infty(x; a)) = \phi_\infty(-r; a)$, since $a > 0$. Hence with

$$r_m = [aq^{m+\delta} + a^{-1}q^{-(m+\delta)}]/2, \quad -1 < \delta < 0, \quad m = 0, 1, \dots \tag{3.1}$$

we find

$$\begin{aligned}
 M(r_n; \phi_\infty(x; a)) &= \phi_\infty(-r_n; a) \\
 &= (-q^{-n-\delta}; q)_n (-q^{-\delta}, -a^2 q^{n+\delta}; q)_\infty \\
 &= q^{[\delta^2 - n - (n+\delta)^2]/2} (-q^{\delta+1}; q)_n (-q^{-\delta}, -a^2 q^{n+\delta}; q)_\infty
 \end{aligned} \tag{3.2}$$

so that

$$\lim_{n \rightarrow \infty} \frac{\ln M(r_n; \phi_\infty)}{\ln^2 r_n} = \frac{1}{2 \ln q^{-1}}.$$

The fact that $\phi_\infty(x; a)$ vanishes at x_{2n} for all n motivates our next theorem.

Theorem 3.1. *An entire function f satisfying (1.2) with $c < 1/(2 \ln q^{-1})$ has a convergent expansion*

$$f(x) = \sum_{k=0}^\infty f_{k,\phi} \phi_k(x; a),$$

with $\{f_{k,\phi}\}$ defined in Theorem 1.1. Moreover, any such f is uniquely determined by its values on $\{x_{2n} : n \geq 0\}$.

Note that $(\mathcal{G}_q^k f)(x_k)$ is a linear combination of $f(x_0), \dots, f(x_{2k})$, so that the coefficients $f_{k,\phi}$ in Theorem 1.1 also depend on the points $\{x_{2n} : n \geq 0\}$.

The proof of Theorem 3.1 relies on a lemma which we now state and prove.

Lemma 3.2. *Let f be entire and satisfy the condition in Theorem 3.1. Then if r_n is defined by (3.1), we have*

$$\lim_{n \rightarrow \infty} \oint_{|y|=r_n} \frac{f(y)}{y-x} \frac{dy}{\phi_\infty(y;a)} = 0.$$

Moreover, the same conclusion holds if

$$\lim_{n \rightarrow \infty} q^{n(n+2\delta+1)/2} \sup\{|f(r_n e^{i\theta})| : 0 \leq \theta < 2\pi\} = 0.$$

Proof. It is clear that $\inf\{|\phi_\infty(y;a)| : |y|=r\} = |\phi_\infty(r;a)|$. Hence for $|y|=r_n$, we have

$$\begin{aligned} |\phi_\infty(y;a)| &\geq |(q^{-n-\delta}; q)_n| (q^{-\delta}, a^2 q^{n+\delta}; q)_\infty \\ &= q^{-n(n+2\delta+1)/2} (q^{\delta+1}; q)_n (q^{-\delta}, a^2 q^{n+\delta}; q)_\infty. \end{aligned}$$

Therefore,

$$\begin{aligned} \ln M(r_n; f(y)/\phi_\infty(y;a)) &\leq \frac{1}{2} [n + (n + \delta)^2] \ln q + \ln M(r_n; f) + O(1) \\ &= \ln M(r_n; f) - \frac{1}{2} \frac{\ln^2 r_n}{\ln q^{-1}} + O(\ln r_n), \end{aligned}$$

and the lemma follows. \square

Instead of proving the expansion in Theorem 3.1 in the basis $\{\phi_n(x;a)\}$ we shall prove the following equivalent result.

Theorem 3.3. *The expansion formula*

$$f(x) = \sum_{n=0}^{\infty} q^n f_{n,\phi} \phi_n(x;a),$$

with

$$f_{n,\phi} = \sum_{k=0}^n \frac{(-1)^k q^{k(k-1)/2} (1 - a^2 q^{2k})}{(q; q)_k (q; q)_{n-k} (a^2 q^k; q)_{n+1}} f(x_{2k}),$$

holds for functions f satisfying the assumptions of Theorem 3.1.

Proof. In Theorem 2.2 we choose C to be C_m , a circle centered at $y = 0$ and radius r_m . Lemma 3.2 shows that the first integral in Theorem 2.2 is small if m is large. We split the remaining sum in Theorem 2.2 into tail terms with $n > m$, and initial terms with $n \leq m$. We will show that the tail is small, leaving the initial terms. Then a residue calculation establishes the expression for $f_{n,\phi}$, because the poles of $f(y)/\phi_{n+1}(y;a)$ are at $y = x_{2k}$, $k = 0, 1, \dots, n$. The details of the residue calculation

are omitted since they are similar to the calculation at the end of the proof of Theorem 3.4.

Note that if $n > m$ then

$$\begin{aligned} & \min\{|\phi_{n+1}(y)| : y \in C_m\} \\ &= |\phi_{n+1}(r_m; a)| = |(q^{-m-\delta}, a^2 q^{m+\delta}; q)_{n+1}| \\ &= (q^{-m-\delta}; q)_m (-1)^m (q^{-\delta}; q)_{n+1-m} (a^2 q^{m+\delta}; q)_{n+1} \\ &= q^{-m(m+2\delta+1)/2} (q^{\delta+1}; q)_m (q^{-\delta}; q)_{n+1-m} (a^2 q^{m+\delta}; q)_{n+1} \\ &\geq q^{-m(m+2\delta+1)/2} A = q^{-((m+\delta)^2+1-\delta^2)/2} A, \end{aligned}$$

where A is a positive constant independent of n and m . Therefore, for sufficiently large m , and $y \in C_m$,

$$\ln[M(r_m; f/\phi_{n+1})] \leq [c_1 + 1/(2 \ln q)] \ln^2 r_m + O(m)$$

for some $c_1, c \leq c_1 < 1/(2 \ln q^{-1})$.

This is a uniform bound of $e^{-D(\ln r_m)^2}$, $D > 0$, for each integral for $n > m$. Since $\phi_n(x; a) \rightarrow \phi_\infty(x; a)$, there is a uniform bound B for $\phi_n(x; a)$ on compact sets. Thus the tail is bounded by

$$\sum_{n=m+1}^\infty Bq^n e^{-D(\ln r_m)^2} \leq Bq^{m+1} e^{-D(\ln r_m)^2} / (1 - q),$$

which is small for m large. \square

For polynomials f we equate the coefficients $f_{n,\phi}$ in the expansions of f in $\{\phi_n(x; a)\}$ in Theorems 3.3 and 1.1 and discover the identity

$$\begin{aligned} & \mathcal{D}_q^n f(x_n) \\ &= \frac{(2a)^n q^{n(n+3)/4}}{(q-1)^n} \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q \frac{(-1)^k q^{k(k-1)/2}}{(a^2 q^k; q)_{n+1}} (1 - a^2 q^{2k}) f(x_{2k}). \end{aligned} \tag{3.3}$$

Since (3.3) holds for arbitrary polynomials it must hold for all continuous functions. Using the notation

$$\eta_q^{\pm 1} f(x) = \check{f}(q^{\pm 1/2} z), \tag{3.4}$$

and noting that a , is a general parameter and $x_n = \eta^n x_0$, we can rewrite (3.3) in the form

$$\mathcal{D}_q^n f(x) = \frac{(2z)^n q^{n(3-n)/4}}{(q-1)^n} \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q \frac{(-1)^k q^{k(k-1)/2} \eta^{2k-n} f(x)}{(z^2 q^{k-n}; q)_k (z^2 q^{2k+1-n}; q)_{n-k}}, \tag{3.5}$$

with $x = (z + z^{-1})/2$. Eq. (3.5) can be shown to be equivalent to Cooper’s (1.16).

Theorem 3.4. *Let f be an entire function satisfying (1.2) and assume that $c < 1/\ln q^{-1}$. Then f has the expansion*

$$f(x) = \sum_{n=0}^{\infty} f_{n,\rho} \rho_n(x),$$

where

$$f_{n,\rho} = i^n \sum_{k=0}^n (-1)^k \frac{(q^k + q^{n-k})q^{(k^2+(n-k)^2)/2}}{2(q^2; q^2)_k (q^2; q^2)_{n-k}} f(u_{n-2k}),$$

and $\{u_n\}$ is given by (1.4).

For general entire functions not necessarily satisfying (1.2), we note that the property $u_j = -u_{-j}$, allows one to conclude that for even functions f , $f_{2n+1,\rho} = 0$ for all $n \geq 0$, while for odd functions f , $f_{2n,\rho} = 0$, for all $n \geq 0$, confirming that f and its formal expansion $\sum_{n=0}^{\infty} f_{n,\rho} \rho_n(x)$ have the same parity.

Proof. We basically repeat the proof of Theorem 3.3, with some changes in the technical details. We will use

$$\frac{\rho_{2N}(x)}{\rho_{2N}(y)} = \frac{(-e^{2i\theta}, -e^{-2i\theta}; q^2)_N}{(-e^{2i\phi}, -e^{-2i\phi}; q^2)_N} \tag{3.6}$$

$$\frac{\rho_{2N+1}(x)}{\rho_{2N+1}(y)} = \frac{x(-qe^{2i\theta}, -qe^{-2i\theta}; q^2)_N}{y(-qe^{2i\phi}, -qe^{-2i\phi}; q^2)_N} \tag{3.7}$$

Let C_m be a circle centered at $y = 0$ with radius r_m ,

$$r_m = [q^{-(m+\delta)/2} - q^{(m+\delta)/2}]/2, \quad -1 < \delta < 0, \tag{3.8}$$

and m is even. We use the form of the Cauchy kernel in (2.10) and show that the first integral in (2.10) with $C = C_m$ tends to zero as $m \rightarrow \infty$. Thus we minimize the modulus of the denominators in order to give an upper bound for the integral. The first denominator is minimized by choosing $y^2 = -r_m^2$,

$$\begin{aligned} & \min\{|(-qe^{2i\phi}, -qe^{-2i\phi}; q^2)_{\infty}| : y \in C_m\} \\ &= |(q^{1-m-\delta}, q^{m+\delta+1}; q^2)_{\infty}| = |(q^{1-m-\delta}; q^2)_{m/2}| (q^{1-\delta}, q^{m+\delta+1}; q^2)_{\infty} \\ &\geq Aq^{-(m+\delta)^2/4} \end{aligned}$$

for some positive constant A independent of m .

Similarly,

$$\begin{aligned} & \min\{|(-e^{2i\phi}, -e^{-2i\phi}; q^2)_{\infty}| : y \in C_m\} \\ &= |(q^{-m-\delta}, q^{m+\delta}; q^2)_{\infty}| = |(q^{-m-\delta}; q^2)_{m/2}| (q^{-\delta}, q^{m+\delta}; q^2)_{\infty} \\ &\geq Aq^{-(m+\delta)^2/4-m/2}. \end{aligned}$$

So the sum of the first two integrals, $|I_m|$, is bounded by

$$|I_m| \leq B M(r_m; f) q^{(m+\delta)^2/4},$$

for some B independent of m . Thus,

$$\ln |I_m| \leq \ln M(r_m; f) + \frac{(\ln r_m)^2}{\ln q} + O(\ln r_m)$$

which proves that $|I_m| \rightarrow 0$ as $m \rightarrow \infty$, m even.

Next we show that $\sum_{n=m+1}^{\infty} q^n |I_{n,m}|$ tends to zero as $m \rightarrow \infty$ and for x in compact sets, where $\{I_{n,m}\}$ are the integrals

$$I_{n,m} = \oint_{C_m} \frac{y \rho_n(x) f(y) dy}{\rho_n(y) [(1 - q^n)^2 + 4y^2 q^n]}. \tag{3.9}$$

For $y \in C_m$, and $n > m$, we have

$$\begin{aligned} |(1 - q^n)^2 + 4y^2 q^n| &= |(1 + q^n e^{2i\phi})(1 + q^n e^{-2i\phi})| \\ &\geq |(1 - q^{n-m-\delta})(1 - q^{n+m+\delta})| \\ &\geq (1 - q^{1-\delta})(1 - q^{1+\delta}). \end{aligned}$$

Moreover after applying (3.6) and (3.7), we get for $y \in C_m$ and $n > m$,

$$\begin{aligned} \left| \frac{\rho_{2n}(x)}{\rho_{2n}(y)} \right| &\leq \frac{|(-e^{2i\theta}, -e^{-2i\theta}; q^2)_n|}{|(q^{-m-\delta}, q^{m+\delta}; q^2)_n|} \\ &\leq \frac{|(-e^{2i\theta}, -e^{-2i\theta}; q^2)_n|}{|(q^{-m-\delta}, q^2)_{m/2} (q^{-\delta}, q^2)_{n-m/2} (q^{m+\delta}, q^2)_n|} \\ &\leq A_1 q^{(m+\delta)^2/4}, \end{aligned}$$

for some constant A_1 , and

$$\begin{aligned} \left| \frac{\rho_{2n+1}(x)}{\rho_{2n+1}(y)} \right| &\leq \frac{|x(-qe^{2i\theta}, -qe^{-2i\theta}; q^2)_n|}{|y(q^{1-m-\delta}, q^{1+m+\delta}; q^2)_n|} \\ &\leq \frac{|x(-qe^{2i\theta}, -qe^{-2i\theta}; q^2)_n|}{|y(q^{1-m-\delta}, q^2)_{m/2} (q^{1-\delta}, q^2)_{n-m/2} (q^{1+m+\delta}, q^2)_n|} \\ &\leq A_2 q^{(m+\delta)^2/4}. \end{aligned}$$

The constants A_1 and A_2 depend on the compact set x to which x is restricted but do not depend on y , m or n .

As before this shows that

$$\begin{aligned} \ln |I_{n,m}| &\leq \ln M(f; r_m) + \frac{(\ln r_m)^2}{\ln q} + O(\ln r_m) \\ &\leq (c_1 + 1/\ln q)(\ln r_m)^2, \end{aligned}$$

for some c_1 , $0 < c_1 < 1/\ln q^{-1}$ which shows that $\sum_{n=m+1}^{\infty} q^n |I_{n,m}|$ tends to zero as $m \rightarrow \infty$.

Next we evaluate the sum $\sum_{n=0}^m q^n I_{m,n}$ by residues then let $m \rightarrow \infty$. From (1.10) it follows that

$$\rho_{n+2}(y) = q^{-n}[(1 - q^n)^2 + 4y^2 q^n] \rho_n(y),$$

hence we need to evaluate

$$\oint_{C_m} \frac{y f(y) dy}{\rho_{n+2}(y)}.$$

The poles $\{y_k\}$ of $y/\rho_{n+2}(y)$ are

$$i[q^{-k+n/2} - q^{k-n/2}]/2,$$

$k = 0, 1, \dots, n$. Let $y_k = \cos \phi_k$, hence

$$e^{i\phi_k} = \begin{cases} -iq^{k-n/2}, & 0 \leq k \leq n/2, \\ iq^{-k+n/2}, & n/2 < k \leq n. \end{cases} \tag{3.10}$$

It is routine to use (3.10) and find that the residue of $y/\rho_{n+2}(y)$ at $i[q^{-k+n/2} - q^{k-n/2}]/2$ is

$$i^n \frac{(-1)^k (q^{n-k} + q^k)}{8(q^2; q^2)_k (q^2; q^2)_{n-k}} q^{k(k-n)+n^2/2}, \quad k = 0, \dots, n,$$

and the theorem follows. \square

Remark. The first part of the proof can be replaced by estimating the first integral in Theorem 2.3 directly. Let $|e^{i\theta}| \leq A$ for all x in a compact set. Hence for $y \in C_m$ and fixed x , we have

$$\begin{aligned} & \ln \left(\left| \frac{y(-qe^{i(\theta+\phi)}, -qe^{i(\theta-\phi)}, -qe^{i(\phi-\theta)}, -qe^{-i(\theta+\phi)}; q)_\infty}{(-e^{2i\phi}, -e^{-2i\phi}; q)_\infty} \right| \right) \\ & \leq \ln((-Aq^{1-(m+\delta)/2}, -q^{1-(m+\delta)/2}/A; q)_\infty) \\ & \quad - \ln(|(q^{-m-\delta}; q)_\infty|) + O(1) \\ & \leq \ln((-Aq^{1-(m+\delta)/2}, -q^{1-(m+\delta)/2}/A; q)_{m/2}) \\ & \quad - \ln(|(q^{-m-\delta}; q)_m|) + O(m) \\ & = \frac{m(m+2\delta)}{4} \ln q + O(m) = -\frac{\ln^2 r_m}{\ln q^{-1}} + O(\ln(r_m)). \end{aligned}$$

Therefore, the first integral on the right-hand side of the equation in Theorem 2.3 tends to zero as $m \rightarrow \infty$.

4. A Mittag–Leffler expansion

It is tempting to substitute for f_n in the first formula in Theorem 3.3 then rearrange the sum and find the coefficient of $f(x_{2k})$. The formal interchange of sums gives

$$f(x) = \sum_{k=0}^{\infty} \frac{(-1)^k q^{k(k+1)/2} (1 - a^2 q^{2k})}{(q; q)_k (a^2 q^k; q)_{k+1}} (ae^{i\theta}, ae^{-i\theta}; q)_k \times {}_2\phi_1 \left(\begin{matrix} aq^k e^{i\theta}, aq^k e^{-i\theta} \\ a^2 q^{2k+1} \end{matrix} \middle| q, q \right) f(x_{2k}). \tag{4.1}$$

The ${}_2\phi_1$ can be summed by the q -analogue of Gauss’ theorem [7, (II.8)] and its sum is $(aq^{k+1}e^{i\theta}, aq^{k+1}e^{-i\theta}; q)_{\infty} / (a^2 q^{2k+1}, q; q)_{\infty}$ and (4.1) becomes

$$\frac{f(x)}{\phi_{\infty}(x; a)} = \sum_{k=0}^{\infty} \frac{(-1)^k q^{k(k+1)/2} (1 - a^2 q^{2k})}{(q; q)_k (q, a^2 q^k; q)_{\infty}} \frac{f(x_{2k})}{1 - 2axq^k + a^2 q^{2k}}. \tag{4.2}$$

Theorem 4.1. *Formula (4.2) holds for entire functions f satisfying (1.2) with $c < 1/(2 \ln q^{-1})$.*

Proof. Let r_m be as in (3.1) and C_m be a circle centered at the origin and have radius r_m . Let x be fixed and m be large enough so that x is interior to C_m . Consider

$$I_m := \frac{1}{2\pi i} \oint_{C_m} \frac{f(y)}{\phi_{\infty}(y; a)} \frac{dy}{y - x}. \tag{4.3}$$

From Lemma 3.2, $I_m \rightarrow 0$ as $m \rightarrow \infty$. On the other hand, the residue theorem implies

$$I_m = \frac{f(x)}{\phi_{\infty}(x; a)} - \sum_{k=0}^m \frac{(-1)^k q^{k(k+1)/2} (1 - a^2 q^{2k})}{(q; q)_k (q, a^2 q^k; q)_{\infty}} \frac{f(x_{2k})}{1 - 2axq^k + a^2 q^{2k}},$$

and the theorem follows. \square

Clearly Theorem 4.1 is a Mittag–Leffler expansion.

5. Applications

Recall that the conclusion of Lemma 3.2 holds provided that

$$\lim_{n \rightarrow \infty} M(r_n; f) q^{n(n+2\delta+1)/2} = 0. \tag{5.1}$$

where r_n is defined in (3.1). By examining the proof of Theorem 3.3 we see that it continues to hold under assumption (5.1). In fact we can replace the sequence $\{r_n\}$ in (5.1) by any subsequence $\{r_{n_k}\}$.

As a first application of the above observation we let

$$g(z) = (be^{i\theta}, be^{-i\theta}; p)_{\infty}, \quad p \leq q, \quad z := \cos \theta. \tag{5.2}$$

To verify (5.1) we employ

$$\begin{aligned} M_n(r_n; g) &\leq (-|b/a|q^{-n-\delta}, -|ab|q^{n+\delta}; p)_{\infty} \\ &\leq (-|b/a|q^{-n-\delta}, -|ab|q^{n+\delta}; q)_{\infty} \\ &\leq |b/a|^n q^{-n(n+2\delta+1)/2} (-|b/a|q^{-\delta}, -q^{1+\delta}|ab|; q)_{\infty}. \end{aligned}$$

Thus (5.1) holds when $|b/a| < 1$, and (4.2) will then hold for $|b| < |a|$ and we have established the series summation

$$\begin{aligned} &\frac{(be^{i\theta}, be^{-i\theta}; p)_{\infty}}{(qae^{i\theta}, qae^{-i\theta}; q)_{\infty}} \\ &= \sum_{k=0}^{\infty} \frac{(-1)^k q^{k(k+1)/2} (a^2, aq, -aq; q)_k}{(q, a, -a; q)_k (q, a^2q; q)_{\infty}} \\ &\quad \times \frac{(ae^{i\theta}, ae^{-i\theta}; q)_k}{(aqe^{i\theta}, aqe^{-i\theta}; q)_k} (abq^k, bq^{-k}/a; p)_{\infty}, \end{aligned} \tag{5.3}$$

valid for $0 < p < q$, or $p = q$ and $|b| < |a|$.

Mizan Rahman pointed out that (5.3) follows from a result of George Gasper. Gasper’s formula is (5.13) on p. 68 in [6] and can be stated as

$$\begin{aligned} &{}_{6+2m}W_{5+2m} \left(A; B, \frac{A}{B}, d, e_1, \dots, e_m, \frac{Aq^{n_1+1}}{e_1}, \dots, \frac{Aq^{n_m+1}}{e_m}; q, \frac{q}{d} q^{-\sum_{j=1}^m n_j} \right) \\ &= \frac{(q, Aq, Aq/Bd, Bq/d; q)_{\infty}}{(Bq, Aq/B, Aq/d, q/d; q)_{\infty}} \prod_{j=1}^m \frac{(Aq/Be_j, Bq/e_j; q)_{n_j}}{(Aq/e_j, q/e_j; q)_{n_j}}. \end{aligned}$$

We put

$$A = a^2, \quad B = ae^{i\theta}, \quad d = q^{-m}, \quad e_j = aqp^{1-j}/b, \quad n_j = 1, \quad 1 \leq j \leq m.$$

Write the ${}_{6+2m}W_{5+2m}$ as a sum over $k, k \geq 0$. The contribution to the k th term of the factors containing e_1, \dots, e_m is

$$\begin{aligned} &\prod_{r=0}^{m-1} \frac{(aqp^{-r}/b, abqp^r; q)_k}{(abp^r, ap^{-r}/b; q)_k} \\ &= \prod_{r=0}^{m-1} \frac{(1 - aq^k p^{-r}/b)(1 - abp^r q^k)}{(1 - ap^{-r}/b)(1 - abp^r)} = q^{km} \frac{(abq^k, bq^{-k}/a; p)_m}{(ab, b/a; q)_m}. \end{aligned}$$

Now (5.3) follows by letting $m \rightarrow \infty$.

When $p = q$ in (5.3), a simple calculation using

$$\begin{aligned} (abq^k, bq^{-k}/a; q)_\infty &= (ab, b/a; q)_\infty \frac{(bq^{-k}/a; q)_k}{(ab; q)_k} \\ &= \frac{(-b/a)^k (aq/b; q)_k}{q^{k(k+1)/2} (ab; q)_k} (ab, b/a; q)_\infty, \end{aligned}$$

shows that the right-hand side of (5.1) is $(ab, b/a; q)_\infty / (q, a^2q; q)_\infty$ times a ${}_6\phi_5$ function. Thus (5.3) with $p = q$ is equivalent to

$$\begin{aligned} {}_6\phi_5 \left(\begin{matrix} a^2, aq, -aq, aq/b, ae^{i\theta}, ae^{-i\theta} \\ a, -a, ab, aqe^{-i\theta}, aqe^{i\theta} \end{matrix} \middle| \begin{matrix} b \\ q, \frac{b}{a} \end{matrix} \right) \\ = \frac{(q, a^2q, be^{i\theta}, be^{-i\theta}; q)_\infty}{(aqe^{i\theta}, aqe^{-i\theta}, ab, b/a; q)_\infty}. \end{aligned} \tag{5.4}$$

Formula (5.4) is the sum of a very well poised ${}_6\phi_5$, [7, (II.20)]. The most general ${}_6\phi_5$ has four free parameters, but our (5.4) has only three free parameters.

Another application of (4.2) is to choose

$$f(z) = \prod_{j=1}^m f_j(z), \quad f_j(\cos \theta) := (b_j e^{i\theta}, b_j e^{-i\theta}; p_j)_\infty. \tag{5.5}$$

Here we will only mention the case when the $p_j = p$ for all j . In this case we choose a positive integer l such that $q^{l+1} < p \leq q^l$. It suffices to take $n = ls$ in (5.1) and for sufficiently large s , we get

$$\begin{aligned} M(r_{ls}; f_j) &\leq (-|b_j/a|q^{-ls-\delta}, -|ab_j|q^{ls+\delta}; p)_\infty \\ &\leq (-|b_j/a|q^{-ls-\delta}; p)_s (-|b_j/a|q^{-\delta}, -|ab_j|^{ls+\delta}; p)_\infty \\ &\leq |b_j/a|^s q^{-s(ls+\delta)} p^{s(s-1)/2} A_j, \end{aligned} \tag{5.6}$$

where A_j is a constant depending only on $a, b_1, \dots, b_m, \delta$ but not on s . With r defined through

$$p = q^r, \quad 1 \leq r < 1 + 1/l, \tag{5.7}$$

we obtain

$$\begin{aligned} M(r_{ls}; f) &q^{ls(ls+2\delta+1)/2} \\ &\leq Bq^{-(ls+\delta)sm+ls(ls+2\delta+1)/2} p^{ms(s-1)/2} \prod_{j=1}^m |b_j/a|^s, \end{aligned} \tag{5.8}$$

for some constant B . Substitute for p in (5.8) from (5.7) to see that the coefficient of s^2 in the exponent of q is nonnegative if and only if

$$l \geq m(2 - r). \tag{5.9}$$

If $l = m(2 - r)$ then the coefficient of s in the exponent of q is $m[\delta + 1 - r\delta - r(1 + m(2 - r))/2]$. Thus we have established the following theorem.

Theorem 5.1. Let f be defined by (5.5) with $p_j = p$ for all $j, 1 \leq j \leq m$, and let $l \geq 1$ be defined by $q^{l+1} < p \leq q^l$. Set $B = |b_1 \cdots b_m|^{1/m}$. If

- (i) $l > m(2 - r)$, or
- (ii) $l = m(2 - r)$ and $Bq^{1-r(1+m(2-r))/2} < |a|$,

holds then f has the Mittag-Leffler expansion (4.2).

The details of consequences of Theorem 5.1 will be explored elsewhere. We just mention the case $p_j = p = q^m$, so $r = 1$ and $m = l$. Thus (4.2) holds if

$$\prod_{j=1}^m |b_j/a| < q^{m(m-1)/2}. \tag{5.10}$$

Thus (4.2) gives

$$\begin{aligned} & \frac{\prod_{j=1}^m (b_j e^{i\theta}, b_j e^{-i\theta}; q^m)_\infty}{(ae^{i\theta}, ae^{-i\theta}; q)_\infty} \\ &= \sum_{k=0}^\infty \frac{(-1)^k q^{k(k+1)/2} (1 - a^2 q^{2k}) \prod_{j=1}^m (b_j q^{-k}/a, ab_j q^k; q^m)_\infty}{(q; q)_k (q, a^2 q^k; q)_\infty (1 - aq^k e^{i\theta})(1 - aq^k e^{-i\theta})}. \end{aligned} \tag{5.11}$$

The special case $m = 2$ of (5.13) follows from [7, (III.38)]. To see this we first write upper case letters for the parameters a, b, \dots, q in [7, (III.38)]. We make the choices

$$\begin{aligned} Q &= q^2, & A &= B = e^{i\theta}, & C &= e^{-i\theta}, \\ D &= q^2/b_1, & E &= q^2/b_2, & F &= a, & G &= aq. \end{aligned} \tag{5.12}$$

The resulting ${}_8\psi_8$ on the left-hand side of (III.38) in [7] reduces to 1 because its numerator parameter AC is 1, while a denominator parameter $(= AQ/B)$ is q^2 , which is the base of the ${}_8\psi_8$. The even and odd sums in (5.11) are the ${}_8\phi_7$'s on the right-hand side of (III.38).

6. Remarks

In [12] we pointed out the importance of the polynomial basis

$$\phi_n(\cos \theta) = (q^{1/4} e^{i\theta}, q^{1/4} e^{-i\theta}; q^{1/2})_n, \tag{6.1}$$

in the theory of basic hypergeometric functions. We also established the q -Taylor series

$$f(x) = \sum_{k=0}^n f_k \phi_k(x), \tag{6.2}$$

for polynomials f , where

$$f_k = \frac{(q - 1)^k}{2^k q^{k/4} (q; q)_k} (\mathcal{D}_q^k f)(\zeta_0) \tag{6.3}$$

and

$$\zeta_n = [q^{(n+1/2)/2} + q^{-(n+1/2)/2}]/2. \tag{6.4}$$

The proof of (6.2) uses

$$\mathcal{D}_q \phi_n(x) = -2q^{1/4} \frac{1 - q^n}{1 - q} \phi_{n-1}(x). \tag{6.5}$$

One can also extend (6.2) to entire functions satisfying (1.2) with $c < 1/\ln q^{-1}$ using Theorem 3.1 because the interpolation points used in (6.4) correspond to replacing q by $q^{1/2}$ and a by $q^{1/4}$ in (1.1).

The q -exponential function of [14] is

$$\begin{aligned} \mathcal{E}_q(\cos \theta; t) &= \frac{(t^2; q^2)_\infty}{(qt^2; q^2)_\infty} \sum_{n=0}^\infty \frac{(-it)^n}{(q; q)_n} q^{n^2/4} \\ &\quad \times (-iq^{(1-n)/2} e^{i\theta}, -iq^{(1-n)/2} e^{-i\theta}; q)_n. \end{aligned} \tag{6.6}$$

The function $\mathcal{E}_q(x; t)$ is entire in x for all t , $|t| < 1$. Corollary 2.5 of [12] is

$$\mathcal{E}_q(\cos \theta; t) = \frac{(-t; q^{1/2})_\infty}{(qt^2; q^2)_\infty} {}_2\phi_1 \left(\begin{matrix} q^{1/4} e^{i\theta}, q^{1/4} e^{-i\theta} \\ -q^{1/2} \end{matrix} \middle| q^{1/2}, -t \right). \tag{6.7}$$

We now show that (6.7) enables us to determine the exact limiting behavior of the maximum modulus of the \mathcal{E}_q function. Let $r = \cosh u$, $u > 0$. Thus (6.7) implies

$$\ln M(\cosh u; \mathcal{E}_q) \leq \ln((-q^{1/4} e^u, -q^{1/4} e^{-u}; q^{1/2})_\infty) + O(1),$$

as $u \rightarrow \infty$. It is clear that for any sequence $\{u_m\}$ tending to infinity, e^{u_m} can be written in the form $q^{-(n_m + \delta_m)/2}$, with positive integers n_m such that $-1/2 \leq \delta_m < 1/2$. From here it is not difficult to see that

$$\limsup_{r \rightarrow \infty} \frac{\ln M(r; \mathcal{E}_q)}{\ln^2 r} \leq \frac{1}{\ln q^{-1}}. \tag{6.8}$$

On the other hand the sequence $r_m = [q^{-(m+1/2)/2} + q^{(m+1/2)/2}]/2$ makes

$$\frac{(qt^2; q^2)_\infty}{(-t; q^{1/2})_\infty} \mathcal{E}_q(r_m; t) = {}_2\phi_1 \left(\begin{matrix} q^{-m/2}, q^{(m+1)/2} \\ -q^{1/2} \end{matrix} \middle| q^{1/2}, -t \right). \tag{6.9}$$

The right-hand side of (6.9) is a little q -Jacobi polynomial $\Phi_m^{(\alpha, \beta)}(x)$, with $\alpha = \beta = -1$ and $x = -t/q^{1/2}$, [13], hence the ${}_2\phi_1$ in (6.9) is asymptotically equal to

$$t^m q^{-m(m+1)/4} (-q^{1/2}/t; q)_\infty / (-q^{1/2}; q^{1/2})_\infty,$$

by (1.5) in [13]. Therefore

$$\limsup_{r \rightarrow \infty} \frac{\ln M(r; \mathcal{E}_q)}{\ln^2 r} \geq \lim_{m \rightarrow \infty} \frac{\ln M(r_m; \mathcal{E}_q)}{\ln^2 r_m} = \frac{1}{\ln q^{-1}}. \tag{6.10}$$

Therefore (6.8) and (6.10) establish the following theorem.

Theorem 6.1. For $|t| < 1$ the maximum modulus of \mathcal{E}_q has the property

$$\limsup_{r \rightarrow \infty} \frac{\ln M(r; \mathcal{E}_q)}{\ln^2 r} = \frac{1}{\ln q^{-1}}.$$

It is worth mentioning that Theorem 6.1 shows that (6.7) does not follow from the general approach developed here. It is of interest to find a function theoretic approach to development of identities like (6.7). A simple proof of (6.7) using ideas from the Sheffer classification [18] is in [15].

There is extensive literature on entire functions of exponential order when $f(n)$ takes integer values at $n, n = 0, 1, \dots$. One such theorem is due to Polya [17] and states that an entire function of exponential order $< \log 2$ which takes integer values at $n, n = 0, 1, \dots$, is a polynomial. Wallisser [20] mentions the following q -analogue, due to Gelfond [8].

Theorem 6.2. Assume that f is an entire function such that

$$\ln M(r; f) \leq \frac{(\ln(\sigma r))^2}{4 \ln q},$$

and q is an integer such that $2 \leq q, \sigma < 1/q$. If f takes integer values at q^n then f is a polynomial.

Wallisser [20] then raises the question of finding the form of the entire functions taking integer values at the points $(q^n - 1)/(q - 1), n = 0, 1, \dots$, with some restrictions on q in addition to $q > 1$. In fact our work raises the question of describing a class of functions f such that if $f(x_{2n})$ (or $f(u_n)$) is an integer for all $n = 0, 1, \dots$, then f is a polynomial.

Acknowledgments

Part of this work was done while the first author was visiting the Liu Bie Ju Center of Mathematical Sciences of City University of Hong Kong and the Hong Kong University of Science and Technology. He gratefully acknowledges the financial support and the hospitality of both universities. Both authors thank Yik-Man Chiang and Mizan Rahman for many discussions and for their interest in this work. Last but not least we thank two very kind referees for many useful comments and remarks on the first version of this paper.

References

- [1] G.E. Andrews, R.A. Askey, R. Roy, Special Functions, Cambridge University Press, Cambridge, 1999.
- [2] R.A. Askey, J.A. Wilson, Some basic hypergeometric orthogonal polynomials that generalize Jacobi polynomials, *Memoirs of the American Mathematical Society*, 319. Providence RI, 1985.

- [3] R.P. Boas, *Entire Functions*, Academic Press, New York, 1954.
- [4] R.P. Boas, R.C. Buck, *Polynomial Expansions of Analytic Functions*, Springer, Berlin, 1964.
- [5] S. Cooper, The Askey–Wilson operator and the ${}_6\phi_5$ summation formula, preprint, March 1996.
- [6] G. Gasper, Elementary derivations of summation and transformation formulas for q -series, in: M.E.H. Ismail, D.R. Masson, M. Rahman (Eds.), *Special Functions, q -Series and Related Topics*, Fields Institute Communication, Vol. 14, American Mathematical Society, Providence, RI, 1997, pp. 55–70.
- [7] G. Gasper, M. Rahman, *Basic Hypergeometric Series*, Cambridge University Press, Cambridge, 1990.
- [8] A.O. Gelfond, *Calculus of Finite Differences*, Translated from the Russian, Hindustan Publishing Corporation, Delhi, 1971.
- [9] G.H. Hardy, *Ramanujan: Twelve Lectures on Subjects Suggested by his Life and Work*, Chelsea, New York, 1978.
- [10] P. Hartman, Difference equations: disconjugacy, principal solutions, Green's functions, complete monotonicity, *Trans. Amer. Math. Soc.* 246 (1978) 1–30.
- [11] M.E.H. Ismail, The Askey–Wilson operator and summation theorems, in: M. Ismail, M.Z. Nashed, A. Zayed, A. Ghaleb (Eds.), *Mathematical Analysis, Wavelets, and Signal Processing*, Contemporary Mathematics, Vol. 190, American Mathematical Society, Providence, RI, 1995, pp. 171–178.
- [12] M.E.H. Ismail, D. Stanton, Applications of q -Taylor theorems, *J. Comp. Appl. Math.* 153 (2003) 259–272.
- [13] M.E.H. Ismail, J. Wilson, Asymptotic and generating relations for the q -Jacobi and the ${}_4\phi_3$ polynomials, *J. Approx. Theory* 36 (1982) 43–54.
- [14] M.E.H. Ismail, R. Zhang, Diagonalization of certain integral operators, *Adv. Math.* 109 (1994) 1–33.
- [15] M.E.H. Ismail, R. Zhang, New proofs of some q -series results, in: M.E.H. Ismail, H. Koelink (Eds.), *Kluwer*, to appear.
- [16] B.H. Makar, *The theory of coordinates*, a book manuscript, 1973.
- [17] G. Polya, Über ganzwertige ganze Funktionen, *Ren. Circ. Mat. Palermo* 40 (1915) 1–16.
- [18] E.D. Rainville, *Special Functions*, Chelsea, the Bronx, 1971.
- [19] J.P. Ramis, About the growth of entire functions solutions of linear algebraic q -difference equations, *Ann. Fac. Sc. Toulouse* 1 (6) (1992) 53–94.
- [20] R.V. Wallisser, On entire functions assuming integer values in a geometric sequence, in: *Théorie des Nombres* (Quebec, PQ, 1987), de Gruyter, Berlin, 1989, pp. 981–989.
- [21] E.T. Whittaker, G.N. Watson, *Modern Analysis*, 4th Edition, Cambridge University Press, Cambridge, 1927.
- [22] J.M. Whittaker, *Interpolatory Function Theory*, Cambridge Tracts in Mathematics and Mathematical Physics, No. 33, Cambridge, 1935.
- [23] J.M. Whittaker, *Sur les Séries de Base de Polynomes Quelconques*, Gauthier-Villars, Paris, 1949.