

Letter to the Editor

Am. J. Hum. Genet. 75:523–524, 2004

The Phylogeography of Mitochondrial DNA Haplogroup L3g in Africa and the Atlantic Slave Trade

To the Editor:

From the 16th to the 19th century, ~4 million slaves were transported from sub-Saharan Africa to Brazil. With the use of historical records, it is possible to estimate that ~65% of them were Bantus from west-central Africa, ~30% originated from western Africa, and ~5% came from southeastern Africa (Klein 2002).

Salas et al. (2004) have compared the phylogeography of mtDNA haplogroups in Africa with available data on Brazilians and have concluded that their results agree with these historical estimates. However, they were careful to point out that the west-central African contribution to Brazil, signaled by a high frequency of haplogroups L1c and L3e (Bortolini et al. 1997; Alves-Silva et al. 2000; Bandelt et al. 2001), derives largely from an area (the Congo basin) that, thus far, has not been thoroughly analyzed for mtDNA variation. Also, Salas et al. (2002, 2004) called attention to the presence of the haplogroup L3g—which they had only encountered in eastern African populations—in three Brazilians (among the 92 African mtDNA haplotypes that were characterized), one Colombian, and one African American individual. On the basis of this observation, they proposed the occurrence of either direct slave trade from eastern Africa to America or hitherto undetected gene flow from eastern Africa into western or southeastern Africa and then into America.

Intrigued by this proposal, we tried to identify historical evidence of direct slave trade from eastern Africa to Brazil and, indeed, found some anecdotal reports but certainly not enough to explain the significant frequency of L3g seen among African mtDNA haplogroups in Brazil (proportion, 3.3%; 95% CI, 0.7%–9.2%). We then tried to identify other potential sources for the Brazilian L3g mtDNAs. While studying the mtDNA haplotypes of 10 individuals from Cameroon (described in Da Silva et al. 1999), we identified 1 individual of undisclosed ethnic origin whose mtDNA unequivocally belonged to the L3g haplogroup (fig. 1). This finding stimulated us

to search for further mtDNA data from Cameroon, and we came across an article by Destro-Bisol et al. (2004) in which they reported 4 instances of the L3g haplogroup among 53 Ewondo individuals. Moreover, the same authors described (on the Laboratory of Molecular Anthropology Web site) another 11 instances of the L3g haplogroup in several ethnic groups (Bakaka, Bassa, Ewondo, Daba, Fali, Podowko, and Mandara) from different geographical regions of Cameroon. The 14 mtDNA sequences from Cameroon belonged to only four different haplotypes.

We incorporated the four L3g lineages from Cameroon with those from eastern Africa and obtained the median-joining network shown in figure 1. One Brazilian haplotype was identical to the most common Cameroones haplotype (seen in 8 of the 14 cases), whereas another was closely related. Both the third Brazilian and the single African American haplotype also clustered with Cameroones sequences. A noteworthy feature of the network was that there did not appear to exist any clear segregation of the Cameroones L3g haplotypes from the eastern African counterparts. This feature, plus the fact that the putative ancestral haplotype was seen in an individual from Sudan and that there is a much smaller haplotype diversity in Cameroon (0.602) as compared with eastern Africa (0.911, excluding the Hadza), suggests that the Cameroones L3g lineages might have originated from eastern Africa by transcontinental gene flow, as put forward by Salas et al. (2004) in one of their two possible scenarios. At any rate, it appears that the L3g lineages seen in America probably have their immediate origin in Cameroon or in neighboring regions and not in eastern Africa.

Acknowledgments

We are grateful to Dr. Giovanni Destro-Bisol, Dr. Valentina Coia, and Dr. Gabriella Spedini from the Laboratory of Molecular Anthropology, Department of Animal and Human Biology, University of Rome “La Sapienza,” who very kindly gave us access to their data and allowed us to use their data in this letter. Our research was supported by the Conselho Nacional de Pesquisas of Brazil and the Coordenação de Aperfeiçoamento de Pessoal de Ensino Superior.




Figure 1 Median network of 20 different L3g lineages identified in populations from sub-Saharan Africa and America. The ancestral haplotype (16223T-16293T-16311C-16355T-16362C) was identified in a single Sudanese individual by Salas et al. (2002). The branches of the network show the additional mutations that identify each haplotype. Positions are indicated as variants from the reference sequence, minus 16,000; a letter next to the position indicates a transversion. The data were compiled from the following groups: Ethiopians, Colombians, and African Americans (Salas et al. 2002); Hadzas (Vigilant et al. 1991); Turkanans (Watson et al. 1997); Nubians (Krings et al. 1999); Ewondos (Destro-Bisol et al. 2004); Bakaka, Bassa, Daba, Falis, Podowkos, and Mandara (see Web site of the Laboratory of Molecular Anthropology, University of Rome “La Sapienza”); Brazilian whites (Alves-Silva et al. 2000); Brazilian blacks (Bortolini et al. 1997); and Camerooneese (present study).

MARIA CÁTIRA BORTOLINI,¹

WILSON A. DA SILVA, JR.,²

MARCO ANTÔNIO ZAGO,³ JACQUES ELION,⁴

RAJAGOPAL KRISHNAMOORTHY,⁴

VANESSA F. GONÇALVES,⁵ AND SÉRGIO D. J. PENA⁵

¹*Departamento de Genética, Universidade Federal do Rio Grande do Sul, Porto Alegre, Brazil;*

²*Departamento de Genética and* ³*Departamento de Clínica Médica, Universidade de São Paulo, Ribeirão Preto, Brazil;*

⁴*INSERM UMR 458, Biochimie Génétique, Hôpital Robert Debré, Paris; and*

⁵*Departamento de Bioquímica e Imunologia, Universidade Federal de Minas Gerais, Belo Horizonte, Brazil*

Electronic-Database Information

The URL for data presented herein is as follows:

Laboratory of Molecular Anthropology, University of Rome

“La Sapienza,” <http://www.scienzemfn.uniroma1.it/labantro/>

References

- Alves-Silva J, da Silva Santos M, Guimarães PEM, Ferreira ACS, Bandelt H-J, Pena SDJ, Prado VF (2000) The ancestry of Brazilian mtDNA lineages. *Am J Hum Genet* 67:444–461
- Bandelt H-J, Alves-Silva J, Guimarães PEM, Santos MS, Brehm A, Pereira L, Coppa A, Larruga JM, Rengo C, Scozzari R, Torroni A, Prata MJ, Amorim A, Prado VF, Pena SDJ (2001) Phylogeography of the human mitochondrial haplogroup L3e: a snapshot of African prehistory and Atlantic slave trade. *Ann Hum Genet* 65:549–563
- Bortolini MC, Zago MA, Salzano FM, Silva-Junior WA, Bonatto SL, da Silva MC, Weimer TA (1997) Evolutionary and anthropological implications of mitochondrial DNA variation in African Brazilian populations. *Hum Biol* 69:141–159
- Da Silva WA Jr, Bortolini MC, Meyer D, Salzano FM, Elion J, Krishnamoorthy R, Schneider MPC, De Guerra DC,

- Layrisse Z, Castellano HM, Weimer TA, Zago MA (1999) Genetic diversity of two African and sixteen South American populations determined on the basis of six hypervariable loci. *Am J Phys Anthropol* 109:425–437
- Destro-Bisol G, Coia V, Boschi I, Verginelli F, Caglia A, Pascali V, Spedini G, Calafell F (2004) The analysis of variation of mtDNA hypervariable region 1 suggests that eastern and western pygmies diverged before the Bantu expansion. *Am Nat* 163:212–226
- Klein HS (2002) As origens Africanas dos escravos brasileiros. In: Pena SDJ (ed) *Homo brasilis: aspectos genéticos, linguísticos, históricos e socioantropológicos da formação do povo brasileiro*. FUNPEC, Ribeirão Preto, Brazil, pp 93–112
- Krings M, Salem AH, Bauer K, Geisert H, Malek AK, Chaix L, Simon C, Welsby D, Di Rienzo A, Utermann G, Sajantila A, Pääbo S, Stoneking M (1999) mtDNA analysis of Nile River Valley populations: a genetic corridor or a barrier to migration? *Am J Hum Genet* 64:1166–1176
- Salas A, Richards M, De la Fe T, Lareu M-V, Sobrino B, Sánchez-Diz P, Macaulay V, Carracedo Á (2002) The making of the African mtDNA landscape. *Am J Hum Genet* 71:1082–1111
- Salas A, Richards M, Lareu M-V, Scozzari R, Coppa A, Torroni A, Macaulay V, Carracedo Á (2004) The African diaspora: mitochondrial DNA and the Atlantic slave trade. *Am J Hum Genet* 74:454–465
- Vigilant L, Stoneking M, Harpending H, Hawkes K, Wilson AC (1991) African populations and the evolution of mitochondrial DNA. *Science* 253:1503–1507
- Watson E, Forster P, Richards M, Bandelt H-J (1997) Mitochondrial footprints of human expansions in Africa. *Am J Hum Genet* 61:691–704

Address for correspondence and reprints: Dr. Sérgio D. J. Pena, Departamento de Bioquímica, ICB-UFMG, Caixa Postal 486, 30161-970 Belo Horizonte, MG, Brazil. E-mail: spena@dcc.ufmg.br

© 2004 by The American Society of Human Genetics. All rights reserved.
0002-9297/2004/7503-0021\$15.00