



Contents lists available at ScienceDirect

Journal of Mathematical Analysis and Applications

www.elsevier.com/locate/jmaa


Mixed modulation spaces and their application to pseudodifferential operators

Shannon Bishop¹

School of Mathematics, Georgia Institute of Technology, Atlanta, GA 30332, USA

ARTICLE INFO

Article history:

Received 23 January 2009

Available online 26 August 2009

Submitted by L. Grafakos

Keywords:

Frames

Gabor transform

Integral operator

Modulation space

Pseudodifferential operator

Schatten class

ABSTRACT

This paper uses frame techniques to characterize the Schatten class properties of integral operators. The main result shows that if the coefficients $\{\langle k, \Phi_{m,n} \rangle\}$ of certain frame expansions of the kernel k of an integral operator are in $\ell^{2,p}$, then the operator is Schatten p -class. As a corollary, we conclude that if the kernel or Kohn–Nirenberg symbol of a pseudodifferential operator lies in a particular mixed modulation space, then the operator is Schatten p -class. Our corollary improves existing Schatten class results for pseudodifferential operators and the corollary is sharp in the sense that larger mixed modulation spaces yield operators that are not Schatten class.

© 2009 Elsevier Inc. All rights reserved.

1. Introduction

Integral operators arise naturally in many areas of mathematics and science. Pseudodifferential operators, which are a particular type of integral operator, have appeared widely in the literature of physics, signal processing and differential equations. An overview of pseudodifferential operators is given in Chapter 14 of [9], while more detailed expositions are found in [6,13,17]. Because of the role of pseudodifferential operators in partial differential equations, the smoothness of the Weyl and Kohn–Nirenberg symbols of a pseudodifferential operator has traditionally been used to characterize properties of the operator, with the Hörmander symbol classes playing key roles.

More recently, pseudodifferential operators have been studied from a time–frequency perspective. Every pseudodifferential operator is a superposition of time–frequency shifts, and the properties of pseudodifferential operators have been well described by time–frequency analysis. Results with this flavor appear in [3,18,22]. In particular the classical modulation spaces $M_w^{p,q}(\mathbb{R}^d)$, which are Banach spaces characterized by time–frequency shifts and mixed norms, have been useful symbol spaces for studying continuity and Schatten class properties of pseudodifferential operators. (See [16] for applications of mixed norms in other areas of harmonic analysis.) Using Gabor frames, elements in these spaces can be decomposed into a superposition of time–frequency shifts, and this Gabor frame decomposition of the symbol of a pseudodifferential operator can be used to characterize the properties of the operator. Results of this type appear in [2,8,10,14,19,21], while modulation spaces appear implicitly in [11,20,5,12,15].

In this paper we develop a technique for analyzing the kernel of an integral operator which both generalizes and improves existing techniques of time–frequency analysis of pseudodifferential operators, and in particular yields larger non-smooth classes of Kohn–Nirenberg symbols which ensure that a given pseudodifferential operator is Schatten p -class.

E-mail address: sbishop@math.gatech.edu.

¹ Partially supported by NSF Grant DMS-0806532.

To obtain our main result, we analyze the slices of the kernel of an integral operator with a frame. If these decomposed slices have a certain decay, then the operator is Schatten p -class. As a special case, we obtain the following theorem.

Theorem 1.1. *Suppose $\{\phi_m\}_{m \in \Lambda}$ is a frame for $L^2(\mathbb{R}^d)$. Let $\Phi_{m,n} = \phi_m \otimes \overline{\phi_n}$. If A is an integral operator with kernel k and $p \in [1, 2]$, then A is Schatten p -class on $L^2(\mathbb{R}^d)$ if*

$$\left(\sum_{n \in \Lambda} \left(\sum_{m \in \Lambda} |\langle k, \Phi_{m,n} \rangle|^2 \right)^{\frac{p}{2}} \right)^{\frac{1}{p}} < \infty.$$

Analyzing the slices of the kernel as in Theorem 1.1 with a Gabor frame in particular gives a time–frequency condition on the kernel which ensures the operator is Schatten p -class. We show that this condition holds for kernels belonging to certain Banach spaces $M(c)_{W}^{p_1, p_2, \dots, p_{2d}}$ that we call mixed modulation spaces, which are natural generalizations of the traditional modulation spaces $M_{W}^{p,q}(\mathbb{R}^d)$. In this paper we show that many of the interesting properties of traditional modulation spaces also hold for mixed modulation spaces. Furthermore, inclusion of the Kohn–Nirenberg symbol in an appropriate mixed modulation space ensures the corresponding operator is Schatten p -class. The relationship between mixed modulation spaces and the kernels and Kohn–Nirenberg symbols of Schatten p -class operators is summarized in the following theorem.

Theorem 1.2. *Let A be a pseudodifferential operator with kernel k and Kohn–Nirenberg symbol τ . Assume $p \in [1, 2]$ and set $2 = p_1 = \dots = p_{2d}$ and $p = p_{2d+1} = \dots = p_{4d}$. For suitable c , if one of k, τ lies in $M(c)^{p_1, p_2, \dots, p_{4d}}$, then so does the other. In this case A is Schatten p -class on $L^2(\mathbb{R}^d)$.*

The strongest known Schatten class result for pseudodifferential operators obtained by time–frequency analysis is Theorem 1.2 of [8], which states that if the Weyl symbol or Kohn–Nirenberg symbol of a pseudodifferential operator is in $M_{v_s}^{2,2}(\mathbb{R}^{2d})$, then the operator is Schatten p -class if $p > \frac{2d}{d+s}$ and $s \geq 0$. Although the crux of both Theorem 1.2 and [8, Theorem 1.2] is time–frequency analysis with Gabor frames, our Theorem 1.2 is obtained by analyzing the slices of the kernel with a Gabor frame, thus permitting a finer control on the properties of the kernel (and, consequently, the symbol). As a result, we can show that Theorem 1.2 is stronger than [8, Theorem 1.2], in the sense that the mixed modulation space described by Theorem 1.2 strictly contains the space $M_{v_s}^{2,2}(\mathbb{R}^{2d})$.

The mixed modulation space $M(c)^{p_1, p_2, \dots, p_{4d}}$ is characterized by $4d$ decay parameters p_1, p_2, \dots, p_{4d} , while the mixed modulation space described by Theorem 1.2 essentially only has two decay parameters. This disparity suggests that Theorem 1.2 may be extended to a larger mixed modulation space by a more subtle analysis of the kernel of a pseudodifferential operator. However, this is not the case. In fact, we show that Theorem 1.2 is sharp in the sense that larger mixed modulation spaces contain kernels and symbols of pseudodifferential operators that are not Schatten p -class.

The paper is organized as follows. Section 2 contains preliminary and background information. Section 3 is devoted to the proof of Theorem 1.1. In Section 4, the definition of mixed modulation spaces $M(c)^{p_1, p_2, \dots, p_{2d}}$ is given and the properties of these spaces are described. In Section 5, we apply the results of Sections 3 and 4 to pseudodifferential operators.

2. Preliminaries

2.1. Weight functions

Definition 2.1. A locally integrable function $v : \mathbb{R}^d \rightarrow [0, \infty)$ is called a *weight function*. A weight function $v : \mathbb{R}^d \rightarrow [0, \infty)$ is *submultiplicative* if

$$v(z_1 + z_2) \leq v(z_1)v(z_2) \quad \text{for all } z_1, z_2 \in \mathbb{R}^d.$$

A weight function v has *polynomial growth* if there are $C, s \geq 0$ such that $v(z) \leq C(1 + |z|)^s$ for all $z \in \mathbb{R}^d$.

For each $s \geq 0$, the function $v_s(z) = (1 + |z|)^s$ is a submultiplicative weight function with polynomial growth.

Definition 2.2. Suppose $w : \mathbb{R}^d \rightarrow [0, \infty)$ is a weight function and $v : \mathbb{R}^d \rightarrow [0, \infty)$ is submultiplicative. If there is a constant C such that

$$w(z_1 + z_2) \leq Cv(z_1)w(z_2) \quad \text{for all } z_1, z_2 \in \mathbb{R}^d,$$

then we call w a *v -moderate weight*.

We will assume throughout this paper that $v : \mathbb{R}^d \rightarrow [0, \infty)$ is a submultiplicative weight function of polynomial growth symmetric in each coordinate, i.e. $v(x_1, \dots, -x_i, \dots, x_d) = v(x_1, \dots, x_i, \dots, x_d)$ for each $i = 1, 2, \dots, d$. We also assume throughout that w is a v -moderate weight.

2.2. Mixed norm spaces

Definition 2.3. Given measure spaces (X_i, μ_i) and given $p_i \in [1, \infty]$ for $i = 1, 2, \dots, d$, we let $L_w^{p_1, p_2, \dots, p_d}(X_1, X_2, \dots, X_d, \mu_1, \mu_2, \dots, \mu_d)$ consist of all of the measurable functions $F : X_1 \times X_2 \times \dots \times X_d \rightarrow \mathbb{C}$ for which the following norm is finite:

$$\|F\|_{L_w^{p_1, p_2, \dots, p_d}} = \left(\int_{X_d} \dots \left(\int_{X_1} |F(x_1, \dots, x_d) w(x_1, \dots, x_d)|^{p_1} d\mu_1(x_1) \right)^{\frac{p_2}{p_1}} \dots d\mu_d(x_d) \right)^{\frac{1}{p_d}},$$

with the usual modifications for indices p_i which equal ∞ .

If the measures μ_i for all $i = 1, 2, \dots, d$ are clear from context we simply write $L_w^{p_1, p_2, \dots, p_d}(X_1, X_2, \dots, X_d)$. If $X_i = \mathbb{R}$ and μ_i is Lebesgue measure on \mathbb{R} for all $i = 1, 2, \dots, d$, then we simply write $L_w^{p_1, p_2, \dots, p_d}$. If each X_i is countable and μ_i is counting measure on X_i we simply write $\ell_w^{p_1, p_2, \dots, p_d}(X_1, X_2, \dots, X_d)$.

The mixed norm spaces $L_w^{p_1, p_2, \dots, p_d}(X_1, X_2, \dots, X_d, \mu_1, \mu_2, \dots, \mu_d)$ are generalizations of the classical spaces L^p and ℓ^p , and the proof that L^p and ℓ^p are Banach spaces can be extended to the mixed norm spaces (see [1]).

The following technical lemma will be useful in later sections.

Lemma 2.4. If $p > \frac{2d}{d+s}$ and $s \geq 0$, then $\ell_{v_s}^{2,2}(\mathbb{Z}^{2d}, \mathbb{Z}^{2d}) \subsetneq \ell^{2,p}(\mathbb{Z}^{2d}, \mathbb{Z}^{2d})$.

2.3. Schatten class operators

Definition 2.5. Suppose H is a Hilbert space and $A : H \rightarrow H$ is a linear operator. We say A is Schatten p -class and write $A \in \mathcal{I}_p(H)$ if

$$\|A\|_{\mathcal{I}_p} = \sup \left(\sum_{n \in \mathbb{N}} |(Af_n, g_n)|^p \right)^{\frac{1}{p}} < \infty,$$

where the supremum is taken over all pairs of orthonormal sequences $\{f_n\}_{n \in \mathbb{N}}, \{g_n\}_{n \in \mathbb{N}}$ in H .

Equivalently, an operator is Schatten p -class if its singular values constitute an ℓ^p sequence. Consequently, trace-class operators are exactly the Schatten 1-class operators and Hilbert–Schmidt operators are the Schatten 2-class operators. Schatten ∞ -class operators are bounded operators.

2.4. Gabor transform

Suppose $f : \mathbb{R}^d \rightarrow \mathbb{C}$ is measurable. For $x, \xi \in \mathbb{R}^d$ define the translation operator T_x and modulation operator M_ξ by

$$T_x f(t) = f(t - x) \quad \text{and} \quad M_\xi f(t) = e^{2\pi i t \cdot \xi} f(t),$$

and define the time–frequency shift $\pi_{(x,\xi)}$ by $\pi_{(x,\xi)} = M_\xi T_x$.

Definition 2.6. Fix $\phi \in S(\mathbb{R}^d)$. Given $f \in S'(\mathbb{R}^d)$, the Gabor transform of f with respect to ϕ is

$$V_\phi f(x, \xi) = \langle f, M_\xi T_x \phi \rangle, \quad x, \xi \in \mathbb{R}^d.$$

The function ϕ is called the window function of the Gabor transform.

The value of $V_\phi f(x, \xi)$ gives information about the time–frequency content of f around x in time and ξ in frequency. See [9] for background and information about the Gabor transform.

2.4.1. Gabor frames

Definition 2.7. A frame for a Hilbert space H is a sequence of elements $\{\phi_x\}_{x \in \Lambda}$ in H such that there are $A, B > 0$ with

$$A \|f\|^2 \leq \sum_{x \in \Lambda} |\langle f, \phi_x \rangle|^2 \leq B \|f\|^2$$

for all $f \in H$. In this case A, B are frame bounds. If we can take $A = B$, then $\{\phi_x\}_{x \in \Lambda}$ is a tight frame. A tight frame is Parseval if $A = B = 1$. A Gabor frame for $L^2(\mathbb{R}^d)$ is a sequence $\{M_\xi T_x \phi\}_{(x,\xi) \in \Lambda}$ that is a frame for $L^2(\mathbb{R}^d)$.

Frames give non-orthogonal expansions of elements of H in terms of the frame elements. In particular, if $\{\phi_x\}_{x \in \Lambda}$ is a tight frame for H with frame bound B , we have

$$f = B^{-1} \sum_{x \in \Lambda} \langle f, \phi_x \rangle \phi_x \quad \forall f \in H.$$

See [4] for general background on frames and [9] for examples and properties of Gabor frames. In particular, there is a tight Gabor frame for $L^2(\mathbb{R}^d)$ whose generator ϕ is a nice function, e.g., $\phi \in C_c^\infty(\mathbb{R}^d)$. However, the different statements of the Balian-Low Theorem show that the elements of a Gabor frame which offers unique expansions (i.e. a Gabor Riesz basis) necessarily have poor time–frequency localization.

2.4.2. Wilson bases

Wilson bases are orthonormal bases similar to Gabor Riesz bases in that they allow for unique, discrete expansions of the elements of $L^2(\mathbb{R}^d)$ in terms of time–frequency “molecules.” However, in contrast with Gabor Riesz bases, the elements of a Wilson bases may be well localized in time and frequency.

For each $k \in \mathbb{Z}^d, n \in (\mathbb{Z}^+)^d$ let

$$\Psi_{k,n}(t) = \psi_{k_1,n_1}(t_1) \psi_{k_2,n_2}(t_2) \cdots \psi_{k_d,n_d}(t_d),$$

where

$$\psi_{k_i,n_i}(t_i) = \begin{cases} T_{k_i} \psi(t_i) & \text{if } n_i = 0, \\ \frac{1}{\sqrt{2}} T_{\frac{k_i}{2}} (M_{n_i} + (-1)^{k_i+n_i} M_{-n_i}) \psi(t_i) & \text{if } n_i > 0. \end{cases}$$

For suitable $\psi \in L^2(\mathbb{R})$, the sequence $\{\Psi_{k,n}\}_{k \in \mathbb{Z}^d, n \in (\mathbb{Z}^+)^d}$ constitutes an orthonormal basis for $L^2(\mathbb{R}^d)$. In this case we call $\{\Psi_{k,n}\}_{k \in \mathbb{Z}^d, n \in (\mathbb{Z}^+)^d}$ the Wilson basis generated by ψ (see [9] for details).

2.4.3. Modulation spaces

Fix $\phi \in S(\mathbb{R}^d)$ and $p, q \in [1, \infty]$. Define

$$\|f\|_{M_w^{p,q}(\mathbb{R}^d)} = \|V_\phi f\|_{L_w^{p_1, p_2, \dots, p_{2d}}},$$

where $p = p_1 = p_2 = \dots = p_d$ and $q = p_{d+1} = p_{d+2} = \dots = p_{2d}$. Let

$$M_w^{p,q}(\mathbb{R}^d) = \{f \in S'(\mathbb{R}^d) : \|f\|_{M_w^{p,q}(\mathbb{R}^d)} < \infty\}.$$

Each $M_w^{p,q}(\mathbb{R}^d)$ is a modulation space. For $w = 1$ we write $M_w^{p,q}(\mathbb{R}^d) = M^{p,q}(\mathbb{R}^d)$.

The modulation space $M_w^{p,q}(\mathbb{R}^d)$ consists of functions with a particular time–frequency decay controlled by the parameters p, q and weight w .

2.5. Integral operators and pseudodifferential operators

An operator A of the form

$$Af(t) = \int_{\mathbb{R}^d} k(t, y) f(y) dy$$

is an integral operator. The function k is the kernel of A .

A pseudodifferential operator with Kohn–Nirenberg symbol τ is an operator having the form

$$K_\tau f(t) = \iint_{\mathbb{R}^{2d}} \hat{\tau}(\xi, x) M_\xi T_{-x} f(t) dx d\xi.$$

Suitable K_τ can be realized as integral operators. In particular, if we let \mathcal{F}_2 denote the partial Fourier transform on the last d variables of a function of $2d$ variables, i.e.

$$(\mathcal{F}_2 F)(x, w) = \int_{\mathbb{R}^d} F(x, y) e^{-2\pi i y \cdot w} dy \quad \text{for all } x, w \in \mathbb{R}^d,$$

then K_τ is an integral operator with kernel $k = \mathcal{F}_2^{-1} \tau \circ N$, where $N(x, y) = (x, x - y)$ for $x, y \in \mathbb{R}^d$.

In general, the time–frequency properties of the symbol of a pseudodifferential operator determine if the operator is Schatten p -class. Results of this type can be found in [8,10,15,20,21]. The strongest of these results is found in [8], in which the authors obtain estimates on the singular values of pseudodifferential operators. The following theorem is a special case of Theorem 1.2 in [8].

Theorem 2.8. Suppose A is a pseudodifferential operator with Kohn–Nirenberg symbol τ . If $\tau \in M_{V_s}^{2,2}(\mathbb{R}^{2d})$ with $p > \frac{2d}{d+s}$ and $s \geq 0$, then $A \in \mathcal{I}_p(L^2(\mathbb{R}^d))$.

3. A Schatten class result for integral operators

In this section we find a general condition on the kernel of an integral operator which ensures the operator is Schatten p -class.

Lemma 3.1. Assume $\{f_j\}_{j \in \mathbb{N}}, \{g_j\}_{j \in \mathbb{N}}$ are orthonormal sequences in $L^2(\mathbb{R}^d)$. Suppose $\{\phi_n\}_{n \in \Lambda}$ is a Parseval frame for $L^2(\mathbb{R}^d)$. For $G \in L^{2,p}(\mathbb{R}^d, \Lambda)$ define

$$T(G) = \left\{ \sum_{n \in \Lambda} \langle f_j, \phi_n \rangle \langle G(\cdot, n), g_j \rangle \right\}_{j \in \mathbb{N}}.$$

Then for all $p \in [1, 2]$, $T : L^{2,p}(\mathbb{R}^d, \Lambda) \rightarrow \ell^p(\mathbb{N})$ is bounded with $\|T\| \leq 1$.

Proof. Since $\{\phi_n\}_{n \in \Lambda}$ has frame bounds $A = B = 1$, we have $\|\phi_n\|_{L^2(\mathbb{R}^d)} \leq 1$ for all $n \in \Lambda$. Therefore

$$\begin{aligned} \|T(G)\|_{\ell^1} &= \sum_{j \in \mathbb{N}} \left| \sum_{n \in \Lambda} \langle f_j, \phi_n \rangle \langle G(\cdot, n), g_j \rangle \right| \leq \sum_{n \in \Lambda} \sum_{j \in \mathbb{N}} |\langle f_j, \phi_n \rangle| |\langle G(\cdot, n), g_j \rangle| \\ &\leq \sum_{n \in \Lambda} \left(\sum_{j \in \mathbb{N}} |\langle f_j, \phi_n \rangle|^2 \right)^{\frac{1}{2}} \left(\sum_{j \in \mathbb{N}} |\langle G(\cdot, n), g_j \rangle|^2 \right)^{\frac{1}{2}} \leq \sum_{n \in \Lambda} \|\phi_n\|_{L^2(\mathbb{R}^d)} \|G(\cdot, n)\|_{L^2(\mathbb{R}^d)} \leq \|G\|_{L^{2,1}(\mathbb{R}^d, \Lambda)} \end{aligned}$$

and

$$\begin{aligned} \|T(G)\|_{\ell^2} &= \left(\sum_{j \in \mathbb{N}} \left| \sum_{n \in \Lambda} \langle f_j, \phi_n \rangle \langle G(\cdot, n), g_j \rangle \right|^2 \right)^{\frac{1}{2}} \leq \left(\sum_{j \in \mathbb{N}} \left(\sum_{n \in \Lambda} |\langle f_j, \phi_n \rangle|^2 \right) \left(\sum_{n \in \Lambda} |\langle G(\cdot, n), g_j \rangle|^2 \right) \right)^{\frac{1}{2}} \\ &= \left(\sum_{n \in \Lambda} \sum_{j \in \mathbb{N}} |\langle g_j, G(\cdot, n) \rangle|^2 \right)^{\frac{1}{2}} \leq \left(\sum_{n \in \Lambda} \|G(\cdot, n)\|_{L^2(\mathbb{R}^d)}^2 \right)^{\frac{1}{2}} = \|G\|_{L^{2,2}(\mathbb{R}^d, \Lambda)}. \end{aligned}$$

Hence the theorem holds for $p = 1$ and $p = 2$. The Riesz–Thorin Interpolation Theorem gives the result for $p \in (1, 2)$. \square

Theorem 3.2. Suppose $\{\phi_m\}_{m \in \Lambda}$ is a Parseval frame for $L^2(\mathbb{R}^d)$. Define $\Phi_{m,n}(t, y) = \phi_m(t) \overline{\phi_n(y)}$. If A is an integral operator with kernel k , then for all $p \in [1, 2]$

$$\|A\|_{\mathcal{I}_p} \leq \left(\sum_{n \in \Lambda} \left(\sum_{m \in \Lambda} |\langle k, \Phi_{m,n} \rangle|^2 \right)^{\frac{p}{2}} \right)^{\frac{1}{p}}.$$

Proof. Suppose $\{f_j\}_{j \in \mathbb{N}}, \{g_j\}_{j \in \mathbb{N}}$ are orthonormal sequences in $L^2(\mathbb{R}^d)$. Let $G(y, n) = A\phi_n(y)$. Notice that $\langle \phi_n, A^*g_j \rangle = \langle G(\cdot, n), g_j \rangle$. Expanding f_j with the frame $\{\phi_m\}_{m \in \Lambda}$ and using the previous lemma, we have

$$\begin{aligned} \left(\sum_{j \in \mathbb{N}} |\langle Af_j, g_j \rangle|^p \right)^{\frac{1}{p}} &= \left(\sum_{j \in \mathbb{N}} |\langle f_j, A^*g_j \rangle|^p \right)^{\frac{1}{p}} = \left(\sum_{j \in \mathbb{N}} \left| \sum_{n \in \Lambda} \langle f_j, \phi_n \rangle \langle G(\cdot, n), g_j \rangle \right|^p \right)^{\frac{1}{p}} \leq \|G\|_{L^{2,p}(\mathbb{R}^d, \Lambda)} \\ &= \left(\sum_{n \in \Lambda} \|A\phi_n\|_{L^2(\mathbb{R}^d)}^p \right)^{\frac{1}{p}} = \left(\sum_{n \in \Lambda} \left(\sum_{m \in \Lambda} |\langle A\phi_n, \phi_m \rangle|^2 \right)^{\frac{p}{2}} \right)^{\frac{1}{p}} = \left(\sum_{n \in \Lambda} \left(\sum_{m \in \Lambda} |\langle k, \Phi_{m,n} \rangle|^2 \right)^{\frac{p}{2}} \right)^{\frac{1}{p}}. \end{aligned}$$

Taking the supremum of $(\sum_{j \in \mathbb{N}} |\langle Af_j, g_j \rangle|^p)^{\frac{1}{p}}$ over all such orthonormal sequences $\{f_j\}_{j \in \mathbb{N}}, \{g_j\}_{j \in \mathbb{N}}$ gives the result. \square

The proofs of Lemma 3.1 and Theorem 3.2 can be generalized to prove Theorem 1.1.

4. Mixed modulation spaces

In this section we introduce a generalization of the modulation spaces $M_w^{p,q}(\mathbb{R}^d)$. Throughout this section, we assume c is a permutation of the set $\{1, 2, \dots, 2d\}$. To simplify some notation we identify c with the bijection $c : \mathbb{R}^{2d} \rightarrow \mathbb{R}^{2d}$ given by $c(x_1, \dots, x_{2d}) = (x_{c(1)}, \dots, x_{c(2d)})$.

Definition 4.1. Suppose $\phi \in S(\mathbb{R}^d)$ and c is a permutation of $\{1, 2, \dots, 2d\}$ corresponding to the map c . Let $M(c)_w^{p_1, p_2, \dots, p_{2d}}$ be the mixed modulation space consisting of all $f \in S'(\mathbb{R}^d)$ for which

$$\|f\|_{M(c)_w^{p_1, p_2, \dots, p_{2d}}} = \|V_\phi f \circ c\|_{L_w^{p_1, p_2, \dots, p_{2d}}} < \infty.$$

When $w = 1$ we write $M(c)_w^{p_1, p_2, \dots, p_{2d}} = M(c)^{p_1, p_2, \dots, p_{2d}}$.

Notice that if c is the identity permutation and $p = p_1 = p_2 = \dots = p_d$ and $q = p_{d+1} = \dots = p_{2d}$, then $M(c)_w^{p_1, p_2, \dots, p_{2d}} = M_w^{p,q}(\mathbb{R}^d)$. Hence the mixed modulation spaces are indeed generalizations of modulation spaces. Also notice that if $p = p_1 = p_2 = \dots = p_d = p_{d+1} = \dots = p_{2d}$, then $M(c)_w^{p_1, p_2, \dots, p_{2d}} = M_w^{p,p}(\mathbb{R}^d)$ for any permutation c .

The most interesting properties of modulation spaces carry over to the mixed modulation spaces. As the proofs are basic generalizations of the proofs for modulation spaces, we state these properties without proof. See [9] for a detailed account of the properties of modulation spaces.

Definition 4.2. Suppose c is a permutation of $\{1, 2, \dots, 2d\}$. For each $x \in \mathbb{R}^{2d}$ let $\pi_x = M_{(x_{d+1}, \dots, x_{2d})} T_{(x_1, \dots, x_d)}$. For measurable $\psi : \mathbb{R}^d \rightarrow \mathbb{C}$ define an operator Υ_ψ by

$$\Upsilon_\psi F(t) = \int_{\mathbb{R}^{2d}} F(x) \pi_{c(x)} \psi(t) dx.$$

Theorem 4.3. Suppose $\psi, \gamma \in M(c)_v^{1, \dots, 1}$.

- (a) For any $f \in M(c)_w^{p_1, p_2, \dots, p_{2d}}$, we have $\Upsilon_\psi (V_\gamma f \circ c) = \langle \psi, \gamma \rangle f$.
- (b) $\|f\| = \|V_\psi f \circ c\|_{L_w^{p_1, p_2, \dots, p_{2d}}}$ is an equivalent norm on $M(c)_w^{p_1, p_2, \dots, p_{2d}}$.

Theorem 4.3(b) implies that the definition of the mixed modulation spaces is independent of the choice of window $\phi \in S(\mathbb{R}^d)$, with different windows ϕ giving equivalent norms. Furthermore, this fact also holds for ϕ in the larger space $M(c)_v^{1, \dots, 1}$. Theorem 4.3(a) shows that for Gabor window functions in $M(c)_v^{1, \dots, 1}$, there is an inversion formula valid on each $M(c)_w^{p_1, p_2, \dots, p_{2d}}$.

Corollary 4.4. For any $p_1, p_2, \dots, p_{2d} \in [1, \infty]$, $M(c)_w^{p_1, p_2, \dots, p_{2d}}$ is a Banach space.

Theorem 4.5. If $p_1, p_2, \dots, p_{2d} \in [1, \infty)$, then $M(c)_w^{p'_1, p'_2, \dots, p'_{2d}}$ is the dual space of $M(c)_w^{p_1, p_2, \dots, p_{2d}}$, where $p'_i \in [1, \infty]$ satisfies $\frac{1}{p_i} + \frac{1}{p'_i} = 1$.

The next theorem states that if the window function is nice, then a Gabor frame for $L^2(\mathbb{R}^d)$ gives bounded decompositions for all mixed modulation spaces.

Theorem 4.6. Suppose $p_1, p_2, \dots, p_{2d} \in [1, \infty]$ and $\psi \in M(c)_v^{1, \dots, 1}$. Further suppose that $\{\pi_{\alpha m} \psi\}_{m \in \mathbb{Z}^{2d}}$ is a frame for $L^2(\mathbb{R}^d)$ with dual frame $\{\pi_{\alpha m} \gamma\}_{m \in \mathbb{Z}^{2d}}$. Then

- (a) $\{\pi_{\alpha m} \psi\}_{m \in \mathbb{Z}^{2d}}$ is a Banach frame for $M(c)_w^{p_1, p_2, \dots, p_{2d}}$ and there exist $0 < A \leq B < \infty$ independent of p_1, p_2, \dots, p_{2d} with

$$A \|f\|_{M(c)_w^{p_1, p_2, \dots, p_{2d}}} \leq \|V_\psi f \circ c\|_{\alpha \mathbb{Z}^{2d}} \|e^{p_1, p_2, \dots, p_{2d}}\|_{e^{p_1, p_2, \dots, p_{2d}}} \leq B \|f\|_{M(c)_w^{p_1, p_2, \dots, p_{2d}}}$$

for all $f \in M(c)_w^{p_1, p_2, \dots, p_{2d}}$.

- (b) If $p_1, p_2, \dots, p_{2d} \in [1, \infty)$, then

$$f = \sum_{m \in \mathbb{Z}^{2d}} \langle f, \pi_{\alpha m} \psi \rangle \pi_{\alpha m} \gamma = \sum_{m \in \mathbb{Z}^{2d}} \langle f, \pi_{\alpha m} \gamma \rangle \pi_{\alpha m} \psi$$

for all $f \in M(c)_w^{p_1, p_2, \dots, p_{2d}}$ with unconditional convergence in $M(c)_w^{p_1, \dots, p_{2d}}$.

(c) If $p_1, p_2, \dots, p_{2d} \in [1, \infty]$, then

$$f = \sum_{m \in \mathbb{Z}^{2d}} \langle f, \pi_{\alpha m} \psi \rangle \pi_{\alpha m} \gamma = \sum_{m \in \mathbb{Z}^{2d}} \langle f, \pi_{\alpha m} \gamma \rangle \pi_{\alpha m} \psi$$

for all $f \in M(c)_W^{p_1, p_2, \dots, p_{2d}}$ with weak* convergence in $M(c)_{\frac{1}{v}}^{\infty, \dots, \infty}$.

Theorem 4.6 can be used to prove embeddings among the mixed modulation spaces.

Lemma 4.7. If $s \geq t$ and $p_i, r_i \in [1, \infty]$ with $p_i \leq r_i$ for all $i = 1, 2, \dots, 2d$, then $M(c)_{v_s}^{p_1, p_2, \dots, p_{2d}} \subset M(c)_{v_t}^{r_1, r_2, \dots, r_{2d}}$.

The following theorem states that Wilson bases are bases for the mixed modulation spaces.

Theorem 4.8. Let $v : \mathbb{R}^{2d} \rightarrow [0, \infty)$ be a weight and w a v -moderate weight. Define $\tilde{v}(t) = \max\{v(t, 0, \dots, 0), v(0, t, 0, \dots, 0), \dots, v(0, \dots, 0, t)\}$ for each $t \in \mathbb{R}$. Assume $\psi \in M_{\tilde{v} \otimes \tilde{v}}^{1,1}(\mathbb{R})$ generates an orthonormal Wilson basis $\{\Psi_{k,n}\}_{n \in (\mathbb{Z}^+)^d, k \in \mathbb{Z}^d}$ for $L^2(\mathbb{R}^d)$. Then $\{\Psi_{k,n}\}_{n \in (\mathbb{Z}^+)^d, k \in \mathbb{Z}^d}$ is an unconditional basis for $M(c)_W^{p_1, p_2, \dots, p_{2d}}$ for each $p_1, p_2, \dots, p_{2d} \in [1, \infty)$.

Corollary 4.9. Let $X_1 = X_2 = \dots = X_d = \mathbb{Z}$ and $X_{d+1} = X_{d+2} = \dots = X_{2d} = \mathbb{Z}^+$. Then the map

$$f \rightarrow \left\{ \langle f, \Psi_{(n_{c(1)}, n_{c(2)}, \dots, n_{c(d)}), (n_{c(d+1)}, \dots, n_{c(2d)})} \rangle \right\}_{n_1 \in X_{c^{-1}(1)}, n_2 \in X_{c^{-1}(2)}, \dots, n_{2d} \in X_{c^{-1}(2d)}}$$

is an isomorphism from $M(c)_W^{p_1, p_2, \dots, p_{2d}}$ to $\ell_W^{p_1, p_2, \dots, p_{2d}}(X_{c^{-1}(1)}, \dots, X_{c^{-1}(2d)})$.

5. Pseudodifferential operators and Schatten classes

In this section we will use Theorem 3.2 to find conditions on the kernel and Kohn–Nirenberg symbol of a pseudodifferential operator that guarantee the operator is Schatten p -class.

We are particularly interested in permutations c of $\{1, 2, \dots, 4d\}$ satisfying the following:

- (a) c maps $\{1, 2, \dots, d, 2d + 1, 2d + 2, \dots, 3d\}$ to $\{1, 2, \dots, 2d\}$ bijectively and
- (b) c maps $\{d + 1, d + 2, \dots, 2d, 3d + 1, 3d + 2, \dots, 4d\}$ to $\{2d + 1, \dots, 4d\}$ bijectively.

We call such permutations *slice permutations* because they relate nicely to the slice analysis of Section 3.

Corollary 5.1. Assume c is a slice permutation. Let $2 = p_1 = p_2 = \dots = p_{2d}$ and $p = p_{2d+1} = \dots = p_{4d}$. If $p \in [1, 2]$, $k \in M(c)^{p_1, \dots, p_{4d}}$ and A is an integral operator with kernel k , then $A \in \mathcal{I}_p(L^2(\mathbb{R}^d))$.

Proof. Let $\{\pi_{\alpha m} \phi\}_{m \in \mathbb{Z}^{2d}} = \{\phi_m\}_{m \in \mathbb{Z}^{2d}}$ be a Parseval Gabor frame for $L^2(\mathbb{R}^d)$ with $\phi \in M^{1,1}(\mathbb{R}^d)$ and let $\Phi(t, y) = \phi(t) \overline{\phi(y)}$. Then $\Phi \in M^{1,1}(\mathbb{R}^{2d})$. Let $\Phi_{m,n}(t, y) = \phi_m(t) \overline{\phi_n(y)}$. By Lemma 3.2 in [11], $\{\Phi_{m,n}\}_{m,n \in \mathbb{Z}^{2d}}$ is a Parseval frame for $L^2(\mathbb{R}^{2d})$. For $m_1, m_2, n_1, n_2 \in \mathbb{Z}^d$, with $m = (m_1, m_2)$ and $n = (n_1, n_2)$, we have

$$\langle k, \Phi_{m,n} \rangle = V_\phi k(\alpha m_1, \alpha n_1, \alpha m_2, \alpha n_2).$$

For each slice permutation c , we see that

$$\begin{aligned} \left(\sum_{n \in \mathbb{Z}^{2d}} \left(\sum_{m \in \mathbb{Z}^{2d}} |\langle k, \Phi_{m,n} \rangle|^2 \right)^{\frac{p}{2}} \right)^{\frac{1}{p}} &= \left(\sum_{n_1, n_2 \in \mathbb{Z}^d} \left(\sum_{m_1, m_2 \in \mathbb{Z}^d} |V_\phi k(\alpha m_1, \alpha n_1, \alpha m_2, \alpha n_2)|^2 \right)^{\frac{p}{2}} \right)^{\frac{1}{p}} \\ &= \left(\sum_{n_1, n_2 \in \mathbb{Z}^d} \left(\sum_{m_1, m_2 \in \mathbb{Z}^d} |V_\phi k(c(\alpha m_1, \alpha m_2, \alpha n_1, \alpha n_2))|^2 \right)^{\frac{p}{2}} \right)^{\frac{1}{p}} \\ &\leq B \|k\|_{M(c)^{p_1, p_2, \dots, p_{4d}}}, \end{aligned}$$

where B is the constant ensured by Theorem 4.6(a). Hence if $k \in M(c)^{p_1, \dots, p_{4d}}$, then, by Theorem 3.2, $A \in \mathcal{I}_p(L^2(\mathbb{R}^d))$. \square

We can extend Corollary 5.1 to conditions on the symbol of a pseudodifferential operator.

Theorem 5.2. Let A be a pseudodifferential operator with kernel k and Kohn–Nirenberg symbol τ . Assume $p \in [1, 2]$ and set $2 = p_1 = \dots = p_{2d}$ and $p = p_{2d+1} = \dots = p_{4d}$. If c is a slice permutation and one of k, τ lies in $M(c)^{p_1, p_2, \dots, p_{4d}}$, then so does the other. In this case $A \in \mathcal{I}_p(L^2(\mathbb{R}^d))$.

Proof. Using the fact that $k = \mathcal{F}_2^{-1}\tau \circ N$ we can show that $|\langle k, M_{(z,t)}T_{(x,y)}\Phi \rangle| = |\langle \tau, M_{(z+t,-y)}T_{(x,-t)}\mathcal{F}_2(\Phi \circ N^{-1}) \rangle|$ for all $x, y, z, t \in \mathbb{R}^d$. Hence

$$\begin{aligned} \|k\|_{M(c)^{p_1,p_2,\dots,p_{4d}}} &= \left(\iint \left(\iint | \langle k, M_{(z,t)}T_{(x,y)}\Phi \rangle |^2 dx dz \right)^{\frac{p}{2}} dy dt \right)^{\frac{1}{p}} \\ &= \left(\iint \left(\iint | \langle \tau, M_{(z+t,-y)}T_{(x,-t)}\mathcal{F}_2(\Phi \circ N^{-1}) \rangle |^2 dx dz \right)^{\frac{p}{2}} dy dt \right)^{\frac{1}{p}} \\ &= \left(\iint \left(\iint | \langle \tau, M_{(z,y)}T_{(x,t)}\mathcal{F}_2(\Phi \circ N^{-1}) \rangle |^2 dx dz \right)^{\frac{p}{2}} dy dt \right)^{\frac{1}{p}} \\ &\asymp \|\tau\|_{M(c)^{p_1,p_2,\dots,p_{4d}}}. \quad \square \end{aligned}$$

Theorem 5.2 is stronger than the previously known Theorem 2.8, as the following lemma shows.

Lemma 5.3. *Let c be a slice permutation and let $2 = p_1 = \dots = p_{2d}, p = p_{2d+1} = \dots = p_{4d}$. If $p > \frac{2d}{d+s}$ with $s \geq 0$, then $M_{v_s}^{2,2}(\mathbb{R}^{2d}) \subsetneq M(c)^{p_1,p_2,\dots,p_{4d}}$.*

Proof. Let $X_1 = X_2 = \dots = X_{2d} = \mathbb{Z}$ and $X_{2d+1} = X_{2d+2} = \dots = X_{4d} = \mathbb{Z}^+$ and define

$$S(f) = \left\{ \langle f, \Psi_{(n_{c(1)}, \dots, n_{c(2d)}), (n_{c(2d+1)}, \dots, n_{c(4d)})} \rangle \right\}_{n_1 \in X_{c^{-1}(1)}, n_2 \in X_{c^{-1}(2)}, \dots, n_{4d} \in X_{c^{-1}(4d)}}.$$

Since $M_{v_s}^{2,2}(\mathbb{R}^{2d}) = M(c)^{2,2,\dots,2}$, Corollary 4.9 implies that

$$S : M_{v_s}^{2,2}(\mathbb{R}^{2d}) \rightarrow \ell_{v_s}^{2,2,\dots,2}(X_{c^{-1}(1)}, \dots, X_{c^{-1}(4d)})$$

and

$$S : M(c)^{p_1,p_2,\dots,p_{4d}} \rightarrow \ell^{p_1,p_2,\dots,p_{4d}}(X_{c^{-1}(1)}, \dots, X_{c^{-1}(4d)})$$

are isomorphisms. Furthermore, by Lemma 2.4, we have

$$\ell_{v_s}^{2,2,\dots,2}(X_{c^{-1}(1)}, \dots, X_{c^{-1}(4d)}) \subsetneq \ell^{p_1,p_2,\dots,p_{4d}}(X_{c^{-1}(1)}, \dots, X_{c^{-1}(4d)})$$

for $p > \frac{2d}{d+s}$ with $s \geq 0$. Hence we obtain the following diagram

$$\begin{array}{ccc} M_{v_s}^{2,2}(\mathbb{R}^{2d}) & & M(c)^{p_1,p_2,\dots,p_{4d}} \\ \downarrow S & & \downarrow S \\ \ell_{v_s}^{2,2,\dots,2}(X_{c^{-1}(1)}, \dots, X_{c^{-1}(4d)}) & \hookrightarrow & \ell^{p_1,p_2,\dots,p_{4d}}(X_{c^{-1}(1)}, \dots, X_{c^{-1}(4d)}) \end{array}$$

Since S is an isomorphism, the result follows. \square

By Lemma 4.7, increasing any one of the exponent parameters p_1, \dots, p_{4d} or decreasing the weight parameter s yields a mixed modulation space larger than $M(c)_{v_s}^{p_1,p_2,\dots,p_{4d}}$. The next theorem shows Theorem 5.2 is sharp in the following sense: increasing the exponent parameters or decreasing the weight parameter of the mixed modulation space in Theorem 5.2 gives a larger mixed modulation space, but pseudodifferential operators with kernels or Kohn–Nirenberg symbols in this larger space need not be Schatten class.

Theorem 5.4. *Assume $s \leq 0, p_1, \dots, p_{2d} \in [2, \infty], p_{2d+1}, \dots, p_{4d} \in [p, \infty]$ and c is a slice permutation. Assume at least one of the following is true:*

- (a) $s < 0$.
- (b) At least one of p_1, \dots, p_{2d} is larger than 2.
- (c) At least one of p_{2d+1}, \dots, p_{4d} is larger than p .

If $1 \leq p \leq 2$, then there are pseudodifferential operators with kernels in $M(c)_{v_s}^{p_1,p_2,\dots,p_{4d}}$ and pseudodifferential operators with Kohn–Nirenberg symbols in $M(c)_{v_s}^{p_1,p_2,\dots,p_{4d}}$ that are not in $\mathcal{I}_p(L^2(\mathbb{R}^d))$.

Proof. Suppose A is a pseudodifferential operator with kernel k and Kohn–Nirenberg symbol τ . Since

$$|(k, M_{(z,t)}T_{(x,y)}\Phi)| = |(\tau, M_{(z+t,-y)}T_{(x,-t)}\mathcal{F}_2(\Phi \circ N^{-1}))|,$$

it follows that for each slice permutation c , there is a slice permutation \tilde{c} with $\|\tau\|_{M(c)_{V_s}^{p_1, p_2, \dots, p_{4d}}} \asymp \|k\|_{M(\tilde{c})_{V_s}^{p_1, p_2, \dots, p_{4d}}}$. Hence it suffices to show that for each slice permutation c , there are pseudodifferential operators with kernels in $M(c)_{V_s}^{p_1, p_2, \dots, p_{4d}}$ that are not Schatten p -class.

To avoid complicated notation, we prove the theorem only for the permutation $c(x_1, \dots, x_{4d}) = (x_{d+1}, \dots, x_{2d}, x_{3d+1}, \dots, x_{4d}, x_1, \dots, x_d, x_{2d+1}, \dots, x_{3d})$. The result is proven similarly for other slice permutations.

In the case that (a) or (b) holds, we can adapt some of the arguments in [7] to complete the proof. In particular, if $k(t, y) = k_1(t)k_2(y)$ is the kernel of an integral operator A , then $Af = \langle f, \overline{k_2} \rangle k_1$. Hence if $k_1 \notin L^2(\mathbb{R}^d)$, then A does not map into $L^2(\mathbb{R}^d)$, and if $k_2 \notin L^2(\mathbb{R}^d)$, then $A : L^2(\mathbb{R}^d) \rightarrow L^2(\mathbb{R}^d)$ is not bounded. Let c' be the permutation with associated bijection $c'(n_1, \dots, n_{2d}) = (n_{d+1}, \dots, n_{2d}, n_1, \dots, n_d)$. If (a) holds, choose $k_1 \in M_{V_s}^{2,2}(\mathbb{R}^d) \setminus L^2(\mathbb{R}^d)$ and $k_2 \in M^{p,p}(\mathbb{R}^d)$. If (b) holds, choose $k_1 \in M(c')^{p_1, \dots, p_{2d}} \setminus L^2(\mathbb{R}^d)$ and $k_2 \in M(c')^{p_{2d+1}, \dots, p_{4d}}$. In either case $k(t, y) = k_1(t)k_2(y) \in M(c)_{V_s}^{p_1, p_2, \dots, p_{4d}}$, but the integral operator with kernel k is not a bounded operator on $L^2(\mathbb{R}^d)$.

Hence we assume (c) is true. Choose $\lambda \in \ell^{p_{2d+1}, \dots, p_{3d}, p_{3d+1}, \dots, p_{4d}}((\mathbb{Z}^+)^d, \mathbb{Z}^d) \setminus \ell^{p,p}((\mathbb{Z}^+)^d, \mathbb{Z}^d)$. Assume $\{\psi_{j,l}\}_{j \in \mathbb{Z}^d, l \in (\mathbb{Z}^+)^d}$ is a Wilson basis for $L^2(\mathbb{R}^d)$ generated by $\psi \in M^{1,1}(\mathbb{R})$. Then

$$\{\Psi_{(j_1, j_2), (l_1, l_2)}\}_{j_1, j_2 \in \mathbb{Z}^d, l_1, l_2 \in (\mathbb{Z}^+)^d} = \{\psi_{j_1, l_1} \otimes \psi_{j_2, l_2}\}_{j_1, j_2 \in \mathbb{Z}^d, l_1, l_2 \in (\mathbb{Z}^+)^d}$$

is a Wilson basis for $L^2(\mathbb{R}^{2d})$ generated by $\psi \in M^{1,1}(\mathbb{R})$. Set

$$k(t, y) = \sum_{j \in \mathbb{Z}^d} \sum_{l \in (\mathbb{Z}^+)^d} \lambda_{l,j} \psi_{j,l}(t) \psi_{j,l}(y).$$

Then

$$\begin{aligned} \Psi_{(n_{c(1)}, n_{c(2)}, \dots, n_{c(2d)}), (n_{c(2d+1)}, \dots, n_{c(4d)})} &= \Psi_{(n_{d+1}, \dots, n_{2d}, n_{3d+1}, \dots, n_{4d}), (n_1, \dots, n_d, n_{2d+1}, \dots, n_{3d})} \\ &= \psi_{(n_{d+1}, \dots, n_{2d}), (n_1, \dots, n_d)} \otimes \psi_{(n_{3d+1}, \dots, n_{4d}), (n_{2d+1}, \dots, n_{3d})}. \end{aligned}$$

By Corollary 4.9

$$\begin{aligned} \|k\|_{M(c)_{V_s}^{p_1, p_2, \dots, p_{4d}}} &\asymp \left(\sum_{n_{4d} \in X_{c^{-1}(4d)}} \dots \left(\sum_{n_1 \in X_{c^{-1}(1)}} |k, \Psi_{(n_{c(1)}, \dots, n_{c(2d)}), (n_{c(2d+1)}, \dots, n_{c(4d)})}|^{p_1} \right)^{\frac{p_2}{p_1}} \dots \right)^{\frac{1}{p_{4d}}} \\ &= \left(\sum_{n_{4d} \in \mathbb{Z}} \left(\dots \left(\sum_{n_{2d+1} \in \mathbb{Z}^+} |\lambda_{(n_{2d+1}, \dots, n_{3d}), (n_{3d+1}, \dots, n_{4d})}|^{p_{2d+1}} \right)^{\frac{p_{4d}+2}{p_{2d+1}}} \dots \right)^{\frac{p_{4d}}{p_{4d-1}}} \right)^{\frac{1}{p_{4d}}} \\ &= \|\lambda\|_{\ell^{p_{2d+1}, \dots, p_{3d}, p_{3d+1}, \dots, p_{4d}}((\mathbb{Z}^+)^d, \mathbb{Z}^d)} \end{aligned}$$

so $k \in M(c)_{V_s}^{p_1, p_2, \dots, p_{4d}} \subset M(c)_{V_s}^{p_1, p_2, \dots, p_{4d}}$. The pseudodifferential operator A with kernel k has singular values equal to the elements of the sequence λ . Hence $A \notin \mathcal{I}_p(L^2(\mathbb{R}^d))$. \square

Notice that the proof of the previous theorem shows that Theorem 5.2 does not hold for $p > 2$. That is, if $p > 2$ and $k \in M(c)^{2, 2, \dots, 2, p, \dots, p}$, the corresponding integral operator may not even be bounded on $L^2(\mathbb{R}^d)$.

Acknowledgment

The author thanks Christopher Heil for helpful discussions.

References

- [1] A. Benedek, R. Panzone, The spaces L^p , with mixed norm, Duke Math. J. 28 (1961) 301–324.
- [2] W. Czaja, Boundedness of pseudodifferential operators on modulation spaces, J. Math. Anal. Appl. 284 (2003) 389–396.
- [3] W. Czaja, Z. Rzeszutnik, Pseudodifferential operators and Gabor frames: Spectral asymptotics, Math. Nachr. 233/234 (2002) 77–88.
- [4] O. Christensen, An Introduction to Frames and Riesz Bases, Birkhäuser, Boston, 2002.
- [5] I. Daubechies, On the distributions corresponding to bounded operators in the Weyl quantization, Comm. Math. Phys. 75 (1980) 229–238.
- [6] G. Folland, Harmonic Analysis in Phase Space, Princeton Univ. Press, Princeton, NJ, 1989.
- [7] K. Gröchenig, C. Heil, Counterexamples for boundedness of pseudodifferential operators, Osaka J. Math. 41 (2004) 681–691.
- [8] K. Gröchenig, C. Heil, Modulation spaces and pseudodifferential operators, Integral Equations Operator Theory 34 (1999) 439–457.
- [9] K. Gröchenig, Foundations of Time–Frequency Analysis, Birkhäuser, Boston, 2001.
- [10] C. Heil, Integral operators, pseudodifferential operators and Gabor frames, in: Advances in Gabor Analysis, Birkhäuser, Boston, 2003.
- [11] C. Heil, J. Ramanathan, P. Topiwala, Singular values of compact pseudodifferential operators, J. Funct. Anal. 150 (1997) 426–452.

- [12] L. Hörmander, The Weyl calculus of pseudodifferential operators, *Comm. Pure Appl. Math.* 32 (1979) 360–444.
- [13] L. Hörmander, *The Analysis of Partial Differential Equations*, Springer-Verlag, Berlin, 1990.
- [14] D. Labate, Pseudodifferential operators on modulation spaces, *J. Math. Anal. Appl.* 262 (2001) 242–255.
- [15] R. Rochberg, K. Tachizawa, Pseudodifferential operators, Gabor frames and local trigonometric bases, in: *Gabor Analysis and Algorithms*, Birkhäuser, Boston, 1998.
- [16] A. Stefanov, R. Torres, Calderón–Zygmund operators on mixed Lebesgue spaces and applications to null forms, *J. Lond. Math. Soc.* 70 (2004) 447–462.
- [17] E. Stein, *Harmonic Analysis*, Princeton Univ. Press, Princeton, NJ, 1993.
- [18] K. Tachizawa, The pseudodifferential operators and Wilson bases, *J. Math. Pures Appl.* 9 (1996) 509–529.
- [19] K. Tachizawa, The boundedness of pseudodifferential operators on modulation spaces, *Math. Nachr.* 168 (1994) 263–277.
- [20] J. Sjöstrand, An algebra of pseudodifferential operators, *Math. Res. Lett.* 1 (1994) 185–192.
- [21] J. Toft, Continuity and Schatten Properties for Pseudo-Differential Operators on Modulation Spaces, *Oper. Theory Adv. Appl.*, vol. 172, 2007, pp. 173–206.
- [22] M. Wong, Trace class Weyl transforms, in: *Pseudo-Differential Operators and Related Topics*, Birkhäuser, Basel, 2005.