

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

Topology and its Applications 143 (2004) 15–26

Topology
and its
Applicationswww.elsevier.com/locate/topol

On Kontsevich integral of torus knots

Julien Marché

*Institut de Mathématiques de Jussieu, Équipe “Topologie et Géométries Algébriques” Case 7012,
Université Paris VII, 75251 Paris cedex 05, France*

Received 13 October 2003; accepted 2 January 2004

Abstract

We study the unwheeled rational Kontsevich integral of torus knots. We give a precise formula for these invariants up to loop degree 3 and show that they appear as colorings of simple diagrams. We show that they behave under cyclic branched coverings in a very simple way. Our proof is combinatorial: it uses the results of Wheels and Wheelings and new decorations of diagrams.

© 2004 Elsevier B.V. All rights reserved.

MSC: 57M27

Keywords: Finite type invariants; Kontsevich integral; Torus knots; Wheels and Wheeling; Rationality

1. Introduction and notations

This article is divided in 4 parts: the first one explains the notations used in the sequel and some facts known about rationality, the second part is a computation of the unwheeled Kontsevich integral of torus knots up to loop degree 3 using formal series which encode series of diagrams. Then, in the third part, we compute a rational form of the preceding expression and show that it appears as a coloring of chain diagrams as it is suggested by Fig. 1. As a consequence of this computation, we show that the operator Lift_r which corresponds to cyclic branched coverings of S^3 along the knot simply acts on a diagram D of loop degree lower than 3 by multiplying it by $r^{-\chi(D)}$ where χ is the Euler characteristic.

The computation of the 2-loop part of torus knots has been done independently by Tomotada Ohtsuki in [8]. We would like to thank Marcos Marino for pointing out a mistake in the last formula of this article.

E-mail address: marche@math.jussieu.fr (J. Marché).

0166-8641/\$ – see front matter © 2004 Elsevier B.V. All rights reserved.

doi:10.1016/j.topol.2004.01.006

$$Z(K_{p,q}) = \exp(\text{---} \bigcirc \text{---} \dots + \bigcirc \text{---} \bigcirc \text{---} \dots + \bigcirc \text{---} \bigcirc \text{---} \bigcirc \text{---} \dots + \dots)$$

Fig. 1. Diagrams appearing in the unweheled Kontsevich integral of torus knots.

1.1. Normalizations of the Kontsevich integral

Let K be a knot in S^3 and suppose that K has a banded structure with self-linking 0. We will note $Z(K)$ the Kontsevich integral of K in the algebra \mathcal{A} of trivalent diagrams lying on a circle.

Let \mathcal{B} be the algebra of uni-trivalent diagrams. It is well known that the Poincaré–Birkhoff–Witt map $\chi : \mathcal{B} \rightarrow \mathcal{A}$ is an isomorphism but not an algebra isomorphism. We will note σ its inverse.

If U is the trivial knot, we define $\Omega = \sigma Z(U)$. The map $\Upsilon = \chi \circ \partial_\Omega : \mathcal{B} \rightarrow \mathcal{A}$ defined by instance in [11] is known to be an algebra isomorphism. The quantity $Z^{\text{un}}(K) = \Upsilon^{-1} Z(K)$ will be called unweheled Kontsevich integral and behaves better than $\sigma Z(K)$ under connected sum and cyclic branched coverings.

For each knot K , the quantities $Z(K)$, $\sigma Z(K)$ and $Z^{\text{un}}(K)$ are group-like, which means that they are exponentials of a series of connected diagrams. We will note with a small z the logarithm of these quantities.

1.2. Loop degree and rationality

If D is a connected diagram of \mathcal{B} , its first Betti number defines a degree called loop degree. The loop degree 1 part of $\sigma Z(K)$ or $Z^{\text{un}}(K)$ is well known: it only depends on the Alexander polynomial of K . For the higher degrees, very few is known. There are formulas for the 2-loop part of small knots in Rozansky’s table (see [10]), and we can find in [2] a formula for the 2-loop part of untwisted whitehead doubles. In the sequel, we give a formula for the 2-loop and 3-loop parts of the Kontsevich integral of torus knots.

In order to make precise computations, we will need the following formalism: let \mathcal{H} be a cocommutative Hopf algebra (or a Hopf algebra up to completion) and \mathcal{M} be an algebra over \mathcal{H} . We consider a space of diagrams noted $\mathcal{D}(\mathcal{H}, \mathcal{M})$ which was defined in [12]. It is roughly obtained by decorating the edges of a diagram by elements of \mathcal{M} and allowing elements of \mathcal{H} to slide through vertices thanks to the coproduct law.

We know that the space \mathcal{B} is isomorphic to $\mathcal{D}(\mathbb{Q}[[h]], \mathbb{Q}[[h]])$. In particular, the diagrams of loop degree 1 appear as colorings of the circle by an even power series without constant term. In the following, we will call wheels such diagrams and identify a power series with the wheel series it represents.

Let $f(x)$ be the power series defined by $\frac{1}{2} \log \frac{\sinh x/2}{x/2}$. The famous wheel formula (see [11]) states that $\sigma z(U) = f(x)$. Further, it was shown in [5] that the loop degree 1 part of $\sigma z(K)$ is $f(x) + Wh_K(x)$ where $Wh_K(x) = -\frac{1}{2} \log \Delta(e^x)$ and Δ is the Alexander polynomial of K .

As we are interested in the higher loop degree part, we need to recall the rationality theorem which was proved in [4]:

Let $\Lambda = \mathbb{Q}[t, t^{-1}]$ and Λ_{loc} the localization of Λ with respect to elements f satisfying $f(1) = 1$. The substitution $t = \exp(h)$ gives a morphism between couples $(\Lambda, \Lambda_{\text{loc}})$ and $(\mathbb{Q}[[h]], \mathbb{Q}[[h]])$, hence a morphism between $\mathcal{D}(\Lambda, \Lambda_{\text{loc}})$ and \mathcal{B} .

We call Hair this application: the rationality theorem tells us that the series $\sigma_Z(K)$ minus the loop degree 1 part lies in the image of the Hair map. More precisely, there is an element $Z^{\text{rat}}(K) \in \mathcal{D}(\Lambda, \Lambda_{\text{loc}})$ whose denominators on each edge is at most $\Delta(K)$ such that $\sigma Z(K) = \exp(f(x) + Wh_K(x)) \text{Hair } Z^{\text{rat}}(K)$. Respectively, there is such an element for the unwheeled invariant but we will give the formula later.

A construction, developed in [4] gives $Z^{\text{rat}}(K)$ as an invariant of K which is not automatic because the Hair map is not injective, although it is in small degrees (see [9]).

2. Computation of the torus knot integral up to loop degree 3

Let p and q be two coprime integers such that $p > 0$. We note $K_{p,q}$ the torus banded knot with parameters p and q and self-linking 0, and $L_{p,q}$ the torus banded knot with banding parallel to the torus on which it lies. This knot has self-linking pq , and his Kontsevich integral is a bit easier to compute.

The method of computation is inspired from [6]: we first compute the Kontsevich integral of the following braid.

Let p points be lying on the vertices of a regular p -gone. We note γ be the braid obtained by rotating the whole picture by an angle $2\pi \frac{q}{p}$.

Let associate to any one-dimensional manifold Γ the space $\mathcal{A}(\Gamma)$ of trivalent diagrams lying on Γ . It defines a contravariant functor with respect to continuous maps relative to boundaries. Let ϕ_p^* be the map induced by the trivial projection $[0, 1]^p \rightarrow [0, 1]$ and \curvearrowright be the only degree 1 diagram in $\mathcal{A}([0, 1])$. Then a direct computation of monodromy of the K - Z connection shows that γ has a Kontsevich integral equal to $\phi_p^*(\exp_{\#}(\frac{q}{2p} \curvearrowright))$.

The banded knot $L_{p,q}$ is obtained by closing the previous braid: this translates diagrammatically to the following: let ψ_p be the map from S^1 to itself defined by $\psi_p(z) = z^p$. Then, $Z(L_{p,q}) = \psi_p^*(v \# \exp_{\#}(\frac{q}{2p} \curvearrowright))$, where $v = Z(U)$.

By Lemma 4.10 of [11], the map ψ_p^* viewed in \mathcal{B} has the following form: if $D \in \mathcal{B}$ has k legs (i.e., univalent vertices) then $\sigma \psi_p^* \chi D = p^k D$. We will note more simply D_p the result of this operation which looks like a change of variable.

Then, to compute $Z(K_{p,q})$ from $Z(L_{p,q})$, we only need to change the framing, that is $Z(K_{p,q}) = \exp_{\#}(-\frac{pq}{2} \curvearrowright) \# Z(L_{p,q})$. We will transform this product in the usual one by applying the unwheeling map Υ^{-1} . As a result, we will have a formula for $Z^{\text{rat}}(K_{p,q})$.

We now sum up the steps of the computation:

- (1) Computation of $\sigma(v \# \exp_{\#}(\frac{q}{2p} \curvearrowright))$;
- (2) Change of variables $x \mapsto px$;
- (3) Unwheeling.

2.1. Step 1

From this point, we will be interested only in diagrams with loop degree 1, 2 or 3 as we were not able till now to perform a computation for general loop degree.

It is clear that all the operations considered below as #, ∂, etc., . . . , cannot decrease the loop degree. Then it is licit to quotient the spaces of diagrams by all diagrams with loop degree greater than 3.

There is a less licit simplification we make: we also quotient by all diagrams without legs or not lying on a one-dimensional manifold. Indeed, the unwheeled invariants may have such elements and we will justify this choice in the Section 3.3.

We recall that $\Upsilon = \chi \circ \partial_\Omega$ is an algebra isomorphism and that $\Upsilon^{-1}\nu = \Omega$ and $\Upsilon^{-1}\smile = \frown$ (modulo closed diagrams).

Then, what we have to compute is

$$\sigma\left(\nu \# \exp_{\#}\left(\frac{q}{2p} \frown\right)\right) = \partial_\Omega\left(\Omega \exp\left(\frac{q}{2p} \frown\right)\right).$$

As this expression is group-like, we just have to find all connected diagrams appearing in it. To do so, we choose a system of notations to name these diagrams (or combinations of them) as it is shown in Fig. 2.

- We recall that x^n is a wheel with n legs (n is even).
- Let $x^n y^m$ represent two wheels glued on one edge, with n remaining legs on the left and m remaining legs on the right. In particular, n and m are odd and $x^n y^m = x^m y^n$.
- The expression $z_1^{m_1} x^n z_2^{m_2}$ represents the coloring of a diagram with three wheels joined by two edges, with n legs on the central wheel and m_1 and m_2 legs on the other wheels. By convention, we put the variable associated to the middle wheel between the two others.
- Finally, we note $[x^n, x^m]$ the sum of all diagrams obtained by gluing two wheels of size n and m in two points.

In the computation of $\partial_\Omega(\Omega \exp(\frac{q}{2p} \frown))$, we will concentrate on the wheels of the derived term and make the following observation: such a wheel can only be glued to another wheel, and each gluing increase the loop degree by one. Hence, there can be at most two gluings. We want to separate the diagrams obtained by gluing these wheels in two points. To compute these contributions, we state the following very useful lemma.




Fig. 2. Examples of diagrams.

Lemma 1.

- Let $f(x)$ and $g(y)$ be two power series of wheels. The series obtained by gluing them in one point is $f'(x)g'(y)$.
- Let $f(x)$ and $g(y)h(z)$ be two series of diagrams. The series obtained by gluing them in one point is $f'(x)g'(y)h(z) + f'(x)h'(z)g(y)$.

The proof is straightforward and has a first application in the following lemma:

Lemma 2. The expression obtained by gluing the derived wheels in two points is

$$\frac{p^2}{q^2} \left[f\left(\frac{q}{p}x\right), f(x) \right] + \frac{1}{2} f'\left(\frac{q}{p}z_1\right) f''(x) f'\left(\frac{q}{p}z_2\right).$$

Proof. Recall that $\Omega = \exp(f(x))$. The first term comes from the gluing of a right wheel to the same left wheel. We have to fill the remaining left legs with the $\frac{q}{2p}$ struts in the only way such that it does not increase the loop degree. Finally, we just multiply by $\frac{q}{p}$ to the power the number of remaining legs. There are as many legs as the left wheel, minus two, which justifies the first part of the formula.

For the second term, we just apply twice the Lemma 1. \square

We can suppose now that each wheel can be glued in at most one leg. Let us examine precisely the following expression:

$$\partial_\Omega \left(\Omega \exp\left(\frac{q}{2p}\curvearrowright\right) \right) = \left\langle \Omega_x, \Omega_{x+y} \exp\left(\frac{q}{2p}x+y \curvearrowright_{x+y}\right) \right\rangle_x.$$

We can replace $\exp(\omega_{x+y})$ by the series $\exp(\omega_y + \hat{\omega}_x)$. The series $\hat{\omega}_x$ is obtained by replacing a wheel by the sum of the coloring of its legs by one x (and the other ones y).

We use a famous trick concerning the doubling of the strut part to get:

$$\begin{aligned} \partial_\Omega \left(\exp\left(\omega + \frac{q}{2p}\curvearrowright\right) \right) &= \left\langle \Omega_x, \exp\left(\omega_y + \hat{\omega}_x + \frac{q}{2p}x \curvearrowright_x + \frac{q}{p}x \curvearrowright_y + \frac{q}{2p}y \curvearrowright_y\right) \right\rangle_x \\ &= \left\langle \partial_{\exp(\hat{\omega}_x)} \partial_{\exp(\frac{q}{2p}x \curvearrowright_x)} \Omega_x, \exp\left(\frac{q}{p}x \curvearrowright_y\right) \right\rangle_x \left\langle \exp\left(\frac{q}{2p}y \curvearrowright_y\right) \Omega_y \right\rangle_x \\ &= \left\langle \partial_{\exp(\hat{\omega}_x)} \Omega_x, \exp\left(\frac{q}{p}x \curvearrowright_y\right) \right\rangle_x \exp\left(\frac{q}{2p}y \curvearrowright_y\right) \Omega_y. \end{aligned}$$

But as $\hat{\omega}_{x+x'} = \hat{\omega}_x + \hat{\omega}_{x'}$, the operator $\partial_{\exp(\hat{\omega}_x)}$ commutes with exponentials and then

$$\partial_{\exp(\hat{\omega}_x)} \Omega_x = \exp(\partial_{\exp(\hat{\omega}_x)} \omega_x) = \exp\left(f(x) + f'(y)f'(x) + \frac{1}{2}f'(z_1)f''(x)f'(z_2)\right).$$

The final connected contribution is $\frac{q}{2p}\curvearrowright + f(x) + f(\frac{q}{p}x) + f'(y)f'(\frac{q}{p}x) + \frac{1}{2}f'(z_1) \times f''(\frac{q}{p}x)f'(z_2)$.

Putting all terms together, we find the following expression for $\partial_\Omega(\Omega \exp(\frac{q}{2p}\curvearrowright))$:

$$\begin{aligned} \exp\left(\frac{q}{2p}\curvearrowright + f(x) + f\left(\frac{q}{p}x\right) + f'(y)f'\left(\frac{q}{p}x\right) + \frac{1}{2}f'(z_1)f''\left(\frac{q}{p}x\right)f'(z_2) \\ + \frac{1}{2}f'\left(\frac{q}{p}z_1\right)f''(x)f'\left(\frac{q}{p}z_2\right) + \frac{p^2}{q^2}\left[f\left(\frac{q}{p}x\right), f(x)\right] \right). \end{aligned}$$

2.2. Step 2

This step is very simple. In order to compute $\sigma Z(L_{p,q})$, we just have to multiply each term by as many factors p as legs.

$$\begin{aligned} \sigma Z(L_{p,q}) = \exp\left(\frac{pq}{2}\curvearrowright + f(px) + f(qx) + f'(py)f'(qx) \\ + \frac{1}{2}f'(pz_1)f''(qx)f'(pz_2) + \frac{1}{2}f'(qz_1)f''(px)(qz_2) \\ + \frac{1}{p^2q^2}[f(qx), f(px)] \right). \end{aligned}$$

2.3. Step 3

In this section, we want to correct the framing defect. The only way we know is to unwheel the preceding expression. Let us note $c(x) = f(px) + f(qx) - f(pqx)$ and factorize it. We then compute

$$Z^{\text{fr}}(K_{p,q}) = \partial_{\Omega^{-1}} \exp\left(\frac{pq}{2}\curvearrowright + f(pqx) + c(x) + R\right) \exp\left(-\frac{pq}{2}\curvearrowright\right).$$

Here,

$$\begin{aligned} R = f'(py)f'(qx) + f'(pz_1)f''(qx)f'(pz_2) \\ + \frac{1}{2}f'(qz_1)f''(px)f'(qz_2) + \frac{1}{p^2q^2}[f(qx), f(px)] \end{aligned}$$

is a series of loop degree 2 and 3.

As before, we first compute the contribution coming from the gluing of c in two points of the same wheel. Lemma 2 give us the following expression, noted U :

$$U = -\frac{1}{p^2q^2}[f(pqx), c(x)] + \frac{1}{2}f'(pqz_1)c''(x)f'(pqz_2).$$

If we suppose that c can be glued to only one leg, we see that the following diagram is commutative:

$$\begin{array}{ccc} \mathcal{B} & \xrightarrow{\times \exp(-c)} & \mathcal{B} \\ \downarrow \partial_\Omega & & \downarrow \partial_{\bar{\Omega}} \\ \mathcal{B} & \xrightarrow{\times \exp(-c)} & \mathcal{B} \end{array}$$

Where $\tilde{\Omega} = \partial_{\exp(\hat{c})}\Omega = \exp(f(x) + c'(x)f'(y) + \frac{1}{2}c'(z_1)f''(x)c'(z_2))$.

But

$$\partial_{\tilde{\Omega}} \exp\left(\frac{pq}{2} \curvearrowright\right) = \left\langle \partial_{\exp(\hat{c}_x)}\Omega_x, \exp\left(\frac{pq}{2} \curvearrowright_{x+y}\right) \right\rangle_x = \tilde{\Omega}_{pq} \exp\left(\frac{pq}{2} \curvearrowright\right).$$

If we factorize this element in the expression of $Z^{\mathfrak{R}}(K_{p,q})$ we get:

$$\begin{aligned} &\partial_{\tilde{\Omega}^{-1}} \left[\partial_{\tilde{\Omega}} \exp\left(\frac{pq}{2} \curvearrowright\right) \exp\left(R - c'(x)f'(pqy) - \frac{1}{2}c'(z_1)f''(pqx)c'(z_2)\right) \right] \\ &\quad \times \exp\left(-\frac{pq}{2} \curvearrowright + c + U\right). \end{aligned}$$

Now, we just have to take care of the action by derivation of $\tilde{\Omega}^{-1}$ on the 2-loop part of the right member, that is:

$$-f'(pqz_1)[pf''(px)f'(qz_2) + qf''(qx)f'(pz_2) - pqf''(pqz_2)c'(x)].$$

Finally, we collect all terms in increasing loop-degree order and conclude the computation.

Proposition 1. *Up to loop degree 3, the unwheeled Kontsevich integral of torus knot can be expressed by the following power series:*

$$\begin{aligned} &c(x) \quad \text{1-loop part} \\ &f'(py)f'(qx) - c'(x)f'(pqy) \quad \text{2-loop part} \\ &\frac{1}{2}f'(pz_1)f''(qx)f'(pz_2) + \frac{1}{2}f'(qz_1)f''(px)f'(qz_2) \quad (*) \\ &+ \frac{1}{2}f'(pqz_1)c''(x)f'(pqz_2) - \frac{1}{2}c'(z_1)f''(pqx)c'(z_2) \\ &- f'(pqz_1)[pf''(px)f'(qz_2) + qf''(qx)f'(pz_2) - pqf''(pqz_2)c'(x)] \\ &\frac{1}{p^2q^2}[f(qx), f(px)] - \frac{1}{p^2q^2}[f(pqx), c(x)]. \end{aligned}$$

3. Rationality

In this part, we will express the diagrams of formula (*) in a rational form and show the following proposition:

Proposition 2. *Let X_n be the connected diagram consisting of n wheels connected by $n - 1$ edges. Then for $n \leq 3$, the unwheeled rational invariant of $K_{p,q}$ of loop degree n appear as a coloring of X_n .*

3.1. Loop degree 1 and 2

It is easy to check that the 1-loop part is as expected:

$$\begin{aligned}
 c(x) - f(x) &= f(px) + f(qx) - f(pqx) - f(x) \\
 &= \frac{1}{2} \log \frac{\sinh(px/2)}{px/2} + \frac{1}{2} \log \frac{\sinh(qx/2)}{qx/2} \\
 &\quad - \frac{1}{2} \log \frac{\sinh(pqx/2)}{pqx/2} - \frac{1}{2} \log \frac{\sinh(x/2)}{x/2} \\
 &= \frac{1}{2} \log \frac{\sinh(px/2) \sinh(qx/2)}{\sinh(pqx/2) \sinh(x/2)} = -\frac{1}{2} \log D_{p,q}(e^x).
 \end{aligned}$$

Here, $D_{p,q}(t) = t^{-\frac{1}{2}(p-1)(q-1)} \frac{(t^{pq}-1)(t-1)}{(t^p-1)(t^q-1)}$ is the Alexander polynomial of $K_{p,q}$.

In the following, we will need a small extension of the spaces of diagrams, because it happens that we need to use rational expressions with poles at the unity. Let us note Λ' the field of fractions of $\Lambda = \mathbb{Q}[t, t^{-1}]$. This is an algebra over Λ and we have an injective Λ -morphism from Λ_{loc} to Λ' .

Then we have a map from $\mathcal{D}(\Lambda, \Lambda_{\text{loc}}) \rightarrow \mathcal{D}(\Lambda, \Lambda')$. There is a corresponding Hair map which fits in the following diagram:

$$\begin{array}{ccc}
 \mathcal{D}(\Lambda, \Lambda_{\text{loc}}) & \longrightarrow & \mathcal{D}(\Lambda, \Lambda') \\
 \downarrow \text{Hair} & & \downarrow \text{Hair} \\
 \mathcal{D}(\mathbb{Q}[[h]], \mathbb{Q}[[h]]) & \longrightarrow & \mathcal{D}(\mathbb{Q}[[h]], \mathbb{Q}[[h]][[h^{-1}]])
 \end{array}$$

Although none of the previous maps is injective, we will identify all diagrams with their image in the latter space (these maps happen to be injective in low degrees).

We can now compute the 2-loop term which we call z_2 , using the formula $f'(x) = \frac{1}{4} \coth(x/2) - \frac{1}{2x}$, all terms containing $\frac{1}{x}$ simplify after symmetrization. If we note $t = e^x$ and $r = e^y$, then

$$\begin{aligned}
 z_2 &= \frac{1}{32} \left[2pq \frac{t^{pq} + 1}{t^{pq} - 1} \frac{s^{pq} + 1}{s^{pq} - 1} - p \frac{t^{pq} + 1}{t^{pq} - 1} \frac{s^p + 1}{s^p - 1} - q \frac{t^{pq} + 1}{t^{pq} - 1} \frac{s^q + 1}{s^q - 1} \right. \\
 &\quad \left. - p \frac{s^{pq} + 1}{s^{pq} - 1} \frac{t^p + 1}{t^p - 1} - q \frac{s^{pq} + 1}{s^{pq} - 1} \frac{t^q + 1}{t^q - 1} + \frac{t^p + 1}{t^p - 1} \frac{s^q + 1}{s^q - 1} + \frac{t^q + 1}{t^q - 1} \frac{s^p + 1}{s^p - 1} \right].
 \end{aligned}$$

This formula appears as a coloring of the dumb-bell graph with denominators dividing $t^{pq} - 1$. If we want to write it with denominators dividing $D_{p,q}$, we are forced to write z_2 as a coloring of \ominus . We can show that there is such a factorization but we were not able to find a close formula for the numerators.

3.2. Loop degree 3

Let us study the 3-loop term, noted z_3 . We decompose it in two parts, z_3^1 and z_3^2 . The z_3^2 part is just the part of z_3 expressed with brackets (the last line of the formula (*)).

In the expression of z_3^1 , if we write $f''(x) = -\frac{1}{8} \frac{1}{\sinh(x/2)^2} + \frac{1}{2x^2}$ and develop, we get a sum of two terms. The first does not contain any fractional term and is obtained from z_3^1 by forgetting all of them (and hence is rational). Concerning the second term, a computation with MAPLE shows that it reduces to:

$$z_3' = -\frac{1}{2pq} \left[f'(py) \frac{1}{x^2} f'(qz) - c'(y) \frac{1}{x^2} f'(pqz) \right].$$




Fig. 3. Diagram containing a $\frac{1}{x}$ term.




Fig. 4. Gluing a leg and a wheel.

Proposition 3. *The series z^3 and z_3^2 cancels.*

Proof. Let us compute the term z_3^2 :

We recall that $z_3^2 = \frac{1}{p^2qz}[f(qx), f(px)] - \frac{1}{p^2qz}[f(pqx), c(x)]$. It is obtained from z_2 by summing all gluings of a left leg on a right leg. There is a normalization factor $\frac{1}{pq}$ and $\frac{1}{2}$ which counterbalance the order given to the two gluings.

We will need the following two lemma which interpret some diagrams with inverse legs.

Lemma 3. *Let D be a diagram consisting of a circle glued on a segment by an edge. Imagine that the circle is colored by a series $g(x)$, and the segment by $\frac{1}{x}$. Using IHX relations we can make the moves suggested by Fig. 3. Making the series sliding, we can cancel the $\frac{1}{x}$ term except for one term which is just the opposite of the initial term.*

This shows that the initial diagram can be expressed by a coloring of another diagram without inverse legs.

Lemma 4. *Let g be a series coloring a circle attached to an edge.*

Consider the sum of the diagrams obtained by gluing the end of a free edge to the legs defined by g . Then this series is obtained by the diagram of Fig. 4.

Proof. We just have to check it for $g(x) = x^n$. In the left-hand side figure, note y a leg lying on the right part of the circle and z a leg on the vertical segment ($z = x - y$).

The left-hand side figure is just obtained by the coloring $\sum_{i+j=n-1} x^i y^j$. Concerning the right part, as in Lemma 3, we make x^n slide to get $(z + y)^n$, eliminate the term y^n , and divide by z . We compute $[(z + y)^n - y^n]/z = (x^n - y^n)/(x - y) = \sum_{i+j=n-1} x^i y^j$. This proves the lemma. \square

If we apply Lemma 4 to both sides of the 2-loop part z_2 , we show that z_3^2 and z_3' cancel. \square

We saw in this computation that both 2-loop terms and 3-loop terms appear as colorings of very special diagrams. The following question is quite natural:

Question 1. *Which kind of diagram are needed in higher loop degrees to express the rational Kontsevich invariant of torus knot? It is unlikely that the degree n part is a coloring of X_n , but we can certainly reduce the number of diagrams needed.*

3.3. The case of closed diagrams

We recall that in all these computations we neglected closed diagrams, although they exist in the expression of $Z^{\text{rat}}(K_{p,q})$. We propose to show that we do not need to add any closed diagram to the expression of $Z^{\text{rat}}(K_{p,q}) = \exp(c + z_2 + z_3)$.

First, there are no closed diagram in the 1-loop or 2-loop terms. The only one comes from z_3 , and precisely from the diagram colored by $z_1 \frac{1}{x^2} z_2$. A direct computation show that this term is $\frac{(p^2-1)(q^2-1)}{1152} \Theta_2$. In the sequel, Θ_n is the graph Θ with the middle edge replaced by n parallel copies.

Following [3], the invariant Z^{rat} is normalized such that for loop degree greater than 1,

$$\text{Hair}_{\langle \Omega, \Omega \rangle} \frac{1}{Z^{\text{rat}}(K)} = Z^{\text{rat}}(K).$$

As $\sigma Z(K_{p,q})$ does not contain any closed diagram, we know that $\langle Z^{\text{rat}}(K), \Omega \rangle = 1$. We must then show that $\langle \text{Hair}_{Z^{\text{rat}}(K_{p,q})}, \Omega \rangle = \langle \Omega, \Omega \rangle$.

But in the left-hand side, the only possible gluing comes from the 1-loop part $\frac{1}{48}(p^2 + q^2 - p^2q^2) \circ$ of c and $\frac{1}{48} \circ$ of Ω . Summed with the closed diagram coming from z_3 , we get $\frac{1}{1152} \Theta_2 = \langle \Omega, \Omega \rangle$.

4. Branched coverings

4.1. Main formula

A great interest for rational expression of Kontsevich integral comes from its relation with branched coverings. More precisely, if $K_{p,q}$ is the torus knot of parameters p and q , and r is an integer, let us note $\Sigma^r(K_{p,q})$ be the pair formed of the cyclic branched covering of S^3 of order r over $K_{p,q}$ and the ramification link.

If r is coprime with p and q , the ramification locus is a knot, and the underlying 3-manifold is a rational homology sphere, the Brieskorn manifold $\Sigma(p, q, r)$.

In [3], a map Lift_r is described which intertwines rational invariant of the cyclic branched coverings and rational invariant of the initial knot in the following way:

$$Z^{\text{rat}}(\Sigma^r(K)) = \exp\left(\frac{\sigma_r(K)}{16} \Theta\right) \text{Lift}_r Z^{\text{rat}}(K).$$

We now study this map in the case of torus knots and loop degree lower than 3 and prove the following proposition:

Proposition 4. Note $Z^{\text{rat}}(K_{p,q}) = \exp(\sum_{k>0} z_k)$ decomposing by loop degree as before, then for all $k \leq 3$,

$$\text{Lift}_r z_k = r^{k-1} z_k.$$

Proof. For $k = 1$, the Alexander polynomial is invariant by these branched coverings. The same will be true for the 1-loop part z_1 .

Let us look to the other cases. The Lift_r map was defined only for diagrams decorated by fractions without poles at r -roots of unity. As we extended the decorations to all fractions, the definition of Lift_r makes sense for any diagram. In the definition of the Lift_r map, we need to express all denominators as polynomials of t^r . Then, we look to the numerators as a coloring by monomials, which is the same as a linear combination of 1-cohomology classes of the underlying graph. We keep only the classes divisible by r and divide them, then we put back denominators replacing t^r by t . Finally we multiply the result by r . This construction is very easy in our case because every edge colored is part of a circle without other colorings and the colorings are the following:

- let n be coprime with r ($n = p, q$ or pq)
- $f_1 = \frac{t^n+1}{t^n-1}$ comes from the first derivative $f'(x) = \frac{1}{4} \coth(x/2) - \frac{1}{2x}$;
- $f_2 = \frac{t^n}{(t^n-1)^2}$ comes from the second derivative $f''(x) = -\frac{1}{8} \frac{1}{\sinh(x/2)^2} + \frac{1}{2x^2}$.

In order to express f_1 with the right denominator, we multiply numerator and denominator by $1 + t^n + \dots + (t^n)^{r-1}$ to get $\frac{1+2t^n+\dots+2t^{n(r-1)}+t^{nr}}{t^{nr}-1}$. The only r -divisible numerators are 1 and t^{nr} because n and r are coprime. The result of the Lift_r map is then the identity (before multiplying by r).

We do the same operation with f_2 and multiply numerator and denominator by $(1 + t^n + \dots + (t^n)^{r-1})^2$ to get $\frac{t^n(1+t^n+\dots+(t^n)^{r-1})^2}{(t^{nr}-1)^2}$. The numerator is

$$t^n(1 + t^n + \dots + (t^n)^{r-1})^2 = \sum_{i,j=0}^{r-1} t^{n(i+j+1)}.$$

The indices for which the order of the monomial is r -divisible are such that $i + j = n - 1$. There are r such terms, and the result of the Lift_r map is then the multiplication by r (before multiplying by r at the end).

For the z_2 term, we have two terms f_1 , and then $\text{Lift}_r z_2 = rz_2$. For the z_3 term, we have two terms f_1 and one term f_2 and then $\text{Lift}_r z_3 = r^2 z_3$. This ends the proof of the proposition. \square

Question 2. Can we extend this proposition for larger values of k ? Do we have any conceptual interpretation of this formula?

4.2. Application to LMO invariant of Brieskorn spheres

The LMO invariant of $\Sigma(p, q, r)$ is just the closed part of $\sigma Z(\Sigma^r(K_{p,q}))$, i.e., $\langle Z^{\text{rat}}(\Sigma^r(K_{p,q})), \Omega \rangle$. Using the map Lift we have the following formula:

$$\text{LMO}(\Sigma(p, q, r)) = \exp\left(\frac{\sigma_r(K)}{16} \ominus\right) \frac{\langle \text{Lift}_r Z^{\text{rat}}(K_{p,q}), \Omega \rangle}{\langle \Omega, \Omega \rangle}.$$

From this formula, we can prove that the degree two term is $\frac{(p^2-1)(q^2-1)(r^2-1)}{1152} \Theta_2$.

Supposing the proposition is true for $n = 4$ we deduced the following formula for the degree 3 term: $\text{LMO}(\Sigma(p, q, r))_3 = \frac{-pqr(p^2-1)(q^2-1)(r^2-1)}{13824} \Theta_3$.

These computations agree with formulas for the LMO invariant of Seiferts spaces which can be found in [1,7].

References

- [1] D. Bar-Natan, R. Lawrence, A rational surgery formula for the LMO invariant, math.GT/0007045, Israel J. Math., submitted for publication.
- [2] S. Garoufalidis, Whitehead doubling persists, math.GT/0003189.
- [3] S. Garoufalidis, A. Kriker, Finite type invariants of cyclic branched covers, math.GT/0107220.
- [4] S. Garoufalidis, A. Kriker, A rational noncommutative invariant of boundary links, math.GT/0105028.
- [5] A. Kriker, The lines of the Kontsevich integral and Rozansky's rationality conjecture, math.GT/0005284.
- [6] C. Lescop, Introduction to the Kontsevich integral of framed tangles, Technical Report, Grenoble Summer School, June 1999.
- [7] M. Mariño, Chern-simons theory, matrix integrals, and perturbative three-manifold invariants, hep-th/0207096.
- [8] T. Ohtsuki, A cabling formula for the 2-loop polynomial of knots, Preprint.
- [9] B. Patereau-Mirand, Non-injectivity of the hair map, math.GT/0202065.
- [10] L. Rozansky, A rationality conjecture about Kontsevich integral of knots and its implications to the structure of the colored jones polynomial, *Topology Appl.* 127 (2003) 47–76.
- [11] D. Thurston, Wheeling: A diagrammatic analogue of the Duflo isomorphism, Ph.D. Thesis, UC Berkeley, math.QA/0006083, 2000.
- [12] P. Vogel, Vassiliev theory, Technical Report, MaPhySto, February 2000.