


Available online at www.sciencedirect.com

Discrete Applied Mathematics 154 (2006) 980–994

DISCRETE
APPLIED
MATHEMATICSwww.elsevier.com/locate/dam

Update schedules of sequential dynamical systems

Reinhard Laubenbacher^a, Bodo Pareigis^b^aVirginia Bioinformatics Institute, 1880 Pratt Drive, Blacksburg, VA 24061, USA^bMath. Inst. d. Universität München, Theresienstr. 39, 80333 München, Germany

Received 21 April 2003; received in revised form 6 October 2005; accepted 11 October 2005

Available online 10 January 2006

Abstract

Sequential dynamical systems have the property, that the updates of states of individual cells occur sequentially, so that the global update of the system depends on the order of the individual updates. This order is given by an order on the set of vertices of the dependency graph. It turns out that only a partial suborder is necessary to describe the global update. This paper defines and studies this partial order and its influence on the global update function.

© 2005 Elsevier B.V. All rights reserved.

Keywords: Sequential dynamical system; Morphism of pographs; Poset models of graphs; Update schedules

1. Introduction

The theory of sequential dynamical systems (SDS) was first introduced in [1–3], with the goal of providing a mathematical foundation for computer simulations. Such a foundation will allow a rigorous mathematical analysis of a variety of questions that arise in simulation practice. Many computer simulations can be represented in terms of SDS for computational purposes. By design SDS carry more internal structure than, say, cellular automata. As a result it is possible to prove general results about SDS relating their structural properties to the dynamics they generate. This represents an important first step toward an understanding of how local properties of a system affect global dynamics.

In [6] we generalized the notion of a SDS, and defined transformations of SDS. Such transformations are compatible with the internal structure and induce a transformation of the associated state spaces, that is, are compatible with the dynamics generated by the systems. One important role such transformations can play is as mathematical formalizations of a simulation of one SDS by another. Such important practical questions as how to reduce the dimension of a simulation can be phrased in this way. Transformations also allow the study of the relationship between structural changes in a simulation to the resulting changes in the dynamics.

A second role for transformations is in a structure theory of SDS as the first step toward a classification. For instance, in [6] it was shown that every SDS can be uniquely decomposed into a product of indecomposable SDS, which can then be studied individually.

Finally, a third role is in comparing SDS with other interesting objects in computer science used for simulations and as computational devices. This can be done very well in a categorical framework, and one goal of this research is to develop a good categorical setting for the study of SDS.

E-mail addresses: reinhard@vbi.vt.edu (R. Laubenbacher), pareigis@lmu.de (B. Pareigis).

Recall from [6] that a SDS is a function

$$f : \prod_{i=1}^n k_i \longrightarrow \prod_{i=1}^n k_i,$$

on strings of length n , with entries in the i th component coming from a specified set k_i . We simply write $k^n := \prod_{i=1}^n k_i$. This function $f : k^n \rightarrow k^n$ is obtained from the following data:

- (1) a *dependency graph* F on n vertices,
- (2) a collection of *local update functions* $f_i : k^n \rightarrow k^n, i = 1, \dots, n$, which change only the i th coordinate, and whose inputs are controlled by the graph F ,
- (3) an *update schedule* α , consisting of a word in n letters a_1, \dots, a_n from the set of vertices V_F of F .

The function f is the composition of the f_i , in the order specified by the update schedule α . (More details can be found in the next section.)

It was shown in [4] that the graph F is implicit in the rest of the data, using a Galois correspondence between collections of local functions and graphs, constructed in [5]. Thus, if expedient, this part of the structure of an SDS may be ignored.

The focus in the present paper is on the update schedule α . Recall that the global update function of the SDS is generated by composing the local update functions in the order prescribed by α . Part of the work reported in [1,2] concerned the extent to which changes in the update schedule affect the global update function, respectively, the resulting dynamic structure. (In contrast to [6] and the present paper the update schedule in [1,2] is taken to be a permutation of the indices of the nodes, rather than a general finite word in a subset of those indices.) It was shown that some changes in the update schedule leave the global update function unchanged. This is similar to a distributed computation in which certain steps can be carried out in various orders, whereas others need to be done according to a prescribed schedule so that the end result remains unchanged. We want to find properties of an update schedule on which the global update function depends and other properties that can freely be changed without changing the global update function and thus the dynamic behavior of the whole system.

We show in this paper that this degree of freedom in the update schedule is a very important aspect of an SDS that deserves to be studied as part of the explicit structure of the SDS. This leads us to propose a new definition of SDS which incorporates this dichotomy of the update schedule. We will see in a subsequent paper that such a change leads to a notion of transformations of SDS with very desirable properties. In particular, we obtain a categorical framework for SDS that is rich in transformations and structure.

In order to describe and study the ordering of the vertices given by the update schedule we introduce the notion of a *poset model of a graph*, an interesting connection between posets and graphs. Let F be a finite graph (e.g., the dependency graph of an SDS), and let

$$\alpha : \{1, \dots, n\} \longrightarrow V_F$$

be an update schedule, i.e. a function into the set of vertices of F . Any update schedule a_1, \dots, a_n of an SDS can be represented as such a function. The key result of this paper is that there exists a unique poset \mathcal{O}_F and poset model $\beta : \mathcal{O}_F \rightarrow V_F$ of the graph F together with a function $\gamma : \mathcal{O}_F \rightarrow \{1, \dots, n\}$, i.e.

$$\{1, \dots, n\} \xleftarrow{\gamma} \mathcal{O}_F \xrightarrow{\beta} V_F,$$

where γ is invertible and order preserving, such that $\alpha = \gamma^{-1} \circ \beta$. That is, we can decompose the update schedule α into a poset model β of the graph F (called *pograph*) and an *update schedule* γ for the pograph $\beta : \mathcal{O}_F \rightarrow V_F$.

Surprisingly, it turns out that the pograph $\beta : \mathcal{O}_F \rightarrow V_F$ completely determines the dynamical behavior of the SDS. Consequently, we can define SDS on a pograph $\mathcal{O}_F \rightarrow V_F$ alone instead of on a graph F together with an update schedule α .

It will turn out that the definition of morphisms of SDS defined on pographs becomes very natural and straightforward.

If the map β is bijective, and we view the poset \mathcal{O}_F as a directed graph via its Hasse diagram, then β becomes a graph map which induces an acyclic orientation on F . In [7, Proposition 1] it is shown that there is a one-to-one correspondence

between acyclic orientations on a graph F on n vertices and the set of equivalence classes of permutations on n letters for a certain equivalence relation. The equivalence relation is generated by making two permutations equivalent if they differ by transpositions of adjacent elements whose corresponding vertices are not connected by an edge in F . The SDS defined by Barrett et al. called permutation SDS in [6], use an update schedule given by a permutation, that is, every local update function is used exactly once in the composition that defines the global update function of the SDS. It was shown in [2] that from an acyclic orientation of F one can construct different update schedules α , which all produce the same global update function. That is, an acyclic orientation of F contains enough information to construct the global update function. This result was used in [2] to derive a sharp upper bound on the number of different global update functions that can be generated by varying the update schedule, for a fixed graph F and fixed local update functions.

In the more general setting considered here, the pograph $\beta : \mathcal{O}_F \rightarrow V_F$ is taking the place of the acyclic orientation of F . We will show that for a fixed β it does not matter what choice we make for the corresponding γ in order to study the dynamic behavior. The global update function is independent of the choice of γ . As a corollary we obtain a one-to-one correspondence between pographs with m elements on a graph F and equivalence classes of update schedules $\{1, \dots, m\} \rightarrow V_F$. The equivalence relation is generated by setting two update schedules α and α' equivalent if they generate the same global update function $f_\alpha = f_{\alpha'}$ from any family of local update functions $f_{\alpha(i)}$.

2. Sequential dynamical systems and graphs with update schedule

For the convenience of the reader we repeat the definition of SDS as used in [6]. We will rephrase that definition under some new points of view. Some of the basic notions are explained in the following:

Remark 2.1. Let X be a set and let $\mathcal{P}(X)$ be its power set. Let $\mathcal{P}_2(X) \subseteq \mathcal{P}(X)$ be the subset of all two-element subsets of X .

A (loop free, undirected) graph $F = (V_F, E_F)$ consists of a set V_F of vertices and a subset $E_F \subseteq \mathcal{P}_2(V_F)$ of edges. Let F be a graph. A 1-neighborhood $N(a)$ of a vertex $a \in V_F$ is the set

$$N(a) := \{b \in V_F \mid \{a, b\} \in E_F \text{ or } a = b\}.$$

Let \mathcal{L} be a subcategory of the category of sets. Let $(k[a] \mid a \in V_F)$ be a family of sets in \mathcal{L} , e.g. finite sets. The set $k[a]$ will be called the set of local states at a . Define

$$k^{V_F} := \prod_{a \in V_F} k[a],$$

the set of (global) states of F . In case V_F is finite with r elements we write

$$k^r := k[a_1] \times \dots \times k[a_r].$$

We use the following notation. For a state $x \in k^r$ and a vertex $a \in V_F$ we write $x[a]$ for the state of the vertex a or the a th component of x so that

$$x = (x[a] \mid a \in V_F) \quad \text{or} \quad x = (x[a_1], \dots, x[a_r]).$$

In case that all $k[a]$ are equal to a set k , this definition reduces to the usual definition of k^{V_F} resp. k^r .

A function $f : k^{V_F} \rightarrow k^{V_F}$ is called local at $a_i \in V_F$ if

$$f(x)[a_j] = \begin{cases} x[a_j] & \text{if } a_j \neq a_i, \\ f^i(x) & \text{if } a_j = a_i, \end{cases}$$

where $f^i(x) = f^i((x[a] \mid a \in V_F)) \in k[a_i]$ depends only on the states $x[a]$ of those variables a that are in the 1-neighborhood $N(a_i)$ of the vertex a_i .

If V_F is finite this means the following. A function $f : k^r \rightarrow k^r$ is local at $a_i \in V_F$ if

$$f(x[a_1], \dots, x[a_r]) = (x[a_1], \dots, x[a_{i-1}], f^i(x[a_1], \dots, x[a_r]), x[a_{i+1}], \dots, x[a_r]),$$

where $f^i(x[a_1], \dots, x[a_r]) \in k[a_i]$ depends only on the states $x[a_j]$ of those vertices a_j that are in the 1-neighborhood $N(a_i)$ of the vertex a_i .

One of the fundamental observations is the following easy fact. If $a, b \in V_F$ such that $\{a, b\} \notin E_F$ then $f_a \circ f_b = f_b \circ f_a$ if f_a and f_b are local functions.

Definition 2.2. Let F be a graph. A map $\alpha : \{1, \dots, n\} \rightarrow V_F$ is called an *update schedule of length n for F* . A pair (F, α) is called a *graph with update schedule* or a *ugraph F* .

Definition 2.3. A *sequential dynamical system (SDS)¹* on a ugraph F

$$\mathcal{F} = (F, (k[a] \mid a \in V_F), (f_a \mid a \in V_F))$$

consists of

- (1) a finite ugraph F ,
- (2) a family of sets $(k[a] \mid a \in V_F)$ in \mathcal{L} ,
- (3) a family of local functions $(f_a : k^r \rightarrow k^r \mid a \in V_F, f_a \text{ local at } a)$ in \mathcal{L} .

Remark 2.4. The update schedule $\alpha : \{1, \dots, n\} \rightarrow V_F$ of a ugraph F of an SDS \mathcal{F} defines an associated *global update function* of the SDS \mathcal{F}

$$f_\alpha := f_{\alpha(1)} \circ \dots \circ f_{\alpha(n)} : k^r \rightarrow k^r.$$

For the moment we consider an update schedule just as an additional structure of the graph used in the definition of an SDS. We will call the graph F with this (or possibly another) additional structure on which an SDS is defined *the basis* of the given SDS. Surprisingly a map $\alpha : \{1, \dots, n\} \rightarrow V_F$ induces an interesting additional structure on the graph F that we will subsequently study.

3. Poset models of graphs

In this section we show the relationship between a graph with update schedule and an associated partially ordered set together with a certain map into the graph which we call a *poset model of F* .

Let \mathcal{O} be a poset (partially ordered set) with order relation \leq . We define

$$i \triangleleft j : \iff (i < j \text{ and } \forall k \in \mathcal{O} : i \leq k \leq j \implies i = k \text{ or } k = j),$$

i.e., j covers i .

Theorem 3.1. *Let F be a ugraph with update schedule $\alpha : \{1, \dots, n\} \rightarrow V_F$. Then there are maps*

$$\{1, \dots, n\} \xleftarrow{\gamma} \mathcal{O}_F \xrightarrow{\beta} V_F$$

with

- a finite poset \mathcal{O}_F ,
- a bijective order preserving map $\gamma : \mathcal{O}_F \rightarrow \{1, \dots, n\}$ and
- a map $\beta : \mathcal{O}_F \rightarrow V_F$

such that

- (1) $\forall i, j \in \mathcal{O}_F : i \triangleleft j \implies \{\beta(i), \beta(j)\} \in E_F$,
- (2) $\forall i, j \in \mathcal{O}_F : \{\beta(i), \beta(j)\} \in E_F \implies i \leq j \vee j \leq i$,
- (3) $\alpha = \beta\gamma^{-1}$.

¹ Subsequently, we will use the acronym SDS for plural as well as singular instances.

Proof. The poset \mathcal{O}_F is constructed as follows. Define $\mathcal{O}_F := \{1, \dots, n\}$ with the following partial order. Define

$$i \dot{\prec} j : \iff (i \leq j \wedge \{\alpha(i), \alpha(j)\} \in E_F).$$

Let $i \preceq j$ be the reflexive and transitive closure of the relation $\dot{\prec}$.

Then (\mathcal{O}_F, \preceq) is a poset since $i \preceq j$ and $j \preceq i$ implies that there are chains $i = i_1 \dot{\prec} i_2 \dot{\prec} \dots \dot{\prec} i_r = j$ and $j = j_1 \dot{\prec} j_2 \dot{\prec} \dots \dot{\prec} j_s = i$. From this we get $i = i_1 \leq i_2 \leq \dots \leq i_r = j$ and $j = j_1 \leq j_2 \leq \dots \leq j_s = i$ hence $i = j$.

Furthermore $\beta := \alpha : \mathcal{O}_F = \{1, \dots, n\} \rightarrow V_F$ satisfies (1), for let $i \triangleleft j$ then j covers i in (\mathcal{O}_F, \preceq) , so we must have $i \dot{\prec} j$ since these pairs generate the order on \mathcal{O}_F . But that implies $\{\beta(i), \beta(j)\} \in E_F$. Assume now that $\{\beta(i), \beta(j)\} \in E_F$ holds. Since $i \leq j$ or $j \leq i$ in $\{1, \dots, n\}$ we get $i \dot{\prec} j$ or $j \dot{\prec} i$ hence $i \preceq j$ or $j \preceq i$. So (2) is satisfied.

By construction $\gamma := \text{id} : \mathcal{O}_F \rightarrow \{1, \dots, n\}$ is bijective. It is order preserving since $i \preceq j$ implies that there is a chain $i = i_1 \dot{\prec} i_2 \dot{\prec} \dots \dot{\prec} i_r = j$. From this we get $i = i_1 \leq i_2 \leq \dots \leq i_r = j$.

Obviously $\alpha = \beta\gamma^{-1}$. \square

Observe that $i \triangleleft j$ in \mathcal{O}_F implies $i \dot{\prec} j$, but the converse does not hold in general.

Definition 3.2. Let F be a graph and \mathcal{O}_F be a poset. Let $\beta : \mathcal{O}_F \rightarrow V_F$ be a map satisfying $\forall i, j \in \mathcal{O}_F$:

- (1) $i \triangleleft j \implies \{\beta(i), \beta(j)\} \in E_F$,
- (2) $\{\beta(i), \beta(j)\} \in E_F \implies i \leq j \vee j \leq i$.

Then $\beta : \mathcal{O}_F \rightarrow V_F$ is called a *poset model of the graph F* or a *pograph*.

Remark 3.3. Theorem 3.1 says that we can construct a pograph $(F, \mathcal{O}_F, \beta)$ together with a bijective order preserving map $\gamma : \mathcal{O}_F \rightarrow \{1, \dots, n\}$ from a ugraph (F, α) . Conversely if we have a pograph $(F, \mathcal{O}_F, \beta)$ together with a bijective order preserving map $\gamma : \mathcal{O}_F \rightarrow \{1, \dots, n\}$ then we get a ugraph with update schedule $\alpha := \beta\gamma^{-1}$.

If β in the definition of a pograph is bijective then this is the same as an acyclic orientation of the graph F as discussed in [7]. In this case the map β and the order of the poset \mathcal{O}_F defines an orientation on each edge of F by condition (2). Since \mathcal{O}_F is a poset the orientation on all of F (as it is defined) will be acyclic, i.e. there are no cycles. Condition (1) means that the order of \mathcal{O}_F is “generated” by the graph. If α is injective then some of the edges will be oriented, the orientation of the edges will be transitive, and F with this orientation will be acyclic. Else it may happen that edges are oriented in both directions.

Example 3.4. An example is $\alpha : \{1, 2, 3, 4\} \rightarrow \{a, b, c\} = V_F$ where all vertices are connected by an edge and $\alpha(1) = \alpha(4) = a, \alpha(2) = b, \alpha(3) = c$. Then $1 \dot{\prec} 2 \dot{\prec} 3 \dot{\prec} 4, 1 \dot{\prec} 3$, and $2 \dot{\prec} 4$, and the edges $\{a, b\}$ and $\{a, c\}$ are directed in both directions. Furthermore $\mathcal{O}_F = \{1, 2, 3, 4\}$ as posets; in particular $1 \preceq 4$ but $1 \dot{\prec} 4$ does not hold.

Definition 3.5. Let $(F, \mathcal{O}_F, \beta)$ be a pograph. A bijective, order preserving map $\gamma : \mathcal{O}_F \rightarrow \{1, \dots, n\}$ is called an *update schedule* for $(F, \mathcal{O}_F, \beta)$. The pair $((F, \mathcal{O}_F, \beta), \gamma)$ is called a *pograph with update schedule*.

Now Theorem 3.1 says that we can construct a pograph with update schedule out of every ugraph. Conversely we can construct a ugraph out of every pograph with update schedule in the obvious way. These two constructions are almost inverses of each other.

Proposition 3.6. Let $F = (F, \mathcal{O}_F, \beta, \gamma)$ and $(F, \mathcal{O}'_F, \beta', \gamma')$ be finite pographs with update schedules $\gamma : \mathcal{O}_F \rightarrow \{1, \dots, n\}$ and $\gamma' : \mathcal{O}'_F \rightarrow \{1, \dots, n\}$ resp. Assume that $\beta\gamma^{-1} = \beta'\gamma'^{-1}$. Then there is a unique isomorphism of posets $\delta : \mathcal{O}_F \rightarrow \mathcal{O}'_F$ such that $\gamma = \gamma'\delta$ and $\beta = \beta'\delta$.

Proof. Obviously $\delta := \gamma'^{-1}\gamma$ is the only choice for this map and δ satisfies $\gamma = \gamma'\delta$ and $\beta = \beta'\delta$. We only have to show that δ is order preserving, since the inverse map will also be order preserving by the symmetry of the situation.

Let $i, j \in \mathcal{O}_F$. Since \mathcal{O}_F is finite we only have to show

$$i \triangleleft j \implies \delta(i) \leq \delta(j).$$

Since $i \triangleleft j$ implies $\{\beta(i), \beta(j)\} \in E_F$ we get $\{\beta' \delta(i), \beta' \delta(j)\} \in E_F$ hence $\delta(i) \leq \delta(j) \vee \delta(j) \leq \delta(i)$. Assume that $\delta(j) \leq \delta(i)$ holds. Then we have $\gamma(j) = \gamma' \delta(j) \leq \gamma' \delta(i) = \gamma(i)$, a contradiction to $i \triangleleft j$. Thus $\delta(i) \leq \delta(j)$. \square

So we see that the pograph with update schedule $(F, \mathcal{O}_F, \beta, \gamma)$ constructed from a ugraph (F, α) by Theorem 3.1 is unique up to an isomorphism (of posets), compatible with the update schedule and with the pograph map β .

Definition 3.7. Let $(F, \mathcal{O}_F, \beta, \gamma)$ and $(F, \mathcal{O}'_F, \beta', \gamma')$ be finite pographs with update schedule. A *strong isomorphism* between these pographs with update schedule is an isomorphism of posets $\delta : \mathcal{O}_F \rightarrow \mathcal{O}'_F$ such that $\gamma = \gamma' \delta$ and $\beta = \beta' \delta$.

Observe that for a strong isomorphism we have $\beta \gamma^{-1} = \beta' \delta \gamma^{-1} = \beta' \gamma'^{-1}$. Obviously strong isomorphisms define an equivalence relation and the above observations give:

Corollary 3.8. *The constructions given in Theorem 3.1 and Proposition 3.6 define a bijection between ugraphs and strong isomorphism classes of pographs with update schedules.*

The next proposition relates the notion of pograph with update schedule to the permutation update schedules used by Barrett et al. For a given graph F on n vertices $\{1, \dots, n\}$ define an equivalence relation \sim_F on the permutations in S_n as follows. Let $\pi, \pi' \in S_n$, denoted by $\pi = (i_1, \dots, i_n)$ and $\pi' = (i'_1, \dots, i'_n)$. (That is, $\pi(j) = i_j$, etc.) Then $\pi \sim_F \pi'$ if there is a $k \in \{1, \dots, n\}$ such that $i_j = i'_j$ for all $j \neq k, k + 1$ and there is no edge in F connecting vertices i_k and i_{k+1} . Let \sim_F be the equivalence relation generated by this relation. Denote by S_n / \sim_F the set of equivalence classes. Let \mathcal{A}_F denote the set of acyclic orientations on F .

Proposition 3.9 (Reidys, [7, Proposition 1]). *There is a one-to-one correspondence between \mathcal{A}_F and S_n / \sim_F .*

The next result shows how the new concepts of poset model and pograph with update schedule reduce to the case of permutation update schedules used in permutation SDS. Let \mathcal{I}_F denote the set of strong isomorphism classes of bijective pographs with update schedule on F (that is, β is bijective).

Proposition 3.10. *There is a one-to-one correspondence between \mathcal{I}_F and \mathcal{A}_F , hence a one-to-one correspondence between the set of strong isomorphism classes of bijective poset models and S_n / \sim_F .*

Proof. Let $\beta : \mathcal{O}_F \rightarrow F$ be a poset model of F , and assume that β is bijective. We define an acyclic orientation \mathcal{O} of F as follows. Let $\{u, v\}$ be an edge of F . Then $u = \beta(i)$ and $v = \beta(j)$ for some $i, j \in \mathcal{O}_F$. Since β is a poset model, we have that $i \leq j$ or $j \leq i$. If $i \leq j$, then orient the edge (u, v) from u to v . Since β is onto, every edge of F is oriented in this way. It is straightforward to see that this orientation is acyclic, since \mathcal{O}_F is a partial order.

If $\beta' : \mathcal{O}'_F \rightarrow F$ is another poset model of F which is strongly isomorphic to β via an isomorphism $\varphi : \mathcal{O}_F \rightarrow \mathcal{O}'_F$, then it is clear that β' induces the same orientation on F .

Conversely, let \mathcal{O} be an acyclic orientation of F . Then \mathcal{O} defines a partial order on the vertices of F , by setting $u < v$ if there is an oriented path in \mathcal{O} from u to v . Then $\text{id} : \mathcal{O} \rightarrow F$ is a poset model of F . Together with Proposition 3.9 this completes the proof of the first statement. \square

As outlined in the introduction, from the point of view of the global update function of an SDS, there is a certain amount of freedom on the order prescribed by an update schedule. There are those order relations that need to be kept fixed if the global update function is not to be changed, and then there are order relations that can be reversed without affecting the global update function. The previous proposition shows that these two aspects of an update schedule are neatly separated in a pograph with update schedule, the poset model β which encodes those order relations that are

required, and the update schedule γ which contains all choices that are available in the update schedule that do not affect the global update function.

4. Morphisms of pographs and of ugraphs

In this section we define morphisms of ugraphs and of pographs with update schedules. The two resulting categories are equivalent. This result will form the basis for a shift from considering SDS on ugraphs to SDS on pographs. The definition of morphisms of pographs is very natural, and, as a consequence, the definition of morphisms of SDS on pographs will be very natural as well.

Definition 4.1. Let $\mathcal{F} = (F, \mathcal{O}_F, \beta_F)$ and $\mathcal{G} = (G, \mathcal{O}_G, \beta_G)$ be two pographs. A *morphism of pographs* $\varphi : \mathcal{F} \rightarrow \mathcal{G}$ is a pair of morphisms $(\varphi_g, \tilde{\varphi})$, where

- $\varphi_g : F \rightarrow G$ is a morphism of graphs and
- $\tilde{\varphi} : \mathcal{O}_F \rightarrow \mathcal{O}_G$ is a morphism of posets such that the diagram

$$\begin{array}{ccc}
 \mathcal{O}_F & \xrightarrow{\beta_F} & V_F \\
 \tilde{\varphi} \downarrow & & \downarrow \varphi_g \\
 \mathcal{O}_G & \xrightarrow{\beta_G} & V_G
 \end{array}$$

commutes.

The composition of morphisms of pographs is again a morphism of pographs. So we obtain a category *Pograph* of pographs.

Next we define morphisms of ugraphs. The following discussion will compare our new definition of a morphism of pographs with the definition we used in the discussion of SDS in [6]. Let $(F, \alpha : \{1, \dots, m\} \rightarrow V_F)$ be a ugraph. For each connected component $F_{(l)}$ of F let $|\alpha_{(l)}| \subseteq \{1, \dots, m\}$ denote the preimage of $F_{(l)}$ under the map α . Consider $|\alpha_{(l)}|$ as an ordered set, the order being induced by the natural order of $\{1, \dots, m\}$. We also define $|\alpha| := \dot{\bigcup} |\alpha_{(l)}|$ with the order induced by the components $|\alpha_{(l)}|$. Then the identity map $\text{id} : |\alpha| \rightarrow \{1, \dots, n\}$ is a morphism of posets, but not an isomorphism, since certain pairs may be unordered in $|\alpha|$ whereas all pairs are ordered in $\{1, \dots, n\}$. We say that $|\alpha|$ has a *coarser partial order* than $\{1, \dots, n\}$.

Observe that $|\alpha|$ is a disjoint union of totally ordered sets, one for each connected component of F , which is equal to $\{1, \dots, n\}$ if F is connected.

Definition 4.2. Let $(F, \alpha_F : \{1, \dots, m\} \rightarrow V_F)$ and $(G, \alpha_G : \{1, \dots, n\} \rightarrow V_G)$ be ugraphs. A *morphism of ugraphs* $\varphi : F \rightarrow G$ consists of

- a morphism of graphs $\varphi_g : F \rightarrow G$ and
- a morphism of posets $\tilde{\varphi} : |\alpha_F| \rightarrow |\alpha_G|$,

such that

$$\begin{array}{ccc}
 |\alpha_F| & \xrightarrow{\alpha_F} & V_F \\
 \tilde{\varphi} \downarrow & & \downarrow \varphi_g \\
 |\alpha_G| & \xrightarrow{\alpha_G} & V_G
 \end{array}$$

commutes. Ugraphs together with these morphisms form a category *Ugraph*.

This definition is equivalent to what we used in [6]. There we studied a pair $(\varphi_g, (\tilde{\varphi}_{(l)}))$ satisfying the conditions:

- $\varphi_g : F \rightarrow G$ is a graph morphism,
- the family

$$(*) \quad \tilde{\varphi}_{(l)} : |(\alpha_F)_{(l)}| \rightarrow \{1, \dots, n\}$$

for each connected component $F_{(l)}$ of F is a family of order preserving maps,

- $\forall l \forall j \in |(\alpha_F)_{(l)}| : \varphi_g(\alpha_F(j)) = \alpha_G(\tilde{\varphi}_{(l)}(j))$, i.e. all $\tilde{\varphi}_{(l)}$ are compatible with the given graph morphism φ_g .

Indeed we have:

Proposition 4.3. *Let $(F, \alpha_F : \{1, \dots, m\} \rightarrow V_F)$ and $(G, \alpha_G : \{1, \dots, n\} \rightarrow V_G)$ be ugraphs.*

(1) *Let $(\varphi_g, (\tilde{\varphi}_{(l)}))$ satisfy conditions (*). Then $\tilde{\varphi} : |\alpha_F| \rightarrow |\alpha_G|$ with $\tilde{\varphi}(i) := \tilde{\varphi}_{(l)}(i)$ for all $i \in |(\alpha_F)_{(l)}|$ is a morphism of posets and $\varphi = (\varphi_g, \tilde{\varphi})$ is a morphism of ugraphs.*

(2) *Conversely let $\varphi = (\varphi_g, \tilde{\varphi})$ be a morphism of ugraphs. Then φ_g together with the family of maps*

$$\tilde{\varphi}_{(l)} : |(\alpha_F)_{(l)}| \subseteq |\alpha_F| \xrightarrow{\tilde{\varphi}} |\alpha_G| \xrightarrow{\text{id}} \{1, \dots, n\}$$

satisfy conditions (*).

Proof. (1) Since $|\alpha_G|$ has a coarser order than $\{1, \dots, n\}$ we have to show that $\tilde{\varphi} : |\alpha_F| \rightarrow |\alpha_G|$ is order preserving. Let $i, j \in |\alpha_F|$ be given with $i \leq j$. Then by definition of the partial order on $|\alpha_F|$ there is a unique connected component $F_{(l)}$ with $\alpha_F(i), \alpha_F(j) \in F_{(l)}$ so that $i, j \in |(\alpha_F)_{(l)}|$. Thus $\tilde{\varphi}_{(l)}(i) \leq \tilde{\varphi}_{(l)}(j)$ and hence $\tilde{\varphi}(i) \leq \tilde{\varphi}(j)$ in $\{1, \dots, n\}$. Since $\varphi_g(F_{(l)}) \subseteq G_{(l)}$ for a unique connected component of G we get $\tilde{\varphi}(i), \tilde{\varphi}(j) \in |(\alpha_G)_{(l)}|$ hence $\tilde{\varphi}(i) \leq \tilde{\varphi}(j)$ in $|(\alpha_G)_{(l)}|$ and also in $|\alpha_G|$. By the compatibility of $\tilde{\varphi}_{(l)}$ with φ_g we get the commutativity of the square.

(2) Obviously $\tilde{\varphi}_{(l)}$ is order preserving and satisfies the compatibility condition of $\tilde{\varphi}_{(l)}$ with φ_g . \square

The definition of morphisms of pographs and of ugraphs are very similar. A ugraph, however, contains the complete information on an update schedule, whereas a pograph contains only part of this information. The main point is that \mathcal{O}_F and $|\alpha_F|$ are different posets!

Lemma 4.4. *Let $\mathcal{F} = (F, \alpha_F : \{1, \dots, m\} \rightarrow V_F)$ and $\mathcal{G} = (G, \alpha_G : \{1, \dots, n\} \rightarrow V_G)$ be ugraphs and let $\varphi : \mathcal{F} \rightarrow \mathcal{G}$ with $\varphi = (\varphi_g, \tilde{\varphi})$ be a morphism of ugraphs. Let $(F, \mathcal{O}_F, \beta_F : \mathcal{O}_F \rightarrow V_F, \gamma_F : \mathcal{O}_F \rightarrow \{1, \dots, m\})$ and $(G, \mathcal{O}_G, \beta_G : \mathcal{O}_G \rightarrow V_G, \gamma_G : \mathcal{O}_G \rightarrow \{1, \dots, n\})$ resp. be the pographs with update schedule as constructed in Theorem 3.1. Then φ induces a morphism of pographs such that*

$$\begin{array}{ccccc} \{1, \dots, m\} & \xleftarrow{\text{id}} & |\alpha_F| & \xleftarrow{\gamma_F} & \mathcal{O}_F & \xrightarrow{\beta_F} & V_F \\ & & \tilde{\varphi} \downarrow & & \tilde{\varphi} \downarrow & & \varphi_g \downarrow \\ \{1, \dots, n\} & \xleftarrow{\text{id}} & |\alpha_G| & \xleftarrow{\gamma_G} & \mathcal{O}_G & \xrightarrow{\beta_G} & V_G \end{array}$$

commutes.

Proof. Since the underlying sets of \mathcal{O}_F and of $|\alpha_F|$ are equal to $\{1, \dots, m\}$ and the map γ_F is the identity, the left square commutes. The same set theoretic argument shows that the right square commutes. We have to show that γ_F and $\tilde{\varphi} : \mathcal{O}_F \rightarrow \mathcal{O}_G$ are order preserving.

Let $i, j \in \mathcal{O}_F$ with $i \prec j$ be given. Then $i \leq j$ in $\{1, \dots, m\}$ and $\{\beta_F(i), \beta_F(j)\} \in E_F$. Thus $\beta_F(i)$ and $\beta_F(j)$ are contained in a common connected component $F_{(l)}$ of F and hence $i \leq j$ in $|\alpha_F|$. This shows that γ_F is order preserving.

Furthermore $\tilde{\varphi}(i) \leq \tilde{\varphi}(j)$ in $|\alpha_G|$ and in $\{1, \dots, n\}$. Since $\{\beta_G \tilde{\varphi}(i), \beta_G \tilde{\varphi}(j)\} \in E_G$ we get $\tilde{\varphi}(i) \prec \tilde{\varphi}(j)$ in \mathcal{O}_G . So $\tilde{\varphi} : \mathcal{O}_F \rightarrow \mathcal{O}_G$ is also order preserving. \square

We have now proved the following:

Theorem 4.5. *The constructions given in Theorem 3.1 and Lemma 4.4 define an embedding functor*

$$\mathcal{Q} : \mathcal{Ugraph} \rightarrow \mathcal{Pograph}$$

by

$$\begin{aligned} \mathcal{Q}((F, \alpha_F : \{1, \dots, n\} \rightarrow V_F)) &= (F, \mathcal{O}_F, \beta_F : \mathcal{O}_F \rightarrow V_F), \\ \mathcal{Q}((\varphi_g, \tilde{\varphi} : |\alpha_F| \rightarrow |\alpha_G|)) &= (\varphi_g, \tilde{\varphi} : \mathcal{O}_F \rightarrow \mathcal{O}_G). \end{aligned}$$

Remark 4.6. From the point of view of SDS, pographs and their morphisms are useless since in the course of the construction of the global update function we need the full total order on the poset. Recall that the local update functions are composed in this total order to give the global update function. However, we will show that we always may complete the partial order of \mathcal{O} to a linear order.

Now two questions arise in this context. First, is the construction of the global update function independent of the choice of this total completion of the order? And second, can the completion of the order be chosen in such a way, that it is compatible with morphisms of pographs, so that the construction of the global update function defines a functor? In the end both questions will be answered to the affirmative.

In the following example we will show the following. Given two ugraphs (F, α_F) and (G, α_G) there may be a morphism from the associated pograph $(F, \mathcal{O}_F, \beta_F)$ to $(G, \mathcal{O}_G, \beta_G)$ that does not arise from a morphism of ugraphs $(\varphi_g, \tilde{\varphi}) : (F, \alpha_F) \rightarrow (G, \alpha_G)$. This shows that the functor $\mathcal{Q} : \mathcal{Ugraph} \rightarrow \mathcal{Pograph}$ is not full. This is almost a counterexample to the second question. The example seems to show that there can be morphisms of pographs that do not arise from morphisms of ugraphs. Compare, however, Theorem 4.15.

Example 4.7. We define two ugraphs. Let $F = G$ with $V_F = \{a, b, c\}$ and $E_F = \{\{a, b\}, \{a, c\}\}$. Let $\alpha_F : \{1, 2, 3\} \rightarrow V_F$ be given by $\alpha_F(1) = a, \alpha_F(2) = b, \alpha_F(3) = c$. Let $\alpha_G : \{1, 2, 3\} \rightarrow V_G$ be given by $\alpha_G(1) = a, \alpha_G(2) = c, \alpha_G(3) = b$.

Then $\mathcal{O}_F = \mathcal{O}_G = \{1, 2, 3\}$ with $1 \preceq 2$ and $1 \preceq 3$. Furthermore $\beta_F(1) = a = \beta_G(1), \beta_F(2) = b = \beta_G(3), \beta_F(3) = c = \beta_G(2)$.

Hence $(\varphi_g, \tilde{\varphi}) : (F, \mathcal{O}_F, \beta_F) \rightarrow (G, \mathcal{O}_G, \beta_G)$ with $\varphi_g = \text{id}$ and $\tilde{\varphi}(1) = 1, \tilde{\varphi}(2) = 3, \tilde{\varphi}(3) = 2$ is a morphism of pographs.

Since both graphs have only one connected component we have $|\alpha_F| = \{1, 2, 3\} = |\alpha_G|$ with the natural total order. Hence $\tilde{\varphi} : |\alpha_F| \rightarrow |\alpha_G|$ as defined above is not order preserving.

So the morphism $(\varphi_g, \tilde{\varphi}) : (F, \mathcal{O}_F, \beta_F) \rightarrow (G, \mathcal{O}_G, \beta_G)$ of pographs is not induced by any morphism of ugraphs from (F, α_F) to (G, α_G) .

If a pograph $(F, \mathcal{O}_F, \beta_F)$ has an update schedule $\gamma : \mathcal{O}_F \rightarrow V_F$ then $(F, \mathcal{O}_F, \beta_F) \cong \mathcal{Q}((F, \beta_F \gamma_F^{-1}))$. So different ugraphs can have isomorphic images under \mathcal{Q} . They clearly differ only in their update schedules. The question if \mathcal{Q} is a representative functor, i.e. if for every object $Y \in \mathcal{Pograph}$ there is an $X \in \mathcal{Ugraph}$ with $\mathcal{Q}(X) \cong Y$, is answered in the following.

We use the following proposition about ordered sets:

Proposition 4.8. *Let (\mathcal{O}, \leq) be a poset and (\mathcal{T}, \leq) be a totally ordered set. Let $\varphi : (\mathcal{O}, \leq) \rightarrow (\mathcal{T}, \leq)$ be a poset map. Then there is a total order \leq' on \mathcal{O} extending \leq such that $\varphi : (\mathcal{O}, \leq') \rightarrow (\mathcal{T}, \leq)$ is a poset map.*

Proof. We consider the set of pairs (U, \leq_U) with $U \subseteq \mathcal{O}$ and \leq_U a total order on U , that is an extension of $\leq|_U$, the order \leq on \mathcal{O} restricted to U , such that $\varphi|_U : (U, \leq_U) \rightarrow (\mathcal{T}, \leq)$ is a poset map. The set \mathcal{S} of these pairs is inductively ordered by $(U, \leq_U) \sqsubseteq (V, \leq_V)$ iff $U \subseteq V$ and $(\leq_V)|_U = \leq_U$. Thus \mathcal{S} has a maximal element (U, \leq') by Zorn's Lemma.

Assume $U \neq \mathcal{O}$. Let $x \in \mathcal{O} \setminus U$. Define $U_x := \{u \in U \mid \varphi(u) < \varphi(x) \text{ or } \exists w \in U : u \leq' w \leq x\}$. Define the following relation \leq'' on $U \cup \{x\}$, where $u, v \in U$

$$\begin{aligned} u \leq'' v &\iff u \leq' v, \\ u \leq'' x &\iff u \in U_x, \\ x \leq'' v &\iff v \notin U_x, \\ x \leq'' x. & \end{aligned}$$

It is easy to show that this is an element in \mathcal{S} . Reflexivity and symmetry of \leq'' are clear from the definition. Furthermore it is clear by definition that this is a total order as soon as we have proved transitivity. The only important cases for transitivity are $u \leq'' x \wedge x \leq'' v, x \leq'' u \leq'' v$, and $u \leq'' v \leq'' x$. These are easy exercises in the axioms for the new order. So is the fact that \leq'' extends \leq .

Then it is clear that \leq'' is a continuation of \leq' . This is a contradiction to the maximality of (U, \leq') . Hence $U = \mathcal{O}$. □

Corollary 4.9 (Trotter, [8]). *Let (\mathcal{O}, \leq) be a poset. Then there is a total order \leq' on \mathcal{O} extending \leq .*

Proof. In the theorem take \mathcal{T} as the one-element totally ordered set and φ the only possible map. □

Observe that the proof of Proposition 4.8 is non-constructive, but that it has sufficient constructive ingredients, in particular the construction of $U \cup \{x\}$ and its order, to define an algorithm in case the sets of interest (e.g. SDS) are finite.

Now we return to the discussion of pographs.

Corollary 4.10. *Let (F, \mathcal{O}, β) be a pograph. Then there exists an update schedule $\gamma : \mathcal{O} \rightarrow \{1, \dots, n\}$.*

Corollary 4.11. *The embedding functor*

$$\mathcal{Q} : \mathcal{U}graph \rightarrow \mathcal{P}ograph$$

is a representative functor.

Definition 4.12. Let $(F, \gamma_F : \mathcal{O}_F \rightarrow \{1, \dots, m\})$ and $(G, \gamma_G : \mathcal{O}_G \rightarrow \{1, \dots, n\})$ be pographs with update schedules. A morphism of pographs with update schedule consists of

- a morphism of graphs $\varphi_g : F \rightarrow G$,
- a morphism of posets $\tilde{\varphi} : \mathcal{O}_F \rightarrow \mathcal{O}_G$,

such that

(1) the diagram

$$\begin{array}{ccc} \mathcal{O}_F & \xrightarrow{\beta_F} & V_F \\ \tilde{\varphi} \downarrow & & \downarrow \varphi_g \\ \mathcal{O}_G & \xrightarrow{\beta_G} & V_G \end{array}$$

commutes and

(2) for all $i, j \in \mathcal{O}_F$ with $\beta_F(i)$ and $\beta_F(j)$ contained in a common connected component of F

$$\gamma_F(i) \leq \gamma_F(j) \implies \gamma_G \tilde{\varphi}(i) \leq \gamma_G \tilde{\varphi}(j).$$

Remark 4.13. It is clear that the composition of two morphisms of pographs with update schedule is again a morphism of pographs with update schedule. Thus pographs with update schedule (F, γ_F) as objects and their morphisms form a category $\mathcal{U}pograph$.

Theorem 4.14. *There is an equivalence between the category of ugraphs and the category of pographs with update schedule*

$$\mathcal{P} : \mathcal{U}graph \simeq \mathcal{U}pograph.$$

Proof. The construction given in Theorem 3.1 defines an pograph with update schedule $\mathcal{P}((F, \alpha_F)) = (F, \mathcal{O}_F, \beta_F, \gamma_F)$ for every ugraph (F, α_F) . Conversely every pograph $(F, \mathcal{O}_F, \beta_F)$ with update schedule $\gamma_F : \mathcal{O}_F \rightarrow \{1, \dots, n\}$ defines a ugraph $\mathcal{P}'((F, \mathcal{O}_F, \beta_F, \gamma_F)) = (F, \beta_F \gamma_F^{-1} : \{1, \dots, n\} \rightarrow V_F)$.

Given a morphism $(\varphi_g, \tilde{\varphi})$ of ugraphs we have seen in Lemma 4.4 that we get a morphism $\mathcal{P}((\varphi_g, \tilde{\varphi})) = (\varphi_g, \tilde{\varphi})$ of pographs.

We have to show that condition (2) is satisfied. We have $\beta_F := \alpha_F$ and $\gamma_F := \text{id}$ from the construction of \mathcal{P} . Let $i, j \in \mathcal{O}_F$ be given with $\beta_F(i) = \alpha_F(i)$ and $\beta_F(j) = \alpha_F(j)$ contained in a common connected component of F . Assume that $i = \gamma_F(i) \leq \gamma_F(j) = j$ in $\{1, \dots, m\}$. Then $\tilde{\varphi}(i) \leq \tilde{\varphi}(j)$ in $|\alpha_G|$ hence also in $\{1, \dots, n\}$. This means $\gamma_G \tilde{\varphi}(i) = \tilde{\varphi}(i) \leq \tilde{\varphi}(j) = \gamma_G \tilde{\varphi}(j)$ in $\{1, \dots, n\}$.

Given a morphism $(\varphi_g, \tilde{\varphi}) : (F, \mathcal{O}_F, \beta_F, \gamma_F) \rightarrow (G, \mathcal{O}_G, \beta_G, \gamma_G)$ in $\mathcal{U}pograph$. By the construction of \mathcal{P}' we have $\alpha_F = \beta_F \gamma_F^{-1}$ and $\alpha_G = \beta_G \gamma_G^{-1}$. Define $\tilde{\varphi}' := \gamma_G \tilde{\varphi} \gamma_F^{-1} : |\alpha_F| \rightarrow |\alpha_G|$. Then $\varphi_g \alpha_F = \varphi_g \beta_F \gamma_F^{-1} = \beta_G \tilde{\varphi} \gamma_F^{-1} = \beta_G \gamma_G^{-1} \tilde{\varphi}' = \alpha_G \tilde{\varphi}'$.

So it remains to show that $\tilde{\varphi}' : |\alpha_F| \rightarrow |\alpha_G|$ is order preserving. Let $i, j \in |\alpha_F|$ with $i \leq j$ in $|\alpha_F|$ be given. Then $i \leq j$ in $\{1, \dots, m\}$, and $\alpha_F(i)$ and $\alpha_F(j)$ are in the same connected component in F . Let $i' := \gamma_F^{-1}(i)$ and $j' := \gamma_F^{-1}(j)$. Then $i', j' \in \mathcal{O}_F$ with $\beta_F(i')$ and $\beta_F(j')$ contained in a common connected component of F . Furthermore we have $\gamma_F(i') = i \leq j = \gamma_F(j')$ hence $\gamma_G \tilde{\varphi}(i') \leq \gamma_G \tilde{\varphi}(j')$. Obviously $\alpha_G \tilde{\varphi}'(i) = \beta_G \gamma_G^{-1} \tilde{\varphi}'(i) = \beta_G \tilde{\varphi}(i') = \varphi_g \beta_F \gamma_F^{-1}(i) = \varphi_g \alpha_F(i)$ and $\alpha_G \tilde{\varphi}'(j) = \varphi_g \alpha_F(j)$ are contained in a common connected component of G . Finally we have $\tilde{\varphi}'(i) = \gamma_G \tilde{\varphi}(i') \leq \gamma_G \tilde{\varphi}(j') = \tilde{\varphi}'(j)$ in $\{1, \dots, n\}$ since γ_G is order preserving so $\tilde{\varphi}'(i) \leq \tilde{\varphi}'(j)$ in $|\alpha_G|$.

It is clear that \mathcal{P} and \mathcal{P}' are functors. Furthermore we have $\mathcal{P}' \mathcal{P} = \text{Id}$ and $\mathcal{P} \mathcal{P}' \cong \text{Id}$ given by the construction of the strong isomorphism in Proposition 3.6 which is an isomorphism in $\mathcal{U}pograph$. \square

The construction of Corollary 4.10 can be extended to morphisms as follows.

Theorem 4.15. *Let $(F, \mathcal{O}_F, \beta_F)$ and $(G, \mathcal{O}_G, \beta_G)$ be pographs and let (φ_g, \tilde{g}) be a morphism of pographs.*

Then there exist update schedules $\gamma_F : \mathcal{O}_F \rightarrow \{1, \dots, m\}$ and $\gamma_G : \mathcal{O}_G \rightarrow \{1, \dots, n\}$ and a morphism of ugraphs $(\varphi_g, \tilde{\varphi}) : (F, \alpha_F) \rightarrow (G, \alpha_G)$ that restricts to the given morphism of pographs (φ_g, \tilde{g}) .

Proof. This is a direct consequence of Proposition 4.8 and Corollary 4.9. \square

5. Sequential dynamical systems on pographs

We have proved that every ugraph defines a pograph with update schedule unique up to strong isomorphism and conversely. Furthermore morphisms of ugraphs turn out to be special morphisms of pographs. We apply this now to SDS.

Definition 5.1. *A sequential dynamical system or an SDS on a pograph F*

$$\mathcal{F} = (F, (k[a] \mid a \in V_F), (f_a \mid a \in V_F))$$

consists of

- (1) a finite pograph F ,
- (2) a family of sets $(k[a] \mid a \in V_F)$ in \mathcal{L} ,
- (3) a family of local functions $(f_a : k^r \rightarrow k^r \mid a \in V_F, f_a \text{ local at } a)$ in \mathcal{L} .

Given an SDS on a pograph and assume that we have an update schedule $\gamma : \mathcal{O} \rightarrow \{1, \dots, n\}$ for the pograph. Then we can define a *global update function* as we did for an SDS over a ugraph:

$$f_{\beta\gamma^{-1}} := f_{\beta\gamma^{-1}(1)} \circ \dots \circ f_{\beta\gamma^{-1}(n)} : k^r \rightarrow k^r.$$

For an SDS on a pograph without a given update schedule, however, it is not clear how to construct a global update function. So the following proposition and its consequences are surprising and important.

Proposition 5.2. *Let \mathcal{F} be a finite SDS on a pograph F . Let $\gamma, \eta : \mathcal{O}_F \rightarrow \{1, \dots, n\}$ be update schedules. Then*

$$f_{\beta\gamma^{-1}} = f_{\beta\eta^{-1}}.$$

Proof. For all $i \in \{1, \dots, n\}$ let $a_i := \beta\gamma^{-1}(i)$ and $b_i := \beta\eta^{-1}(i)$. We want to show $f_{a_1} \circ \dots \circ f_{a_n} = f_{b_1} \circ \dots \circ f_{b_n}$. For each j there is an $i (= \gamma\eta^{-1}(j))$ such that $b_j = a_i$ and conversely.

Claim: Given $j, k \in \{1, \dots, n\}$ such that $k < j$ and $\eta\gamma^{-1}(j) < \eta\gamma^{-1}(k)$, then $f_{\beta\gamma^{-1}(j)} \circ f_{\beta\gamma^{-1}(k)} = f_{\beta\gamma^{-1}(k)} \circ f_{\beta\gamma^{-1}(j)}$. Let $u := \gamma^{-1}(j)$ and $v := \gamma^{-1}(k)$. Then $\gamma(v) < \gamma(u)$ and $\eta(u) < \eta(v)$. Since both maps γ and η are order preserving we get that $u \not\leq v$ and $v \not\leq u$ in the poset \mathcal{O}_F hence $\{\beta(u), \beta(v)\} \notin E_F$. So we get that $f_{\beta(u)} \circ f_{\beta(v)} = f_{\beta(v)} \circ f_{\beta(u)}$ by Remark 2.1.

Assume now that we have already arranged a reordering of the update function such that

$$f_{a_1} \circ \dots \circ f_{a_n} = f_{b_1} \circ \dots \circ f_{b_{j-1}} \circ f_{a_l} \circ \dots \circ f_{a_m},$$

where the f_{a_l}, \dots, f_{a_m} are those factors among the f_{a_1}, \dots, f_{a_n} that do not occur as factors $f_{b_1}, \dots, f_{b_{j-1}}$ and where their product is taken in the same order as in f_{a_1}, \dots, f_{a_n} .

Let $\eta^{-1}(j) = \gamma^{-1}(i)$ and thus $b_j = a_i$ and $f_{b_j} = f_{a_i}$. We want to shift the local update function $f_{b_j} = f_{a_i}$ in the right hand side of the equation toward the left. Given k with $l \leq k < i$ then $\eta\gamma^{-1}(i) = j < \eta\gamma^{-1}(k)$ (because $f_{b_{\eta\gamma^{-1}(k)}}$ in the update function $f_{b_1} \circ \dots \circ f_{b_n}$ does not occur in the partial product $f_{b_1} \circ \dots \circ f_{b_{j-1}}$). Thus $f_{b_j} \circ f_{a_k} = f_{a_k} \circ f_{b_j}$. So we can rearrange the update function to

$$f_{a_1} \circ \dots \circ f_{a_n} = f_{b_1} \circ \dots \circ f_{b_j} \circ f_{a_{l'}} \circ \dots \circ f_{a_{m'}}.$$

By induction this completes the proof. \square

Theorem 5.3. Let \mathcal{F} be an SDS on a pograph. Then \mathcal{F} has a well-defined global update function $f_\beta := f_{\beta\gamma^{-1}} : k^r \rightarrow k^r$.

Proof. By Corollary 4.10 there is an update schedule γ . By Proposition 5.2 the global update function is independent of the choice of the update schedule γ . \square

In a subsequent paper where we will define and study morphisms of SDS on pographs we will use this theorem and Theorem 4.15 to prove the following:

The construction of global update functions of SDS on pographs and its associated state graphs define a functor to the category of dynamical systems and to the category of graphs.

Remark 5.4. What we have proved is that any two update schedules of pographs give the same global update function. Furthermore, the number of different update schedules for a ugraph giving the same global update function is greater than or equal to the number of bijective order preserving maps $\mathcal{O}_F \rightarrow \{1, \dots, n\}$.

Actually we can prove more.

Proposition 5.5. Let F be a graph and let $\alpha : \{1, \dots, n\} \rightarrow V_F$ be an update schedule with canonical decomposition $\{1, \dots, n\} \xleftarrow{\gamma} \mathcal{O}_F \xrightarrow{\beta} V_F$. If $\tilde{\alpha} : \{1, \dots, n\} \rightarrow V_F$ is an update schedule for F that does not factor through β then there is a structure of an SDS on F (in particular a family of state spaces and a family of local update functions) such that $f_\alpha \neq f_{\tilde{\alpha}}$.

We first need the following:

Lemma 5.6. Let $\beta : \mathcal{O} \rightarrow V_F$ and $\tilde{\beta} : \tilde{\mathcal{O}} \rightarrow V_F$ be pographs on the same graph F . Assume there is a bijective order preserving map $\delta : \mathcal{O} \rightarrow \tilde{\mathcal{O}}$ such that $\beta\delta = \tilde{\beta}$. Then δ is an isomorphism of posets.

Proof. Let $i < j$ in $\tilde{\mathcal{O}}$. Then $\{\tilde{\beta}(i), \tilde{\beta}(j)\} \in E_F$. Let $u := \delta^{-1}(i)$ and $v := \delta^{-1}(j)$. Then $\{\beta(u), \beta(v)\} = \{\tilde{\beta}\delta(u), \tilde{\beta}\delta(v)\} = \{\tilde{\beta}(i), \tilde{\beta}(j)\} \in E_F$, hence $u \leq v$ or $v \leq u$. Since δ is order preserving we obtain $u \leq v$ hence $\delta^{-1}(i) \leq \delta^{-1}(j)$. Thus δ^{-1} is order preserving. \square

Proof of Proposition. Since $\alpha \neq \tilde{\alpha}$ we can consider three cases.

Case 1: Let $\text{Im}(\alpha) \neq \text{Im}(\tilde{\alpha})$. Without loss of generality assume $a \in \text{Im}(\alpha) \setminus \text{Im}(\tilde{\alpha})$. Then set $f_b = \text{id}$ for all $b \neq a$ in V_F . Then the global update functions are $f_{\tilde{\alpha}} = \text{id}$ and $f_{\alpha} = f_a^r \neq \text{id}$ for a suitable choice of f_a , hence $f_{\alpha} \neq f_{\tilde{\alpha}}$.

Case 2: Let $\text{Im}(\alpha) = \text{Im}(\tilde{\alpha})$ and assume there is an $a \in \text{Im}(\alpha)$ such that f_{α} contains r copies of f_a and $f_{\tilde{\alpha}}$ contains $s \neq r$ copies of f_a . Again set all $f_b = \text{id}$ for $b \neq a$. Then by a suitable choice of f_a we get $f_{\alpha} \neq f_{\tilde{\alpha}}$.

Case 3: Let $\text{Im}(\alpha) = \text{Im}(\tilde{\alpha})$ and let there be the same number of factors f_a in f_{α} resp. $f_{\tilde{\alpha}}$ for all $a \in V_F$. So $f_{\tilde{\alpha}}$ arises from f_{α} by a reordering of the factors. Thus there is a permutation $\sigma : \{1, \dots, n\} \rightarrow \{1, \dots, n\}$ such that $\tilde{\alpha}\sigma = \alpha$. We can choose σ on the preimage of each $a \in E_F$ to be order preserving, since the orderings of factors f_a among each other in the global update function are irrelevant.

We use the decomposition of α and $\tilde{\alpha}$ into $(\mathcal{O}_F, \beta, \gamma = \text{id})$ and $(\tilde{\mathcal{O}}_F, \tilde{\beta}, \gamma = \text{id})$ resp. as constructed in Theorem 3.1. Then σ may be considered as a map $\sigma : \mathcal{O}_F \rightarrow \tilde{\mathcal{O}}_F$. Assume $f_{\alpha} = f_{\tilde{\alpha}}$ for all choices of families of local update functions (f_a) . We claim that $\sigma : \mathcal{O}_F \rightarrow \tilde{\mathcal{O}}_F$ is order preserving.

Let $a, b \in V_F$ with $\{a, b\} \in E_F$. Assume that the subwords of f_{α} and $f_{\tilde{\alpha}}$ consisting of factors f_a and f_b are of the form $f_a^{i_1} f_b^{i_2} f_a^{i_3} \dots f_b^{i_r}$ and $f_a^{j_1} f_b^{j_2} f_a^{j_3} \dots f_b^{j_s}$ (with $i_k, j_k > 0$ except for i_1, i_r, j_1, j_s which may also be zero). We will show further down that there are choices for f_a and f_b such that $f_a^{i_1} f_b^{i_2} f_a^{i_3} \dots f_b^{i_r} = f_a^{j_1} f_b^{j_2} f_a^{j_3} \dots f_b^{j_s}$ iff $(i_1, \dots, i_r) = (j_1, \dots, j_s)$. This is equivalent to σ being order preserving on the preimage of $\{a, b\}$ under α . Since we may choose $f_c = \text{id}$ for all $c \neq a, b$, the assumption $f_{\alpha} = f_{\tilde{\alpha}}$ implies that σ is order preserving on the preimage of $\{a, b\}$ under α .

Now we show under the given assumptions that σ is order preserving. Let $i, j \in \mathcal{O}_F$ with $i \prec j$. Then $i \leq j$ (as numbers) and $\{\alpha(i), \alpha(j)\} \in E_F$. Define $a := \alpha(i)$ and $b := \alpha(j)$. Then $\tilde{\alpha}\sigma(i) = a$ and $\tilde{\alpha}\sigma(j) = b$. Hence $\sigma(i) \prec \sigma(j)$ or $\sigma(j) \prec \sigma(i)$ in $\tilde{\mathcal{O}}_F$. If $\sigma(j) \prec \sigma(i)$ holds, then σ is not order preserving on the preimage of $\{a, b\}$ under α . Hence we have $\sigma(i) \prec \sigma(j)$ and thus σ is order preserving.

By Lemma 5.6 we find that $\tilde{\alpha}$ factors through $\beta : \mathcal{O}_F \rightarrow V_F$. This is a contradiction to the assumption in the proposition. So $f_{\alpha} \neq f_{\tilde{\alpha}}$ for some choice of a family of local update functions (f_a) .

It remains to show that there are choices for f_a and f_b such that $f_a^{i_1} f_b^{i_2} f_a^{i_3} \dots f_b^{i_r} = f_a^{j_1} f_b^{j_2} f_a^{j_3} \dots f_b^{j_s}$ iff $(i_1, \dots, i_r) = (j_1, \dots, j_s)$. Take $k[a] = k[b] = \mathbb{N}$ (or a suitable finite subset thereof). Define

$$f_a(\dots, x[a], \dots, x[b], \dots) = \begin{cases} (\dots, p \cdot x[a], \dots, x[b], \dots) & \text{if } x[a] > x[b], \\ (\dots, q \cdot x[b], \dots, x[b], \dots) & \text{if } x[a] \leq x[b], \end{cases}$$

where p is the largest prime dividing $x[a]$ and q is the smallest prime not dividing $x[b]$. Furthermore let

$$f_b(\dots, x[a], \dots, x[b], \dots) = \begin{cases} (\dots, x[a], \dots, q \cdot x[a], \dots) & \text{if } x[a] \geq x[b], \\ (\dots, x[a], \dots, p \cdot x[b], \dots) & \text{if } x[a] < x[b], \end{cases}$$

where p is the largest prime dividing $x[b]$ and q is the smallest prime not dividing $x[a]$. Then

$$f_a^{i_1} f_b^{i_2} \dots f_a^{i_{r-1}} f_b^{i_r}(\dots, 1, \dots, 1, \dots) = (\dots, 2^{i_r} \cdot 3^{i_{r-1}} \cdot \dots \cdot p_{r-1}^{i_2} \cdot p_r^{i_1}, \dots, 2^{i_r} \cdot 3^{i_{r-1}} \cdot \dots \cdot p_{r-1}^{i_2}, \dots),$$

for $i_k > 0$. A similar argument holds for $i_1 = 0$ and/or $i_r = 0$. This proves the claim and the proposition. \square

Corollary 5.7. *The number of bijective order preserving maps $\mathcal{O}_F \rightarrow \{1, \dots, n\}$ is a sharp lower bound for the number of different update schedules for a graph giving the same global update function.*

Let F be a graph. We call two update schedules $\alpha, \alpha' : \{1, \dots, n\} \rightarrow V_F$ equivalent, iff for all choices of local state spaces $(k[a] \mid a \in V_F)$ and all choices of local update functions $(f_a \mid a \in V_F)$ on the given graph F the global update functions f_{α} and $f_{\alpha'}$ are equal.

Corollary 5.8. *There is a one-to-one correspondence between equivalence classes of update schedules of length n and poset models with n elements of the graph F .*

6. On the posets of pographs

We have seen that the structure of SDS strongly depends on the underlying pographs. In this section we investigate which posets can occur as posets in pographs.

Definition 6.1. A pograph $(F, \mathcal{O}_F, \beta)$ is called *rigid*, if there is only one bijective map of posets $\gamma : \mathcal{O}_F \rightarrow \{1, \dots, n\}$. In other words, a pograph is rigid, if \mathcal{O}_F has a unique extension to a total ordering.

An update schedule $\alpha : \{1, \dots, n\} \rightarrow V_F$ of a graph F is called *rigid* if there is only one bijective map of posets $\gamma : \mathcal{O}_F \rightarrow \{1, \dots, n\}$, where \mathcal{O}_F is induced by α .

Proposition 6.2. α is rigid if and only if \mathcal{O}_F is totally ordered.

Proof. This follows from the fact that a poset has a unique extension to a total ordering if and only if it is already totally ordered. See [8, p. 17]. \square

Thus for pographs with totally ordered poset \mathcal{O}_F we have only one update schedule. The opposite observation is that a pograph with a discrete poset (no two elements can be compared) has n^n update functions $\beta : \mathcal{O}_F \rightarrow \{1, \dots, n\}$. They all have the same global update function.

Examples 6.3.

- (1) A linear graph a_1, \dots, a_n with $E_F = \{\{a_i, a_{i+1}\} \mid i = 1, \dots, n - 1\}$ has a rigid update schedule $\alpha(i) = a_i$. An example for such an SDS is a column of cars on a road a_1, \dots, a_n where each car determines its behavior or its local update function upon the preceding car. This gives a linear graph and the update schedule α cannot be changed without the risk of changing the global update function of the system.
- (2) A linear graph with at least three vertices has a non-rigid update schedule. Let a, b, c be three consecutive vertices in F with edges $\{a, b\}$ and $\{b, c\}$. Consider the rigid update schedule α with $\alpha(i) = a, \alpha(i + 1) = b$, and $\alpha(i + 2) = c$. Define new update schedules $\tilde{\alpha}$ by $\tilde{\alpha}(j) := \alpha(j)$ for $j \neq i + 1, i + 2$ and $\tilde{\alpha}(i + 1) = c, \tilde{\alpha}(i + 2) = b$ and $\bar{\alpha}$ by $\bar{\alpha}(j) := \alpha(j)$ for $j \neq i, i + 1, i + 2$, and $\bar{\alpha}(i) = c, \bar{\alpha}(i + 1) = a, \bar{\alpha}(i + 2) = b$. Then it is easy to see that both update schedules define the same poset \mathcal{O} (as constructed in the proof of Theorem 3.1).
- (3) The hypercube $F = \{(x_1, \dots, x_n) \mid x_i \in \{0, 1\}\}$ with 2^n vertices has a rigid update schedule $\alpha : \{1, \dots, 2^n\} \rightarrow F$. It is well known that F is a Hamiltonian graph. By omitting the last edge in a Hamiltonian path we get a rigid update schedule $\alpha : \{1, \dots, 2^n\} \rightarrow V_F$.
- (4) It is an easy exercise to show that hypercubes of dimension $n \geq 2$ have non-rigid update schedules.
- (5) An interesting example of a rigid update schedule is given in Example 3.4.

Now we show that any finite poset \mathcal{O} can occur as a poset of a pograph.

Proposition 6.4. (1) Let \mathcal{O} be a finite poset. Let V be a set and $\beta : \mathcal{O} \rightarrow V$ be a map such that $i \triangleleft j$ implies $\beta(i) \neq \beta(j)$ for all $i, j \in \mathcal{O}$. Then the Hasse diagram

$$E_F := \{\{\beta(i), \beta(j)\} \mid i, j \in \mathcal{O} : i \triangleleft j\}$$

defines a graph F with vertex set $V_F = V$ such that (F, \mathcal{O}, β) is a pograph.

(2) Let \mathcal{O} be a finite poset. Let V be a set and $\beta : \mathcal{O} \rightarrow V$ be a map such that $i \triangleleft j$ implies $\beta(i) \neq \beta(j)$ for all $i, j \in \mathcal{O}$. Then

$$E_F := \{\{\beta(i), \beta(j)\} \mid i, j \in \mathcal{O} : i < j \text{ and } \beta(i) \neq \beta(j)\}$$

defines a graph F with vertex set $V_F = V$ such that (F, \mathcal{O}, β) is a pograph.

(3) The graph constructed in (1) over $\beta : \mathcal{O} \rightarrow V$ is the smallest subgraph of the complete graph on V such that (F, \mathcal{O}, β) is a pograph. The graph constructed in (2) over $\beta : \mathcal{O} \rightarrow V$ is the largest subgraph of the complete graph on V such that (F, \mathcal{O}, β) is a pograph.

Proof. (1) We have to check that $\beta : \mathcal{O} \rightarrow V_F$ is a pograph. By definition of the graph F we have for all $i, j \in \mathcal{O}$

$$i \triangleleft j \iff \{\beta(i), \beta(j)\} \in E_F.$$

This implies both conditions (1) and (2).

(2) Again we have to check that $\beta : \mathcal{O} \rightarrow V_F$ is a pograph. By definition of the graph F we have for all $i, j \in \mathcal{O}$ such that $\beta(i) \neq \beta(j)$:

$$\{\beta(i), \beta(j)\} \in E_F \iff i < j \vee j < i.$$

This implies also both conditions (1) and (2).

(3) Let F be a graph with $V_F = V$ and assume that (F, \mathcal{O}, β) is a pograph. Let F_{\min} be the graph constructed in (1) from the map β . Let $\{\beta(i), \beta(j)\}$ be in F_{\min} . Then $i \triangleleft j$ hence $\{\beta(i), \beta(j)\}$ is an edge in F by axiom (1).

Let F_{\max} be the graph constructed in (2) from the map β . Let $\{\beta(i), \beta(j)\}$ be in F . Then $i < j$ (or $j < i$) by axiom (2). Hence $\{\beta(i), \beta(j)\}$ is an edge in F_{\max} . \square

It is interesting to note that any graph F' between F_{\min} and F_{\max} as constructed above (subgraph and supergraph on the same set of vertices) gives also a poset model $\beta : \mathcal{O} \rightarrow V_{F'}$ as can be easily checked.

References

- [1] C.L. Barrett, H.S. Mortveit, C.M. Reidys, Elements of a theory of computer simulation II: sequential dynamical systems, *Appl. Math. Comput.* 107 (2–3) (1999) 121–136.
- [2] C.L. Barrett, H.S. Mortveit, C.M. Reidys, Elements of a theory of computer simulation III: equivalence of SDS, *Appl. Math. Comput.* 122 (2001) 325–340.
- [3] C.L. Barrett, C.M. Reidys, Elements of a theory of simulation, I: sequential CA over random graphs, *Appl. Math. Comput.* 98 (1999) 241–259.
- [4] L. Garcia, A. Jarrar, R. Laubenbacher, Sequential dynamical systems over words, *Appl. Math. Comput.*, 2005, to appear.
- [5] R. Laubenbacher, B. Pareigis, Equivalence relations on finite dynamical systems, *Adv. Appl. Math.* 26 (2001) 237–251.
- [6] R. Laubenbacher, B. Pareigis, Decomposition and simulation of sequential dynamical systems, *Adv. Appl. Math.* 30 (2003) 655–678.
- [7] C. Reidys, Acyclic orientations of random graphs, *Adv. Appl. Math.* 21 (1998) 181–192.
- [8] T. Trotter, *Combinatorics and Partially Ordered Sets*, Johns Hopkins University Press, Baltimore, 1992.