

Available online at www.sciencedirect.com
ScienceDirectADVANCES IN
Mathematics

Advances in Mathematics 228 (2011) 97–152

www.elsevier.com/locate/aim

Non-commutative integrability, paths and quasi-determinants

Philippe Di Francesco^{a,b,*}, Rinat Kedem^c^a Department of Mathematics, University of Michigan, 530 Church Street, Ann Arbor, MI 48190, USA^b Institut de Physique Théorique du Commissariat à l'Energie Atomique, Unité de Recherche associée du CNRS, CEA Saclay/IPhT/Bat 774, F-91191 Gif sur Yvette Cedex, France^c Department of Mathematics, University of Illinois MC-382, Urbana, IL 61821, USA

Received 25 June 2010; accepted 10 May 2011

Available online 31 May 2011

Communicated by Andrei Zelevinsky

Abstract

In previous work, we showed that the solution of certain systems of discrete integrable equations, notably Q and T -systems, is given in terms of partition functions of positively weighted paths, thereby proving the positive Laurent phenomenon of Fomin and Zelevinsky for these cases. This method of solution is amenable to generalization to non-commutative weighted paths. Under certain circumstances, these describe solutions of discrete evolution equations in non-commutative variables: Examples are the corresponding quantum cluster algebras (Berenstein and Zelevinsky (2005) [3]), the Kontsevich evolution (Di Francesco and Kedem (2010) [10]) and the T -systems themselves (Di Francesco and Kedem (2009) [8]). In this paper, we formulate certain non-commutative integrable evolutions by considering paths with non-commutative weights, together with an evolution of the weights that reduces to cluster algebra mutations in the commutative limit. The general weights are expressed as Laurent monomials of quasi-determinants of path partition functions, allowing for a non-commutative version of the positive Laurent phenomenon. We apply this construction to the known systems, and obtain Laurent positivity results for their solutions in terms of initial data.

© 2011 Elsevier Inc. All rights reserved.

Keywords: Discrete integrable systems; Noncommutative; Quasi-determinants; Path models; Continued fractions; Cluster algebra; Quantum cluster algebra

* Corresponding author at: Institut de Physique Théorique du Commissariat à l'Energie Atomique, Unité de Recherche associée du CNRS, CEA Saclay/IPhT/Bat 774, F-91191 Gif sur Yvette Cedex, France.

E-mail addresses: philippe.di-francesco@cea.fr (P. Di Francesco), rinat@illinois.edu (R. Kedem).

Contents

| | | |
|------|---|-----|
| 1. | Introduction | 98 |
| 1.1. | Discrete integrable systems and positivity | 98 |
| 1.2. | The A_r T -system and Q -system | 99 |
| 1.3. | Example: Commutative and non-commutative A_1 Q -systems | 100 |
| 1.4. | Plan of the paper | 101 |
| 2. | Continued fraction solutions to Q -systems | 102 |
| 2.1. | The A_r Q -system | 102 |
| 2.2. | Initial data sets | 102 |
| 2.3. | Path partition functions | 103 |
| 2.4. | Mutations | 107 |
| 3. | Non-commutative weights and evolutions | 112 |
| 3.1. | Paths with non-commutative weights and non-commutative continued fractions | 112 |
| 3.2. | Mutations | 115 |
| 3.3. | An explicit expression for the weights | 118 |
| 4. | Application I: T -system | 128 |
| 4.1. | The A_r T -system as non-commutative Q -system | 129 |
| 4.2. | Quasi-Wronskian formulation | 130 |
| 5. | Application II: Quantum Q -system | 131 |
| 5.1. | Quantum cluster algebra | 131 |
| 5.2. | Quantum Q -system | 133 |
| 5.3. | Quasi-Wronskian formulation | 134 |
| 6. | Other non-commutative systems | 137 |
| 6.1. | Affine rank 2 non-commutative systems | 137 |
| 6.2. | Some rank $2k + 1$ cases | 138 |
| 6.3. | A non-coprime rank 2 case related to rank 3 | 139 |
| 7. | Towards a non-commutative Lindström–Gessel–Viennot theorem | 139 |
| 8. | Conclusion: Towards non-commutative cluster algebras | 142 |
| | Acknowledgments | 143 |
| | Appendix A. Relation of weighted graphs $G_{\mathbf{m}}$ to the graphs $\Gamma_{\mathbf{m}}$ of [6] | 143 |
| | A.1. The graphs $\Gamma_{\mathbf{m}}$ | 143 |
| | A.2. Compactified graphs | 146 |
| | A.3. The graph $G_{\mathbf{m}}$ | 147 |
| | Appendix B. Proof of Theorem 6.2 | 149 |
| | References | 151 |

1. Introduction

1.1. Discrete integrable systems and positivity

Discrete integrable systems are evolution equations in a discrete time variable, which possess sufficiently many conserved quantities (discrete integrals of motion) so that they can be solved explicitly in terms of some discrete Cauchy initial data.

Cluster algebra mutations [15] can be viewed as discrete evolution equations for certain commutative variables, where at each step, the evolution is a rational transformation. The evolution is known to result in variables which are Laurent polynomials as functions of the initial data, even though the evolution equations are rational transformations [16].

Cluster algebras are algebras over commutative variables (the cluster variables) in which sets of variables (the clusters) are associated with the nodes of a regular n -tree, and cluster variables on connected nodes are related by an evolution called a mutation. One can choose a cluster at any fixed node to be the initial data for the evolution. Fomin and Zelevinsky conjectured that in terms of any initial data, the cluster variables at any node are positive Laurent polynomials [15]. This is known as the positivity conjecture.

The A_r Q -systems and T -systems are discrete integrable evolution equations which are a subset of a cluster algebra structure [21,6]. These systems were solved, and thus their positivity property proved, in [9,8] (for the T -systems, see [5] for generic boundary conditions). The approach of [9] and [8] involves the interpretation of the solutions of the systems as partition functions for paths on some discrete target space, with weights explicitly related to the (discrete Cauchy) initial data of the evolution equations.

The T -system can be considered as a particular *non-commutative* version of the Q -system, in which we adjoin a non-commuting element to the commutative algebra generated by Q -system solutions [8]. The T -system equations are matrix elements of the cluster algebra mutations twisted by the non-commutative element. The solutions are then expressed in terms of partition functions for paths as for the Q -system, but with non-commutative weights.

In all these examples, the positive Laurent property follows from the fact that the weights are explicit positive Laurent monomials of the initial data.

The aim of the present paper is to formalize this underlying path structure in the more general case of arbitrary non-commutative weights, with the idea of defining non-commutative (integrable) cluster algebra mutations with a built-in non-commutative positive Laurent property.

We note that there exists an example of a non-commutative cluster algebra with these properties: the quantum cluster algebra [3], where the cluster variables q -commute. This is the mildest form of non-commutativity, and in this case, much more can be said about the structure of the solutions.

1.2. The A_r T -system and Q -system

The A_r T -system is a system of discrete evolution equations for the commuting variables $\{T_{i,j,k}\}$. It can be written in the following form¹:

$$T_{i,j,k+1}T_{i,j,k-1} = T_{i,j+1,k}T_{i,j-1,k} + T_{i+1,j,k}T_{i-1,j,k}, \quad 1 \leq i \leq r, j, k \in \mathbb{Z}, \quad (1.1)$$

with the convention that $T_{0,j,k} = T_{r+1,j,k} = 1$.

Eq. (1.1) (with a sign change on the right-hand side) was first introduced [2,24] in the context of the generalized Heisenberg spin chain in statistical mechanics. With appropriate boundary conditions, it is the “fusion relation” satisfied by its transfer matrices. Its solutions are the q -characters of $U_q(\mathfrak{sl}_{r+1})$ [17,27].

A closely related system is the A_r Q -system [22],

$$R_{i,k+1}R_{i,k-1} = R_{i,k}R_{i,k} + R_{i+1,k}R_{i-1,k}, \quad 1 \leq i \leq r, k \in \mathbb{Z}, \quad (1.2)$$

¹ The relative minus sign which appears in the original version [2,24] is normalized away in this paper.

with $R_{0,k} = R_{r+1,k} = 1$. This system is obtained from the T -system (1.1) by dropping the index j , for example, by considering solutions of (1.1) which are 2-periodic in j , $T_{i,j+2,k} = T_{i,j,k}$ for all i, j, k . Then $R_{i,k} = T_{i,i+k \bmod 2,k}$ is a solution to (1.2).

Remark 1.1. Given the appropriate boundary conditions ($R_{i,0} = (-1)^{\lfloor i/2 \rfloor}$) this system is satisfied by the characters $Q_{i,n} = (-1)^{\lfloor i/2 \rfloor} R_{i,n}$ ($n > 0$) of the irreducible A_r -module with special highest weights which are multiples of one of the fundamental weights. Using Eq. (1.2), these are expressed as polynomials in the characters of the fundamental modules, identified as $\{Q_{i,1}: 1 \leq i \leq r\}$.

In reference to the origin of the equation, we refer to Eq. (1.2) as the A_r Q -system throughout this paper, without imposing this special boundary condition.

To solve such a system means to express the variables for any n in terms of the initial data set. For example, valid initial data sets for the Q -system are sets of $2r$ variables $\{Q_{i,m_i}, Q_{i,m_i+1}: 1 \leq i \leq r\}$ such that $|m_i - m_{i+1}| \leq 1$. For fixed values of (i, k) or (i, j, k) , Eqs. (1.1), (1.2) are mutations of cluster variables in a cluster algebra. We will therefore investigate the systems from the point of view of local mutations of their initial data.

1.3. Example: Commutative and non-commutative A_1 Q -systems

The simplest case of the Q -system occurs for $r = 1$. This example was previously considered from a different point of view by several authors [4,28,26,13]. Let us recall the solutions of both the commutative and non-commutative versions of the A_1 Q -system [7,10]. We wish to stress the similarity of their path solutions, which relies on the existence of a sufficient number of conserved quantities and a linear recursion relation involving these.

Consider the recursion relation for the commutative variables $\{R_n\}$,

$$R_{n+1}R_{n-1} = (R_n)^2 + 1. \tag{1.3}$$

The system has a manifest translational invariance. We thus compute the solution $R_n, n \geq 0$ in terms of the initial data (R_0, R_1) .

The system is integrable: The quantity $c = \frac{R_{n+1}}{R_n} + \frac{1}{R_n R_{n+1}} + \frac{R_n}{R_{n+1}}$ is independent of n . Moreover, the solutions of (1.3) obey the linear recursion relation $R_{n+1} - cR_n + R_{n-1} = 0$. Let

$$y_1 = R_1/R_0, \quad y_2 = 1/(R_0R_1), \quad y_3 = 1/y_1 = R_0/R_1$$

and define the generating function

$$F_0(t) = \sum_{n=0}^{\infty} t^n \frac{R_n}{R_0} = \frac{1}{1 - \frac{ty_1}{1 - \frac{ty_2}{1 - ty_3}}}. \tag{1.4}$$

This is the partition function for weighted paths on the integer segment $[1, 4]$, from 1 to 1, and with steps ± 1 , with weights 1 for step $i \rightarrow i + 1$ and ty_i for step $i + 1 \rightarrow i, i = 1, 2, 3$. As mentioned above, this interpretation makes positivity manifest, as the y_i 's are explicit Laurent monomials of the initial data (R_0, R_1) .

The non-commutative A_1 Q -system is in a class of wall crossing formulas introduced by Kontsevich, and involves the variables $\mathbf{R}_n \in \mathcal{A}$, a skew field of rational fractions in the variables $\mathbf{R}_0^{\pm 1}, \mathbf{R}_1^{\pm 1}$. It can be written as [10]

$$\mathbf{R}_{n+1}(\mathbf{R}_n)^{-1}\mathbf{R}_{n-1} = \mathbf{R}_n + (\mathbf{R}_n)^{-1}. \tag{1.5}$$

The goal is to express $\mathbf{R}_n, n \geq 0$ in terms of the initial data $(\mathbf{R}_0, \mathbf{R}_1)$.

The system is integrable, in the sense that there are two conserved quantities [10], independent of n :

$$\mathbf{C} = (\mathbf{R}_{n+1})^{-1}\mathbf{R}_n\mathbf{R}_{n+1}(\mathbf{R}_n)^{-1}, \quad \mathbf{K} = \mathbf{R}_{n+1}(\mathbf{R}_n)^{-1} + (\mathbf{R}_{n+1})^{-1}(\mathbf{R}_n)^{-1} + (\mathbf{R}_{n+1})^{-1}\mathbf{R}_n$$

and the solutions to (1.3) obey the linear recursion relation $\mathbf{R}_{n+1} - \mathbf{K}\mathbf{R}_n + \mathbf{C}\mathbf{R}_{n-1} = 0$. Define $\mathbf{y}_1 = \mathbf{R}_1(\mathbf{R}_0)^{-1}, \mathbf{y}_2 = (\mathbf{R}_1)^{-1}(\mathbf{R}_0)^{-1}$ and $\mathbf{y}_3 = (\mathbf{R}_1)^{-1}\mathbf{R}_0$, with $\mathbf{y}_3\mathbf{y}_1 = \mathbf{C}$. Then the generating function

$$\mathbf{F}_0(t) = \sum_{n=0}^{\infty} t^n \mathbf{R}_n(\mathbf{R}_0)^{-1} = (1 - (1 - (1 - t\mathbf{y}_3)^{-1}t\mathbf{y}_2)^{-1}t\mathbf{y}_1)^{-1}. \tag{1.6}$$

This can be interpreted as the partition function of weighted paths on the integer segment [1, 4], from 1 to 1, and with steps ± 1 , with non-commutative weights 1 for $i \rightarrow i + 1$ and $t\mathbf{y}_i$ for $i + 1 \rightarrow i, i = 1, 2, 3$. In this case, the weight of a path is the product of step weights, ordered according to the order in which they are taken.

We see that the non-commutative A_1 Q -system, despite its complexity, has essentially the *same* solution as the commutative one: it is expressed in terms of paths on the same graph, with similar but non-commutative weights related to the initial data, and with the condition that the order in which weights are multiplied is that in which the corresponding steps are taken.

We recall that in the commutative case, the mutation of initial data in terms of which R_n is expressed may be implemented via local *rearrangements* of the finite continued fraction expression (1.4) for $F_0(t)$ [7]. This property was shown to carry over in the non-commutative case [10], where mutations of the initial data are obtained via local non-commutative rearrangements of the non-commutative finite continued fraction expression (1.6) for $\mathbf{F}_0(t)$.

1.4. Plan of the paper

The example above suggests the following strategy for exploration of the non-commutative evolutions. Instead of starting from an evolution equation, we start from the “solution”, namely some path models on the *same* target graphs as those considered in the solution of commutative systems, but with non-commutative weights. It turns out that the partition functions of these paths have a structure similar to that observed in the commutative case. In particular, we will define non-commutative mutations connecting them via non-commutative finite continued fraction rearrangements. Moreover, by investigating the relation between the coefficients of these continued fractions (i.e. the weights of the path models) and the coefficients of their series expansion, we will be led naturally to express results in terms of the quasi-determinants introduced in [18]. Our main result is the construction of a chain of path models with non-commutative weights, all related via mutations, and whose partition functions involve a basic set of non-commuting variables, in a manifestly positive Laurent polynomial way.

The paper is organized as follows.

We first review in Section 2 the solution of the (commutative) A_r Q -system, in a new path formulation related to that of [7] (the equivalence is described in detail in Appendix A). Each initial data, and therefore each graph and associated set of weights is coded by a Motzkin path of length $r - 1$. The corresponding partition functions of weighted paths display a simple finite continued fraction structure, and the mutations are shown to be expressed through local rearrangements.

Section 3 is devoted to the non-commutative generalization of this structure, and involves paths on the same graphs but with non-commutative weights. Mutations are shown to be expressed again as some non-commutative local rearrangements of the finite continued fraction expressions for the partition functions of paths, giving rise to simple evolution equations for the corresponding non-commutative weights. We then define non-commutative variables by discrete quasi-Wronskian determinants of the coefficients of the series expansion of the partition functions. These variables are shown to satisfy an evolution equation (the discrete non-commutative Hirota equation, Theorem 3.27) playing the role of the Q - or T -system in the non-commutative setting. Finally, we express the weights of all the mutated path models in terms of these discrete quasi-Wronskian variables (Theorem 3.33), as manifestly positive Laurent monomials.

Sections 4 and 5 are applications respectively to the T -system, as an example of non-commutative Q -system, and to the quantum Q -system, in relation with the corresponding quantum cluster algebra [3]. In both cases, we compute the discrete quasi-Wronskian variables in terms of the initial data, and obtain Laurent positivity theorems for the solutions of the corresponding systems.

In Section 6, we give some further examples of non-commutative evolutions which have a Laurent positivity property. Section 7 considers the problem of generalizing of the Lindström–Gessel–Viennot theorem to the non-commutative setting. We conclude with some remarks about the possible generalization of cluster algebra relations and variables to the generic non-commutative setting.

2. Continued fraction solutions to Q -systems

2.1. The A_r Q -system

We consider the set of discrete evolution equations

$$R_{i,k+1}R_{i,k-1} = R_{i,k}^2 + R_{i-1,k}R_{i+1,k}, \quad 1 \leq i \leq r, \quad k \in \mathbb{Z} \tag{2.1}$$

with the boundary conditions $R_{0,k} = R_{r+1,k} = 1$ for all $k \in \mathbb{Z}$.

2.2. Initial data sets

For fixed α and n , the variable $R_{\alpha,n+1}$ is a rational function of $R_{\beta,n}$ with $|\alpha - \beta| \leq 1$ and $R_{\alpha,n-1}$. Therefore the recursion relation (2.1) has a solution for a given set of initial data, which must include the $2r$ variables $R(\mathbf{m}) = \{R_{\alpha,m_\alpha}, R_{\alpha,m_\alpha+1} : 1 \leq \alpha \leq r\}$ where the sequence $\mathbf{m} = (m_1, \dots, m_r)$ is such that $|m_i - m_{i+1}| \leq 1$. That is, initial data sets of the Q -system are in bijection with Motzkin paths of length $r - 1$.

Any variable $R_{\alpha,k}$ which is a solution of (2.1) can then be expressed as a function of the initial data set $R(\mathbf{m})$. Moreover, due to translational invariance of Eq. (2.1), the expression for $R_{\alpha,n}$ as a function of $R(\mathbf{m})$ is the *same* as that of $R_{\alpha,n+k}$ as a function of $R(k + \mathbf{m})$, with the

translated Motzkin path $k + \mathbf{m} = (k + m_1, \dots, k + m_r)$. We may therefore restrict ourselves to a fundamental domain of Motzkin paths modulo translations, $\mathcal{M} = \{\mathbf{m} \mid \text{Min}(m_i) = 0\}$.

In [21], we showed that the relations (2.1) are mutations in a coefficient-free cluster algebra, defined from the seed cluster variables (\mathbf{x}_0, B) , where

$$\mathbf{x}_0 = (R_{1,0}, \dots, R_{r,0}; R_{1,1}, \dots, R_{1,r}), \quad B = \begin{pmatrix} 0 & -C \\ C & 0 \end{pmatrix},$$

where C is the Cartan matrix of the Lie algebra A_r . This seed corresponds to the fundamental Motzkin path, $\mathbf{m} = (0, \dots, 0)$.

Not all mutations in the cluster algebra are of the form (2.1). However, all the variables $\{R_{\alpha,n} : 1 \leq \alpha \leq r, n \in \mathbb{Z}\}$ are contained in the subalgebra made up of the union of seeds parametrized by the set of Motzkin paths of length $r - 1$. Thus, we consider the corresponding sub-cluster algebra, restricting to the nodes of the Cayley tree that are labeled by Motzkin paths, and to the mutations that connect them.

We note that mutations in the sub-cluster algebra under consideration here have the following property: μ_i is given by one of the Q -system equations if and only if it maps $R(\mathbf{m})$ to $R(\mathbf{m}')$, where $m'_j = m_j \pm \delta_{i,j}$. That is, it maps a Motzkin path to another Motzkin path which differs from it in only one node.

2.3. Path partition functions

2.3.1. Positivity from weighted paths on graphs

In [7], we first eliminated from the Q -system (2.1) the variables $R_{i,k}$ $i > 1, k \in \mathbb{Z}$, in terms of the $R_{1,n}$, $n \in \mathbb{Z}$, by noting that the discrete Wronskians

$$W_{i,n} = \det_{1 \leq a,b \leq i} (R_{1,n+i+1-a-b}) \tag{2.2}$$

actually satisfy the same recursion relation (2.1), as a consequence of the Desnanot–Jacobi identity for the minors of any given matrix (also a particular case of the Plücker relations, see [7] for details). As moreover $W_{0,n} = 1$ and $W_{1,n} = R_{1,n}$, we conclude that $W_{i,n} = R_{i,n}$ for all $i \geq 0, n \in \mathbb{Z}$.

The Q -system therefore boils down to a single non-linear recursion relation for $R_{1,n}$, in the form $W_{r+1,n} = 1$. In Ref. [7], we explicitly solved this equation, by using its integrability. The latter implied the existence of a linear recursion relation for the $R_{1,n}$'s, with constant coefficients (independent of n , and expressible explicitly in terms of initial data). Exploiting this, we further showed that $R_{1,n}$ could be related to the partition functions of certain weighted path models on target graphs $\Gamma_{\mathbf{m}}$ (see definition in Appendix A), indexed by Motzkin paths \mathbf{m} of length $r - 1$, the weights being directly expressed as positive Laurent monomials of the corresponding initial data $R(\mathbf{m})$.


Finally, the quantity $R_{i,n}$, expressed as the determinant $W_{i,n}$ (2.2) was shown to also be related to path partition functions, namely that of collections of i strongly non-intersecting paths on the same target graphs $\Gamma_{\mathbf{m}}$, with the same weights. This followed from a slight generalization of the Lindström–Gessel–Viennot theorem [25,20].

Moreover, we obtained an explicit expression for the generating function of the variables $\{R_{1,n} : n \geq 0\}$ as the partition function of paths on the graph $\Gamma_{\mathbf{m}}$, in the form of an explicit finite continued fraction.

In this section, we present a similar but simpler description in terms of paths on another, more compact graph $G_{\mathbf{m}}$, with related weights. The equivalence of the two formulations is proved in Appendix A.

2.3.2. *The weighted graphs $G_{\mathbf{m}}$*

For fixed r , consider the weighted graph $G_{\mathbf{m}}$. The graph itself depends only on r . It has $r + 1$ vertices labeled $1, \dots, r + 1$ and oriented edges connecting vertex i to itself, to $i - 1$ and to $i + 1$. This is pictured below,



where each edge connecting two different vertices is to be considered as a doubly oriented edge. The map $\mathbf{m} \mapsto \Gamma_{\mathbf{m}} \mapsto G_{\mathbf{m}}$ is described in Appendix A.

To make the contact between the $G_{\mathbf{m}}$ path formulation and that on $\Gamma_{\mathbf{m}}$, we need a few definitions. We start from a collection of $2r + 1$ formal variables, $y(\mathbf{m}) = (y_1(\mathbf{m}), \dots, y_{2r+1}(\mathbf{m}))$. These will eventually be identified with the skeleton weights of the graph $\Gamma_{\mathbf{m}}$ (see Appendix A).

Definition 2.1. To $y(\mathbf{m})$ we associate bijectively the collection $\widehat{y}(\mathbf{m}) = (\widehat{y}_1(\mathbf{m}), \dots, \widehat{y}_{2r+1}(\mathbf{m}))$. This is defined according to the local structure of the Motzkin path \mathbf{m} as follows:

$$\widehat{y}_{2i} = ty_{2i}(ty_{2i+1})^{m_{i+1}-m_i},$$

$$\widehat{y}_{2i-1} = \begin{cases} ty_{2i-1} - \frac{y_{2i}}{y_{2i+1}}, & m_{i+1} = m_i - 1; \\ ty_{2i-1} + ty_{2i}, & m_{i+1} = m_i + 1; \\ ty_{2i-1}, & \text{otherwise.} \end{cases} \tag{2.3}$$

Example 2.2. The three basic examples are as follows:

- If $\mathbf{m} = \mathbf{m}_0 = (0, \dots, 0)$, the fundamental Motzkin path, we have $\widehat{y}_i = ty_i, i = 1, 2, \dots, 2r + 1$.
- If $\mathbf{m} = \mathbf{m}_1 = (0, 1, 2, \dots, r - 1)$, the strictly ascending Motzkin path, we have

$$\widehat{y}_{2i-1} = t(y_{2i-1} + y_{2i}), \quad \widehat{y}_{2i} = t^2y_{2i}y_{2i+1} \quad (1 \leq i \leq r - 1),$$

with $\widehat{y}_a = ty_a$ for $a \geq 2r - 1$.

- If $\mathbf{m} = \mathbf{m}_2 = (r - 1, r - 2, \dots, 0)$, the strictly descending Motzkin path,

$$\widehat{y}_{2i-1} = ty_{2i-1} - \frac{y_{2i}}{y_{2i+1}}, \quad \widehat{y}_{2i} = \frac{y_{2i}}{y_{2i+1}} \quad (1 \leq i \leq r - 1),$$

with $\widehat{y}_a = ty_a$ for $a \geq 2r - 1$.

We now attach weights to the edges of the graph $G_{\mathbf{m}}$. The edge connecting vertex i to vertex j carries a weight $w_{i,j} := w_{i,j}(\mathbf{m})$, where

$$w_{i,i+1} = 1, \quad w_{i,i} = \widehat{y}_{2i-1}(\mathbf{m}), \quad w_{i+1,i} = \widehat{y}_{2i}(\mathbf{m}),$$

with $\widehat{y}_i(\mathbf{m})$ as in (2.3).

2.3.3. Partition functions of paths on $G_{\mathbf{m}}$ and continued fractions


Each path p on the graph $G_{\mathbf{m}}$ from node i to node j carries a weight $wt(p)$ which is the product of the weights $w_{i,j}$ of the edges $i \rightarrow j$ traversed by the path. The partition function of paths from i to j is the sum of the weights of all paths $p \in \mathcal{P}_{i,j}(G_{\mathbf{m}})$ which start at vertex i and end at vertex j :

$$Z_{i,j}(G_{\mathbf{m}}) = \sum_{p \in \mathcal{P}_{i,j}(G_{\mathbf{m}})} wt(p).$$

Once expressed in terms of $\{y_i\}$, the weights of Definition 2.1 introduce terms with negative signs. The following lemma shows why this does not introduce minus signs in the generating function of paths.

Lemma 2.3. *The partition function $Z_{i,j}(G_{\mathbf{m}})$ as a power series in t , has coefficients which are positive Laurent polynomials in $\{y_i\}_i$.*

Proof. Let $I_{\pm} = \{i: m_i = m_{i-1} \pm 1\}$. The partition function for paths on $G_{\mathbf{m}}$ is the same as that on the graph $G'_{\mathbf{m}}$, where each loop at vertex i such that $i \in I_-$ is decomposed into two loops as follows:



The corresponding piece of $G'_{\mathbf{m}}$ is depicted on the right. In this case, $a = ty_{2i-1}$ and $b = y_{2i}/y_{2i+1}$. We call the two corresponding loops on the right-hand side ℓ_i^+ (with weight a) and ℓ_i^- (with weight $-b$). Note that the weight of ℓ_i^- is equal to $-w_{i+1,i}$, as shown in the picture.

We define an involution φ on paths $p \in \mathcal{P}_{i,j}(G'_m)$ as follows:

- $\varphi(p) = p$ if p does not pass through any ℓ_i^- ($i \in I_-$) nor contains any sequence of steps $i \rightarrow i + 1 \rightarrow i$ ($i \in I_-$).
- Otherwise, consider s , which is either the first step of p passing through some ℓ_i^- , or the first sequence of the form $i \rightarrow i + 1 \rightarrow i$ ($i \in I_-$) in p , whichever comes first. The path $\varphi(p)$ is obtained from p by replacing s by the other sequence (a step through ℓ_i^- by $i \rightarrow i + 1 \rightarrow i$ and vice versa).

Clearly, φ is an involution on $\mathcal{P}_{i,j}(G'_m)$. Its effect is to reverse the sign of the weight, $wt(\varphi(p)) = -wt(p)$ if $\varphi(p) \neq p$. There is therefore a pairwise cancellation of contributions of paths such that $p \neq \varphi(p)$. The terms which are left in the partition function are the fixed points $p = \varphi(p)$ of the involution. These have no step through any loop ℓ_i^- ($i \in I_-$) nor any sequence $i \rightarrow i + 1 \rightarrow i$ ($i \in I_-$).

The remaining weights are all positive Laurent polynomials of the y_i 's, and the lemma follows. \square

Given the graph G_m with weights $w_{i,j}(\mathbf{m})$, the partition function of paths from vertex 1 to itself can be written as a finite continued fraction as follows.

Definition 2.4. The Jacobi-type (finite) continued fraction is defined inductively as

$$J^{(r)}(x_1, \dots, x_{2r+1}) = \frac{1}{1 - x_1 - x_2 J^{(r-1)}(x_3, \dots, x_{2r+1})}, \quad J^{(1)}(x) = \frac{1}{1 - x}. \quad (2.4)$$

Lemma 2.5. *The generating function of weighted paths on G_m from and to the vertex 1 can be written as*

$$Z_{1,1}(G_m) = J^{(r)}(\widehat{y}(\mathbf{m})). \quad (2.5)$$

Proof. By induction. \square

We regard $J^{(r)}(\widehat{y}(\mathbf{m}))$ as a power series in t . As we have seen in Lemma 2.3, the coefficients of t^n are positive Laurent polynomials in $y(\mathbf{m})$.

Alternatively, the positivity of the generating function $Z_{1,1}(G_m)$ may be recovered by a simple rearrangement of the continued fraction $J^{(r)}(\widehat{y}(\mathbf{m}))$. Starting with the bottom term in the continued fraction and moving upwards, one iteratively rearranges the fraction for each $i \in I_-$ using the identity

$$a - \frac{b}{c} + \frac{\frac{b}{c}}{1 - c - u} = a + \frac{b + \frac{b}{c}u}{1 - c - u} \quad (2.6)$$

with $a = ty_{2i-1}$, $b = ty_{2i}$ and $c = ty_{2i+1}$. (Here, u denotes the already rearranged remainder of the fraction.) This eliminates any negative weights in the fraction, at the cost of a modification of the branching structure of the fraction (the fraction u appears in two places), and the introduction of new, positive weights coming from the term $\frac{b}{c}u$.

The fraction $J^{(r)}(\widehat{y}(m))$ can be re-expressed in a *canonical*, manifestly positive form as follows. Going through the fraction iteratively from bottom to top, whenever $i \in I_-$, apply the identity (2.6). Otherwise, if $m_i = m_{i-1} + 1$, apply the identity

$$a + b + \frac{bc}{1 - c - u} = a + \frac{b}{1 - \frac{c}{1-u}} \tag{2.7}$$

with $a = ty_{2i-1}$, $b = ty_{2i}$ and $c = ty_{2i+1}$. (Here, u is the already rearranged remainder of the fraction.)

The resulting rearranged form of $J^{(r)}(\widehat{y}(\mathbf{m}))$ has the property that all its coefficients are Laurent monomials of $y(\mathbf{m})$ with coefficient t .

We call this form canonical.

Definition 2.6. The canonical form of the fraction $J^{(r)}(\widehat{y}(\mathbf{m}))$ is the fraction resulting from the use of the rearrangements (2.6) and (2.7), such that the final fraction has coefficients which are all Laurent monomials of $y(\mathbf{m})$, with coefficient t .

Another important generating function is the partition function of paths from $1'$ to $1'$ on the graph $\Gamma_{\mathbf{m}}$ of Appendix A. It can be written as follows:

$$F_{\mathbf{m}}(t) = 1 + ty_1(\mathbf{m})J^{(r)}(\widehat{y}(\mathbf{m})). \tag{2.8}$$

Remark 2.7. The canonical form of $F_{\mathbf{m}}(t)$ itself (with the identity (2.7) applied also at the top level, with $a = 0$, $b = 1$ and $c = y_1(\mathbf{m})$) is related directly to the path interpretation of $F_{\mathbf{m}}(t)$ as generating weighted paths on the graph $\Gamma_{\mathbf{m}}$ (see Appendix A, Eq. (A.2)); while the y_i 's are the skeleton weights of $\Gamma_{\mathbf{m}}$, the redundant weights have been produced by the rearrangements (2.6).

Example 2.8. The fraction corresponding to the Motzkin path $(0, 1, 0)$ for $r = 3$ reads:

$$\begin{aligned} F_{0,1,0}(t) &= 1 + t \frac{y_1}{1 - t(y_1 + y_2) - t^2 \frac{y_2 y_3}{1 - ty_3 + \frac{y_4}{y_5} - \frac{\frac{y_4}{y_5}}{1 - ty_5 - t \frac{y_6}{1 - ty_7}}}} \\ &= \frac{1}{1 - t \frac{y_1}{1 - t \frac{y_2}{1 - t \frac{y_3}{1 - t \frac{\frac{y_4 y_6}{y_5}}{1 - t \frac{y_4 + \frac{y_5}{1 - ty_7}}{1 - ty_5 - t \frac{y_6}{1 - ty_7}}}}}}}} \end{aligned}$$

Note the occurrence of a new “redundant” weight $ty_4 y_6 / y_5$.

2.4. Mutations

Without loss of generality we now restrict ourselves to forward mutations of \mathbf{m} , $\mu_i: \mathbf{m} \mapsto \mathbf{m}'$, with $m'_j = m_j + \delta_{j,i}$. We also restrict ourselves to two special cases:

- Case (i): $\mu_i: (m_{i-1}, m_i, m_{i+1}) = (a, a, a + 1) \mapsto (a, a + 1, a + 1)$.
- Case (ii): $\mu_i: (m_{i-1}, m_i, m_{i-1}) = (a, a, a) \mapsto (a, a + 1, a)$.

These two cases are sufficient to obtain any Motzkin path, up to a global translation by an overall integer, from the fundamental Motzkin path $\mathbf{m}_0 = (0, 0, \dots, 0)$.

We also define the action of a mutation on $\mathbb{Z}(y)$:

Definition 2.9. Let $\mu_i(\mathbf{m}) = \mathbf{m}'$. Define $y' := y(\mathbf{m}')$ as a function of $y := y(\mathbf{m})$ as follows:

$$\text{Case (i): } \begin{cases} y'_{2i-1} = y_{2i-1} + y_{2i}; \\ y'_{2i} = y_{2i+1}y_{2i}/(y_{2i-1} + y_{2i}); \\ y'_{2i+1} = y_{2i+1}y_{2i-1}/(y_{2i-1} + y_{2i}); \end{cases} \tag{2.9}$$

$$\text{Case (ii): } \begin{cases} y'_{2i-1} = y_{2i-1} + y_{2i}; \\ y'_{2i} = y_{2i+1}y_{2i}/(y_{2i-1} + y_{2i}); \\ y'_{2i+1} = y_{2i+1}y_{2i-1}/(y_{2i-1} + y_{2i}); \\ y'_{2i+2} = y_{2i+2}y_{2i-1}/(y_{2i-1} + y_{2i}) \end{cases} \tag{2.10}$$

while all other weights are left unchanged,

We then have the natural definitions of $\widehat{y}(\mathbf{m}')$ in terms of $y(\mathbf{m}')$, using Eqs. (2.1).

Theorem 2.10. Under the mutation μ_i , the function $J^{(r)}(\widehat{y}(\mathbf{m}))$ is transformed as

$$J^{(r)}(\widehat{y}(\mathbf{m})) = \begin{cases} J^{(r)}(\widehat{y}(\mathbf{m}')), & i > 1; \\ 1 + ty_1(\mathbf{m}')J^{(r)}(\widehat{y}(\mathbf{m}')), & i = 1. \end{cases} \tag{2.11}$$

This has an immediate corollary.

Corollary 2.11. Under the same mutation μ_i , $F_{\mathbf{m}}(t)$ is transformed as:

$$F_{\mathbf{m}}(t) = \begin{cases} F_{\mathbf{m}'}(t), & i > 1; \\ 1 + ty_1(\mathbf{m})F_{\mathbf{m}'}(t), & i = 1. \end{cases} \tag{2.12}$$

To prove the theorem, we use two rearrangement lemmas in addition to Eqs. (2.6) and (2.7):

Lemma 2.12. For all a, b we have

$$1 + \frac{a}{1 - a - b} = \frac{1}{1 - \frac{a}{1-b}}. \tag{2.13}$$

Lemma 2.13. For all a, b, c, d , with $a + b$ invertible, we have

$$a + \frac{b}{1 - \frac{c}{1-d}} = \frac{a'}{1 - \frac{b'}{1-c'-d}} \tag{2.14}$$

where a', b', c' are defined by

$$a' = a + b, \quad b' = \frac{bc}{a + b}, \quad c' = \frac{ac}{a + b}. \tag{2.15}$$

We now turn to the proof of Theorem 2.10.

Proof. Let $J_i(\mathbf{m})$ be defined inductively as follows:

$$J_i(\mathbf{m}) = \widehat{y}_{2i-1} + \frac{\widehat{y}_{2i}}{1 - J_{i+1}(\mathbf{m})} \quad (i \geq 0), \quad J_{r+1}(\mathbf{m}) = \widehat{y}_{2r+1}(\mathbf{m}),$$

with the convention that $\widehat{y}_{-1} = 0$ and $\widehat{y}_0 = 1$. Note that $J_0(\mathbf{m}) = J^{(r)}(\widehat{y}(\mathbf{m}))$. We refer below to the “level” i of the fraction to denote the part of the fraction $J^{(r)}$ where J_i appears. Let

$$\begin{aligned} a &= ty_{2i-3}(\mathbf{m}), & b &= ty_{2i-2}(\mathbf{m}), & c &= ty_{2i-1}(\mathbf{m}), & d &= ty_{2i}(\mathbf{m}), \\ e &= ty_{2i+1}(\mathbf{m}), & f &= ty_{2i+2}(\mathbf{m}), & u &= \frac{1}{1 - J_{i+2}(\mathbf{m})} \end{aligned}$$

together with their primed counterparts, which are functions of \mathbf{m}' .

We now consider the effects of the mutations of Definition 2.9 on $J^{(r)}(\widehat{y}(\mathbf{m}))$.

- **Case (i):** We first transform the Jacobi-like form into the local canonical form for $J_{i-1}(\mathbf{m})$ by applying (2.7) at the level i , namely:


$$J_{i-1}(\mathbf{m}) = a + \frac{b}{1 - c - d - \frac{de}{1 - e - fu}} = a + \frac{\widehat{b}}{1 - c - \frac{d}{1 - \frac{e}{1 - fu}}}.$$

Next, we apply Lemma 2.13 to the second level of this fraction, namely:

$$J_{i-1}(\mathbf{m}) = a + \frac{b}{1 - \frac{c'}{1 - \frac{d'}{1 - e' - fu}}}, \quad \text{with } c' = c + d, \quad d' = de/c', \quad \text{and } e' = ce/c'.$$

Finally, we put this back into Jacobi-like form, by applying (2.7) backwards at the level $i - 1$. The complete transformation reads:

$$J_{i-1}(\mathbf{m}) = a + \frac{b}{1 - c - d - \frac{de}{1 - e - fu}} = J_{i-1}(\mathbf{m}') = a + b + \frac{bc'}{1 - c' - \frac{d'}{1 - e' - fu}}.$$



- **Case (ii):** We apply directly Lemma 2.13 to the level i , namely:


$$J_{i-1}(\mathbf{m}) = a + \frac{b}{1 - c - \frac{d}{1 - e - fu}} = a + \frac{b}{1 - \frac{c'}{1 - \frac{d' + \frac{d'f'}{e'f'u}}{1 - e' - f'u}}}, \quad \text{with} \quad \begin{cases} c' = c + d; \\ d' = de/c'; \\ e' = ce/c'; \\ f' = cf/c'. \end{cases}$$

We bring this back to Jacobi-like form by applying first (2.6) backward on its first level, and then (2.7) backward on its level zero, namely:

$$J_{i-1}(\mathbf{m}) = a + \frac{b}{1 - \frac{c'}{1 + \frac{d'}{e' - \frac{d'}{1 - e' - f'u}}}} = a + b + \frac{bc'}{1 - c' + \frac{d'}{e'} - \frac{d'}{1 - e' - f'u}}.$$

The complete transformation reads:

$$J_{i-1}(\mathbf{m}) = a + \frac{b}{1 - c - \frac{d}{1 - e - fu}} = J_{i-1}(\mathbf{m}') = a + b + \frac{bc'}{1 - c' + \frac{d'}{e'} - \frac{d'}{1 - e' - f'u}}.$$



Note that for $i > 1$, we always have $J_0(\mathbf{m}') = J_0(\mathbf{m})$, or equivalently $F_{\mathbf{m}}(t) = F_{\mathbf{m}'}(t)$. When $i = 1$, we have $a = 0, b = 1$ and in all cases $c' = y_1(\mathbf{m}')$, so the final formula amounts to $J_0(\mathbf{m}) = 1 + tJ_0(\mathbf{m}')y_1(\mathbf{m}')$, or equivalently $F_{\mathbf{m}}(t) = 1 + tF_{\mathbf{m}'}(t)y_1(\mathbf{m})$.

This proves the theorem and its corollary. □

2.4.1. *Explicit expression for the weights $y_i(\mathbf{m})$*

We can now identify the definitions of the mutations on the initial data variables $\mu_i: R(\mathbf{m}) \mapsto R(\mathbf{m}')$ and the action on weights $\mu_i: y(\mathbf{m}) \rightarrow y(\mathbf{m}')$.

Corollary 2.11 may be used as a definition of the series $F_{\mathbf{m}}(t)$, by using induction over mutations, once we specify the initial condition. We can do so by giving the value of $F_{\mathbf{m}_0}(t)$ as a function of $R(\mathbf{m}_0)$, the variables corresponding to the fundamental Motzkin path. This will define the initial values $y_i(\mathbf{m}_0)$ of the weights. Using Definition 2.9, we can then determine $y(\mathbf{m})$ for any \mathbf{m} inductively.

In Ref. [7], we used the integrability of the Q -system (2.1) to show that the generating function

$$F_0(t) = \sum_{n \geq 0} t^n R_{1,n} / R_{1,0}$$

is expressed explicitly in terms of the variables $R(\mathbf{m}_0)$ as the finite continued fraction:

$$F_0(t) = 1 + t \frac{y_1}{1 - ty_1 - t \frac{y_2}{1 - ty_3 - t \frac{y_4}{\dots \frac{y_{2r}}{1 - ty_{2r-1} - t \frac{y_{2r+1}}{y_{2r+1}}}}}}$$

where

$$y_{2i-1} = \frac{R_{i,1}R_{i-1,0}}{R_{i,0}R_{i-1,1}}, \quad y_{2i} = \frac{R_{i+1,0}R_{i-1,1}}{R_{i,0}R_{i,1}}. \tag{2.16}$$

Comparing this expression with that for $F_{\mathbf{m}}(t)$ (2.8), one sees that the choice of weights $y_i(\mathbf{m}_0) = y_i$ as in (2.16) gives

$$F_{\mathbf{m}_0}(t) = F_0(t).$$

We can now derive explicit expressions for the variables $y(\mathbf{m})$ for any \mathbf{m} using Definition 2.9.

Theorem 2.14. *For any Motzkin path $\mathbf{m} = (m_1, m_2, \dots, m_r)$, the weights $y_i(\mathbf{m})$ defined by (2.9)–(2.10) with the initial conditions $y_i(\mathbf{m}_0)$ as in (2.16) are expressed in terms of the variables $R(\mathbf{m})$ as:*

$$y_{2i-1}(\mathbf{m}) = \frac{c_{i,m_i+1}}{c_{i-1,m_{i-1}+1}}, \quad y_{2i}(\mathbf{m}) = d_{i,m_i+1}L_{i,i+1}(L_{i,i-1})^{-1}, \tag{2.17}$$

where

$$L_{i,i+\varepsilon} = \begin{cases} \frac{c_{i+\varepsilon,m_i+\varepsilon+1}}{c_{i+\varepsilon,m_i+1}}, & \text{if } m_i = m_{i+\varepsilon} + \varepsilon; \\ 1, & \text{otherwise,} \end{cases}$$

and

$$c_{i,m} = \frac{R_{i,m}}{R_{i,m-1}}, \quad d_{i,m} = \frac{R_{i+1,m}R_{i-1,m-1}}{R_{i,m}R_{i,m-1}}.$$

Proof. By induction under mutations, from \mathbf{m}_0 to any Motzkin path \mathbf{m} . We explicitly check the relations (2.9)–(2.10) on the expressions (2.17), by use of the Q -system. \square

Corollary 2.15. *The series $F_{\mathbf{m}}(t)$ is the following generating function for solutions of the Q -system:*

$$F_{\mathbf{m}}(t) = \sum_{n \geq 0} t^n R_{1,n+m_1} / R_{1,m_1}. \tag{2.18}$$

Proof. Whenever a mutation $\mu_i: \mathbf{m} \mapsto \mathbf{m}'$, $i > 1$ is applied, the function $F_{\mathbf{m}}(t) = F_{\mathbf{m}'}(t)$ is preserved, hence its series coefficient remain the same. When $i = 1$ however, we have the transformation $F_{\mathbf{m}}(t) = 1 + ty_1(\mathbf{m})F_{\mathbf{m}'}(t) = 1 + t \frac{R_{1,m_1+1}}{R_{1,m_1}}F_{\mathbf{m}'}(t)$, and therefore the coefficients of the series $F_{\mathbf{m}'}(t)$ change according to (2.18). \square

3. Non-commutative weights and evolutions

In this section, we introduce a non-commutative version of the path partition functions and weight evolutions which appear in the A_r Q -system. Most of the key ideas of the previous section generalize to the non-commutative setting because paths are themselves non-commutative objects, and respect ordering of non-commutative weights.

3.1. Paths with non-commutative weights and non-commutative continued fractions

Let $\{y_i^{\pm 1}; 1 \leq i \leq 2r + 1\}$ be non-commutative variables and let \mathcal{A} be the skew field of rational functions in $\{y_1^{\pm 1}, \dots, y_{2r+1}^{\pm 1}\}$. Let $G_{\mathbf{m}}$ denote the graph used in Section 2.3.2. We associate to each edge of $G_{\mathbf{m}}$ a non-commutative weight, which is an element in $\mathcal{A}[t]$.

A path on a graph with non-commutative weights is defined to have a weight equal to the product of weights of the edges traversed by the path, written *in the order in which the steps are taken*, from left to right. For example, if the path is of the form $p = (i_1, i_2, \dots, i_n)$, then its weight is $\mathbf{wt}(p) = \mathbf{w}_{i_1, i_2} \mathbf{w}_{i_2, i_3} \cdots \mathbf{w}_{i_{n-1}, i_n}$, where $\mathbf{w}_{i, j}$ is the weight of edge from vertex i to vertex j .

3.1.1. Non-commutative weights

For each Motzkin path in the fundamental domain, $\mathbf{m} \in \mathcal{M}$, let $\mathbf{y}(\mathbf{m}) \equiv (y_1(\mathbf{m}), \dots, y_{2r+1}(\mathbf{m})) \in \mathcal{A}^{2r+1}$ be a collection of $2r + 1$ non-commutative variables. Let $I_{\pm} = \{i: m_i = m_{i-1} \pm 1\}$ as before. To $\mathbf{y}(\mathbf{m})$ we associate the variables $\widehat{\mathbf{y}}(\mathbf{m}) = (\widehat{y}_1(\mathbf{m}), \dots, \widehat{y}_{2r+1}(\mathbf{m})) \in \mathcal{A}[t]^{2r+1}$ defined as follows:

$$\begin{aligned} \widehat{y}_{2i-1} &= t(y_{2i-1} + y_{2i}), & \widehat{y}_{2i} &= t^2 y_{2i+1} y_{2i}, & \text{if } i \in I_+; \\ \widehat{y}_{2i-1} &= t y_{2i-1} - y_{2i+1}^{-1} y_{2i}, & \widehat{y}_{2i} &= y_{2i+1}^{-1} y_{2i}, & \text{if } i \in I_-; \\ \widehat{y}_{2i-1} &= t y_{2i-1}, & \widehat{y}_{2i} &= t y_{2i}, & \text{otherwise.} \end{aligned} \tag{3.1}$$

Here, we have omitted the reference to the Motzkin path \mathbf{m} . Note that these are non-commutative versions of the variables \widehat{y} defined in Eq. (2.3).

Finally, we define the weight on the edge from i to j of $G_{\mathbf{m}}$ to be $\mathbf{w}_{i, j}$, where

$$\mathbf{w}_{i, i} = \widehat{y}_{2i-1}(\mathbf{m}), \quad \mathbf{w}_{i, i+1} = 1, \quad \mathbf{w}_{i+1, i} = \widehat{y}_{2i}(\mathbf{m}).$$

3.1.2. Non-commutative partition functions

Let $\mathcal{P}_{a, b}$ be the set of paths on $G_{\mathbf{m}}$ from vertex a to vertex b . Each path $p \in \mathcal{P}_{a, b}$ has a non-commutative weight $\mathbf{wt}(p) \in \mathcal{A}$ defined like in Section 2.3.3 as the product of step weights, in the order in which the steps are taken. The partition function, or generating function, of paths from a to b is defined to be

$$\mathbf{Z}(G_{\mathbf{m}})_{a, b} = \sum_{p \in \mathcal{P}_{a, b}} \mathbf{wt}(p). \tag{3.2}$$

3.1.3. Non-commutative continued fractions

Definition 3.1. For any $\mathbf{x} = (\mathbf{x}_1, \dots, \mathbf{x}_{2r+1})$ with $\mathbf{x}_i \in \mathcal{A}[t]$, we define the non-commutative Jacobi-like finite continued fraction $\mathbf{J}^{(r)}(\mathbf{x})$ inductively by $\mathbf{J}^{(1)}(\mathbf{x}_1) = (\mathbf{1} - \mathbf{x}_1)^{-1}$ and

$$\mathbf{J}^{(k)}(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_{2k+1}) = (\mathbf{1} - \mathbf{x}_1 - \mathbf{J}^{(k-1)}(\mathbf{x}_3, \mathbf{x}_4, \dots, \mathbf{x}_{2k+1})\mathbf{x}_2)^{-1}. \tag{3.3}$$

As in the commutative case, we may consider such a fraction as an element of $\mathcal{A}[[t]]$, that is, as a power series in t .

It is clear that $\mathbf{Z}(\mathbf{m})_{1,1} = \mathbf{J}^{(r)}(\widehat{\mathbf{y}}(\mathbf{m}))$ in terms of the weights (3.1). We also define the generating function $\mathbf{F}_\mathbf{m}(t)$, in analogy with the commutative case, to be

$$\mathbf{F}_\mathbf{m}(t) = 1 + t\mathbf{J}^{(r)}(\widehat{\mathbf{y}}_1(\mathbf{m}), \dots, \widehat{\mathbf{y}}_{2r+1}(\mathbf{m}))\mathbf{y}_1(\mathbf{m}). \tag{3.4}$$

Lemma 3.2. *The coefficient of t^n in the power series expansion of $\mathbf{F}_\mathbf{m}(t)$ is a Laurent polynomial in $\mathbf{y}(\mathbf{m})$ with non-negative integer coefficients.*

Proof. The proof is identical to the proof in the commutative case (cf. Lemma 2.3), since weights respect the order of steps in the paths. Suppose $i \in I_-$, so that $\mathbf{w}_{i,i}$ has a negative term. Decompose the loop $i \rightarrow i$ with weight $\mathbf{w}_{ii} = \widehat{\mathbf{y}}_{2i-1}$ into two separate loops from i to i , one, l_i^+ , with weight $t\mathbf{y}_{2i-1}$ and the other, l_i^- , with weight $-\mathbf{y}_{2i+1}^{-1}\mathbf{y}_{2i} = -\mathbf{w}_{i+1,i}$. Using the involution φ defined in Lemma 2.3 on paths on the resulting graph $G'_\mathbf{m}$, we again conclude that we have a cancellation of all paths $p \neq \varphi(p)$ in the partition function. \square

An alternative proof of Lemma 3.2 uses the structure of non-commutative continued fractions. One may simply rearrange the continued fraction expression (3.4) (from bottom to top) whenever $i \in I_-$, by use the following non-commutative version of the identity (2.6). For any $\mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{u} \in \mathcal{A}$:

$$\mathbf{a} - \mathbf{c}^{-1}\mathbf{b} + (\mathbf{1} - \mathbf{c} - \mathbf{u})^{-1}\mathbf{c}^{-1}\mathbf{b} = \mathbf{a} + (\mathbf{1} - \mathbf{c} - \mathbf{u})^{-1}(\mathbf{b} + \mathbf{u}\mathbf{c}^{-1}\mathbf{b}). \tag{3.5}$$

Once again, this is done at the cost of introducing multiple branching of the new fraction, since \mathbf{u} now appears in two places. Moreover, new “redundant” weights appear, from the new term $\mathbf{u}\mathbf{c}^{-1}\mathbf{b}$. These are, however, positive Laurent monomials in $\mathbf{y}(\mathbf{m})$.

For use below, we note that we also have an analogue of the identity (2.7): For any $\mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{u} \in \mathcal{A}$,

$$\mathbf{a} + \mathbf{b} + (\mathbf{1} - \mathbf{c} - \mathbf{u})^{-1}\mathbf{c}\mathbf{b} = \mathbf{a} + (\mathbf{1} - (\mathbf{1} - \mathbf{u})^{-1}\mathbf{c})^{-1}\mathbf{b}. \tag{3.6}$$

3.1.4. Canonical form of continued fractions

As in the commutative case, the fraction $\mathbf{F}_\mathbf{m}(t)$ can be rearranged into a “canonical” form, a (finite, multiply branching) continued fraction whose coefficients are all Laurent monomials in \mathbf{y} , with coefficients equal to t .

This is done as follows. From the bottom to the top level of the fraction ($i = r, r - 1, \dots, 1$), we rearrange the fraction whenever $m_i \neq m_{i-1}$. Let $(\mathbf{a}, \mathbf{b}, \mathbf{c}) = t(\mathbf{y}_{2i-1}, \mathbf{y}_{2i}, \mathbf{y}_{2i+1})$ and use (3.5) if $i \in I_-$, (3.6) if $i \in I_+$, to rearrange the fraction. This includes a last “level zero” step of the continued fraction as well (case $i = 0$), which corresponds to the conventions $\mathbf{y}_{-1} = 0$ and $\mathbf{y}_0 = 1$.

The resulting “canonical” continued fraction is interpreted as the generating function for paths from and to the root vertex on the graph $\Gamma_{\mathbf{m}}$ (see Appendix A), but now with non-commutative skeleton weights $t\mathbf{y}_i(\mathbf{m})$.

In particular, the non-commutative “redundant” weights $\mathbf{w}_{i,j}$, $i - j > 1$, generalizing (A.1) now read, as a consequence of iterations of (3.5):

$$\mathbf{w}_{i,j} = t \left(\prod_{a=i-1}^{j+1} \mathbf{y}_{2a} \mathbf{y}_{2a-1}^{-1} \right) \mathbf{y}_{2j} \tag{3.7}$$

where the product is taken in decreasing order, $a = i - 1, i - 2, \dots, j + 1$.

Example 3.3. In the case of the fundamental Motzkin path $\mathbf{m}_0 = (0, 0, \dots, 0)$, we have

$$\mathbf{F}_{\mathbf{m}_0}(t) = 1 + t\mathbf{J}^{(r)}(t\mathbf{y}_1, t\mathbf{y}_2, \dots, t\mathbf{y}_{2r+1})\mathbf{y}_1 \tag{3.8}$$

with $\mathbf{y}_i \equiv \mathbf{y}_i(\mathbf{m}_0)$. The canonical form is obtained by applying (3.6) at level zero, with $\mathbf{a} = 0$, $\mathbf{b} = 1$ and $\mathbf{c} = t\mathbf{y}_1$, with the result:

$$\begin{aligned} \mathbf{F}_{\mathbf{m}_0}(t) = & (1 - t(1 - t(1 - t\mathbf{y}_3 - t(\dots(1 - t\mathbf{y}_{2r-1} \\ & - t(1 - t\mathbf{y}_{2r+1})^{-1}\mathbf{y}_{2r})^{-1} \dots)^{-1}\mathbf{y}_4)^{-1}\mathbf{y}_2)^{-1}\mathbf{y}_1)^{-1}. \end{aligned} \tag{3.9}$$

This is the generating function for paths on the graph $\Gamma_{\mathbf{m}_0}$ of Example A.3, from vertex $1'$ to itself, with the non-commutative weights $\mathbf{w}_{i,i'} = \mathbf{w}_{i-1,i} = 1$, $\mathbf{w}_{i',i} = t\mathbf{y}_{2i-1}$, $\mathbf{w}_{i,i-1} = t\mathbf{y}_{2i-2}$ for $i \geq 2$ and $\mathbf{w}_{1',1} = 1$, $\mathbf{w}_{1,1'} = t\mathbf{y}_1$.

Example 3.4. In the case $\mathbf{m} = \mathbf{m}_1 = (0, 1, 2, \dots, r - 1)$, we have $m_i = m_{i-1} + 1$ for all i . The corresponding continued fraction reads, with $\mathbf{y}_i \equiv \mathbf{y}_i(\mathbf{m}_1)$:

$$\begin{aligned} \mathbf{F}_{\mathbf{m}_1}(t) = & 1 + t(1 - t\mathbf{y}_1 - t\mathbf{y}_2 - t^2(\dots(1 - t\mathbf{y}_{2r-1} \\ & - t\mathbf{y}_{2r} - t^2(1 - t\mathbf{y}_{2r+1})^{-1}\mathbf{y}_{2r+1}\mathbf{y}_{2r})^{-1} \dots)^{-1}\mathbf{y}_2\mathbf{y}_1)\mathbf{y}_1 \\ = & (1 - t(1 - t(\dots(1 - t(1 - t\mathbf{y}_{2r+1})^{-1}\mathbf{y}_{2r})^{-1} \dots)^{-1}\mathbf{y}_2)^{-1}\mathbf{y}_1)^{-1} \end{aligned} \tag{3.10}$$

where the first line is the Jacobi-like expression (3.4) and the second line has the canonical form.

The second identity in Example 3.4 is the natural non-commutative generalization of the Stieltjes continued fraction expansion. In the following, we will refer to the case $\mathbf{m} = \mathbf{m}_1$ as the “Stieltjes point”.

The graph $\Gamma_{\mathbf{m}_1}$ associated to this example is a chain of $2r + 2$ vertices labeled $1, 2, \dots, 2r + 2$ (see Appendix A), with weights $\mathbf{w}_{i,i+1} = 1$ and $\mathbf{w}_{i+1,i} = t\mathbf{y}_i$, $i = 1, 2, \dots, 2r + 1$.

Example 3.5. In the case $r = 3$ and $\mathbf{m} = (0, 1, 0)$, we have, with $\mathbf{y}_i \equiv \mathbf{y}_i(0, 1, 0)$:

$$\begin{aligned} \mathbf{F}_{(0,1,0)}(t) &= 1 + t(1 - t(\mathbf{y}_1 + \mathbf{y}_2)) - t^2(1 - t\mathbf{y}_3 + \mathbf{y}_5^{-1}\mathbf{y}_4 \\ &\quad - (1 - t\mathbf{y}_5 - t(1 - t\mathbf{y}_7)^{-1}\mathbf{y}_6)^{-1}\mathbf{y}_5^{-1}\mathbf{y}_4)^{-1}\mathbf{y}_3\mathbf{y}_2)\mathbf{y}_1 \\ &= (1 - t(1 - t(1 - t(1 - t(1 - t\mathbf{y}_5 - t(1 - t\mathbf{y}_7)^{-1}\mathbf{y}_6)^{-1}\mathbf{y}_5)^{-1}\mathbf{y}_4)^{-1}\mathbf{y}_3) \\ &\quad \times (\mathbf{y}_4 + (1 - t\mathbf{y}_7)^{-1}\mathbf{y}_6\mathbf{y}_5^{-1}\mathbf{y}_4))^{-1}\mathbf{y}_3)^{-1}\mathbf{y}_2)^{-1}\mathbf{y}_1)^{-1} \end{aligned}$$

where the first expression is the Jacobi-like form (3.4), and the second is the canonical form, obtained by first applying (3.5) with $\mathbf{a} = t\mathbf{y}_3$, $\mathbf{b} = t\mathbf{y}_4$ and $\mathbf{c} = t\mathbf{y}_5$ while $\mathbf{u} = 1 - t\mathbf{y}_5 - t(1 - t\mathbf{y}_7)^{-1}\mathbf{y}_6$, then (3.6) with $\mathbf{a} = t\mathbf{y}_1$, $\mathbf{b} = t\mathbf{y}_2$, $\mathbf{c} = t\mathbf{y}_3$, and finally (3.6) at “level zero”, with $\mathbf{a} = 0$, $\mathbf{b} = 1$ and $\mathbf{c} = t\mathbf{y}_1$. We see the emergence of the “redundant” weight $t\mathbf{y}_6\mathbf{y}_5^{-1}\mathbf{y}_4$. See the corresponding graph $\Gamma_{(0,1,0)}$ in Appendix A.

3.2. Mutations

In the non-commutative case, we do not have an evolution equation for cluster variables. Instead, we start by defining mutations of the weights \mathbf{y} . We will then show that these mutations have the same effect on the generating functions $F_{\mathbf{m}}(t)$, considered as continued fractions, as the non-commutative analogues of the rearrangement Lemmas 2.12 and 2.13.

We consider the set of Motzkin paths $\mathbf{m} \in \mathcal{M}$. We use the same notion of mutations on Motzkin paths as in the commutative case. Recall that mutations act on Motzkin paths as follows: $\mu_i(\mathbf{m}) = \mathbf{m}'$ where \mathbf{m}' differs from \mathbf{m} only in that $m'_i = m_i + 1$. In order to obtain any Motzkin path $\mathbf{m} \in \mathcal{M}$ via a sequence of mutations, we need only consider the two types of mutations described in cases (i) and (ii) of Section 2.4.

3.2.1. Mutations of weights

We now define the action of mutations on \mathcal{A} , as follows.

Definition 3.6. Let $\mu_i(\mathbf{m}) = \mathbf{m}'$. Define $\mathbf{y}' := \mathbf{y}(\mathbf{m}')$ as a function of $\mathbf{y} := \mathbf{y}(\mathbf{m})$ as follows:

$$\text{Case (i): } \begin{cases} \mathbf{y}'_{2i-1} = \mathbf{y}_{2i-1} + \mathbf{y}_{2i}; \\ \mathbf{y}'_{2i} = \mathbf{y}_{2i+1}\mathbf{y}_{2i}(\mathbf{y}_{2i-1} + \mathbf{y}_{2i})^{-1}; \\ \mathbf{y}'_{2i+1} = \mathbf{y}_{2i+1}\mathbf{y}_{2i-1}(\mathbf{y}_{2i-1} + \mathbf{y}_{2i})^{-1}; \end{cases} \quad (3.11)$$

$$\text{Case (ii): } \begin{cases} \mathbf{y}'_{2i-1} = \mathbf{y}_{2i-1} + \mathbf{y}_{2i}; \\ \mathbf{y}'_{2i} = \mathbf{y}_{2i+1}\mathbf{y}_{2i}(\mathbf{y}_{2i-1} + \mathbf{y}_{2i})^{-1}; \\ \mathbf{y}'_{2i+1} = \mathbf{y}_{2i+1}\mathbf{y}_{2i-1}(\mathbf{y}_{2i-1} + \mathbf{y}_{2i})^{-1}; \\ \mathbf{y}'_{2i+2} = \mathbf{y}_{2i+2}\mathbf{y}_{2i-1}(\mathbf{y}_{2i-1} + \mathbf{y}_{2i})^{-1} \end{cases} \quad (3.12)$$

with all other $\mathbf{y}'_j = \mathbf{y}_j$ unchanged.

Our main goal in this section is the proof of Theorem 3.10, concerning the action of mutations on the generating functions $F_{\mathbf{m}}(t)$. Towards this end, we prove some non-commutative analogues of the continued fraction rearrangement lemmas of Section 2.

3.2.2. *Mutations as non-commutative rearrangements*

In addition to the two identities (3.5) and (3.6), we have the two following local rearrangement lemmas for non-commutative continued fractions.

Lemma 3.7. *For all $\mathbf{a}, \mathbf{b} \in \mathcal{A}$ we have*

$$1 + (1 - \mathbf{a} - \mathbf{b})^{-1} \mathbf{a} = (1 - (1 - \mathbf{b})^{-1} \mathbf{a})^{-1}. \tag{3.13}$$

Proof. By direct calculation. This is the “level zero” application of identity (3.6). \square

Lemma 3.8. *For all $\mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{d} \in \mathcal{A}$, with $\mathbf{a} + \mathbf{b}$ invertible, we have*

$$\mathbf{a} + (1 - (1 - \mathbf{d})^{-1} \mathbf{c})^{-1} \mathbf{b} = (1 - \mathbf{d} - (1 - \mathbf{c}')^{-1} \mathbf{b}')^{-1} \mathbf{a}' \tag{3.14}$$

where $\mathbf{a}', \mathbf{b}', \mathbf{c}'$ are defined by

$$\mathbf{a}' = \mathbf{a} + \mathbf{b}, \quad \mathbf{c}' = \mathbf{c} \mathbf{b} (\mathbf{a} + \mathbf{b})^{-1}, \quad \mathbf{b}' = \mathbf{c} \mathbf{a} (\mathbf{a} + \mathbf{b})^{-1}. \tag{3.15}$$

Proof. By direct calculation. \square

Together with the identities (3.5)–(3.6), these two lemmas may be used to rearrange the non-commutative continued fraction expression of $\mathbf{F}_{\mathbf{m}}(t) \mapsto \mathbf{F}_{\mathbf{m}'}(t)$ (3.4).

Define the non-commutative version of the fractions J_i as in the proof of Lemma 2.13.

Definition 3.9. The non-commutative finite continued fraction $\mathbf{J}_i(\mathbf{m})$ is defined inductively as follows:

$$\mathbf{J}_i(\mathbf{m}) = \widehat{\mathbf{y}}_{2i-1} + (1 - \mathbf{J}_{i+1}(\mathbf{m}))^{-1} \widehat{\mathbf{y}}_{2i}, \quad \mathbf{J}_{r+1}(\mathbf{m}) = \widehat{\mathbf{y}}_{2r+1}(\mathbf{m}).$$

Note that $\mathbf{J}_0(\mathbf{m}) = \mathbf{J}^{(r)}(\widehat{\mathbf{y}}(\mathbf{m}))$, and $\mathbf{F}_{\mathbf{m}}(t) = 1 + t \mathbf{J}_0(\mathbf{m}) \mathbf{y}_1$, with the convention that $\widehat{\mathbf{y}}_{-1} = 0$ and $\widehat{\mathbf{y}}_0 = 1$.

Let us fix i , and define:

$$\begin{aligned} \mathbf{a} &= t \mathbf{y}_{2i-3}, & \mathbf{b} &= t \mathbf{y}_{2i-2}, & \mathbf{c} &= t \mathbf{y}_{2i-1}, & \mathbf{d} &= t \mathbf{y}_{2i}, \\ \mathbf{e} &= t \mathbf{y}_{2i+1}, & \mathbf{f} &= t \mathbf{y}_{2i+2}, & \mathbf{u} &= (1 - \mathbf{J}_{i+2}(\mathbf{m}))^{-1}, \end{aligned}$$

where $\mathbf{y}_i := \mathbf{y}_i(\mathbf{m})$. Similarly, we use the same notation for the primed counterparts $\mathbf{y}'_i := \mathbf{y}_i(\mathbf{m}')$.

Starting from $\mathbf{F}_{\mathbf{m}}(t)$ in Jacobi-like form (3.4), we apply fraction rearrangements on the part \mathbf{J}_{i-1} of $\mathbf{F}_{\mathbf{m}}(t)$.

- **Case (i):** We first transform the Jacobi-like form into the local canonical form for $\mathbf{J}_{i-1}(\mathbf{m})$ by applying (3.6) at the second level, namely:


$$\begin{aligned} \mathbf{J}_{i-1}(\mathbf{m}) &= \mathbf{a} + (1 - \mathbf{c} - \mathbf{d} - (1 - \mathbf{e} - \mathbf{u} \mathbf{f})^{-1} \mathbf{e} \mathbf{d})^{-1} \mathbf{b} \\ &= \mathbf{a} + (1 - \mathbf{c} - (1 - (1 - \mathbf{u} \mathbf{f})^{-1} \mathbf{e})^{-1} \mathbf{d})^{-1} \mathbf{b}. \end{aligned}$$

Next, we apply Lemma 3.8 to the second level, namely:

$$\mathbf{J}_{i-1}(\mathbf{m}) = \mathbf{a} + (1 - (1 - (1 - \mathbf{e}' - \mathbf{uf})^{-1} \mathbf{d}')^{-1} \mathbf{c}')^{-1} \mathbf{b}$$

with $\mathbf{c}' = \mathbf{c} + \mathbf{d}$, $\mathbf{d}' = \mathbf{ed}(\mathbf{c}')^{-1}$ and $\mathbf{e}' = \mathbf{ec}(\mathbf{c}')^{-1}$. Finally, we put this back in Jacobi-like form, by applying (3.6) backwards to the first level, namely:

$$\mathbf{J}_{i-1}(\mathbf{m}') = \mathbf{J}_{i-1}(\mathbf{m}) = \mathbf{a} + \mathbf{b} + (1 - \mathbf{c}' - (1 - \mathbf{e}' - \mathbf{uf})^{-1} \mathbf{d}')^{-1} \mathbf{c}' \mathbf{b}.$$




- **Case (ii):** We apply directly Lemma 3.8 to the second level of $\mathbf{J}_{i-1}(\mathbf{m})$, namely:

$$\begin{aligned} \mathbf{J}_{i-1}(\mathbf{m}) &= \mathbf{a} + (1 - \mathbf{c} - (1 - \mathbf{e} - \mathbf{uf})^{-1} \mathbf{d})^{-1} \mathbf{b} \\ &= \mathbf{a} + (1 - (1 - (1 - \mathbf{e}' - \mathbf{uf}')^{-1} (\mathbf{d}' + \mathbf{uf}'(\mathbf{e}')^{-1} \mathbf{d}'))^{-1} \mathbf{c}')^{-1} \mathbf{b} \end{aligned}$$

with $\mathbf{c}' = \mathbf{c} + \mathbf{d}$, $\mathbf{d}' = \mathbf{ed}(\mathbf{c}')^{-1}$, $\mathbf{e}' = \mathbf{ec}(\mathbf{c}')^{-1}$ and $\mathbf{f}' = \mathbf{fc}(\mathbf{c}')^{-1}$. To bring this back to Jacobi-like form, we first apply (3.5) backward on the third level, and then (3.6) backward on the first one:

$$\begin{aligned} \mathbf{J}_{i-1}(\mathbf{m}) &= \mathbf{a} + (1 - (1 + (\mathbf{e}')^{-1} \mathbf{d}' - (1 - \mathbf{e}' - \mathbf{uf}')^{-1} (\mathbf{e}')^{-1} \mathbf{d}')^{-1} \mathbf{c}')^{-1} \mathbf{b} \\ &= \mathbf{a} + \mathbf{b} + (1 - \mathbf{c}' + (\mathbf{e}')^{-1} \mathbf{d}' - (1 - \mathbf{e}' - \mathbf{uf}')^{-1} (\mathbf{e}')^{-1} \mathbf{d}')^{-1} \mathbf{c}' \mathbf{b}. \end{aligned}$$



Note that in both cases, the relation of the primed variables to the unprimed variables is the same as the relation of the mutated variables y'_j to y_j in Definition 3.6.

Clearly, for $i > 1$, we always have $\mathbf{J}_0(\mathbf{m}') = \mathbf{J}_0(\mathbf{m})$, or equivalently $\mathbf{F}_m(t) = \mathbf{F}_{m'}(t)$. When $i = 1$, we have $\mathbf{a} = 0$, $\mathbf{b} = 1$ and in all cases $\mathbf{c}' = \mathbf{y}_1(\mathbf{m}')$, so the final formula amounts to $\mathbf{J}_0(\mathbf{m}) = 1 + t\mathbf{J}_0(\mathbf{m}')\mathbf{y}_1(\mathbf{m}')$, or equivalently $\mathbf{F}_m(t) = 1 + t\mathbf{F}_{m'}(t)\mathbf{y}_1(\mathbf{m})$.

We therefore have the following theorem:

Theorem 3.10. *Under the mutation μ_i on \mathcal{A} , $\mu_i: \mathbf{y}(\mathbf{m}) \mapsto \mathbf{y}(\mathbf{m}')$ of Definition 3.6, the function $\mathbf{F}_{\mathbf{m}}(t)$ is transformed as:*

$$i = 1: \quad \mathbf{F}_{\mathbf{m}}(t) = 1 + t\mathbf{F}_{\mathbf{m}'}(t)\mathbf{y}_1(\mathbf{m}), \tag{3.16}$$

$$i > 1: \quad \mathbf{F}_{\mathbf{m}}(t) = \mathbf{F}_{\mathbf{m}'}(t). \tag{3.17}$$

3.3. An explicit expression for the weights

Although we do not have evolution equations of the form (2.1) in the non-commutative case, we make a definition of the variables \mathbf{R}_n as follows.

Definition 3.11. Let $R_n R_0^{-1}$ be defined as the coefficients in the power series expansion of $F_{m_0}(t)$, as follows:

$$\mathbf{F}_{m_0}(t) = \sum_{n=0}^{\infty} t^n \mathbf{R}_n \mathbf{R}_0^{-1}. \tag{3.18}$$

In particular, $\mathbf{y}_1(\mathbf{m}_0) = R_1 R_0^{-1}$. To make the contact with the commutative case of Section 2, one should compare \mathbf{R}_n with $R_{1,n}$ of that section.

In this section, we relate the weights $\mathbf{y}(\mathbf{m})$ to the coefficients of the generating series. The main result is Theorem 3.33 below, which gives an explicit expression for all $\mathbf{y}(\mathbf{m})$ in terms of the \mathbf{R}_n 's.

3.3.1. Stieltjes point and quasi-determinants

The generating series $\mathbf{F}_{m_0}(t)$ may be written as the non-commutative ‘‘Stieltjes’’ continued fraction $\mathbf{F}_{\mathbf{m}_1}(t)$ in its canonical form (3.10) at the Stieltjes point $\mathbf{m} = \mathbf{m}_1$. Since the sequence of forward mutations from \mathbf{m}_0 to \mathbf{m}_1 only involves mutations μ_i with $i > 1$, the generating series is preserved at each step, and $\mathbf{F}_{m_0} = \mathbf{F}_{m_1}$.

The coefficients $\mathbf{y}_i(\mathbf{m}_1)$ of the non-commutative Stieltjes continued fraction (3.10) are expressed in terms of the \mathbf{R}_n 's via a non-commutative generalization of the celebrated Stieltjes formula (Theorem 3.17 below). For this, we need a number of definitions. In the following, A denotes a square $n \times n$ matrix with coefficients $a_{i,j} \in \mathcal{A}$.

We recall the notion of quasi-determinant [18].

Definition 3.12 (Quasi-determinant). The quasi-determinant of A with respect to the entry (p, q) is defined inductively by the formula:

$$|A|_{p,q} = a_{p,q} - \sum_{i \neq p, j \neq q} a_{p,j} (|A^{p,q}|_{i,j})^{-1} a_{i,q},$$

$$|(a_{1,1})|_{1,1} = a_{1,1}$$

where $A^{p,q}$ denotes the $(n - 1) \times (n - 1)$ matrix obtained from A by erasing row p and column q , while keeping the row and column labeling of A .

We refer to [18] for a review of the many properties of the quasi-determinant. For now we will only need the following:

Proposition 3.13. (See [18], Proposition 1.4.6.) *The three following statements are equivalent:*

- (i) $|A|_{p,q} = 0$;
- (ii) *the p -th row of A is a left linear combination of the other rows;*
- (iii) *the q -th column of A is a right linear combination of the other columns.*

Note that the quasi-determinant is not a generalization of the determinant, but rather of the ratio of the determinant by a minor:

Proposition 3.14. *If \mathcal{A} is commutative, and the minor $|A^{p,q}| \neq 0$, then the quasi-determinant $|A|_{p,q}$ reduces to a ratio of the determinant of A by its p, q minor:*

$$|A|_{p,q} = \frac{|A|}{|A^{p,q}|}. \tag{3.19}$$

Finally, we also need:

Definition 3.15 (*Quasi-Wronskian*). The quasi-Wronskians of the sequence $(\mathbf{R}_M)_{M \in \mathbb{Z}}$ are defined for $i \geq 1$ and $n \in \mathbb{Z}$ as the following $i \times i$ quasi-determinants:

$$\Delta_{i,n} = \left| \begin{array}{cccccc} \mathbf{R}_{n+i-1} & \mathbf{R}_{n+i-2} & \cdots & \cdots & \mathbf{R}_n & \\ \mathbf{R}_{n+i-2} & \mathbf{R}_{n+i-3} & \cdots & \cdots & \mathbf{R}_{n-1} & \\ \vdots & \vdots & & & \vdots & \\ \mathbf{R}_n & \mathbf{R}_{n-1} & \cdots & \cdots & \mathbf{R}_{n-i+1} & \end{array} \right|_{1,1}. \tag{3.20}$$

Example 3.16. We have for $i = 1, 2, 3$:

$$\begin{aligned} \Delta_{1,n} &= \mathbf{R}_n, \\ \Delta_{2,n} &= \mathbf{R}_{n+1} - \mathbf{R}_n \mathbf{R}_{n-1}^{-1} \mathbf{R}_n, \\ \Delta_{3,n} &= \mathbf{R}_{n+2} - \mathbf{R}_n (\mathbf{R}_{n-2} - \mathbf{R}_{n-1} \mathbf{R}_n^{-1} \mathbf{R}_{n-1})^{-1} \mathbf{R}_n - \mathbf{R}_{n+1} (\mathbf{R}_{n-1} - \mathbf{R}_{n-2} \mathbf{R}_{n-1}^{-1} \mathbf{R}_n)^{-1} \mathbf{R}_n \\ &\quad - \mathbf{R}_n (\mathbf{R}_{n-1} - \mathbf{R}_n \mathbf{R}_{n-1}^{-1} \mathbf{R}_{n-2})^{-1} \mathbf{R}_{n+1} - \mathbf{R}_{n+1} (\mathbf{R}_n - \mathbf{R}_{n-1} \mathbf{R}_{n-2}^{-1} \mathbf{R}_{n-1})^{-1} \mathbf{R}_{n+1}. \end{aligned}$$

We are now ready for the non-commutative Stieltjes formula [19]. We start from the (infinite) non-commutative continued fraction $\mathbf{S}(t; \mathbf{x})$ with coefficients $\mathbf{x}_i \in \mathcal{A}$, $i \in \mathbb{Z}_{>0}$. It is defined iteratively by

$$\begin{aligned} \mathbf{S}(t; \mathbf{x}_1) &= (1 - t\mathbf{x}_1)^{-1}, \\ \mathbf{S}(t; \mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k) &= \mathbf{S}(t; \mathbf{x}_1, \dots, \mathbf{x}_{k-2}, (1 - t\mathbf{x}_k)^{-1} \mathbf{x}_{k-1}), \\ \mathbf{S}(t; \mathbf{x}) &= \lim_{k \rightarrow \infty} \mathbf{S}(t; \mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k). \end{aligned} \tag{3.21}$$

Theorem 3.17. *The coefficients \mathbf{x}_i of the non-commutative continued fraction $S(t; \mathbf{x})$ are related to the coefficients of the corresponding power series $\mathbf{S}(t; \mathbf{x}) = \sum_{n \geq 0} t^n \mathbf{R}_n \mathbf{R}_0^{-1}$ via:*

$$\mathbf{x}_{2i-1} = \Delta_{i,i} \Delta_{i,i-1}^{-1}, \tag{3.22}$$

$$\mathbf{x}_{2i} = \Delta_{i+1,i} \Delta_{i,i}^{-1}, \tag{3.23}$$

for all $i > 0$.

Proof. See Ref. [19], Corollary 8.3. \square

Comparing $\mathbf{S}(t; \mathbf{x})$ (3.21) with the finite canonical continued fraction at the Stieltjes point $\mathbf{F}_{\mathbf{m}_1}(t)$ (3.10), we see that the latter is a *finite truncation* of the former, which may be implemented by imposing that $\mathbf{x}_{2r+2} = 0$, while $\mathbf{x}_i = \mathbf{y}_i(\mathbf{m}_1)$ for $i = 1, 2, \dots, 2r + 1$. From Theorem 3.17, this finite truncation condition is equivalent to the equation $\Delta_{r+2,r+1} = 0$, and Theorem 3.17 gives us an evaluation of the non-commutative weights at the Stieltjes point in terms of the \mathbf{R}_n 's. In fact, we have:

Theorem 3.18. *The coefficients \mathbf{R}_n of the power series expansion of the finite continued fraction $\mathbf{F}_{\mathbf{m}_1}(t)$ satisfy:*

$$\Delta_{r+2,n} = 0 \quad (n > r). \tag{3.24}$$

Proof. As a consequence of the definition (3.10), we may write (with $\mathbf{y}_i \equiv \mathbf{y}_i(\mathbf{m}_1)$):

$$\mathbf{F}_{\mathbf{m}_1}(t) = P(t; \mathbf{y}_1, \dots, \mathbf{y}_{2r+1})^{-1} Q(t; \mathbf{y}_2, \dots, \mathbf{y}_{2r+1}) = P(t)^{-1} Q(t)$$

with polynomials $P, Q \in \mathcal{A}[t]$, such that $P(t), Q(t) = 1 + O(t)$ and $\deg(P) = r + 1$ and $\deg(Q) = r$. Indeed, from the inductive definition (3.21) we get

$$P(t; \mathbf{y}_1, \dots, \mathbf{y}_{s+1}) = P(t; \mathbf{y}_1, \dots, \mathbf{y}_s) - t \mathbf{y}_{s+1} P(t; \mathbf{y}_1, \dots, \mathbf{y}_{s-1}) \tag{3.25}$$

while $P(t; \mathbf{y}_1) = 1 - t \mathbf{y}_1$, and $Q(t; \mathbf{y}_2, \dots, \mathbf{y}_s) = P(t; 0, \mathbf{y}_2, \dots, \mathbf{y}_s)$. Writing $P(t; \mathbf{y}_1, \dots, \mathbf{y}_{2r+1}) = \sum_{i=0}^{r+1} (-t)^i P_i$, we conclude that the \mathbf{R}_n 's satisfy a linear recursion relation of the form

$$\sum_{i=0}^{r+1} (-1)^i P_i \mathbf{R}_{n+r+1-i} = 0 \tag{3.26}$$

for all $n \geq 0$, with $P_0 = 1$ and $P_{r+1} = \mathbf{y}_{2r+1} \mathbf{y}_{2r-1} \cdots \mathbf{y}_1$. This implies that the first row of the Wronskian matrix of (3.20) is a left linear combination of its other rows, and the theorem follows from Proposition 3.13. \square

Remark 3.19 (Conserved quantities). Note that the coefficients $P_m, m = 0, 1, \dots, r + 1$ of the linear recursion relation (3.26) are conserved quantities under the evolution of the weights of Definition 3.6. Indeed, throughout this evolution, the denominator $D_{\mathbf{m}}^{-1}$ of the generating function $\mathbf{F}_{\mathbf{m}} = D_{\mathbf{m}}^{-1} N_{\mathbf{m}}$ (where both factors are equal to 1 modulo t) remains invariant. Therefore,

$D_{\mathbf{m}} = P(t; \mathbf{y}_1, \dots, \mathbf{y}_{2r+1})$ defined above, is the generating function for the $r + 1$ conserved quantities P_m (with $P_0 = 1$). Note, however, that in general these quantities do not commute with each other.

Remark 3.20. The conserved quantities P_m have an interpretation in terms of non-commutative hard particle partition function on a chain, as in [7] in the commutative case. That is, P_m is the sum over configurations of m hard particles on the integer segment $[1, 2r + 1]$, with a weight $y_{i_m} y_{i_{m-1}} \cdots y_{i_1}$ per particle configuration occupying vertices i_1, i_2, \dots, i_m such that $i_{j+1} - i_j > 1$. This interpretation is a consequence of the recursion relation (3.25).

More generally, one can show that for any Motzkin path \mathbf{m} , the conserved quantities, when expressed in terms of the weights $\mathbf{y}_i(\mathbf{m})$, are (non-commutative) hard-particle partition functions on certain graphs $G_{\mathbf{m}}$ introduced in [7].

For later use, it will be convenient to extend the definition of \mathbf{R}_n to $n < 0$. This is easily done by extending the linear recursion relation (3.26) to any $n \in \mathbb{Z}$. For instance, this allows to define $\mathbf{R}_{-1} = P_{r+1}^{-1} \sum_{i=0}^r (-1)^{r+i} P_i \mathbf{R}_{r-i}$. The main equation extends accordingly to

$$\Delta_{r+2,n} = 0 \quad \text{for all } n \in \mathbb{Z}. \tag{3.27}$$

Remark 3.21. If \mathcal{A} is commutative, and $R_{i,n}$ are solutions to the A_r Q -system (2.1), we have $\mathbf{R}_n = R_{1,n} = R_n$. Then according to Proposition 3.14 we have:

$$\Delta_{i,n} = \frac{\det((R_{n+i+1-a-b})_{1 \leq a, b \leq i})}{\det((R_{n+i+1-a-b})_{2 \leq a, b \leq i})} = \frac{R_{i,n}}{R_{i-1,n-1}}.$$

The finite truncation condition (3.27) is equivalent to $R_{r+2,n} = 0$, for all $n \in \mathbb{Z}$.

3.3.2. Properties: From Plücker to Hirota

In this section, we derive a recursion relation for the quasi-Wronskians, which is a non-commutative analogue of the discrete Hirota equation. To this end, we must first review a few useful properties of quasi-determinants [18].

We recall the so-called heredity property of quasi-determinant (see [18], Theorem 1.4.3). This states that we get the same result if we apply Definition 3.12 to any $r \times r$ decomposition of the matrix A into possibly rectangular blocks, viewing the blocks as some new non-commutative matrix elements, and then evaluate the quasi-determinant of the resulting matrix. This holds only if the (p, q) entry of A belongs to a square block, and all expressions (involving inverses) are well defined.

Using this property, we immediately get the following two lemmas:

Lemma 3.22. *Let A be a $k + 1 \times k + 1$ matrix with last column such that $a_{i,k+1} = 0$ for $i = 1, 2, \dots, k$. Then for $1 \leq p, q \leq k$, we have:*

$$|A|_{p,q} = |A^{k+1,k+1}|_{p,q}. \tag{3.28}$$

Proof. By the heredity property, we decompose the matrix A into a first $k \times k$ square block $B = A^{k+1,k+1}$ which contains the (p, q) entry of A , a zero $k \times 1$ column vector z , a $1 \times k$

row $u = (a_{k+1,1}, \dots, a_{k+1,k})$ and the single matrix element $x = a_{k+1,k+1}$, to get: $\begin{vmatrix} B & z \\ u & x \end{vmatrix}_{p,q} = |B - zx^{-1}u|_{p,q} = |B|_{p,q}$. \square

Lemma 3.23. *Let A be a $k + 1 \times k + 1$ matrix with first column such that $a_{i,1} = 0$ for $i = 1, 2, \dots, k$. Then we have:*

$$|A|_{k+1,1} = a_{k+1,1}. \tag{3.29}$$

Proof. By the heredity property, we decompose the matrix A into a zero $k \times 1$ column z , the $k \times k$ block $B = A^{k+1,1}$, the single matrix element $x = a_{k+1,1}$, and the $1 \times k$ row vector $u = (a_{k+1,2}, \dots, a_{k+1,k+1})$:

$$|A|_{k+1,1} = \begin{vmatrix} z & B \\ x & u \end{vmatrix}_{k+1,1} = |x - uB^{-1}z|_{k+1,1} = x = a_{k+1,1}. \quad \square$$

Quasi-determinants and minors are related via the so-called homological relations:

Proposition 3.24 (*Homological relations*). (See [18], Theorem 1.4.2.)

$$(|A^{k,j}|_{i,\ell})^{-1} |A|_{i,j} = -(|A^{i,j}|_{k,\ell})^{-1} |A|_{k,j}. \tag{3.30}$$

Quasi-determinants obey the following non-commutative generalizations of the Plücker relations. Let A be a $k \times m$ matrix with elements in \mathcal{A} , and $m > k$. Let $A(i_1, \dots, i_n)$ denote the $k \times n$ matrix made of the columns of A indexed i_1, \dots, i_n .

Definition 3.25. For any ordered sequence I of column indices of A , we define the left quasi-Plücker coordinates

$$q_{i,j}^I = (|A(i, I)|_{s,i})^{-1} |A(j, I)|_{s,j}. \tag{3.31}$$

Note that the definition is independent of the choice of row index s . Moreover, if j is among I , then $q_{i,j}^I = 0$.

We may now write the generalized Plücker relations:

Proposition 3.26 (*Non-commutative Plücker relations*). (See [18], Theorem 4.4.2.) *Let $M = m_1, m_2, \dots, m_{k-1}$ and $L = \ell_1, \ell_2, \dots, \ell_k$ be two (ordered) sequences of column indices of A , then for any $i \notin L$:*

$$\sum_{j \in L \setminus M} q_{i,j}^M q_{j,i}^{L \setminus \{j\}} = 1. \tag{3.32}$$

With these properties in mind, we now turn to our main theorem:

Theorem 3.27 (*Discrete non-commutative Hirota equation*). *The quasi-Wronskians obey the following relations*

$$\Delta_{i+1,n} = \Delta_{i,n+1} - \Delta_{i,n}(\Delta_{i,n-1}^{-1} - \Delta_{i-1,n}^{-1})\Delta_{i,n} \tag{3.33}$$

for $i \geq 1$ and $n \in \mathbb{Z}$, with the convention $\Delta_{0,n}^{-1} = 0$.

Proof. Note that using the initial values $i = 1, 2$ of Example 3.16, we get $\Delta_{2,n} = \mathbf{R}_{n+1} - \mathbf{R}_n \mathbf{R}_{n-1}^{-1} \mathbf{R}_n = \Delta_{1,n+1} - \Delta_{1,n} \Delta_{1,n-1}^{-1} \Delta_{1,n}$, which agrees with (3.33) for $i = 1$, given the convention for $\Delta_{0,n}^{-1} = 0$. We now turn to the general proof of the identity (3.33) for $i > 1$. Let us apply the Plücker relation (3.32) to the following $i + 1 \times i + 3$ matrix and column index sets:

$$A = \begin{pmatrix} 1 & \mathbf{R}_{n+i} & \mathbf{R}_{n+i-1} & \cdots & \mathbf{R}_n & 0 \\ 0 & \mathbf{R}_{n+i-1} & \mathbf{R}_{n+i-2} & \cdots & \mathbf{R}_{n-1} & 0 \\ \vdots & \vdots & \vdots & & \vdots & \vdots \\ 0 & \mathbf{R}_{n+1} & \mathbf{R}_n & \cdots & \mathbf{R}_{n-i+1} & 0 \\ 0 & \mathbf{R}_n & \mathbf{R}_{n-1} & \cdots & \mathbf{R}_{n-i} & 1 \end{pmatrix}, \quad \begin{cases} M = (3, 4, \dots, i + 1, i + 2); \\ L = (2, 3, \dots, i + 1, i + 3), \end{cases}$$

so that $L \setminus M = 2, i + 3$. We now compute the various Plücker coordinates involved in the sum (3.32). Introducing

$$\begin{aligned} \theta_{i+1,n} &= \begin{vmatrix} \mathbf{R}_{n+i} & \mathbf{R}_{n+i-1} & \cdots & \mathbf{R}_{n+1} & 0 \\ \mathbf{R}_{n+i-1} & \mathbf{R}_{n+i-2} & \cdots & \mathbf{R}_n & 0 \\ \vdots & \vdots & & \vdots & \vdots \\ \mathbf{R}_{n+1} & \mathbf{R}_n & \cdots & \mathbf{R}_{n-i+2} & 0 \\ \mathbf{R}_n & \mathbf{R}_{n-1} & \cdots & \mathbf{R}_{n-i+1} & 1 \end{vmatrix}_{1,1}, \\ \varphi_{i+1,n} &= \begin{vmatrix} 0 & \mathbf{R}_{n+i-1} & \mathbf{R}_{n+i-2} & \cdots & \mathbf{R}_n \\ 0 & \mathbf{R}_{n+i-2} & \mathbf{R}_{n+i-3} & \cdots & \mathbf{R}_{n-1} \\ \vdots & \vdots & \vdots & & \vdots \\ 0 & \mathbf{R}_n & \mathbf{R}_{n-1} & \cdots & \mathbf{R}_{n-i+1} \\ 1 & \mathbf{R}_{n-1} & \mathbf{R}_{n-2} & \cdots & \mathbf{R}_{n-i} \end{vmatrix}_{1,1}, \end{aligned} \tag{3.34}$$

we find

$$q_{1,2}^M = \Delta_{i+1,n}, \quad q_{2,1}^{L \setminus \{2\}} = \theta_{i+1,n}^{-1}, \quad q_{1,i+3}^M = \varphi_{i+1,n}, \quad q_{i+3,1}^{L \setminus \{i+3\}} = \varphi_{i+1,n+1}^{-1}$$

so the Plücker relation turns into

$$\Delta_{i+1,n} \theta_{i+1,n}^{-1} + \varphi_{i+1,n} \varphi_{i+1,n+1}^{-1} = 1. \tag{3.35}$$

By direct application of Lemma 3.22, we compute $\theta_{i+1,n} = \Delta_{i,n+1}$. To compute $\varphi_{i+1,n}$ we apply the homological relation (3.30) with $i = j = 1, k = i + 1, \ell = 2$ and with the matrix A with entries: $a_{x,1} = \delta_{x,i+1}$, and $a_{x,y} = \mathbf{R}_{n-x-y+i}$ for $y > 1$. We get $\varphi_{i+1,n} = |A|_{1,1} = -|A^{i+1,1}|_{1,2} (|A^{1,1}|_{i+1,2})^{-1} |A|_{i+1,1}$. The first factor is $|A^{i+1,1}|_{1,2} = \Delta_{i,n}$ by definition, while the last factor is $|A|_{i+1,1} = a_{i+1,1} = 1$ by applying Lemma 3.23 with $k = i$. We are left with the task of computing $|A^{1,1}|_{i+1,2} = \psi_{i,n-1}$, where

$$\psi_{i,n} = \begin{vmatrix} \mathbf{R}_{n+i-1} & \mathbf{R}_{n+i-2} & \cdots & \cdots & \mathbf{R}_n \\ \mathbf{R}_{n+i-2} & \mathbf{R}_{n+i-3} & \cdots & \cdots & \mathbf{R}_{n-1} \\ \vdots & \vdots & & & \vdots \\ \mathbf{R}_n & \mathbf{R}_{n-1} & \cdots & \cdots & \mathbf{R}_{n-i+1} \end{vmatrix}_{i,1}.$$

Again, we use the homological relation (3.30) with $a_{x,y} = \mathbf{R}_{n-x-y+i+1}$, $i = i$, $j = k = 1$ and $\ell = 2$ to get:

$$\psi_{i,n} = -|A^{1,1}|_{i,2} (|A^{i,1}|_{1,2})^{-1} |A|_{1,1} = -\psi_{i-1,n-1} \Delta_{i-1,n}^{-1} \Delta_{i,n}.$$

Recalling that $\varphi_{i+1,n} = -\Delta_{i,n} \psi_{i,n-1}^{-1}$, this turns into the recursion relation

$$\varphi_{i+1,n} = -(\Delta_{i,n} \Delta_{i,n-1}^{-1}) \varphi_{i,n-1}. \tag{3.36}$$

Collecting all the results, we re-express (3.35) as:

$$\Delta_{i+1,n} = \Delta_{i,n+1} - \varphi_{i+1,n} \varphi_{i+1,n+1}^{-1} \Delta_{i,n+1} \tag{3.37}$$

$$= \Delta_{i,n+1} + \varphi_{i+1,n} \varphi_{i,n}^{-1} \Delta_{i,n}. \tag{3.38}$$

We deduce that

$$\begin{aligned} \Delta_{i,n} (\Delta_{i-1,n}^{-1} - \Delta_{i,n-1}^{-1}) &= \Delta_{i,n} \Delta_{i-1,n}^{-1} (\Delta_{i,n-1} - \Delta_{i-1,n}) \Delta_{i,n-1}^{-1} \\ &= \Delta_{i,n} \Delta_{i-1,n}^{-1} \varphi_{i,n-1} \varphi_{i-1,n-1}^{-1} \Delta_{i-1,n-1} \Delta_{i,n-1}^{-1} \\ &= \varphi_{i+1,n} \varphi_{i,n}^{-1} = (\Delta_{i+1,n} - \Delta_{i,n+1}) \Delta_{i,n}^{-1} \end{aligned}$$

and the theorem follows. \square

Remark 3.28. Theorem 3.27 is a discrete analogue of the continuous formula of Theorem 9.7.2 of [18]. Note that the continuous formula contains a misprint (there should be no -1 power for the term in parentheses).

Remark 3.29. In the case when \mathcal{A} is commutative, we write $\mathbf{R}_n = R_n$, and define for $n \in \mathbb{Z}$: $r_{0,n} = 1$, $r_{1,n} = R_n$ and $r_{i,n} = \det_{1 \leq i, j \leq i} R_{n+i+j-i-1}$. Then, according to Proposition 3.14, we have $\Delta_{i,n} = r_{i,n}/r_{i-1,n-1}$. It is easy to check that in this case the Hirota equation (3.33) is equivalent to the so-called $A_{\infty/2}$ Q -system:

$$r_{i,n+1} r_{i,n-1} = r_{i,n}^2 + r_{i+1,n} r_{i-1,n} \quad (i > 0, n \in \mathbb{Z}).$$

Further restricting to $r_{r+2,n} = 0$ for all $n \in \mathbb{Z}$ amounts to the finite truncation condition (3.27). We may then pick the A_r boundary condition $r_{r+1,n} = 1$ for which $r_{i,n} = R_{i,n}$, the solution of the A_r Q -system.

Remark 3.30. The truncation equation $\Delta_{r+2,n} = 0$ (3.27) implies the existence of a conserved quantity. Indeed, writing the Hirota equation in the form (3.37) for $i = r + 1$, we get that $\varphi_{r+2,n+1} = \varphi_{r+2,n}$ is independent of n , due to the truncation equation.

3.3.3. General expressions for the weights

By applying the non-commutative Hirota equation (3.33), we may compute explicitly all the weights $\mathbf{y}(\mathbf{m})$, $\mathbf{m} \in \mathcal{M}$, as given by Theorem 3.10, in terms of the \mathbf{R}_n 's. Let us first define new variables for $i \geq 0$ and $n \in \mathbb{Z}$:

$$D_{i,n} = \Delta_{i+1,n} \Delta_{i,n}^{-1}, \quad C_{i,n} = (-1)^i \varphi_{i+1,n} \tag{3.39}$$

with the initial conditions $D_{0,n} = 0$ and $C_{0,n} = 1$. Note that (3.36) implies the relation

$$C_{i,n} = \Delta_{i,n} \Delta_{i,n-1}^{-1} C_{i-1,n-1} \tag{3.40}$$

hence we have

$$C_{i,n} = \prod_{j=0}^{i-1} \Delta_{i-j,n-j} \Delta_{i-j,n-j-1}^{-1}. \tag{3.41}$$

Note also that Remark 3.30 implies that

$$C_{r+1,n} = C_{r+1} \tag{3.42}$$

is independent of n .

Remark 3.31. In the commutative case $\mathbf{R}_n = R_n$, with $r_{0,n} = 1$, $r_{1,n} = R_n$ and $r_{i,n}$ solution of the $A_{\infty/2}$ Q -system of Remark 3.29, we have

$$\Delta_{i,n} = \frac{r_{i,n}}{r_{i-1,n-1}}, \quad C_{i,n} = \frac{r_{i,n}}{r_{i,n-1}}, \quad D_{i,n} = \frac{r_{i+1,n} r_{i-1,n+1}}{r_{i,n+1} r_{i,n}}.$$

If we further restrict to the case of the A_r Q -system (by imposing $r_{r+2,n} = 0$ and $r_{r+1,n} = 1$), we note that the variables $C_{i,m}$ and $D_{i,m}$ reduce to the variables $c_{i,n}$ and $d_{i,n}$ of Theorem 2.14.

The variables C, D obey the following relations:

Lemma 3.32. *We have*

$$D_{i,n+1} = C_{i+1,n+1} C_{i,n}^{-1} D_{i,n} C_{i-1,n} C_{i,n+1}^{-1}, \tag{3.43}$$

$$C_{i,n+1} C_{i-1,n}^{-1} = D_{i,n} + C_{i,n} C_{i-1,n}^{-1}, \tag{3.44}$$

$$C_{i+1,n} C_{i,n-1}^{-1} = D_{i,n} + C_{i+1,n} C_{i,n}^{-1}. \tag{3.45}$$

Proof. Eq. (3.43) holds by definition, as a consequence of (3.40). Eq. (3.44) is equivalent to the Hirota equation in the form (3.38), upon multiplication by $\Delta_{i,n}^{-1}$. Using (3.44) and (3.40) we compute

$$\begin{aligned} D_{i,n} + C_{i+1,n} C_{i,n}^{-1} &= \Delta_{i+1,n} \Delta_{i+1,n-1}^{-1} (D_{i,n-1} + C_{i,n-1} C_{i-1,n-1}^{-1}) \Delta_{i,n-1} \Delta_{i,n}^{-1} \\ &= \Delta_{i+1,n} \Delta_{i+1,n-1}^{-1} C_{i,n} C_{i-1,n-1}^{-1} \Delta_{i,n-1} \Delta_{i,n}^{-1} = C_{i+1,n} C_{i,n-1}^{-1} \end{aligned}$$

and the lemma follows. \square

We are now ready for the main theorem of this section:

Theorem 3.33. *The weights $\{y_i(\mathbf{m})\}$ are expressed in terms of the \mathbf{R}_n 's as:*

$$y_{2i-1}(\mathbf{m}) = C_{i,m_i+1} C_{i-1,m_{i-1}+1}^{-1}, \tag{3.46}$$

$$y_{2i}(\mathbf{m}) = \left\{ \begin{array}{l} C_{i+1,m_{i+1}+1} C_{i+1,m_i+1}^{-1} \quad \text{if } m_i = m_{i+1} + 1, \\ 1 \quad \text{otherwise} \end{array} \right\} \\ \times D_{i,m_i+1} \times \left\{ \begin{array}{l} C_{i-1,m_i+1} C_{i-1,m_{i-1}+1}^{-1} \quad \text{if } m_i = m_{i-1} - 1, \\ 1 \quad \text{otherwise} \end{array} \right\}. \tag{3.47}$$

Proof. As already pointed out, the recursions (3.11)–(3.12) determine uniquely the weights $\mathbf{y}(\mathbf{m})$ in terms of the initial data $\mathbf{y}(\mathbf{m}_0)$ at the fundamental point $\mathbf{m}_0 = (0, 0, \dots, 0)$. However, Theorem 3.17 with $\mathbf{x}_{2r+2} = 0$ gives expressions for the weights $\mathbf{x}_i = \mathbf{y}_i(\mathbf{m}_1)$ at the Stieltjes point $\mathbf{m}_1 = (0, 1, 2, \dots, r - 1)$. So we will assume that Eqs. (3.46)–(3.47) hold at the fundamental point, and prove by induction that they hold everywhere. We then simply have to check that (3.46)–(3.47) hold at the Stieltjes point and the theorem will follow.

Using (3.46)–(3.47) for $m_i = i - 1$, we get:

$$y_{2i-1}(\mathbf{m}_1) = C_{i,i} C_{i-1,i-1}^{-1}, \quad y_{2i}(\mathbf{m}_1) = D_{i,i}.$$

These reduce respectively to (3.22) by (3.40) and to (3.23) by (3.39). So the solution (3.46)–(3.47) holds at the Stieltjes point.

We are left with proving the inductive step: assume (3.46)–(3.47) hold at some point \mathbf{m} , with weights $\mathbf{y}_i \equiv \mathbf{y}_i(\mathbf{m})$, then show that it does at $\mathbf{m}' = \mu_i \mathbf{m}$, with weights $\mathbf{y}'_i \equiv \mathbf{y}_i(\mathbf{m}')$. We must distinguish the two usual cases:

- **Case (i):** $m_{i-1} = m_i = m, m_{i+1} = m + 1$. By (3.46)–(3.47) we have:

$$y_{2i-1} = C_{i,m+1} C_{i-1,m+1}^{-1}, \quad y_{2i} = D_{i,m+1}, \quad y_{2i+1} = C_{i+1,m+2} C_{i,m+1}^{-1}.$$

The evolutions (3.11) give:

$$y'_{2i-1} = y_{2i-1} + y_{2i} = C_{i,m+2} C_{i-1,m+1}^{-1} \quad (\text{by Eq. (3.44)}),$$

$$y'_{2i} = y_{2i+1} y_{2i} (y'_{2i-1})^{-1} = D_{i,m+2} \quad (\text{by Eq. (3.43)}),$$

$$y'_{2i+1} = y_{2i+1} y_{2i} (y'_{2i-1})^{-1} = C_{i+1,m+2} C_{i,m+2}^{-1}$$

and (3.46)–(3.47) hold for \mathbf{m}' , with $m'_{i-1} = m$ and $m'_i = m'_{i+1} = m + 1$.

- **Case (ii):** $m_{i-1} = m_i = m_{i+1} = m$. Now we have by (3.46)–(3.47):

$$y_{2i-1} = C_{i,m+1} C_{i-1,m+1}^{-1}, \quad y_{2i} = D_{i,m+1},$$

$$y_{2i+1} = C_{i+1,m+1} C_{i,m+1}^{-1}, \quad y_{2i+2} = A D_{i+1,m+1}$$

where $A = C_{i+1,m+1}C_{i,m+1}^{-1}$ if $m_{i+2} = m - 1$, and $A = 1$ otherwise. The evolutions (3.12) give:

$$\begin{aligned} \mathbf{y}'_{2i-1} &= \mathbf{y}_{2i-1} + \mathbf{y}_{2i} = C_{i,m+2}C_{i-1,m+1}^{-1} \quad (\text{by Eq. (3.44)}), \\ \mathbf{y}'_{2i} &= \mathbf{y}_{2i+1}\mathbf{y}_{2i}(\mathbf{y}'_{2i-1})^{-1} = C_{i,m+1}C_{i+1,m+2}^{-1}D_{i,m+2} \quad (\text{by Eq. (3.43)}), \\ \mathbf{y}'_{2i+1} &= \mathbf{y}_{2i+1}\mathbf{y}_{2i-1}(\mathbf{y}'_{2i-1})^{-1} = C_{i+1,m+1}C_{i,m+2}^{-1}, \\ \mathbf{y}'_{2i+2} &= \mathbf{y}_{2i+2}\mathbf{y}_{2i-1}(\mathbf{y}'_{2i-1})^{-1} = AD_{i+1,m+1}C_{i,m+1}C_{i,m+2}^{-1} \end{aligned}$$

and (3.46)–(3.47) hold for \mathbf{m}' , with $m'_{i-1} = m'_{i+1} = m$ and $m'_i = m + 1$. \square

Example 3.34. The particular set of non-decreasing Motzkin paths with $m_1 \leq m_2 \leq \dots \leq m_r$ is obtained by only applying mutations pertaining to the case (i). The corresponding weights are particularly simple:

$$\mathbf{y}_{2i-1}(\mathbf{m}) = C_{i,m_i+1}C_{i-1,m_{i-1}+1}^{-1}, \quad \mathbf{y}_{2i}(\mathbf{m}) = D_{i,m_i+1}.$$

In particular, for the fundamental point \mathbf{m}_0 , we get

$$\mathbf{y}_{2i-1}(\mathbf{m}_0) = C_{i,1}C_{i-1,1}^{-1}, \quad \mathbf{y}_{2i}(\mathbf{m}_0) = D_{i,1}.$$

This can be compared to the so-called J -fraction expressions of [19], Theorem 8.12.

Example 3.35. In the case $r = 2$, we have the three Motzkin paths $(0, 0)$, $(0, 1)$ and $(1, 0)$. We have $C_{0,n} = 1$ and $C_{3,n} = C_3$. The weights are:

$$\begin{array}{lll} \mathbf{y}_1(0, 0) = C_{1,1}, & \mathbf{y}_1(0, 1) = C_{1,1}, & \mathbf{y}_1(1, 0) = C_{1,2}, \\ \mathbf{y}_2(0, 0) = D_{1,1}, & \mathbf{y}_2(0, 1) = D_{1,1}, & \mathbf{y}_2(1, 0) = C_{2,1}C_{2,2}^{-1}D_{1,2}, \\ \mathbf{y}_3(0, 0) = C_{2,1}C_{1,1}^{-1}, & \mathbf{y}_3(0, 1) = C_{2,2}C_{1,1}^{-1}, & \mathbf{y}_3(1, 0) = C_{2,1}C_{1,2}^{-1}, \\ \mathbf{y}_4(0, 0) = D_{2,1}, & \mathbf{y}_4(0, 1) = D_{2,2}, & \mathbf{y}_4(1, 0) = D_{2,1}C_{1,1}C_{1,2}^{-1}, \\ \mathbf{y}_5(0, 0) = C_3C_{2,1}^{-1}, & \mathbf{y}_5(0, 1) = C_3C_{2,2}^{-1}, & \mathbf{y}_5(1, 0) = C_3C_{2,1}^{-1}. \end{array}$$

In the formulation as paths on $\Gamma_{\mathbf{m}}$ via the canonical form of the continued fractions, the only “redundant” weight occurs in the last case $\mathbf{m} = (1, 0)$, where $\mathbf{w}_{3,1} = t\mathbf{y}_4\mathbf{y}_3^{-1}\mathbf{y}_2 = tD_{2,1}C_{1,1}C_{2,2}^{-1}D_{1,2}$. Note that weights may involve \mathbf{R}_n ’s with $n < 0$ in their definition (e.g. $\Delta_{i,n}$ for $n < i - 1$).

Theorem 3.36. We have

$$F_{\mathbf{m}}(t) = \sum_{n \geq 0} t^n \mathbf{R}_{1,n+m_1} \mathbf{R}_{1,m_1}^{-1}.$$

In other words, the quantity $\mathbf{R}_{1,n+m_1} \mathbf{R}_{1,m_1}^{-1}$ is the partition function of paths from vertex 1 to itself on $\Gamma_{\mathbf{m}}$, and with the non-commutative weights (3.1) associated to (3.46)–(3.47). As such it is a Laurent polynomial of these weights, with non-negative integer coefficients.

Proof. The statement is proved by induction on mutations. It holds for $\mathbf{m} = \mathbf{m}_0$, with $m_1 = (\mathbf{m}_0)_1 = 0$. Any given $\mathbf{m} \in \mathcal{M}$ is obtained by an iterative application of mutations μ_i of type (i) or (ii). Assume the theorem holds for some $\mathbf{m} \in \mathcal{M}$. Let us form $\mathbf{m}' = \mu_i \mathbf{m}$. When $i > 1$, we have $\mathbf{F}_{\mathbf{m}}(t) = \mathbf{F}_{\mathbf{m}'}(t) = \sum_{n \geq 0} t^n \mathbf{R}_{n+m_1} \mathbf{R}_{m_1}^{-1}$ by (3.17): hence the same quantity $\mathbf{R}_{n+m_1} \mathbf{R}_{m_1}^{-1}$ is interpreted as the partition function for paths on $\Gamma_{\mathbf{m}'}$ with the weights associated to $\mathbf{y}_i(\mathbf{m}')$, and the theorem holds for \mathbf{m}' , as $m'_1 = m_1$. If $i = 1$, then we have (3.16). Using $\mathbf{y}_1(\mathbf{m}) = C_{1,m_1+1} = \mathbf{R}_{m_1+1} \mathbf{R}_{m_1}^{-1}$, we get

$$\begin{aligned} \mathbf{F}_{\mathbf{m}}(t) &= \sum_{n \geq 0} t^n \mathbf{R}_{n+m_1} \mathbf{R}_{m_1}^{-1} = 1 + t \mathbf{F}_{\mathbf{m}'}(t) \mathbf{R}_{m_1+1} \mathbf{R}_{m_1}^{-1}, \\ \mathbf{F}_{\mathbf{m}'}(t) &= \sum_{n \geq 0} t^n \mathbf{R}_{n+m_1+1} \mathbf{R}_{m_1+1}^{-1} \end{aligned}$$

and the theorem holds for \mathbf{m}' , with $m'_1 = m_1 + 1$. \square

Example 3.37. Let us apply Theorem 3.36 to the calculation of \mathbf{R}_3 in the case $r = 2$, at the points $\mathbf{m} = (0, 0), (0, 1), (1, 0)$ respectively. We have

$$\begin{aligned} \mathbf{m} = (0, 0): \quad \mathbf{R}_3 \mathbf{R}_0^{-1} &= (\mathbf{y}_1 + \mathbf{y}_2)^2 \mathbf{y}_1 + (\mathbf{y}_3 + \mathbf{y}_4) \mathbf{y}_2 \mathbf{y}_1 \\ &= (C_{1,1} + D_{1,1})^2 C_{1,1} + (C_{2,1} + D_{2,1} C_{1,1}) C_{1,1}^{-1} D_{1,1} C_{1,1}, \\ \mathbf{m} = (0, 1): \quad \mathbf{R}_3 \mathbf{R}_0^{-1} &= (\mathbf{y}_1 + \mathbf{y}_2)^2 \mathbf{y}_1 + \mathbf{y}_3 \mathbf{y}_2 \mathbf{y}_1 \\ &= (C_{1,1} + D_{1,1})^2 C_{1,1} + C_{2,2} C_{1,1}^{-1} D_{1,1} C_{1,1}, \\ \mathbf{m} = (1, 0): \quad \mathbf{R}_3 \mathbf{R}_1^{-1} &= \mathbf{y}_1^2 + \mathbf{y}_2 \mathbf{y}_1 + \mathbf{y}_3 \mathbf{y}_1 \\ &= (C_{1,2})^2 + (C_{2,1} + D_{2,1} C_{1,1}) C_{2,2}^{-1} D_{1,2} C_{1,2} \end{aligned}$$

respectively equal to the partition functions for non-commutative weighted paths from the root to itself on the corresponding graphs $\Gamma_{\mathbf{m}}$, with weights $\mathbf{y}_i = \mathbf{y}_i(\mathbf{m})$ given in Example 3.35, and with respectively 3, 3, 2 steps towards the root. The identities between these expressions are obtained from identities of Lemma 3.32 relating C and D :

$$C_{1,1} + D_{1,1} = C_{1,2}, \quad C_{2,1} + D_{2,1} C_{1,1} = C_{2,2}, \quad C_{1,1}^{-1} D_{1,1} = C_{2,2}^{-1} D_{1,2} C_{1,2}.$$

4. Application I: T -system

Here, we apply the construction of the previous section to the A_r T -system (1.1), which can be written as a non-commutative Q -system as shown in [8]. The A_r boundary condition implies a truncation relation $T_{r+2,j,k} = 0$ for all $j, k \in \mathbb{Z}$, similar to that studied in Section 3, so we can use the results of Section 3 to give an alternative derivation of the T -system positivity results of [8].

4.1. The A_r T -system as non-commutative Q -system

We may view the T -system equations (1.1) as matrix elements of a relation in a non-commutative algebra in a particular representation. We consider the non-commuting variables $\mathbf{T}_{i,k} \in \mathcal{A}$ which have a faithful representation on a Hilbert space \mathcal{H} with basis $\{|i\rangle\}_{i \in \mathbb{Z}}$:

$$\mathbf{T}_{i,k}|j+k+i\rangle = T_{i,j,k}|j-k-i\rangle \quad (i \geq 0; j, k \in \mathbb{Z}). \tag{4.1}$$

Introducing the shift operator \mathbf{d} which acts on the same representation as $\mathbf{d}|t\rangle = |t-1\rangle$ we may rewrite the A_r boundary conditions $T_{0,j,k} = T_{r+1,j,k} = 1$ as:

$$\mathbf{T}_{0,k} = \mathbf{d}^{2k}, \quad \mathbf{T}_{r+1,k} = \mathbf{d}^{2(k+r+1)}, \tag{4.2}$$

while the finite truncation condition implies

$$\mathbf{T}_{r+2,k} = 0 \quad (k \in \mathbb{Z}).$$

Note that all variables $\mathbf{T}_{i,k}$, $0 \leq i \leq r+1$, are invertible for generic non-vanishing matrix elements $T_{i,j,k}$, with

$$(\mathbf{T}_{i,k})^{-1}|j-k-i\rangle = \frac{1}{T_{i,j,k}}|j+k+i\rangle \quad (0 \leq i \leq r+1; j, k \in \mathbb{Z}). \tag{4.3}$$

We have the following:

Proposition 4.1. *The A_r T -system (1.1) is obtained as matrix elements in \mathcal{H} of the following non-commutative Q -system:*

$$\mathbf{T}_{i,k+1}(\mathbf{T}_{i,k})^{-1}\mathbf{T}_{i,k-1} = \mathbf{T}_{i,k} + \mathbf{T}_{i+1,k}(\mathbf{T}_{i,k})^{-1}\mathbf{T}_{i-1,k} \quad (1 \leq i \leq r; k \in \mathbb{Z}) \tag{4.4}$$

with the boundary conditions (4.2).

Proof. Applying (4.4) to the vector $|j+k-1+i\rangle$ and using (4.1) and (4.3) yields (1.1) divided by $T_{i,j+1,k}$ times $|j-k+1-i\rangle$. \square

Remark 4.2. Note that the operators $\mathbf{T}_{i,k}$ are only “mildly” non-commuting, in the sense that the operators $\{\mathbf{d}^m \mathbf{T}_{i,k} \mathbf{d}^{-m-2i-2k}\}_{m,k \in \mathbb{Z}}$ are all diagonal when acting on \mathcal{H} , and therefore commute with each other. In a sense the \mathbf{d} operator is the only source of non-commutation. It was interpreted in [8] as a time-shift operator, and the non-commutation of the variables is due to their time-ordering. Nevertheless, this gives us a first non-trivial example of a non-commutative Q -system.

4.2. Quasi-Wronskian formulation

The A_r T -system is a particular case of the non-commutative finite truncation equation (3.27). We start from the solution $T_{i,j,k}$ of the system (1.1) and the operators $\mathbf{T}_{i,k}$ (4.1)–(4.3). This gives rise to the sequence $\mathbf{R}_n = \mathbf{T}_{1,n} \in \mathcal{A}$, $n \in \mathbb{Z}$. Defining $\Delta_{i,n}$, $n \in \mathbb{Z}$ as the quasi-Wronskians (3.20) of this sequence, we have the following action of the quasi-Wronskians on the basis $\{|t\rangle\}_{t \in \mathbb{Z}}$:

Theorem 4.3.

$$\Delta_{i,n}|j+n+i\rangle = \frac{T_{i,j,n}}{T_{i-1,j,n-1}}|j-n-i\rangle \quad (1 \leq i \leq r+2; j, n \in \mathbb{Z}). \tag{4.5}$$

Proof. By induction on i . For $i = 1$, we have $\Delta_{1,n} = \mathbf{R}_n$, and $\mathbf{R}_n|j+n+1\rangle = T_{1,j,n}|j-n-1\rangle$, which amounts to (4.5) for $i = 1$, as $T_{0,j,k} = 1$ for all j, k . Assume (4.5) holds for all $\beta \leq i$. Using the Hirota equation (3.33), and the recursion hypotheses we get: $\Delta_{i+1,n}|j+n+i+1\rangle = \delta_{i+1,n}|j-n-i-1\rangle$, with

$$\begin{aligned} \delta_{i+1,n} &= \frac{T_{i,j,n+1}}{T_{i-1,j,n}} + \frac{T_{i,j-1,n}}{T_{i-1,j,n-1}} \left(\frac{T_{i-2,j,n-1}}{T_{i-1,j,n}} - \frac{T_{i-1,j,n-2}}{T_{i,j,n-1}} \right) \frac{T_{i,j+1,n}}{T_{i-1,j+1,n-1}} \\ &= \frac{T_{i,j,n+1}}{T_{i-1,j,n}} - \frac{T_{i,j-1,n}T_{i,j+1,n}}{T_{i-1,j,n}T_{i,j,n-1}} \\ &= \frac{T_{i+1,j,n}}{T_{i,j,n-1}} \end{aligned}$$

where the second line follows from (1.1) for $i \rightarrow i - 1$ and $k = n - 1$, while the third follows from (1.1) with $k = n$. \square

Remark 4.4 (*Relation to the known determinant relation of [2]*). The T -system (1.1) may be simplified by first eliminating the $T_{i,j,n}$, $i > 1$ and $j, n \in \mathbb{Z}$ in terms of the $T_{j,n} = T_{1,j,n}$ by use of the Desnanot–Jacobi formula, as the following determinant:

$$T_{i,j,n} = \det(\Theta_{i,j,n}), \quad (\Theta_{i,j,n})_{a,b} = T_{j-a+b,n+i+1-a-b} \quad (1 \leq a, b \leq i).$$

We note that

$$\frac{T_{i,j,n}}{T_{i-1,j,n-1}} = |\Theta_{i,j,n}|_{1,1},$$

so the diagonal operator $\mathbf{d}^{-n-i} \Delta_{i,n} \mathbf{d}^{-n-i}$ has the eigenvalue $|\Theta_{i,j,n}|_{1,1}$ when acting on $|j\rangle$, for all $j \in \mathbb{Z}$.

Applying Theorem 4.3 for $i = r + 2$, we find:

Corollary 4.5. *The T -system solution $(\mathbf{R}_k)_{k \in \mathbb{Z}}$ satisfies the finite truncation equation (3.27).*

We may therefore apply the results of Section 3. The C, D operators (3.39) act on the basis $\{|t\rangle\}_{t \in \mathbb{Z}}$ as:

$$C_{i,n} |t\rangle = \frac{T_{i,t+n-i,n}}{T_{i,t+n-i+1,n-1}} |t - 2i\rangle, \quad C_{i,n}^{-1} |t\rangle = \frac{T_{i,t+n+i+1,n-1}}{T_{i,t+n+i,n}} |t + 2i\rangle,$$

$$D_{i,n} |t\rangle = \frac{T_{i+1,t+n+i-1,n} T_{i-1,t+n+i,n-1}}{T_{i,t+n+i-1,n-1} T_{i,t+n+i,n}} |t - 2\rangle.$$

Using the notation

$$c_{i,t,m} = \frac{T_{i,t,m}}{T_{i,t+1,m-1}}, \quad d_{i,t,m} = \frac{T_{i+1,t,m} T_{i-1,t+1,m-1}}{T_{i,t+1,m} T_{i,t,m-1}} \tag{4.6}$$

and the fact that expressions of the form $C_{i,m} C_{i,p}^{-1}$ act diagonally, Theorem 3.33 yields non-commutative weights $\mathbf{y}_i(\mathbf{m})$ which act on the basis $\{|t\rangle\}_{t \in \mathbb{Z}}$ as:

$$\mathbf{y}_{2i-1}(\mathbf{m}) |t\rangle = \frac{c_{i,t+m_i+i-1,m_i+1}}{c_{i-1,t+m_{i-1}+i,m_{i-1}+1}} |t - 2\rangle, \tag{4.7}$$

$$\mathbf{y}_{2i}(\mathbf{m}) |t\rangle = \left\{ \begin{array}{ll} \frac{c_{i+1,t+m_{i+1}+i,m_{i+1}+1}}{c_{i+1,t+m_i+i,m_i+1}} & \text{if } m_i = m_{i+1} + 1, \\ 1 & \text{otherwise} \end{array} \right\} \times d_{i,t+m_i+i,m_i+1} \times \left\{ \begin{array}{ll} \frac{c_{i-1,t+m_i+i,m_i+1}}{c_{i-1,t+m_{i-1}+i,m_{i-1}+1}} & \text{if } m_i = m_{i-1} - 1, \\ 1 & \text{otherwise} \end{array} \right\} |t - 2\rangle. \tag{4.8}$$

Remark 4.6. To be able to compare the results (4.7)–(4.8) to those of [8], we note that here, in the non-commutative weighted path interpretation, the up steps have weights $\mathbf{1}$, acting as the identity on the basis $\{|t\rangle\}_{t \in \mathbb{Z}}$, while all the down weights (including the redundant ones) act as a diagonal operator times \mathbf{d}^2 . Interpreting t as a discrete time variable, we see that up steps use no time, while down steps use two units of time. In Ref. [8], the time-dependent weighted path formulation of the T -system solutions uses skeleton down steps as well as up steps that all use one unit of time (i.e. are the product of a diagonal operator by \mathbf{d}), while the redundant steps may go back in time (a step $i \rightarrow i - k, k > 1$ uses $2 - k$ units of time, hence the corresponding weight is a diagonal operator times \mathbf{d}^{2-k}). The two descriptions are equivalent up to a “gauge” transformation, namely a conjugation of the weights (4.7)–(4.8): $\mathbf{y}_{2i-1} \rightarrow \mathbf{d}^{-i-1} \mathbf{y}_{2i-1} \mathbf{d}^i$ and $\mathbf{y}_{2i} \rightarrow \mathbf{d}^{-i-1} \mathbf{y}_{2i} \mathbf{d}^i$.

5. Application II: Quantum Q -system

5.1. Quantum cluster algebra

The solutions of Q -system for A_r have been shown to form particular clusters in a suitable cluster algebra [21]. On the other hand, cluster algebras are known to have quantum analogues, the so-called quantum cluster algebras [3]. In particular, there exists a quantum deformation of the Q -system relations.

The cluster algebra for the A_r Q -system is a rank $2r$ cluster algebra with trivial coefficients, and with initial cluster \mathbf{x}_0 and exchange matrix B_0 :

$$\mathbf{x}_0 = (R_{1,0}, R_{2,0}, \dots, R_{r,0}, R_{1,1}, R_{2,1}, \dots, R_{r,1}), \quad B_0 = \begin{pmatrix} 0 & -C^t \\ C & 0 \end{pmatrix}$$

where C is the Cartan matrix for A_r . The subset of (forward) mutations we have considered so far is such that each mutation amounts to one application of the Q -system equation, allowing to update the cluster via a substitution of the form: $R_{i,n-1} \rightarrow R_{i,n+1} = (R_{i,n}^2 + R_{i+1,n}R_{i-1,n})/R_{i,n-1}$ (mutation μ_i if n is odd, μ_{r+i} if n is even). Recall also that the compound mutation $\mu = \mu_{2r}\mu_{2r-1} \cdots \mu_1$ acts on \mathbf{x}_0 as a global translation of all the n indices by $+2$ (i.e. $R_{i,n} \rightarrow R_{i,n+2}$ for all i), preserving B_0 . The “half-translation” of all indices by $+1$ is realized by acting only with $\mu_{\text{even}} = \mu_r\mu_{r-1} \cdots \mu_1$ or $\mu_{\text{odd}} = \mu_{2r}\mu_{2r-1} \cdots \mu_{r+1}$ in alternance, which reverse the sign of the exchange matrix.

A quantum cluster algebra with trivial coefficients corresponding to this involves the *same* exchange matrix, subject to the same evolution under mutation, but the cluster variables are now non-commuting variables $\mathbf{R}_{i,n} \in \mathcal{A}$ whose commutation relations within each cluster are dictated by the exchange matrix as follows. We have for any pair (x_i, x_j) of variables within the same cluster the following q -commutation relation:

$$x_i x_j = q^{\Lambda_{i,j}} x_j x_i \tag{5.1}$$

where Λ is a suitably normalized $2r \times 2r$ matrix proportional to the inverse of the exchange matrix. Note that in a different cluster the commutation relations are different, as they are entirely dictated by the exchange matrix, and may involve other cluster variables.

For the initial cluster, in order for Λ to have only integer entries, we choose $\Lambda = (r+1)B_0^{-1} = \begin{pmatrix} 0 & \lambda \\ -\lambda & 0 \end{pmatrix}$, where $\lambda = (r+1)C^{-1}$. We obtain the following q -commutation relations between variables of the fundamental cluster:

Lemma 5.1. *For $1 \leq i, j \leq r$, we have:*

$$\mathbf{R}_{i,0}\mathbf{R}_{j,1} = q^{\lambda_{i,j}} \mathbf{R}_{j,1}\mathbf{R}_{i,0}, \tag{5.2}$$

$$\mathbf{R}_{i,n}\mathbf{R}_{j,n} = \mathbf{R}_{j,n}\mathbf{R}_{i,n} \quad (n = 0, 1), \tag{5.3}$$

where

$$\lambda_{i,j} = \text{Min}(i, j)(r + 1 - \text{Max}(i, j)) \quad (1 \leq i, j \leq r). \tag{5.4}$$

Proof. By explicitly inverting the Cartan matrix C , with entries $2\delta_{i,j} - \delta_{|i-j|,1}$. \square

Note that the relations (5.2) and (5.3) are preserved under the above-mentioned half-translation of all n indices by $+1$, as $B_0 \rightarrow -B_0$, and therefore $\Lambda \rightarrow -\Lambda$ for μ_{even} and vice versa for μ_{odd} . Hence the relations (5.2)–(5.3) extend to

$$\mathbf{R}_{i,n}\mathbf{R}_{j,n+1} = q^{\lambda_{i,j}} \mathbf{R}_{j,n+1}\mathbf{R}_{i,n}, \tag{5.5}$$

$$\mathbf{R}_{i,n}\mathbf{R}_{j,n} = \mathbf{R}_{j,n}\mathbf{R}_{i,n} \tag{5.6}$$

for all $n \in \mathbb{Z}$. In particular, the initial sequence $\mathbf{R}_n = \mathbf{R}_{1,n}$, $n \in \mathbb{Z}$, has the q -commutation relations:

$$\mathbf{R}_n \mathbf{R}_{n+1} = q^r \mathbf{R}_{n+1} \mathbf{R}_n. \tag{5.7}$$

In the following we consider the *same* subset of cluster mutations as in the commuting A_r - Q -system cluster algebra, leading to a subset of cluster nodes indexed by Motzkin paths $\mathbf{m} \in \mathcal{M}$. Each of these clusters $\mathbf{x}_{\mathbf{m}}$ is now made of $2r$ non-commuting variables $\{\mathbf{R}_{i,m_i}, \mathbf{R}_{i,m_i+1}\}_{i=1}^r$, ordered as in the commuting case. As before, these will play the role of possible initial conditions for the quantum Q -system below.

5.2. Quantum Q -system

The particular form of the quantum Q -system we choose is borrowed from the T -system case (4.4), and compatible up to multiplicative redefinition of the variables with the mutations of the quantum cluster algebra.

Definition 5.2. We define the quantum Q -system for A_r as:

$$\mathbf{R}_{i,k+1} (\mathbf{R}_{i,k})^{-1} \mathbf{R}_{i,k-1} = \mathbf{R}_{i,k} + \mathbf{R}_{i+1,k} (\mathbf{R}_{i,k})^{-1} \mathbf{R}_{i-1,k} \quad (i \geq 1; k \in \mathbb{Z}) \tag{5.8}$$

with the boundary conditions

$$\mathbf{R}_{0,k} = \mathbf{1}, \quad \mathbf{R}_{r+1,k} = \mathbf{1} \quad (k \in \mathbb{Z}). \tag{5.9}$$

Note that the latter imply a relation

$$\mathbf{R}_{r+2,k} = 0 \quad (k \in \mathbb{Z}) \tag{5.10}$$

as $\lambda_{r+1,r+1} = 0$. We also have $\mathbf{R}_{-1,n} = 0$ as $\lambda_{0,0} = 0$.

Using the q -commutations (5.5)–(5.6), we may rewrite the system (5.8) in a simpler form:

Lemma 5.3. *The A_r quantum Q -system is equivalent to:*

$$q^{\lambda_{i,i}} \mathbf{R}_{i,k+1} \mathbf{R}_{i,k-1} = (\mathbf{R}_{i,k})^2 + \mathbf{R}_{i+1,k} \mathbf{R}_{i-1,k} \quad (i \geq 1; k \in \mathbb{Z}) \tag{5.11}$$

with $\lambda_{i,j}$ as in (5.4), and with the same boundary condition (5.9).

Remark 5.4. The relation (5.11) is equivalent to the relevant quantum cluster mutations of [3] up to a scalar renormalization of the cluster variables. Indeed, defining the variables $\mathbf{X}_{i,n} = q^{\frac{r+1}{4} \lambda_{i,i}} \mathbf{R}_{i,n}$, we find:

$$\mathbf{X}_{i,k+1} = q^{\lambda_{i,i}} (\mathbf{X}_{i,k-1})^{-1} (\mathbf{X}_{i,k})^2 + q^{\frac{1}{2}(\lambda_{i,i-1} + \lambda_{i,i+1})} (\mathbf{X}_{i,k-1})^{-1} \mathbf{X}_{i-1,k} \mathbf{X}_{i+1,k} \tag{5.12}$$

in agreement with the definitions of [3]. Note that boundary conditions remain the same: $\mathbf{X}_{0,n} = \mathbf{X}_{r+1,n} = \mathbf{1}$ for all $n \in \mathbb{Z}$.

5.3. *Quasi-Wronskian formulation*

The quasi-Wronskians (3.20) of the variables $\mathbf{R}_n = \mathbf{R}_{1,n}$ may be expressed as simple monomials in terms of the solutions of the quantum Q -system (5.8)–(5.9). We have:

Theorem 5.5.

$$\Delta_{i,n} = q^{-\frac{1}{2}(i-1)(2r+2-i)} \mathbf{R}_{i,n} \mathbf{R}_{i-1,n-1}^{-1} \quad (i > 1, n \in \mathbb{Z}). \tag{5.13}$$

Proof. By induction on i . Define $\Gamma_{i,n} = q^{-\frac{1}{2}(i-1)(2r+2-i)} \mathbf{R}_{i,n} \mathbf{R}_{i-1,n-1}^{-1}$, the r.h.s. of (5.13) and $a_i = \frac{1}{2}(i-1)(2r+2-i)$. For $i = 1$, we have $\Delta_{1,n} = \mathbf{R}_{1,n} = \Gamma_{1,n}$ from the boundary condition (5.9). Assume (5.13) holds for all $1 \leq j \leq i$. We use the Hirota equation (3.33) to express

$$\Delta_{i+1,n} - \Gamma_{i+1,n} = \Gamma_{i,n+1} - \Gamma_{i+1,n} - \Gamma_{i,n} \Gamma_{i,n-1}^{-1} (\Gamma_{i,n-1} - \Gamma_{i-1,n}) \Gamma_{i-1,n}^{-1} \Gamma_{i,n}. \tag{5.14}$$

Moreover, defining $B_{i,n} = (\mathbf{R}_{i,n})^2 + \mathbf{R}_{i+1,n} \mathbf{R}_{i-1,n} - q^{\lambda_{i,i}} \mathbf{R}_{i,n+1} \mathbf{R}_{i,n-1}$ ($= 0$ due to the quantum Q -system equation (5.11)), we find that

$$\Gamma_{i-1,n} - \Gamma_{i,n-1} = q^{-a_i} ((\mathbf{R}_{i-1,n-1})^2 - B_{i-1,n-1}) \mathbf{R}_{i-2,n-1}^{-1} \mathbf{R}_{i-1,n-2}^{-1}$$

due to the q -commutation relations (5.5)–(5.6) and the identity $\lambda_{i-1,i-1} - \lambda_{i-2,i-1} - a_i = -a_{i-1}$. Substituting this into (5.14), we get

$$\begin{aligned} \Delta_{i+1,n} - \Gamma_{i+1,n} &= q^{-a_{i+1}} \mathbf{R}_{i,n}^2 \mathbf{R}_{i-1,n}^{-1} \mathbf{R}_{i,n-1}^{-1} \\ &\quad - q^{-a_i} \Gamma_{i,n} \Gamma_{i,n-1}^{-1} (\mathbf{R}_{i-1,n-1})^2 \mathbf{R}_{i-2,n-1}^{-1} \mathbf{R}_{i-1,n-2}^{-1} \Gamma_{i-1,n}^{-1} \Gamma_{i,n} \\ &= 0 \end{aligned}$$

by use of the q -commutation relations (5.5)–(5.6). \square

Remark 5.6. Theorem 5.5 allows to interpret $\mathbf{R}_{i,n}$ as a “quantum Wronskian”, namely a “quantum determinant” of the Wronskian matrix used in Definition 3.15. Indeed, we may write

$$\mathbf{R}_{i,n} = q^{\frac{i(i-1)}{6}(3r+2-i)} \Delta_{i,n} \Delta_{i-1,n-1} \cdots \Delta_{1,n-i+1} \quad (i \geq 1)$$

which is a product of principal quasi-minors of the Wronskian matrix, like in the standard definitions of quantum determinants [11] (note however that here the quasi-minors do not commute). In the cases $i = 1, 2, 3$, using the q -commutations and the quantum Q -system, we find the following expressions for the quantum Wronskian:

$$\begin{aligned} \mathbf{R}_{1,n} &= \mathbf{R}_n, \\ \mathbf{R}_{2,n} &= q^r \mathbf{R}_{n+1} \mathbf{R}_{n-1} - \mathbf{R}_n^2, \\ \mathbf{R}_{3,n} &= q^{3r-1} \mathbf{R}_{n+2} \mathbf{R}_n \mathbf{R}_{n-2} - q^{2r-1} (\mathbf{R}_{n+2} \mathbf{R}_{n-1}^2 + \mathbf{R}_{n+1}^2 \mathbf{R}_{n-2}) \\ &\quad + q^{r-1} (q^{r+1} + 1) \mathbf{R}_{n+1} \mathbf{R}_n \mathbf{R}_{n-1} - \mathbf{R}_n^3. \end{aligned}$$

Applying Theorem 5.5 for $i = r + 2$, and using Eq. (5.10), we get:

Corollary 5.7. *The solution of the quantum Q -system (5.8)–(5.9) satisfies the finite truncation condition $\Delta_{r+2,n} = 0$ for all $n \in \mathbb{Z}$.*

We may therefore apply the results of Section 3. Assuming $\mathbf{R}_{i,n}$ is a solution of (5.8)–(5.9), we consider the usual subset of clusters \mathbf{x}_m , also viewed as initial data for the quantum Q -system. Recall that the commutation relations between the quantum cluster variables within each cluster \mathbf{x}_m are determined by the exchange matrix B_m . Using the explicit expressions for the exchange matrix B_m of Ref. [7], it is an easy exercise to compute the q -commutation relations between the cluster variables within any given cluster \mathbf{x}_m . We find that:

Lemma 5.8.

$$\mathbf{R}_{i,n} \mathbf{R}_{j,n+p} = q^{p\lambda_{i,j}} \mathbf{R}_{j,n+p} \mathbf{R}_{i,n} \tag{5.15}$$

for all n, p, i, j such that the variables belong to the same cluster \mathbf{x}_m .

Remark 5.9. Lemma 5.8 does not cover for instance the case $j = i, p = 2$, as $\mathbf{R}_{i,n}$ and $\mathbf{R}_{i,n+2}$ never belong to the same cluster \mathbf{x}_m . Using the quantum Q -system and the valid q -commutations, we find for instance

$$\mathbf{R}_{i,n} \mathbf{R}_{i,n+2} = q^{2\lambda_{i,i}} (q^{-r-1} \mathbf{R}_{i,n+2} \mathbf{R}_{i,n} + (q^{-r-1} - 1) \mathbf{R}_{i,n+1}^2).$$

However, we may apply Lemma 5.8 to compute the commutation of $\Delta_{i,n}$ and $\Delta_{j,m}$, using Theorem 5.5, when the four variables $\mathbf{R}_{i,n}, \mathbf{R}_{i-1,n-1}, \mathbf{R}_{j,m}, \mathbf{R}_{j-1,m-1}$ belong to the same cluster \mathbf{x}_m :

$$\Delta_{i,n} \Delta_{j,m} = q^{(n-m)(\lambda_{i,j-1} + \lambda_{j,i-1}) + r(j-i)} \Delta_{j,m} \Delta_{i,n}.$$

Using the q -commutation relations (5.15) we easily get the following:

Lemma 5.10. *The quantities $C_{i,n}$ and $D_{i,n}$ (3.39) are expressed as:*

$$C_{i,n} = q^{\frac{i(i-1)}{2}} \mathbf{R}_{i,n} \mathbf{R}_{i,n-1}^{-1}, \quad D_{i,n} = \mathbf{R}_{i+1,n} \mathbf{R}_{i,n}^{-1} \mathbf{R}_{i,n-1}^{-1} \mathbf{R}_{i-1,n-1}.$$

In particular, the value of the conserved quantity (3.42) follows from Lemma 5.10 and the boundary condition (5.9):

$$C_{r+1} = q^{\frac{r(r+1)}{2}}.$$

Note that in this case, this conserved quantity is central.

Theorem 5.11. *Let $\mathbf{R}_{i,n}$ be a solution of the quantum Q -system (5.8)–(5.9) and $\mathbf{R}_n = \mathbf{R}_{1,n}$. The quantity $\mathbf{R}_{n+m_1} \mathbf{R}_{m_1}^{-1}$ for $n \geq 0$, when expressed in terms of initial data \mathbf{x}_m , is the partition function for non-commutative weighted paths on the graph Γ_m from the root to itself with n steps towards the root and with skeleton weights:*

$$y_{2i-1}(\mathbf{m}) = q^{i-1} \mathbf{R}_{i,m_i+1} \mathbf{R}_{i,m_i}^{-1} \mathbf{R}_{i-1,m_{i-1}} \mathbf{R}_{i-1,m_{i-1}+1}^{-1}, \tag{5.16}$$

$$y_{2i}(\mathbf{m}) = \left\{ \begin{array}{l} \mathbf{R}_{i+1,m_{i+1}+1} \mathbf{R}_{i+1,m_{i+1}}^{-1} \mathbf{R}_{i+1,m_i} \mathbf{R}_{i+1,m_i+1}^{-1} \quad \text{if } m_i = m_{i+1} + 1, \\ \mathbf{1} \quad \text{otherwise} \end{array} \right\} \\ \times \mathbf{R}_{i+1,m_i+1} \mathbf{R}_{i,m_i+1}^{-1} \mathbf{R}_{i,m_i}^{-1} \mathbf{R}_{i-1,m_i} \\ \times \left\{ \begin{array}{l} \mathbf{R}_{i-1,m_i+1} \mathbf{R}_{i-1,m_i}^{-1} \mathbf{R}_{i-1,m_{i-1}} \mathbf{R}_{i-1,m_{i-1}+1}^{-1} \quad \text{if } m_i = m_{i-1} - 1, \\ \mathbf{1} \quad \text{otherwise} \end{array} \right\} \tag{5.17}$$

that are all non-commutative Laurent monomials of the initial data, times some possible integer powers of q . As such it is a Laurent polynomial of the initial data with coefficients in $\mathbb{Z}_+[q, q^{-1}]$.

Proof. We apply Theorem 3.36 with the weights (3.33). We then rewrite all the skeleton weights using Lemma 5.10. Finally, the redundant weights (3.7) are also monomials of the initial data, with a possible integer power of q in factor. Reordering and rearranging the terms may only produce coefficients in $\mathbb{Z}_+[q, q^{-1}]$. \square

Example 5.12. In the case of non-decreasing Motzkin paths (see Example 3.34 above), we have the following expressions for the skeleton weights:

$$y_{2i-1}(\mathbf{m}) = \begin{cases} \mathbf{R}_{i,m_i+1} \mathbf{R}_{i-1,m_{i-1}+1}^{-1} \mathbf{R}_{i,m_i}^{-1} \mathbf{R}_{i-1,m_{i-1}} & \text{if } m_i = m_{i-1}; \\ \mathbf{R}_{i,m_i+1} \mathbf{R}_{i-1,m_{i-1}+1}^{-1} \mathbf{R}_{i-1,m_{i-1}} \mathbf{R}_{i,m_i}^{-1} & \text{if } m_i = m_{i-1} + 1, \end{cases} \\ y_{2i}(\mathbf{m}) = \mathbf{R}_{i+1,m_i+1} \mathbf{R}_{i,m_i+1}^{-1} \mathbf{R}_{i,m_i}^{-1} \mathbf{R}_{i-1,m_i}$$

where we have used the q commutations (5.15) and the identity $\lambda_{i-1,i} - \lambda_{i-1,i-1} + i - 1 = 0$ to simplify the monomials. For the fundamental initial condition $\mathbf{x}_0 = \mathbf{x}_{\mathbf{m}_0}$, this gives:

$$y_{2i-1}(\mathbf{m}_0) = \mathbf{R}_{i,1} \mathbf{R}_{i-1,1}^{-1} \mathbf{R}_{i,0}^{-1} \mathbf{R}_{i-1,0}, \\ y_{2i}(\mathbf{m}_0) = \mathbf{R}_{i+1,1} \mathbf{R}_{i,1}^{-1} \mathbf{R}_{i,0}^{-1} \mathbf{R}_{i-1,0}.$$

For the Stieltjes point $\mathbf{x}_{\mathbf{m}_1}$, we have:

$$y_{2i-1}(\mathbf{m}_1) = \mathbf{R}_{i,i} \mathbf{R}_{i-1,i-1}^{-1} \mathbf{R}_{i-1,i-2} \mathbf{R}_{i,i-1}^{-1}, \\ y_{2i}(\mathbf{m}_1) = \mathbf{R}_{i+1,i} \mathbf{R}_{i,i}^{-1} \mathbf{R}_{i,i-1}^{-1} \mathbf{R}_{i-1,i-1}.$$

Example 5.13. Let us consider the case $r = 2$ of Example 3.35. Setting $\mathbf{R}_n = \mathbf{R}_{1,n}$ and $\mathbf{S}_n = \mathbf{R}_{2,n}$, we have the following commutation relations:

$$\mathbf{R}_n \mathbf{R}_{n+1} = q^2 \mathbf{R}_{n+1} \mathbf{R}_n, \quad \mathbf{S}_n \mathbf{S}_{n+1} = q^2 \mathbf{S}_{n+1} \mathbf{S}_n, \quad \mathbf{R}_n \mathbf{S}_n = \mathbf{S}_n \mathbf{R}_n, \\ \mathbf{S}_n \mathbf{R}_{n+1} = q \mathbf{R}_{n+1} \mathbf{S}_n, \quad \mathbf{R}_n \mathbf{S}_{n+1} = q \mathbf{S}_{n+1} \mathbf{R}_n \quad (n \in \mathbb{Z}). \tag{5.18}$$

The A_2 quantum Q -system reads:

$$\begin{aligned} q^2 \mathbf{R}_{n+1} \mathbf{R}_{n-1} &= (\mathbf{R}_n)^2 + \mathbf{S}_n, \\ q^2 \mathbf{S}_{n+1} \mathbf{S}_{n-1} &= (\mathbf{S}_n)^2 + \mathbf{R}_n \end{aligned}$$

and the weights at the various Motzkin paths may be expressed as:

$$\begin{aligned} \mathbf{y}_1(0, 0) &= \mathbf{R}_1 \mathbf{R}_0^{-1}, & \mathbf{y}_1(0, 1) &= \mathbf{R}_1 \mathbf{R}_0^{-1}, & \mathbf{y}_1(1, 0) &= \mathbf{R}_2 \mathbf{R}_1^{-1}, \\ \mathbf{y}_2(0, 0) &= \mathbf{S}_1 \mathbf{R}_1^{-1} \mathbf{R}_0^{-1}, & \mathbf{y}_2(0, 1) &= \mathbf{S}_1 \mathbf{R}_1^{-1} \mathbf{R}_0^{-1}, & \mathbf{y}_2(1, 0) &= \mathbf{S}_0^{-1} (\mathbf{S}_1)^2 \mathbf{R}_1^{-1} \mathbf{R}_2^{-1}, \\ \mathbf{y}_3(0, 0) &= \mathbf{S}_1 \mathbf{R}_1^{-1} \mathbf{S}_0^{-1} \mathbf{R}_0, & \mathbf{y}_3(0, 1) &= \mathbf{S}_2 \mathbf{R}_1^{-1} \mathbf{R}_0 \mathbf{S}_1^{-1}, & \mathbf{y}_3(1, 0) &= \mathbf{S}_1 \mathbf{R}_1 \mathbf{S}_0^{-1} \mathbf{R}_2^{-1}, \\ \mathbf{y}_4(0, 0) &= \mathbf{S}_1^{-1} \mathbf{S}_0^{-1} \mathbf{R}_0, & \mathbf{y}_4(0, 1) &= \mathbf{S}_2^{-1} \mathbf{S}_1^{-1} \mathbf{R}_1, & \mathbf{y}_4(1, 0) &= \mathbf{S}_0^{-1} \mathbf{S}_1^{-1} (\mathbf{R}_1)^2 \mathbf{R}_2^{-1}, \\ \mathbf{y}_5(0, 0) &= \mathbf{S}_1^{-1} \mathbf{S}_0, & \mathbf{y}_5(0, 1) &= \mathbf{S}_2^{-1} \mathbf{S}_1, & \mathbf{y}_5(1, 0) &= \mathbf{S}_1^{-1} \mathbf{S}_0 \end{aligned}$$

where we have used (5.18) to eliminate the q -dependent prefactors. The redundant weight of the last case $\mathbf{m} = (1, 0)$ is $\mathbf{w}_{3,1} = t \mathbf{y}_4 \mathbf{y}_3^{-1} \mathbf{y}_2 = t q^{-1} \mathbf{R}_2^{-1} \mathbf{S}_0^{-1}$. Note that the conservation law $\mathbf{y}_5(\mathbf{m}) \mathbf{y}_3(\mathbf{m}) \mathbf{y}_1(\mathbf{m}) = C_3 = q^3$ is satisfied for the three Motzkin paths.

Remark 5.14. One can show that the conserved quantities P_m of Remark 3.19 commute with each other in this case. This suggests an analogy with the definition of classical integrability, where the integrals of the motion are in involution. In fact, in the A_1 quantum Q -system case, the Hamiltonian structure and integrability were shown in [13]. The case of the A_1 T -system was treated in [12].

Remark 5.15. The quasi-Wronskians $\Delta_{i,n}$ are “bar-invariant” quantities. Let us define an anti-automorphism (called usually the bar involution), which we denote by $*$, by $(ab)^* = b^* a^*$ for all $a, b \in \mathcal{A}$, $\mathbf{1}^* = \mathbf{1}$ and $q^* = q^{-1}$, while $(\mathbf{R}_n)^* = \mathbf{R}_n$ for all $n \in \mathbb{Z}$. Then, as a trivial consequence of the Hirota equation (3.33) and of the initial condition $\Delta_{1,n} = \mathbf{R}_n$, we have that

$$(\Delta_{i,n})^* = \Delta_{i,n}.$$

6. Other non-commutative systems

In this section, we consider further examples of non-commutative recursions which have the positive Laurent property. In the following, all the variables we consider belong to some skew field of rational fractions \mathcal{A} .

6.1. Affine rank 2 non-commutative systems

The non-commutative A_1 Q -system (1.5) corresponds to one of the so-called rank two affine cluster algebras, namely with initial exchange matrix B_0 related to affine Cartan matrices of rank 2. These read $B_0 = \begin{pmatrix} 0 & -c \\ b & 0 \end{pmatrix}$ with $bc = 4$, $b, c \in \mathbb{Z}_{>0}$. While the A_1 case corresponds to

$b = c = 2$, the two other cases $(b, c) = (1, 4)$ or $(4, 1)$ were shown in Ref. [10] to be related to the following non-commutative system for $\mathbf{R}_n \in \mathcal{A}$, $n \in \mathbb{Z}$:

$$\mathbf{R}_{n+1}(\mathbf{R}_n)^{-1}\mathbf{R}_{n-1}\mathbf{R}_n = \begin{cases} (\mathbf{R}_n)^b + 1 & \text{if } n \text{ odd;} \\ (\mathbf{R}_n)^c + 1 & \text{if } n \text{ even.} \end{cases} \tag{6.1}$$

Let us fix $b = 1, c = 4$.

Theorem 6.1. *The solution \mathbf{R}_n of the system (6.1) is a positive Laurent polynomial of any initial data $(\mathbf{R}_i, \mathbf{R}_{i+1})$, for any $i, n \in \mathbb{Z}$.*

We sketch the proof below. As in the A_1 case, one finds two conserved quantities and a linear recursion relation for \mathbf{R}_n with constant coefficients. Finally, the generating function for $\mathbf{R}_{2n}\mathbf{R}_0^{-1}$ reads:

$$\mathbf{F}_0(t) = \sum_{n \geq 0} t^n \mathbf{R}_{2n} \mathbf{R}_0^{-1} = (1 - t\mathbf{z}_1 - t^2(1 - t\mathbf{z}_3)^{-1}\mathbf{z}_2)^{-1}$$

where

$$\begin{aligned} \mathbf{z}_1 &= \mathbf{u}_1 \mathbf{u}_0^{-1} = (\mathbf{R}_1^{-1} + 1)\mathbf{R}_0^{-1}\mathbf{R}_1\mathbf{R}_0^{-1}, \\ \mathbf{z}_2 &= (\mathbf{K} - \mathbf{z}_1)\mathbf{z}_1 - \mathbf{C} = (\mathbf{R}_1^{-1}\mathbf{R}_0^2\mathbf{R}_1^{-1} + (1 + \mathbf{R}_1^{-1})\mathbf{R}_0^{-2}(1 + \mathbf{R}_1^{-1}))\mathbf{R}_0^{-1}\mathbf{R}_1\mathbf{R}_0^{-1}, \\ \mathbf{z}_3 &= \mathbf{K} - \mathbf{z}_1 = \mathbf{R}_1^{-1}\mathbf{R}_0^2 + (1 + \mathbf{R}_1^{-1})\mathbf{R}_0^{-2} \end{aligned}$$

and the Laurent positivity follows for $(\mathbf{R}_0, \mathbf{R}_1)$. The mutated expressions in terms of $(\mathbf{R}_i, \mathbf{R}_{i+1})$ are again obtained by manifestly positive non-commutative rearrangements of $\mathbf{F}_0(t)$, and the general Laurent positivity follows from symmetries of the system.

6.2. Some rank $2k + 1$ cases

For any fixed $k \in \mathbb{Z}_+$, we consider the “rank $2k + 1$ ” recursion relation for $n \in \mathbb{Z}$, for variables $\mathbf{u}_n \in \mathcal{A}$:

$$\mathbf{u}_{2n+2k+1}\mathbf{u}_{2n} = \mathbf{u}_{2n+1}\mathbf{u}_{2n+2k} + \mathbf{1}, \tag{6.2}$$

$$\mathbf{u}_{2n+1}\mathbf{u}_{2n+2k+2} = \mathbf{u}_{2n+2k+1}\mathbf{u}_{2n+2} + \mathbf{1}. \tag{6.3}$$

Admissible initial data for this system are $\mathbf{x}_i = (\mathbf{u}_i, \mathbf{u}_{i+1}, \dots, \mathbf{u}_{i+2k})$ for all $i \in \mathbb{Z}$. We have the following positivity theorem.

Theorem 6.2. *The general solution \mathbf{u}_n to the system (6.2)–(6.3) is a positive Laurent polynomial of any initial data \mathbf{x}_i , for all $n, i \in \mathbb{Z}$.*

This is proved in Appendix B, by using the integrability of the system and constructing non-commutative weighted path models for u_n . The commutative version of the system (6.2)–(6.3) corresponds to the affine A_{2k} cluster algebra [1]. Note that the exchange matrix of the fundamental seed is singular, so there is no quantum cluster algebra version of the system (6.2)–(6.3).

6.3. A non-coprime rank 2 case related to rank 3

The following non-commutative recursion relation also satisfies the positive Laurent property:

$$\mathbf{R}_{n+1}(\mathbf{R}_n)^{-1}\mathbf{R}_{n-1} = \mathbf{R}_n + 1. \tag{6.4}$$

Note that we still have the quasi-commutation relation $\mathbf{R}_n\mathbf{R}_{n+1} = \mathbf{R}_{n+1}\mathbf{C}\mathbf{R}_n$ from the conserved quantity \mathbf{C} , and that we may equivalently write $\mathbf{R}_{n+1}\mathbf{C}\mathbf{R}_{n-1} = \mathbf{R}_n(\mathbf{R}_n + 1)$, or

$$\mathbf{R}_{n-1}(\mathbf{R}_n)^{-1}\mathbf{R}_{n+1} = \mathbf{R}_n + \mathbf{C}^{-1}. \tag{6.5}$$

In the commuting case, it would correspond to the recursion relation $R_{n+1}R_{n-1} = R_n(R_n + 1)$ which does not correspond to a cluster algebra mutation, as the two terms on the r.h.s. are not coprime. However, this relation may be connected to a non-commutative rank 3 system, with the positive Laurent property, as follows. Introduce the change of variables:

$$\mathbf{R}_{2n+1} = \mathbf{u}_{2n+1}\mathbf{u}_{2n}, \quad \mathbf{R}_{2n} = \mathbf{u}_{2n-1}\mathbf{u}_{2n}$$

for some variables $\mathbf{u}_n, n \in \mathbb{Z}$. We have:

$$\begin{aligned} \mathbf{u}_{2n+1}\mathbf{u}_{2n-2} &= \mathbf{u}_{2n-1}\mathbf{u}_{2n} + \mathbf{1}, \\ \mathbf{u}_{2n-1}\mathbf{u}_{2n+2} &= \mathbf{u}_{2n+1}\mathbf{u}_{2n} + \mathbf{C}^{-1}. \end{aligned}$$

This is nothing but the $k = 1$ system (6.2)–(6.3) with an additional fixed coefficient \mathbf{C}^{-1} . Choosing the initial data $\mathbf{u}_0 = \mathbf{R}_1, \mathbf{u}_1 = \mathbf{1}$ and $\mathbf{u}_2 = \mathbf{R}_2$, with $\mathbf{C}^{-1} = \mathbf{u}_0\mathbf{u}_2^{-1}\mathbf{u}_0^{-1}\mathbf{u}_2$, we may repeat the proof of Appendix B so as to include the fixed coefficient \mathbf{C}^{-1} , and deduce the Laurent positivity property in terms of $(\mathbf{R}_1, \mathbf{R}_2)$.

7. Towards a non-commutative Lindström–Gessel–Viennot theorem

All the cases we have been investigating so far are expressible in terms of non-commutative weighted path models. More precisely, we have been able to represent the generating function $\sum_{n \geq 0} t^n \mathbf{R}_n \mathbf{R}_0^{-1}$ as the partition function for paths with non-commutative weights. In the application of the A_r case to the quantum A_r Q -system of Section 5, we also have variables $\mathbf{R}_{\alpha,n}, n > 1$, expressed as “quantum Wronskians” (see Remark 5.6). This is actually a non-commutative generalization of the “Wronskian” expressions for the Q -system solutions in the commuting case (2.2), which take the form:

$$\frac{R_{\alpha,n+m_1}}{(R_{1,m_1})^\alpha} = \det_{1 \leq i, j \leq \alpha} \left(\frac{R_{1,i+j-\alpha-1+n+m_1}}{R_{1,m_1}} \right).$$

In the cases $\mathbf{m} = \mathbf{m}_0$ or \mathbf{m}_1 for instance (fundamental initial data and Stieltjes point, both with $m_1 = 0$) the Lindström–Gessel–Viennot theorem [25,20] allows to interpret directly the l.h.s. as




Fig. 1. A typical contribution to $\mathbf{P}_{n+1}\mathbf{R}_0\mathbf{C}\mathbf{P}_{n-1}\mathbf{R}_0\mathbf{C}$ for $n = 8$ is made of a pair of paths $A \rightarrow B$ (blue, left to right) and $C \rightarrow D$ (red, right to left), with two intermediate stations at vertex 1, $B \rightarrow C$ and $D \rightarrow A$. When paths intersect (like here), this may also be interpreted as a contribution to $\mathbf{P}_n\mathbf{R}_0\mathbf{P}_n\mathbf{R}_0\mathbf{C}$, upon reversing the direction of travel of the portions of paths to the right of their rightmost intersection X , with a weight \mathbf{R}_0 for the step $C \rightarrow B$. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

the partition function for a family of α non-intersecting lattice paths, namely paths traveled by walkers starting at the root of the graph $\Gamma_{\mathbf{m}}$ at times $0, 2, \dots, 2\alpha - 4, 2\alpha - 2$ and returning at times $2n + 2\alpha - 2, 2n + 2\alpha - 4, \dots, 2n + 2, 2n$, and such that no two walkers meet at any vertex of $\Gamma_{\mathbf{m}}$. In the case of general \mathbf{m} , we have a more subtle application of the theorem, in which the paths must be strongly non-intersecting (see Ref. [7] for details).

It would be extremely interesting to extend the classical Lindström–Gessel–Viennot theorem to the case of paths with non-commutative weights. Such a generalization was introduced in [14] in the context of non-commutative Schur functions, but does not seem to apply directly to the present framework.

So far, we have found such an interpretation only in the fully non-commutative A_1 Q -system case. We wish to interpret the Q -system relation $\mathbf{R}_{n+1}\mathbf{C}\mathbf{R}_{n-1} - \mathbf{R}_n^2 = 1$ in terms of pairs of non-intersecting paths on the graph $\Gamma_{(0)}$, i.e. the integer segment $[1, 4]$. Let $\mathbf{P}_n = \mathbf{R}_n\mathbf{R}_0^{-1}$ denote the partition function of paths from vertex 0 to itself on $\Gamma_{(0)}$ with the non-commutative weights $\mathbf{y}_1, \mathbf{y}_2, \mathbf{y}_3$ respectively for the steps $2 \rightarrow 1, 3 \rightarrow 2$ and $4 \rightarrow 3$, while the up steps $1 \rightarrow 2, 2 \rightarrow 3$ and $3 \rightarrow 4$ receive the trivial weight 1. Multiplying the Q -system relation by \mathbf{C} , we have:

$$\mathbf{P}_{n+1}\mathbf{R}_0\mathbf{C}\mathbf{P}_{n-1}\mathbf{R}_0\mathbf{C} - \mathbf{P}_n\mathbf{R}_0\mathbf{P}_n\mathbf{R}_0\mathbf{C} = \mathbf{C}. \tag{7.1}$$

To proceed, we note the following important relations between the weights $\mathbf{y}_1, \mathbf{y}_2, \mathbf{y}_3$ and \mathbf{R}_0, \mathbf{C} :

$$(\mathbf{y}_3)^n \mathbf{y}_2 \mathbf{y}_1 \mathbf{R}_0 \mathbf{C} (\mathbf{y}_1)^n = \mathbf{y}_2 \mathbf{y}_1 \mathbf{R}_0 \mathbf{C} = \mathbf{R}_0^{-1}, \quad (\mathbf{y}_1)^n \mathbf{R}_0 (\mathbf{y}_3)^n = \mathbf{R}_0 \quad (n \in \mathbb{Z}). \tag{7.2}$$

We now interpret the quantity $\mathbf{P}_{n+1}\mathbf{R}_0\mathbf{C}\mathbf{P}_{n-1}\mathbf{R}_0\mathbf{C}$ as the partition function for a path loop $A \rightarrow B \rightarrow C \rightarrow D \rightarrow A$ (see Fig. 1), where the path $A \rightarrow B$ (partition function \mathbf{P}_{n+1}) is traveled from left to right and assigned the usual descending step weights \mathbf{y}_i for a step $i + 1 \rightarrow i$, while the portions $B \rightarrow C, D \rightarrow A$ receive the weight $\mathbf{R}_0\mathbf{C}$, and the path $C \rightarrow D$ (partition function \mathbf{P}_{n-1}) is traveled from right to left, and receives the usual descending step

weights y_i for a step $i + 1 \rightarrow i$. The contribution for the configuration in the top of Fig. 1 is $y_3^2 y_2^2 y_1 y_3^2 y_2 y_1 \mathbf{R}_0 \mathbf{C} y_1^2 y_3 y_2 y_1^3 \mathbf{R}_0 \mathbf{C}$.

Theorem 7.1. *The quantity*

$$\mathbf{P}_{n+1} \mathbf{R}_0 \mathbf{C} \mathbf{P}_{n-1} \mathbf{R}_0 \mathbf{C} - \mathbf{P}_n \mathbf{R}_0 \mathbf{P}_n \mathbf{R}_0 \mathbf{C} \tag{7.3}$$

is the partition function of loops $A \rightarrow B \rightarrow C \rightarrow D \rightarrow A$ formed by pairs of non-intersecting paths on $[1, 4]$ respectively from $A = (0, 1)$ to $B = (2n + 2, 1)$, then from $C = (2n, 1)$ to $D = (2, 1)$, with the two intermediate stations $B \rightarrow C$ and $D \rightarrow A$ at the root receiving the weight $\mathbf{R}_0 \mathbf{C}$.

Proof. Recall first that the portions of loop from $B \rightarrow C$, and $D \rightarrow A$ receive the weight $\mathbf{R}_0 \mathbf{C}$. We define the weight for the same portions, but traveled in the other direction ($C \rightarrow B$ and $A \rightarrow D$) to be \mathbf{R}_0 . Let us assume the two paths $B \rightarrow C$, and $D \rightarrow A$ intersect, namely that they have at least a common vertex. Let us denote by X the rightmost such common vertex.

We may now decompose the above loop into three pieces: that from $A \rightarrow X$, $X \rightarrow B \rightarrow C \rightarrow X$ and that from $X \rightarrow A$. The central portion $X \rightarrow B \rightarrow C \rightarrow X$, with weight W_{XBCX} , may be “flipped” into $X \rightarrow C \rightarrow B \rightarrow X$, with new weight W_{XCBX} , by reversing the direction of travel, so that the new configuration now corresponds to pairs of paths $A \rightarrow C$ and $B \rightarrow D$ of equal length $2n$, now separated by the two intermediate stations $C \rightarrow B$ and $D \rightarrow A$ with respective weights \mathbf{R}_0 and $\mathbf{R}_0 \mathbf{C}$, thus contributing to $\mathbf{P}_n \mathbf{R}_0 \mathbf{P}_n \mathbf{R}_0 \mathbf{C}$.

We now note that $W_{XCBX} = W_{XBCX}$. Indeed, depending on the height of X , we have the two following possibilities (the first case is illustrated in Fig. 1):

- (i) $X = (2t + 1, 2)$ for some t , then the blue path $X \rightarrow B$ is $3 \rightarrow 3 \rightarrow 4 \rightarrow 3 \rightarrow 4 \rightarrow \dots \rightarrow 4 \rightarrow 3 \rightarrow 2$ (weight $y_3^m y_2 y_1$, $m = n - t$) and the red path $C \rightarrow X$ is $1 \rightarrow 2 \rightarrow 1 \rightarrow 2 \rightarrow \dots \rightarrow 1 \rightarrow 2$ (weight y_1^m). The weight of the central loop $X \rightarrow B \rightarrow C \rightarrow X$ is:

$$W_{XBCX} = y_3^m (y_2 y_1 \mathbf{R}_0 \mathbf{C}) y_1^m = \mathbf{R}_0^{-1} = y_1 \mathbf{R}_0 y_2 = y_1^{m+1} \mathbf{R}_0 y_3^m y_2 = W_{XCBX}$$

by use of (7.2).

- (ii) $X = (2t + 2, 3)$ for some t , then the blue path $X \rightarrow B$ is $3 \rightarrow 4 \rightarrow 3 \rightarrow 4 \rightarrow \dots \rightarrow 4 \rightarrow 3 \rightarrow 2$ (weight $y_3^m y_2 y_1$, $m = n - t$) and the red path $C \rightarrow X$ is $1 \rightarrow 2 \rightarrow 1 \rightarrow 2 \rightarrow \dots \rightarrow 1 \rightarrow 2 \rightarrow 3$ (weight y_1^{m-1}). The weight of the central loop $X \rightarrow B \rightarrow C \rightarrow X$ is:

$$W_{XBCX} = y_3^m (y_2 y_1 \mathbf{R}_0 \mathbf{C}) y_1^{m-1} = y_3 \mathbf{R}_0^{-1} = \mathbf{R}_1^{-1} = y_2 \mathbf{R}_0 = y_2 y_1^m \mathbf{R}_0 y_3^m = W_{XCBX}$$

by use of (7.2).

The above flipping procedure is a bijection between pairs of paths $A \rightarrow B$ and $C \rightarrow D$ that intersect and pairs of paths $A \rightarrow C$ and $B \rightarrow D$. They cancel each other in the formula (7.3), leaving us only with non-intersecting paths, and the theorem follows. \square

Note that there is a unique configuration of non-intersecting paths, where $A \rightarrow B$ is $1 \rightarrow 2 \rightarrow 3 \rightarrow 4 \rightarrow 3 \rightarrow \dots \rightarrow 4 \rightarrow 3 \rightarrow 2 \rightarrow 1$ and $C \rightarrow D$ is $1 \rightarrow 2 \rightarrow 1 \rightarrow 2 \rightarrow \dots \rightarrow 2 \rightarrow 1$. The corresponding weight is $\mathbf{y}_3^{n-1}(\mathbf{y}_2\mathbf{y}_1\mathbf{R}_0\mathbf{C})\mathbf{y}_1^{n-1}\mathbf{R}_0\mathbf{C} = \mathbf{C}$ by (7.2), and we recover (7.1).

8. Conclusion: Towards non-commutative cluster algebras

The rank 2 non-commutative cluster algebras are defined as follows. Let us consider a bipartite chain \mathbb{T}_2 (Cayley tree of degree 2), with alternating black and white vertices indexed by $v \in \mathbb{Z}$ (black if v is even, white otherwise), and edges with alternating labels 1, 2, and a fixed element $\mathbf{C} \in \mathcal{A}$. To each vertex v of \mathbb{T}_2 , we attach a cluster, i.e. a vector of the form $\mathbf{x}_v = (\mathbf{a}, \mathbf{b})$ in \mathcal{A}^2 , where the cluster variables \mathbf{a}, \mathbf{b} obey the quasi-commutation relations

$$\begin{aligned} \mathbf{ab} &= \mathbf{bCa} & \text{if } v \text{ is black,} \\ \mathbf{ba} &= \mathbf{aCb} & \text{if } v \text{ is white.} \end{aligned}$$

To each (doubly oriented) edge labeled i connecting a black and a white vertex, we associate the two following mutations: (i) the forward mutation μ_i^+ corresponding to the oriented edge from the black vertex to the white (ii) the backward mutation $\mu_i^- = (\mu_i^+)^{-1}$ corresponding to the oriented edge from the white vertex to the black. Depending on the color of the vertex v , the forward and backward mutations of $x_v = (\mathbf{a}, \mathbf{b})$ are defined as follows:

$$\begin{aligned} v \text{ black: } & \mu_1^+ : (\mathbf{a}, \mathbf{b}) \mapsto (\mathbf{a}', \mathbf{b}) \quad \text{with} \quad \mathbf{a}'\mathbf{Ca} = (\mathbf{b})^b + \mathbf{1}, \\ & \mu_2^- : (\mathbf{a}, \mathbf{b}) \mapsto (\mathbf{a}, \mathbf{b}') \quad \text{with} \quad \mathbf{bCb}' = (\mathbf{a})^c + \mathbf{1}, \\ v \text{ white: } & \mu_2^+ : (\mathbf{a}, \mathbf{b}) \mapsto (\mathbf{a}, \mathbf{b}') \quad \text{with} \quad \mathbf{b}'\mathbf{Cb} = (\mathbf{a})^c + \mathbf{1}, \\ & \mu_1^- : (\mathbf{a}, \mathbf{b}) \mapsto (\mathbf{a}', \mathbf{b}) \quad \text{with} \quad \mathbf{aCa}' = (\mathbf{b})^b + \mathbf{1}. \end{aligned}$$

These definitions are compatible with an exchange matrix equal to $\mathbf{B}_\bullet = \begin{pmatrix} 0 & -c \\ b & 0 \end{pmatrix}$ for black vertices and $\mathbf{B}_\circ = -\mathbf{B}_\bullet$ for white ones, evolving according to the rules of the usual rank 2 cluster algebras. The general conjecture of [23] may be rephrased as follows: *for any positive integers b, c , the cluster variables at any node of \mathbb{T}_2 may be expressed as positive Laurent polynomials of the cluster variables at any other node of \mathbb{T}_2 .* This is easily proved in the “finite cases” for $bc < 4$ (leading to only finitely many distinct cluster variables in the commuting case), where one can show explicitly that there exists a positive integer $m_{b,c}$ such that $\mathbf{x}_{v+m_{b,c}} = \mathbf{C}\mathbf{x}_v\mathbf{C}^{-1}$, with $m_{1,1} = 5$, $m_{1,2} = 6$, and $m_{1,3} = 8$, leaving us with only finitely many cluster variables to inspect. In the affine case $bc = 4$, this was proved in [10] (see the A_1 case $b = c = 2$ in the introduction, and the case $b = 1, c = 4$ Theorem 6.1 above), while the case $bc > 4$ is still open.

To generalize this construction, we may think of using the results of the present paper in the following way. Assuming that the A_r Q -system cluster algebra introduced in [21] may be generalized to a non-commutative setting, we would expect the relevant cluster variables to still be expressible in terms of non-commutatively weighted paths on target graphs, in terms of given initial data. In this paper, we have shown how to introduce mutations of non-commutative weights via non-commutative continued fractions. Concentrating first on the Stieltjes point, we may trace

the expressions for $\Delta_{1,n} = \mathbf{R}_{1,n}$ back to the initial variables $\Delta_{\alpha,\alpha}$ and $\Delta_{\alpha,\alpha-1}$, $\alpha = 1, 2, \dots, r+1$ (see Theorem 3.17). This seems to indicate, in this case at least, that the quasi-Wronskians are good non-commutative cluster variables. However, the general expressions of Theorem 3.33 for $\mathbf{R}_{1,n}$ in terms of quasi-Wronskians or the more compact C, D variables (see (3.39) and (3.41)) do not clearly display which initial data they are related to. We clearly need another definition of the non-commutative cluster variables.

We hope to report on this issue in a later publication.

Acknowledgments

We would like to thank S. Fomin for discussions, and L. Faddeev for pointing out Refs. [12, 13] to us. We thank the Mathematisches Forschungsinstitut Oberwolfach, Germany, Research in Pairs program (Aug. 2009) during which this work was initiated. P.D.F. also thanks S. Fomin for hospitality at the Dept. of Mathematics of the University of Michigan, Ann Arbor (Spring 2010). R.K. thanks the Institut Henri Poincaré, Paris, France, for hospitality during the semester “Statistical Physics, Combinatorics and Probability” (Fall 2009), and the Institut de Physique Théorique du CEA Saclay, France. P.D.F. received partial support from the ANR Grant GranMa, the ENIGMA research training network MRTN-CT-2004-5652, and the ESF program MISGAM. R.K. is supported by NSF grant DMS-0802511.

Appendix A. Relation of weighted graphs $G_{\mathbf{m}}$ to the graphs $\Gamma_{\mathbf{m}}$ of [6]

In [6], we introduced a family of weighted graphs $\Gamma_{\mathbf{m}}$ constructed from a Motzkin path \mathbf{m} . Here we recall the definition of these graphs, as well as the sequence of bijections which brings them to form of the weighted graphs $G_{\mathbf{m}}$ which we use in the main body of the paper. It is important to understand this connection, because the most basic proof of positivity relies on the fact that the graphs $\Gamma_{\mathbf{m}}$ are positively weighted with weights which are monomials in the initial data $R(\mathbf{m})$.

In fact, we describe a sequence of simple bijections

$$\mathbf{m} \leftrightarrow \Gamma_{\mathbf{m}} \leftrightarrow \Gamma'_{\mathbf{m}} \leftrightarrow G_{\mathbf{m}}.$$

The graph $\Gamma'_{\mathbf{m}}$ is the “compactified” graph of [9]. Note that in general, only the graph $\Gamma_{\mathbf{m}}$ has manifestly positive weights.

Below, let \mathbf{m} be a fixed Motzkin path of length r .


A.1. The graphs $\Gamma_{\mathbf{m}}$

(1) Decompose the Motzkin path \mathbf{m} into a disjoint union of maximal “descending” segments, including those of length one, such that $\mathbf{m} = \bigsqcup_i (m_i, m_{i+1}, \dots, m_{i+k_i-1})$. Here, $m_{i+a} = m_i - a$, $0 \leq a < k$ in the segment beginning with index i and having length k .

Example A.1. If $\mathbf{m} = (2, 1, 2, 2, 3, 2, 1, 1)$ then $\mathbf{m} = (2, 1) \sqcup (2) \sqcup (2) \sqcup (3, 2, 1) \sqcup (1)$.

For notational purposes, we call the maximal descending segment starting with \mathbf{m}_i $\mathbf{m}^{(i)}$, so that $\mathbf{m} = \bigsqcup_i \mathbf{m}^{(i)}$.

(2) To each maximal descending segment $\mathbf{m}^{(i)}$ of length, say, k , is associated a graph $\Gamma_{\mathbf{m}^{(i)}}$. It has $2(k + 1)$ vertices, labeled $(i, i + 1, \dots, i + k)$ and $(i', (i + 1)', \dots, (i + k)')$. It has the following list of oriented edges: $(a, a + 1), (a + 1, a), (a, a'), (a', a)$ (for all a), and (b, a) for all $k + i \geq b > a + 1 \geq 2$. Below we give a picture of the graph $\Gamma_{\mathbf{m}^{(1)}}$ for the example of the descending subpath $\mathbf{m}^{(1)} = (3, 2, 1, 0)$.



(In this picture, an unoriented edge on the graph Γ is considered to be doubly-oriented, for ease of reading.)

(3) Consecutive maximal descending segments are separated by either a “flat” step of the form $m_a = m_{a+1}$, or an “ascending” step, $m_a + 1 = m_{a+1}$.


Example A.2. For the Motzkin path in Example A.1, the step separating $\mathbf{m}^{(1)} = (2, 1)$ and $\mathbf{m}^{(3)} = (2)$ is ascending, and the step separating $\mathbf{m}^{(3)} = (2)$ from $\mathbf{m}^{(4)} = (2)$ is flat.

We “glue” consecutive graphs $\Gamma_1 = \Gamma_{\mathbf{m}^{(i)}}$ and $\Gamma_2 = \Gamma_{\mathbf{m}^{(j)}}$ where $m^{(i)}$ and $m^{(j)}$ are neighboring descending segments in \mathbf{m} with $j > i$. The gluing procedure is dictated by whether the segments are separated by an ascending or flat step. We assume $\mathbf{m}^{(i)}$ has length k_i and $\mathbf{m}^{(j)}$ has length k_j .


- (a) If the separation between the segments is flat, we identify the edge and vertices $(j', j) \in \Gamma_2$ with $((k_i + i)', k_i + i) \in \Gamma_1$.
- (b) If the separation between the segments is “ascending”, we identify the vertices and edges $(j, j') \in \Gamma_2$ with $((k_i + i)', k_i + i) \in \Gamma_1$ (i.e., the identification is in the opposite sense).

We define $\Gamma_{\mathbf{m}}$ to be the result of gluing the graphs corresponding to consecutive strictly descending pieces of \mathbf{m} according to the procedure given in (4) above.

Example A.3. The purely ascending Motzkin path $\mathbf{m} = \mathbf{m}_1 = (0, 1, \dots, k)$ maps to the chain with $2k + 2$ vertices, connected by doubly-oriented edges. The flat Motzkin path $\mathbf{m}_0 = (0, \dots, 0)$ of length k maps to the graph



As an example of the gluing procedure, in the case of $r = 3$ and $\mathbf{m} = (0, 1, 0)$, we have



$\Gamma_{\mathbf{m}}$ has a root vertex, which is the vertex $1'$ in the picture of $\Gamma_{\mathbf{m}_0}$, or the bottom-most vertex in $\Gamma_{\mathbf{m}}$ in general.

The weights attached to the graph $\Gamma_{\mathbf{m}}$ are as follows. Each edge connecting neighboring vertices and pointing towards the root vertex $1'$ carries weight ty_a , with the edges numbered $a = 1, 2, \dots, 2r + 1$ from the root vertex on (with edge (i', i) preceding $(i + 1, i)$ in cases where it exists). The weights y_a are called the *skeleton weights* of $\Gamma_{\mathbf{m}}$.

Let $w_{a,b}$ be the weight corresponding to the edge connecting vertex a to b . Then $w_{a,a+1} = w_{a,a'} = 1$, $w_{a+1,a} = ty_{2a}$ and $w_{a',a} = ty_{2a-1}$. The weights $w_{a,b}$ of the edges (a, b) with $(a - b > 1)$ are called *redundant weights*, and are expressed solely in terms of skeleton weights between the vertices a, b as:

$$w_{a,b} = t \prod_{i=b+1}^a y_{2i-2} \prod_{i=b+1}^{a-1} y_{2i-1}^{-1} \quad (a - b \geq 1). \tag{A.1}$$

These graphs have manifestly positive weights. Let \mathbf{m} be a Motzkin path with first component m_1 . Then:

Theorem A.4. (See [6].) *The ratio $R_{1,n+m_1}/R_{1,m_1}$ is the coefficient of t^n in the partition function of paths on $\Gamma_{\mathbf{m}}$ from the vertex $1'$ to itself.*

This theorem can be stated in terms of the resolvent of the transfer matrix associated with the graph in the following way. Let $T_{\Gamma_{\mathbf{m}}}$ be the matrix with entries $(T_{\Gamma_{\mathbf{m}}})_{a,b} = w_{a,b}$. Then the generating function for paths from vertex $1'$ to itself on $\Gamma_{\mathbf{m}}$ is the resolvent

$$F_{\mathbf{m}}(t) = ((\mathbb{I} - T_{\Gamma_{\mathbf{m}}})^{-1})_{1',1'} = \sum_{n \geq 0} t^n R_{1,n+m_1} / R_{1,m_1}. \tag{A.2}$$

A.2. Compactified graphs

We now define the graphs $\Gamma'_{\mathbf{m}}$, which have the property that

$$((\mathbb{I} - T_{\Gamma_{\mathbf{m}}})^{-1})_{1,1} = ((\mathbb{I} - T_{\Gamma'_{\mathbf{m}}})^{-1})_{1,1}.$$

Note that $Z_{1',1'}(t) = 1 + ty_1((\mathbb{I} - T_{\Gamma'_{\mathbf{m}}})^{-1})_{1,1}$.

The compactified graphs $\Gamma'_{\mathbf{m}}$ are obtained from $\Gamma_{\mathbf{m}}$ as explained in Section 6 of [9]. This process involves an identification of pairs of vertices in the graph $\Gamma_{\mathbf{m}}$, such that the final number of vertices is reduced to $r + 1$. This produces loops and also, in certain cases, extra directed edges.

- (1) For subgraphs arising from descending or flat segments of \mathbf{m} , compactified graphs are obtained simply from identification of the vertices i and i' for each i , resulting in a loop at vertex i with weight $w_{i',i}$. All other edges retain their weights.
- (2) Consider the vertices and edges arising from each ascending segment of length k in \mathbf{m} . This is a chain of $2k + 2$ vertices.
 - (a) Name the vertices from bottom to top, $1, \dots, 2k + 2$ and identify the vertices $2a - 1, 2a$ ($a = 1, \dots, k + 1$). This leaves $k + 1$ vertices.
 - (b) Rename the vertices from bottom to top again, $1, \dots, k + 1$.
 - (c) Identification of vertices results in a loop at each vertex a with weight inherited from the weight associated with the edge connecting $2a$ to $2a - 1$, $w_{2a,2a-1}$. The spine retains weights $w_{2a+1,2a}$ along each edge $(a + 1, a)$.
 - (d) In addition we adjoin ascending directed edges, connecting vertex a to b for all $1 \leq a < b \leq k + 1$, with weight $(-1)^{b-a+1}$.

Remark A.5. In fact, the construction of the compactified graph directly from the Motzkin path is easy to describe. The Motzkin path is decomposed into maximal ascending, descending and flat segments, the corresponding compactified graphs and their weights are constructed as above, and the graphs are glued: The top two vertices corresponding to a segment are identified with the bottom two vertices corresponding to the next segment. Incident edges between these vertices are identified. (See pictures below.)

Lemma A.6. (See [9].) *We have the equality of partition functions*

$$((\mathbb{I} - T_{\Gamma_{\mathbf{m}}})^{-1})_{1,1} = ((\mathbb{I} - T_{\Gamma'_{\mathbf{m}}})^{-1})_{1,1}.$$

In light of Remark A.5, we can focus on graph segments corresponding to segments of \mathbf{m} which do not change direction.

A.3. The graph $G_{\mathbf{m}}$

We now turn to the equivalence of resolvents of the graph $G_{\mathbf{m}}$, defined in Section 2, to those on the graph $\Gamma'_{\mathbf{m}}$. The difference is the presence of long directed edges in $\Gamma'_{\mathbf{m}}$ (connecting vertices along the spine of the graph with indices which differ by 2 or more).

On the level of the transfer matrix $T_{\Gamma'_{\mathbf{m}}}$, these edges are represented by non-zero elements above or below the second diagonals. These entries can be eliminated using row (column) elimination for descending (ascending) edges, respectively. Conveniently,

Lemma A.7. *The matrices E and F such that $E(\mathbb{I} - T_{\Gamma'_{\mathbf{m}}})F$ has no non-trivial entries away from the second diagonals are such that E is upper unitriangular and F is lower unitriangular.*

The $(1, 1)$ -resolvent

$$((\mathbb{I} - T_{\Gamma'_{\mathbf{m}}})^{-1})_{1,1}$$

is obviously unchanged under such transformations.

Proof. It is sufficient to show this for each block of $T' = T_{\Gamma'_{\mathbf{m}}}$ corresponding to maximal descending and maximal ascending pieces of \mathbf{m} . The case of flat pieces is trivial as it has no entries off the second diagonals.

(1) *Descending pieces.* The matrix $\mathbb{I} - T'$ related to the vertices in the graph $\Gamma'_{\mathbf{m}}$ corresponding to a descending path $(k - 1, k - 2, \dots, 0)$ is the $(k + 1) \times (k + 1)$ -matrix

$$\mathbb{I} - T' = \begin{pmatrix} 1 - x_1 & -1 & 0 & \cdots & & 0 \\ -x_2 & 1 - x_3 & -1 & 0 & \cdots & \\ -\frac{x_2x_4}{x_3} & -x_4 & 1 - x_5 & -1 & 0 & \\ -\frac{x_2x_4x_6}{x_3x_5} & -\frac{x_4x_6}{x_5} & -x_6 & 1 - x_7 & -1 & \vdots \\ \vdots & & \ddots & \ddots & \ddots & \\ \vdots & \cdots & & -x_{2k-2} & 1 - x_{2k-1} & -1 \\ \vdots & \cdots & & -\frac{x_{2k-2}x_{2k}}{x_{2k-1}} & -x_{2k} & 1 - x_{2k+1} \end{pmatrix}.$$

Here, $x_{2a-1} = w_{a',a}$ and $x_{2a} = w_{a+1,a}$. Clearly, there is a lower triangular matrix which eliminates all of the entries below the second diagonal when multiplied on the right:

$$F_{i,i} = 1 \quad (1 \leq i \leq k + 1), \quad F_{i+1,i} = -\frac{x_{2i}}{x_{2i+1}} \quad (1 \leq k - 1), \quad F_{i,j} = 0 \quad \text{otherwise.}$$

The resulting matrix $M = (\mathbb{I} - T')F$ has non-vanishing entries $M_{i,i+1} = -1$, and for $i \leq k - 1$,

$$M_{ii} = 1 - \left(x_{2i-1} - \frac{x_{2i}}{x_{2i+1}} \right), \quad M_{i+1,i} = -\frac{x_{2i}}{x_{2i+1}}. \tag{A.3}$$

In addition, the last two columns of the matrix M are those of $\mathbb{I} - T'$.

Note that we have lost the positivity property of the weights in this case, although the generating function of paths is positive. Note also that the weights attached to the edges connecting the last two vertices have not changed under this transformation.

- (2) *Ascending pieces.* The matrix T' , attached to the vertices in the graph Γ'_m corresponding to an ascending path $(0, \dots, k - 1)$ is the $(k + 1) \times (k + 1)$ -matrix

$$\mathbb{I} - T' = \begin{pmatrix} 1 - x_1 & -1 & 1 & -1 & 1 & \dots & \pm 1 \\ -x_2 & 1 - x_3 & -1 & 1 & -1 & \dots & \\ 0 & -x_4 & 1 - x_5 & -1 & 1 & & \\ \vdots & 0 & -x_6 & 1 - x_7 & -1 & \dots & \\ & & \ddots & \ddots & \ddots & \ddots & \vdots \\ \vdots & & & & -x_{2k-2} & 1 - x_{2k-1} & -1 \\ 0 & \dots & & & 0 & -x_{2k} & 1 - x_{2k+1} \end{pmatrix}.$$


There is obviously an upper triangular matrix E , corresponding to row operations on this matrix, or multiplication on the left, such that $M = E(\mathbb{I} - T')$ has no entries away from the second off-diagonals. The resolvent is equal to the resolvent of the matrix with non-vanishing entries $M_{i,i+1} = -1$, and if $i \leq k - 1$,

$$M_{ii} = 1 - (x_i + x_{i+1}), \quad M_{i+1,i} = -x_{2i}x_{2i+1}. \tag{A.4}$$


The last two rows of M are those of $\mathbb{I} - T'$. Note that in this case, we retain the positivity property of the weights. \square

To summarize, we can draw the graphs corresponding to the sequence of transformations from \mathbf{m} (an ascending, descending or flat path) to G_m in graphical notation:


- (1) For ascending paths,



(2) For descending paths,



(3) For flat paths,



In the final graphs $G_{\mathbf{m}}$ for ascending and descending Motzkin paths, the symbols \times and $=$ along certain edges indicates that the weights have been renormalized from the initial values x_{2i-1} and x_{2i} to the values $x_{2i-1} - x_{2i}/x_{2i+1}$ and x_{2i}/x_{2i+1} for $=$ (see Eq. (A.3)) and to the values $x_{2i-1} + x_{2i}$ and $x_{2i}x_{2i+1}$ for \times (see Eq. (A.4)). For $x_i = ty_i$ with the appropriate labeling of skeleton weights, these are precisely the weights \hat{y} of Definition 2.1 of Section 2.

Gluing two path pieces, \mathbf{m} followed by \mathbf{m}' into a longer Motzkin path corresponds to identifying the two top vertices (and incident edges) of $G_{\mathbf{m}}$, the lower graph, with the bottom two of $G_{\mathbf{m}'}$, the top graph. In the cases where the weights have been renormalized in the lower part of the upper path, the renormalized weights appear in the glued graph.

Appendix B. Proof of Theorem 6.2

Due to the invariances of the system (6.2)–(6.3), we only need to express the solutions \mathbf{u}_n in terms of the initial data $\mathbf{x}_0 = (\mathbf{u}_0, \mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_{2k})$. Indeed, introducing the anti-automorphism φ such that $\varphi(\mathbf{1}) = \mathbf{1}$, $\varphi(ab) = \varphi(b)\varphi(a)$ for all $a, b \in \mathcal{A}$, and $\varphi(\mathbf{u}_i) = \mathbf{u}_{i+1}$, for $i = 0, 1, \dots, 2k$, we easily find that $\varphi(\mathbf{u}_n) = \mathbf{u}_{n+1}$ for all $n \in \mathbb{Z}$. So if we have an expression $\mathbf{u}_n = f_n(\mathbf{u}_0, \mathbf{u}_1, \dots, \mathbf{u}_{2k})$ for the general solution of (6.2)–(6.3) in terms of the initial data \mathbf{x}_0 , we may apply φ iteratively on it to get $\mathbf{u}_n = \varphi^i(f_{n-i}(\mathbf{u}_0, \mathbf{u}_1, \dots, \mathbf{u}_{2k})) = g_{n,i}(\mathbf{u}_i, \mathbf{u}_{i+1}, \dots, \mathbf{u}_{i+2k})$,

namely an expression for \mathbf{u}_n in terms of any other initial data \mathbf{x}_i , for all $n, i \in \mathbb{Z}$. We may also restrict ourselves to \mathbf{u}_n with $n \in \mathbb{Z}_+$. Indeed, let ψ be the anti-automorphism such that $\psi(\mathbf{u}_i) = \mathbf{u}_{2k-1-i}$, $i = 0, 1, \dots, 2k$, then we have $\psi(\mathbf{u}_n) = \mathbf{u}_{2k-1-n}$ for all $n \in \mathbb{Z}$. Hence if we have an expression $\mathbf{u}_n = f_n(\mathbf{u}_0, \mathbf{u}_1, \dots, \mathbf{u}_{2k})$ for $n \geq 2k + 1$, then applying $\varphi\psi$ to it gives $\mathbf{u}_{2k-n} = h_n(\mathbf{u}_0, \mathbf{u}_1, \dots, \mathbf{u}_{2k})$, namely an expression for \mathbf{u}_n , $n \leq 0$, in terms of \mathbf{x}_0 . Moreover all of the above expressions are positive Laurent polynomials iff f_n is a Laurent polynomial for $n \geq 0$.

By the above remarks, we only need prove the statement of the theorem for $n \geq 0$ and $i = 0$. We first need the following lemma, expressing the integrability of the system (6.2)–(6.3).

Lemma B.1. *There exist two linear recursion relations of the form:*

$$\begin{aligned} \mathbf{u}_{2n+2k+1} - \mathbf{u}_{2n+1}\mathbf{K} + \mathbf{u}_{2n-2k+1} &= 0, \\ \mathbf{u}_{2n+2k} - \mathbf{K}\mathbf{u}_{2n} + \mathbf{u}_{2n-2k} &= 0 \end{aligned}$$

for all $n \in \mathbb{Z}$, and with K expressed in terms of \mathbf{x}_0 as:

$$\mathbf{K} = \mathbf{u}_0\mathbf{u}_{2k}^{-1} + \mathbf{u}_{2k}\mathbf{u}_0^{-1} + \sum_{j=1}^k \mathbf{u}_{2j-1}^{-1}(\mathbf{u}_{2j}^{-1} + \mathbf{u}_{2j+2}^{-1}). \tag{B.1}$$

Proof. Introduce

$$\mathbf{K}_n = \mathbf{u}_{2n+1}^{-1}(\mathbf{u}_{2n+2k+1} + \mathbf{u}_{2n-2k+1}), \quad \mathbf{L}_n = (\mathbf{u}_{2n+2k} + \mathbf{u}_{2n-2k})\mathbf{u}_{2n}^{-1}.$$

Subtracting (6.2) for $n \rightarrow n - k$ from that for n , we get $(\mathbf{u}_{2n+2k+1} + \mathbf{u}_{2n-2k+1})\mathbf{u}_{2n} = \mathbf{u}_{2n+1}(\mathbf{u}_{2n-2k} + \mathbf{u}_{2n+2k})$, hence $\mathbf{K}_n = \mathbf{L}_n$. Moreover,

$$\begin{aligned} \mathbf{u}_{2n+1}\mathbf{K}_n\mathbf{u}_{2n+2} &= \mathbf{u}_{2n+1}\mathbf{u}_{2n+2k+2} - \mathbf{1} + \mathbf{u}_{2n-2k+1}\mathbf{u}_{2n+2} \\ &= \mathbf{u}_{2n+1}(\mathbf{u}_{2n+2k+2} + \mathbf{u}_{2n-2k+2}) = \mathbf{u}_{2n+1}\mathbf{L}_{n+1}\mathbf{u}_{2n+2} \end{aligned}$$

hence $\mathbf{K}_n = \mathbf{L}_{n+1} = \mathbf{K}_{n+1} = \mathbf{K}$ is a conserved quantity of the system (6.2)–(6.3). Eq. (B.1) is easily derived by induction on k , and the lemma follows. \square

Note that $\varphi(\mathbf{L}_n) = \mathbf{K}_n$, hence $\varphi(\mathbf{K}) = \mathbf{K}$. Analogously, $\psi(\mathbf{L}_n) = \mathbf{K}_{k-1-n}$, hence $\psi(\mathbf{K}) = \mathbf{K}$ as well.

Introducing the generating functions

$$\mathbf{F}_i(t) = \sum_{n=0}^{\infty} t^n \mathbf{u}_{2i+2kn}, \quad \mathbf{G}_i(t) = \sum_{n=0}^{\infty} t^n \mathbf{u}_{2i+1+2kn}$$

for $i = 0, 1, \dots, k - 1$ and the weights:

$$\begin{aligned} \mathbf{x}_{2i} &= \mathbf{u}_{2i+2k}\mathbf{u}_{2i}^{-1}, \\ \mathbf{z}_{2i} &= \mathbf{K} - \mathbf{x}_{2i} = \mathbf{u}_{2i-2k}\mathbf{u}_{2i}^{-1}, \\ \mathbf{w}_{2i} &= \mathbf{z}_{2i}\mathbf{x}_{2i} - \mathbf{1}, \end{aligned}$$

$$\begin{aligned} \mathbf{x}_{2i+1} &= \mathbf{u}_{2i+1}^{-1} \mathbf{u}_{2i+1+2k}, \\ \mathbf{z}_{2i+1} &= \mathbf{K} - \mathbf{x}_{2i+1} = \mathbf{u}_{2i+1}^{-1} \mathbf{u}_{2i+1-2k}, \\ \mathbf{w}_{2i+1} &= \mathbf{x}_{2i+1} \mathbf{z}_{2i+1} - \mathbf{1}. \end{aligned}$$

Lemma B.1 implies the following continued fraction expressions:

$$\begin{aligned} \mathbf{F}_i(t) &= (\mathbf{1} - (\mathbf{x}_{2i} + \mathbf{z}_{2i})t + t^2)^{-1} (\mathbf{1} - \mathbf{z}_{2i}t) \mathbf{u}_{2i} \\ &= (\mathbf{1} - \mathbf{x}_{2i}t - (\mathbf{1} - \mathbf{z}_{2i}t)^{-1} \mathbf{w}_{2i}t^2)^{-1} \mathbf{u}_{2i}, \\ \mathbf{G}_i(t) &= \mathbf{u}_{2i+1} (\mathbf{1} - \mathbf{z}_{2i+1}t) (\mathbf{1} - (\mathbf{x}_{2i+1} + \mathbf{z}_{2i+1})t + t^2)^{-1} \\ &= \mathbf{u}_{2i+1} (\mathbf{1} - t\mathbf{x}_{2i+1} - t^2 \mathbf{w}_{2i+1} (\mathbf{1} - \mathbf{z}_{2i+1}t)^{-1})^{-1} \end{aligned}$$

whose series expansions in t have coefficients that are manifestly positive Laurent polynomials of $(\mathbf{x}_i, \mathbf{z}_i, \mathbf{w}_i)$. Finally, we see that the \mathbf{x} 's are defined recursively by $\mathbf{x}_0 = \mathbf{u}_{2k} \mathbf{u}_0^{-1}$ and $\mathbf{x}_{2j} = \mathbf{x}_{2j-1} + \mathbf{u}_{2j-1}^{-1} \mathbf{u}_{2j}^{-1}$, $\mathbf{x}_{2j+1} = \mathbf{x}_{2j} + \mathbf{u}_{2j+1}^{-1} \mathbf{u}_{2j}^{-1}$, hence they form subsums of \mathbf{K} , and so do the \mathbf{z} 's. Also, all \mathbf{x} 's contain the term $\mathbf{u}_{2k} \mathbf{u}_0^{-1}$, while all \mathbf{z} 's contain $\mathbf{u}_0 \mathbf{u}_{2k}^{-1}$, hence both $\mathbf{x}_i \mathbf{z}_i$ and $\mathbf{z}_i \mathbf{x}_i$ contain $\mathbf{1}$: we conclude that the $\mathbf{x}, \mathbf{z}, \mathbf{w}$'s are all positive Laurent polynomials of \mathbf{x}_0 . This completes the proof of Theorem 6.2.

The existence of the linear recursion relations of Lemma B.1 implies by Proposition 3.13 that all 3×3 quasi-Wronskians $\mathbf{W}_{2kn+i} = |(\mathbf{u}_{2kn+i-2a-2b+8})_{1 \leq a, b \leq 3}|_{1,1}$ must vanish. Each function $\mathbf{F}_i(t)$, $i = 0, 1, \dots, k - 1$, is actually a particular instance of the general non-commutative case with $r = 1$, discussed in Section 3. It is the generating function of paths on the graph G_1 , with respective weights $\mathbf{w}_{1,2} = \mathbf{1}$ and:

$$(\mathbf{w}_{1,1}, \mathbf{w}_{2,1}, \mathbf{w}_{2,2}) = (t(\mathbf{y}_1 + \mathbf{y}_2), t^2 \mathbf{y}_3 \mathbf{y}_2, t \mathbf{y}_3) = (t \mathbf{x}_{2i}, t^2 \mathbf{w}_{2i}, t \mathbf{z}_{2i}).$$

As we also have a condition $\mathbf{w}_{2i} = \mathbf{z}_{2i} \mathbf{x}_{2i} - \mathbf{1}$, we deduce that the conserved quantity $C_{2,n} = \mathbf{C} = \mathbf{y}_3 \mathbf{y}_1 = \mathbf{1}$. This provides us with examples in the general A_1 setting that are different from the non-commutative A_1 Q -system of Section 1.3. In the latter indeed, \mathbf{C} is kept arbitrary, but the value of $\Delta_{2,n} = \mathbf{R}_{n+1} - \mathbf{R}_n \mathbf{R}_{n-1}^{-1} \mathbf{R}_n$ is fixed by the Q -system relation $\Delta_{2,n} = \mathbf{R}_n^{-1} \mathbf{R}_{n-1} \mathbf{R}_n$. Here we have $\mathbf{C} = \mathbf{1}$, hence the recursion relation $\Delta_{2,n+1} = \mathbf{R}_{n-1} \mathbf{R}_n^{-1} \Delta_{2,n}$ (with $\mathbf{R}_n = \mathbf{u}_{2i+2kn}$), but there is no such additional relation. Note that $\mathbf{C} = \mathbf{1}$ also implies that there is no quantum version in which \mathbf{C} would be a non-trivial central element of \mathcal{A} . This is in agreement with the above-mentioned fact that the exchange matrix of the associated cluster algebra is singular.

References

[1] I. Assem, C. Reutenauer, D. Smith, Frises, preprint, arXiv:0906.2026 [math.RA].
 [2] V. Bazhanov, N. Reshetikhin, Restricted solid-on-solid models connected with simply laced algebras and conformal field theory, *J. Phys. A* 23 (1990) 1477–1492.
 [3] A. Berenstein, A. Zelevinsky, Quantum cluster algebras, *Adv. Math.* 195 (2005) 405–455, arXiv:math/0404446 [math.QA].
 [4] P. Caldero, A. Zelevinsky, Laurent expansions in cluster algebras via quiver representations, *Mosc. Math. J.* 6 (3) (2006) 411–429, arXiv:math/0604054 [math.RT].
 [5] P. Di Francesco, The solution of the A_r T -system with arbitrary boundary, preprint, arXiv:1002.4427 [math.CO], 2010.

- [6] P. Di Francesco, R. Kedem, Q -systems as cluster algebras II: Cartan matrix of finite type and the polynomial property, *Lett. Math. Phys.* 89 (3) (2009) 183–216, arXiv:0803.0362 [math.RT].
- [7] P. Di Francesco, R. Kedem, Q -systems, heaps, paths and cluster positivity, *Comm. Math. Phys.* 293 (3) (2009) 727–802, doi:10.1007/s00220-009-0947-5, arXiv:0811.3027 [math.CO].
- [8] P. Di Francesco, R. Kedem, Positivity of the T -system cluster algebra, *Electron. J. Combin.* 16 (1) (2009) R140, Oberwolfach preprint OWP 2009-21, arXiv:0908.3122 [math.CO].
- [9] P. Di Francesco, R. Kedem, Q -system cluster algebras, paths and total positivity, *SIGMA Symmetry Integrability Geom. Methods Appl.* 6 (014) (2010), 36 pp., arXiv:0906.3421 [math.CO].
- [10] P. Di Francesco, R. Kedem, Discrete non-commutative integrability: proof of a conjecture by M. Kontsevich, *Int. Math. Res. Not. IMRN* (2010), doi:10.1093/imrn/rnq024, arXiv:0909.0615 [math-ph].
- [11] P. Etingof, V. Retakh, Quantum determinants and quasideterminants, *Asian J. Math.* 3 (1999) 345–351.
- [12] L.D. Faddeev, R.M. Kashaev, A.Y. Volkov, Strongly coupled quantum discrete Liouville theory, I. Algebraic approach and duality, *Comm. Math. Phys.* 219 (2001) 199–219, arXiv:hep-th/0006156.
- [13] L.D. Faddeev, A.Y. Volkov, Discrete evolution for the zero modes of the quantum Liouville model, *J. Phys. A* 41 (19) (2008) 194008, 12 pp., arXiv:0803.0230 [hep-th].
- [14] S. Fomin, C. Green, Noncommutative Schur functions and their applications, *Discrete Math.* 306 (10–11) (2006) 1080–1096.
- [15] S. Fomin, A. Zelevinsky, Cluster algebras, I, *J. Amer. Math. Soc.* 15 (2) (2002) 497–529, arXiv:math/0104151 [math.RT].
- [16] S. Fomin, A. Zelevinsky, The Laurent phenomenon, *Adv. in Appl. Math.* 28 (2) (2002) 119–144, arXiv:math/0104241 [math.CO].
- [17] E. Frenkel, N. Reshetikhin, The q -characters of representations of quantum affine algebras and deformations of W -algebras, in: *Recent Developments in Quantum Affine Algebras and Related Topics*, Raleigh, NC, 1998, in: *Contemp. Math.*, vol. 248, 1999, pp. 163–205.
- [18] I. Gelfand, S. Gelfand, V. Retakh, R.L. Wilson, Quasideterminants, *Adv. Math.* 193 (1) (2005) 56–141, arXiv:math/0208146v4 [math.QA].
- [19] I. Gelfand, D. Krob, A. Lascoux, B. Leclerc, V. Retakh, J.-Y. Thibon, Noncommutative symmetric functions, *Adv. Math.* 112 (2) (1995) 218–348, arXiv:hep-th/9407124.
- [20] I.M. Gessel, X. Viennot, Binomial determinants, paths and hook formulae, *Adv. Math.* 58 (1985) 300–321.
- [21] R. Kedem, Q -systems as cluster algebras, *J. Phys. A* 41 (2008) 194011, 14 pp., arXiv:0712.2695 [math.RT].
- [22] A.N. Kirillov, N.Yu. Reshetikhin, Representations of Yangians and multiplicity of occurrence of the irreducible components of the tensor product of representations of simple Lie algebras, *J. Sov. Math.* 52 (1990) 3156–3164.
- [23] M. Kontsevich, private communication.
- [24] A. Kuniba, T. Nakanishi, J. Suzuki, Functional relations in solvable lattice models, I. Functional relations and representation theory, *Internat. J. Modern Phys. A* 9 (1994) 5215–5266.
- [25] B. Lindström, On the vector representations of induced matroids, *Bull. Lond. Math. Soc.* 5 (1973) 85–90.
- [26] G. Musiker, J. Propp, Combinatorial interpretations for rank-two cluster algebras of affine type, *Electron. J. Combin.* 14 (2007), R15, arXiv:math/0602408.
- [27] H. Nakajima, t -Analogues of q -characters of Kirillov–Reshetikhin modules of quantum affine algebras, *Represent. Theory* 7 (2003) 259–274.
- [28] P. Sherman, A. Zelevinsky, Positivity and canonical bases in rank 2 cluster algebras of finite and affine types, *Mosc. Math. J.* 4 (4) (2004) 947–974, arXiv:math/0307082 [math.RT].