

CtBP1 Overexpression in Keratinocytes Perturbs Skin Homeostasis

Hui Deng^{1,2,3,6}, Fulun Li^{2,4,6}, Hong Li^{1,6}, Yu Deng¹, Jing Liu¹, Donna Wang², Gangwen Han², Xiao-Jing Wang^{1,2} and Qinghong Zhang^{1,2,5}

Carboxyl-terminal-binding protein-1 (CtBP1) is a transcriptional corepressor with multiple *in vitro* targets, but its *in vivo* functions are largely unknown. We generated keratinocyte-specific CtBP1 transgenic mice with a keratin-5 promoter (K5.CtBP1) to probe the pathological roles of CtBP1. At transgene expression levels comparable to endogenous CtBP1 in acute skin wounds, the K5.CtBP1 epidermis displayed hyperproliferation, loss of E-cadherin, and failed terminal differentiation. Known CtBP1 target genes associated with these processes, e.g., *p21*, *Brca1*, and *E-cadherin*, were downregulated in K5.CtBP1 skin. Surprisingly, K5.CtBP1 pups also exhibited a hair loss phenotype. We found that expression of the *Distal-less 3 (Dlx3)*, a critical regulator of hair follicle differentiation and cycling, was decreased in K5.CtBP1 mice. Molecular studies revealed that CtBP1 directly suppressed *Dlx3* transcription. Consistently, K5.CtBP1 mice displayed abnormal hair follicles with decreased expression of *Dlx3* downstream targets *Gata3*, *Hoxc13*, and *hair keratins*. In summary, this CtBP1 transgenic model provides *in vivo* evidence for certain CtBP1 functions predicted from *in vitro* studies, reveals—to our knowledge—previously unreported functions and transcriptional activities of CtBP1 in the context of epithelial-mesenchymal interplay, and suggests that CtBP1 has a pathogenic role in hair follicle morphogenesis and differentiation.

Journal of Investigative Dermatology (2014) **134**, 1323–1331; doi:10.1038/jid.2013.504; published online 2 January 2014

INTRODUCTION

CtBP was originally identified on the basis of its ability to bind with the carboxyl terminus of the E1A oncoprotein (Boyd *et al.*, 1993; Schaeper *et al.*, 1995). Subsequently, carboxyl-terminal-binding protein (CtBP) was found as a transcriptional corepressor involved in a variety of biological processes, including proliferation and anti-apoptosis (Chinnadurai, 2002). CtBP indirectly binds DNA with various DNA-binding partners at multiple DNA sequences; thus, CtBP-mediated transcriptional repression is context-specific. For instance, CtBP represses *E-cadherin* in epithelial cells (Grootclaes and Frisch, 2000; Grootclaes *et al.*, 2003; Zhang *et al.*, 2006), *IL-4* in human T cells (Kitamura *et al.*,

2009), and *dll4*, *sprouty*, and *ve-cadherin* for endothelial sprouting (Roukens *et al.*, 2010). Recently, we and others found that CtBP downregulates DNA damage repair by directly suppressing the transcription of *breast cancer type 1 susceptibility protein (Brca1)* in cancer cells (Deng *et al.*, 2010; Di *et al.*, 2010).

In mammals, there are two isoforms, carboxyl-terminal-binding protein-1 (CtBP1) and CtBP2. Both isoforms are expressed in the wild-type mouse embryo and have overlapping and unique roles (Hildebrand and Soriano, 2002). CtBP1 knockout mice are small and 23% die of an unknown cause by 20 days postpartum. CtBP2-null mice are small in size, have axial truncations, delayed neural, muscular, and skeletal development, and defects in heart morphogenesis; they die by E10.5 because of defects in both yolk sac and placental vascularization. In most human and mouse adult tissues, CtBP expression is low. Reactivation of CtBP expression has been shown in pathological conditions, e.g., cancer (Nadauld *et al.*, 2006; Deng *et al.*, 2010), but its *in vivo* role in adult tissue is virtually unknown.

To determine the effects of aberrant CtBP1 expression in keratinocytes *in vivo*, we targeted CtBP1 overexpression to the basal layer of the epidermis and hair follicle using a keratin-5 promoter (K5.CtBP1) (He *et al.*, 2002). The K5.CtBP1 transgenic epidermis displayed loss of E-cadherin, hyperproliferation, and decreased differentiation. Unexpectedly, K5.CtBP1 mice also exhibited defective hair morphogenesis starting at postnatal stage. Molecular analyses revealed that CtBP1 directly repressed transcription of the *Distal-less 3 (Dlx3)* gene, a

¹Department of Dermatology, University of Colorado Denver, Aurora, Colorado, USA; ²Department of Pathology, University of Colorado Denver, Aurora, Colorado, USA; ³Department of Dermatology, The Sixth People's Hospital of Shanghai, Shanghai Jiaotong University, Shanghai, China; ⁴Department of Dermatology, Yueyang Hospital Affiliated to Shanghai University of TCM, Shanghai, China and ⁵Department of Biochemistry and Molecular Genetics, University of Colorado Denver, Aurora, Colorado, USA
⁶Co-first authors.

Correspondence: Xiao-Jing Wang, Department of Pathology, University of Colorado Denver, Aurora, Colorado 80045, USA.
E-mail: xj.wang@UCDenver.edu or Qinghong Zhang, Department of Dermatology, University of Colorado Denver, Aurora, Colorado 80045, USA.
E-mail: Qinghong.Zhang@UCDenver.edu

Abbreviations: *Brca1*, breast cancer type 1 susceptibility protein; CtBP, carboxyl-terminal-binding protein; *Dlx3*, *Distal-less 3*

Received 22 April 2013; revised 24 September 2013; accepted 8 October 2013; accepted article preview online 26 November 2013; published online 2 January 2014

homeobox transcription factor that has a critical role in hair development by orchestrating the differentiation of the inner root sheath and hair shaft (Hwang *et al.*, 2008). Supporting this link, we found a hair follicle formation defect in K5.CtBP1 mice, with decreased expression of *Dlx3* and its target genes. Our study provides *in vivo* model for CtBP1 overexpression and reveals that CtBP1 overexpression perturbs epidermal and hair follicle homeostasis.

RESULTS

Generation of CtBP1 transgenic mice

In normal mouse skin, CtBP1 is barely detectable. Acute skin wound by punch biopsy induced CtBP1 expression ~6-fold higher than in non-wounded skin (Figure 1a). To evaluate the role of CtBP1 overexpression in the skin, we generated K5.CtBP1 transgenic mice by inserting human CtBP1 cDNA (99% amino acid homology to mouse CtBP1 protein) into a K5 vector (He *et al.*, 2002). Three K5.CtBP1 transgenic founders

(K1, K2, and K3) were generated. Their CtBP1 transgene expression levels were 7–10-fold higher than that of endogenous CtBP1 but were comparable to wounds at the mRNA level (Figure 1a). Overall, K5.CtBP1 phenotype severity correlated with transgene expression levels, suggesting that CtBP1 overexpression causes the phenotype. Results from the representative line K2 were shown in this study. K5.CtBP1 pups were born without gross abnormality (not shown) but began to exhibit thickened skin at 1 week postpartum when wild-type mice developed their first coat of hair and K5.CtBP1 pups had no hair growth (Figure 1b). After weaning, K5.CtBP1 mice exhibited hair loss on their dorsal and ventricle sides (Figure 1c). These transgenic mice died between 3 and 6 weeks of age due to severe hyperplasia in the esophagus (Figure 1d) and forestomach (not shown) where CtBP1 transgene was also expressed, compromising food intake. Compared with the wild-type control, these K5.CtBP1 mice also displayed abnormal epithelium in their tongues (Figure 1d).




Figure 1. Generation of K5.CtBP1 mice and phenotypes. (a) *CtBP1* mRNA expression in skin of wild-type (WT) mice and K5.CtBP1 transgenic mice (K1, K2, K3), and acutely wounded WT skin (wound). The mRNA level in WT skin was arbitrarily set as “1”. Error bars indicate SD ($n = 3$); significance was determined using Student’s *t* test. ** $P < 0.01$; * $P < 0.05$. (b) Hyperplasia/hyperkeratotic gross appearance of a K5.CtBP1 transgenic pup 8 days postpartum, in comparison with WT littermates. (c) Thin and patchy hair in K5.CtBP1 mice 3 weeks postpartum, in comparison with WT littermates. (d) Hematoxylin and eosin staining of WT and K5.CtBP1 tongue (top panels) and esophagus (bottom panels). Scale bar = 15 μm (top panels) and 40 μm (bottom panels).

K5.CtBP1 mice displayed epidermal hyperproliferation, with decreased expression of p21 and Brca1

Reactivation of CtBP1 expression has been shown in cancer (Nadauld *et al.*, 2006; Deng *et al.*, 2010) and during wound healing (Figure 1a), presumably contributing to the hyperproliferative process in these conditions. We biopsied the K5.CtBP1 skin on day 21 postpartum to examine morphology changes (Figure 2a) and expression of CtBP1 (Figure 2b and c). Hyperplasia in the epidermis of transgenic mice became obvious at the microscopic level (Figure 2a). Although the K5 promoter targets CtBP1 expression to basal keratinocytes, nuclear CtBP1 staining was apparent throughout the entire epidermis and hair follicles in K5.CtBP1 transgenic mice

(Figure 2c), possibly because of protein retention in differentiated layers. In contrast, weak CtBP1 staining was detected in wild-type skin (Figure 2b and c). Interestingly, although the CtBP1 transgene is targeted to keratinocytes, CtBP1-positive cells were also observed in the K5.CtBP1 dermis but not in wild-type dermis (Figure 2c). This is likely increased endogenous CtBP1 in stromal cells in response to changes in the epidermis and hair follicles. The origin of the cells overexpressing endogenous CtBP1 in the transgenic dermis remains to be determined.

In vitro studies show that CtBP1 has both anti-apoptotic and proliferative effects (Grooteclaes *et al.*, 2003; Mroz *et al.*, 2008), potentially causing epidermal hyperplasia. Apoptosis,




Figure 2. Microscopic characteristics of K5.CtBP1 skin. (a) Hematoxylin and eosin (H&E) staining of 21-day-old mice shows hyperplasia of the epidermis in K5.CtBP1 skin compared with wild-type (WT) skin. (b) Western blotting of CtBP1 in WT and K5.CtBP1 skin. Keratin 14 (K14) was used as loading control. Immunohistochemistry using antibodies specific for CtBP1 (c) and PCNA (d). Scale bar = 40 μ m (all panels). qRT-PCR analysis shows downregulation of CtBP1 target genes *p21* (e) and *Brca1* (f) in K5.CtBP1 skins. *p21* mRNA was decreased to 0.17 ± 0.09 and *Brca1* mRNA was decreased to 0.09 ± 0.03 in transgenic skin. (g) Keratinocytes from K5.CtBP1 mice displayed an increased BrdU index. Keratin 14 (red) was used as a counterstain. Scale bar = 15 μ m.

as determined by cleaved Caspase-3, was not changed in K5.CtBP1 skin (not shown). Proliferative cells, identified by PCNA staining, were sporadic in the basal layer of the epidermis and hair follicles of wild-type skin, but were expanded to suprabasal layers of K5.CtBP1 transgenic epidermis and hair follicles (Figure 2d). Furthermore, we found that mRNA levels of *p21* and *Brca1* were decreased in skin with CtBP1 overexpression, with a 5- and 11-fold reduction, respectively, in K5.CtBP1 skin compared with control skin (Figure 2e and f), supporting the notion that CtBP1 positively contributes to proliferation in keratinocytes. We cultured keratinocytes from K5.CtBP1 mice. Consistent with the hyperplastic phenotype observed in K5.CtBP1 mice, keratinocytes derived from these animals displayed a higher BrdU index than the wild-type controls (Figure 2g), suggesting that the proliferative property is cell autonomous. No major difference was detected in the cell cycle analysis and migration assay (not shown).

CtBP1 transgene expression reduced differentiation and caused loss of E-cadherin in the epidermis and hair follicles

Previously, we and others showed that CtBP1 represses the transcription of *E-cadherin* by directly targeting the *E-cadherin* promoter (Grootclaes and Frisch, 2000; Grootclaes et al., 2003; Zhang et al., 2006). To determine whether CtBP1 expression in epithelial cells leads to *E-cadherin* reduction, we performed qRT-PCR on K5.CtBP1 skin on day 21 postpartum and found that *E-cadherin* mRNA was decreased when compared with that of wild-type control littermates (Figure 3a). Consistent with the mRNA decrease, E-cadherin protein was largely lost in K5.CtBP1 epidermis and hair follicles as seen by immunostaining (Figure 3b). The E-cadherin loss in K5.CtBP1 mice is progressive, as only a small decrease in E-cadherin was detected in day 1 pup skin compared with wild-type littermates (not shown).

Next, we examined the epidermal differentiation of K5.CtBP1 mice. Early epidermal differentiation markers, keratins K1 and K10, were not altered (not shown), but epidermal terminal differentiation markers loricrin and filaggrin were largely diminished by immunostaining in the K5.CtBP1 epidermis (Figure 3c) and significantly reduced by western blotting (Figure 3d). Because CtBP1 is a classic transcriptional corepressor and potentially contributes to the decreased differentiation by transcriptional repression of the terminal differentiation players, we assayed the mRNA levels of *loricrin* and *filaggrin*, important physical barrier components in the epidermis against the environment (Kalinin et al., 2001; Candi et al., 2005). Different from the protein loss observed in the K5.CtBP1 epidermis, neither *loricrin* nor *filaggrin* showed a decrease in their mRNA levels (Figure 3e), suggesting that the defect in epidermal terminal differentiation is a secondary effect.

CtBP1 transgene suppressed *Dlx3*, a critical regulator of hair follicle differentiation

We biopsied neonatal skin and found no obvious hair follicle changes at birth (not shown), suggesting that CtBP1 did not affect hair development. Consistently, similar expression of LEF-1, β -catenin, and pSmad1/5/8 was detected in the CtBP1

transgenic skin hair follicles (not shown), implying that Wnt and bone morphogenic protein signaling pathways are not affected by CtBP1 overexpression in the epidermis. Structural hair follicle abnormalities were observed by day 9 postpartum. Hair follicles had formed large hair bulbs and differentiated hair shafts in wild-type mice. In contrast, CtBP1 transgenic hair follicles displayed smaller hair bulbs, with reduced keratinized medulla and defects in the inner root sheath and hair shaft (Figure 4a), phenotypes reminiscent of hair follicle abnormalities observed in *Dlx3*-null mice (Hwang et al., 2008). Therefore, we used the *Dlx3* hair follicle differentiation marker to explore the molecular mechanism associated with hair loss in K5.CtBP1 mice. On day 9 postpartum, *Dlx3* was expressed in hair matrix cells, inner root sheath, as well as in hair-forming compartments such as the cortex, medulla, and cuticle in hair follicles in wild-type mice (Figure 4b, left panel). However, expression of *Dlx3* was largely reduced in hair follicles in K5.CtBP1 mice (Figure 4b, right panel), suggesting that CtBP1 overexpression in hair follicles down-regulates *Dlx3* to induce the differentiation defect of the inner root sheath and hair shaft. Cross-sectional co-immunofluorescence staining of *Dlx3* and CtBP1 revealed that in wild-type hair follicles the *Dlx3* expression is in the inner root sheath, with CtBP1 expression mainly in the outer root sheath (Figure 4c, left panel). In K5.CtBP1 hair follicles, expression of *Dlx3* was decreased, with CtBP1 expansion to the inner root sheath (Figure 4c, right panel). To evaluate the *Dlx3* changes at the mRNA level, *in situ* hybridization analysis was performed. *Dlx3* mRNA was detected in the wild-type mice hair follicles (Supplementary Figure S1 online, left panel). In contrast, there was very little *Dlx3* detected in the K5.CtBP1 hair follicles (Supplementary Figure S1 online, right panel).

Next, we assayed the mRNA level of *Dlx3* in K5.CtBP1 skin by qRT-PCR and compared it with that in wild-type littermates. A significant decrease in *Dlx3* mRNA was detected in CtBP1 transgenic skin (Figure 5a). To determine whether CtBP1 has a direct role in regulation of the *Dlx3* gene, we performed chromatin immunoprecipitation (ChIP) to determine whether CtBP1 is recruited to the *Dlx3* promoter in mouse keratinocytes. Mouse keratinocytes were transfected with a vector expressing CtBP1, either wild-type or the PLDLX-binding deficient mutant tagged with the FLAG epitope, and the cross-linked chromatin was immunoprecipitated with an anti-FLAG antibody. Wild-type CtBP1 bound the *Dlx3* promoter region, whereas the PLDLX-binding deficient mutant did not (Figure 5b). This binding is limited to the promoter region, as no signal was detected with PCR amplification of the ChIP material using primers either 5' or 3' 2 kb to the *Dlx3* promoter (Figure 5b). Consistent with mRNA changes observed during CtBP1 knockdown, the luciferase activity of the *Dlx3* promoter decreased by 60% with CtBP1 overexpression in mouse keratinocytes (Figure 5c), indicating that CtBP1 regulates *Dlx3* transcription, at least partially, by binding to its promoter. To investigate whether the CtBP1-mediated repression of *Dlx3* gene occurs in human cells, we knocked down CtBP1 in Fadu cells, a human SCC cell line exhibiting high endogenous CtBP1 (Deng et al., 2010), and assayed the *Dlx3*-luciferase reporter. CtBP1 knockdown increased the *Dlx3*-luc




Figure 3. Aberrant differentiation in the epidermis and hair follicles of K5.CtBP1 skin. (a) Downregulation of CtBP1 target gene *E-cadherin* in K5.CtBP1 skin. $**P < 0.01$. (b) Immunofluorescence staining for E-cadherin (green) shows significant loss of E-cadherin in K5.CtBP1 epidermis and hair follicles. Keratin 14 (red) was used as a counterstain. Scale bar = 40 μm . (c) Immunofluorescence staining (green) for differentiation markers loricrin and filaggrin shows loss of these proteins in the K5.CtBP1 epidermis. Keratin 14 (red) was used as a counterstain. Scale bar = 40 μm . (d) Western blotting of loricrin and filaggrin in wild-type (WT) and K5.CtBP1 skin. Keratin 14 (K14) serves as a loading control. (e) qRT-PCR analysis of the differentiation marker genes *loricrin* and *filaggrin* in K5.CtBP1 skins. NS, no statistical significance by Student's *t* test.

reporter activity (Figure 5d), suggesting that CtBP1's repressive role in *Dlx3* transcription is conserved between mouse and human cells.

To further study the functional consequence of CtBP1-mediated repression of the *Dlx3* gene, we examined mRNA levels of *Dlx3* transcriptional targets in K5.CtBP1 skin. *Gata3* and *Hoxc13* are transcription factors affecting hair differentiation (Godwin and Capecchi, 1998; Kurek et al., 2007). Decreased *Gata3* and *Hoxc13* expression was detected in genetically engineered K14-*Dlx3*^{-/-} mice (Hwang et al., 2008). As shown in Figure 6a, CtBP1 transgene expression decreased mRNA levels of *Gata3* to 60% of those seen in control skin, a small but significant change. In contrast, expression of *Hoxc13* was decreased to 10% of that seen in control skin by CtBP1 transgene expression (Figure 6a).

Figure 6b illustrates the decrease in *Gata3* and *Hoxc13* expression in hair follicles of K5.CtBP1 mice.

A recent study shows that *Dlx3* upregulates the expression of the inner root sheath-forming keratins (Kim et al., 2012). Therefore, we measured the levels of Type I inner root sheath keratin genes *Krt25*, *Krt27*, and *Krt28* and Type II inner root sheath keratin gene *Krt71* in wild-type and K5.CtBP1 skin. Similar to the decrease in *Hoxc13* expression observed in K5.CtBP1 skin, the mRNA levels of *Krt25*, *Krt27*, and *Krt28* decreased significantly when compared with control skin (Figure 6c). The expression of *Krt71* was decreased by 4-fold as well (Figure 6c). Taken together, these findings indicate that CtBP1 overexpression triggered a downregulation of transcription factors and hair keratins critical for the maintenance of hair follicle homeostasis.




Figure 4. CtBP1 suppresses *Dlx3* gene expression in K5.CtBP1 skins. (a) Hematoxylin and eosin (H&E) staining of wild-type (WT) and K5.CtBP1 hair shaft and bulb. Scale bar = 15 μ m. Note the reduced hair bulbs in the transgenic mice. (b) Immunofluorescence staining of *Dlx3* (green) shows its aberrant expression pattern in K5.CtBP1 hair follicles. Keratin 14 (red) was used as a counterstain. Scale bar = 15 μ m. (c) Cross-sectional immunofluorescence staining of *Dlx3* (green) and CtBP1 (blue) shows the decreased *Dlx3* expression with CtBP1 expansion to the inner root sheath (IRS) in K5.CtBP1 hair follicles. ORS, outer root sheath. Keratin 14 (red) was used as a counterstain. Scale bar = 5 μ m.

DISCUSSION

Epidermal E-cadherin loss, hyperproliferation, and poor differentiation are caused by keratinocyte-specific CtBP1 overexpression

We and others have shown that CtBP1 expression can be reinitiated in cancers (Nadauld *et al.*, 2006; Deng *et al.*, 2010); thus, we studied whether pathologically induced CtBP1 overexpression perturbs skin homeostasis. Consistent with previous *in vitro* studies, we found hyperproliferation, down-regulated differentiation, and loss of E-cadherin in K5.CtBP1 mouse skin. Among potential CtBP1 targets associated with proliferation, we found downregulation of *p21* and *Brca1*. Increased proliferation can be, but is not always, associated with reduced differentiation. Both *p21* and *Brca1* inhibit cell cycle progression and induce differentiation in the epidermis (Missero *et al.*, 1996; Berton *et al.*, 2003). Therefore, reduced *p21* and *Brca1* could contribute to reduced differentiation of K5.CtBP1 epidermis. We also found gradual E-cadherin loss in the transgenic epidermis and hair follicles. Keratinocyte-specific *E-cadherin* knockout mice do not show epidermal blisters but display epidermal hyperproliferation and poor




Figure 5. CtBP1 suppresses the *Dlx3* gene transcription. (a) Downregulation of the *Dlx3* gene in K5.CtBP1 skins from 9-day-old mice. (b) CtBP1 binding to the *Dlx3* promoter. ChIP assay was performed in keratinocytes after transfection with FLAG-tagged CtBP1-expressing vector, either wild-type or the PLDLX-binding deficient mutant. ***P* < 0.01 vs. the PLDLX-binding deficient mutant (mt). (c) CtBP1 transfection represses the *Dlx3* reporter. The pGL4.26 *Dlx3* promoter reporter was generated by cloning the -286 to 0 bp fragment of the *Dlx3* promoter. (d) siCtBP1 increases activity of the *Dlx3* reporter. Fadu cells were transfected with scrambled siRNA (SC) or siRNA to CtBP1 (siCtBP1) and luciferase activity was measured.

differentiation in both the epidermis and hair follicles (Young *et al.*, 2003; Tinkle *et al.*, 2004). Thus, E-cadherin loss in K5.CtBP1 transgenic mice could significantly contribute to progressive hyperplasia and poor terminal differentiation.

Dlx3 ablation in keratinocytes has been shown to induce hyperplasia of the epidermis (Hwang *et al.*, 2008). In the current study, we found that CtBP1 suppresses transcription of the *Dlx3* gene, which may shift the balance between proliferation and differentiation and contribute to the overly proliferative phenotype of the K5.CtBP1 mice. In addition, the resultant downregulation of *Gata3* may further facilitate overproliferation because *Gata3* ablation in the epidermis has been shown to induce epidermal hyperplasia (Kurek *et al.*,




Figure 6. Downregulation of *Dlx3* target genes in K5.CtBP1 skins. (a) Wild-type (WT) and K5.CtBP1 skins were used for qRT-PCR. *Gata3* mRNA decreased to 0.35 ± 0.07 and *Hoxc13* mRNA decreased to 0.11 ± 0.06 in transgenic skin. $**P < 0.01$. (b) Immunofluorescence staining (green) of *Gata3* and *Hoxc13* shows decreased expression in K5.CtBP1 hair follicles compared with WT follicles. Keratin 14 (red) was used as a counterstain. Scale bar = 15 μ m. (c) WT and K5.CtBP1 skins were used for qRT-PCR *Keratin 25* (*KRT25*), *keratin 27* (*KRT27*), *keratin 28* (*KRT28*), and *keratin 71* (*KRT71*) mRNA expression levels. Error bars indicate SD ($n = 3$), and significance was determined using Student's *t* test. $**P < 0.01$.

2007). Therefore, keratinocyte proliferation and reduced differentiation of cells in the epidermis and hair follicles appear to be critically regulated by CtBP1, as seen from the synergistic actions of p21, E-cadherin, and terminal differentiation regulators of the epidermis and hair follicles such as *Dlx3* and *Gata3*.

CtBP1 transcriptionally represses the *Dlx3* gene, a critical regulator of hair follicle differentiation

In our K5.CtBP1 transgenic model, we unexpectedly found defective hair morphogenesis caused by CtBP1 overexpression in keratinocytes. Previous studies have shown that *Dlx3* is a transcriptional activator and has a critical role in the

development of the epidermis, hair, bone, and placenta (Morasso *et al.*, 1996; Feledy *et al.*, 1999; Morasso *et al.*, 1999; Hassan *et al.*, 2004; Hwang *et al.*, 2008), and *Dlx3* mutations are responsible for the defects in hair, teeth, and bone development called the Tricho-Dento-Osseous syndrome (Price *et al.*, 1998a, b). Our transgenic mice displayed hair follicle differentiation abnormalities similar to *K14.Dlx3*^{-/-} mice, with smaller hair bulbs, reduced keratinized medulla, and defects in the inner root sheath and hair shaft (Hwang *et al.*, 2008). Although the hair loss seems regional, abnormal hair follicles were also observed where hair was retained (not shown), suggesting that mechanical triggers, such as rubbing, facilitate hair loss. Unlike the genetically engineered mice with *Dlx3* ablation in the epidermis in which the Wnt and bone morphogenic protein signaling pathways are disturbed, the Wnt and bone morphogenic protein signaling pathways were not perturbed by CtBP1 overexpression in keratinocytes, presumably reflecting an incomplete shutdown of *Dlx3* in K5.CtBP1 mice. Preservation of the bone morphogenic protein and Wnt signaling is consistent with the ability of K5.CtBP1 mice to regenerate hair despite their abnormal hair formation.

Consistent with *Dlx3*-mediated effects in hair follicles, keratinocyte-specific overexpression of CtBP1 caused a decrease in *Gata3* and *Hoxc13*, transcription factors involved in hair differentiation (Godwin and Capocchi, 1998; Tkatchenko *et al.*, 2001), and in Type I/II keratins. A previous study has shown that HOXC13 regulates human hair keratin gene expression (Jave-Suarez *et al.*, 2002). The inner root sheath forms its structure by obligate heterodimerization of the specified keratins. Type I inner root sheath keratin genes *Krt25*, *Krt27*, and *Krt28* and Type II inner root sheath keratin gene *Krt71* are specifically expressed in all three layers of the inner root sheath and support the structure (Runkel *et al.*, 2006; Tanaka *et al.*, 2007). We observed that expression of these inner root sheath keratins was downregulated by CtBP1. As a result of keratin loss, the heterodimer may be reduced, hindering inner root sheath formation and resulting in further hair abnormality. For instance, mutations in the helix termination motif of mouse Type I inner root sheath keratin genes have been shown to impair the assembly of keratin intermediate filament (Tanaka *et al.*, 2007). The ability to downregulate both Type I and Type II keratins may further impair the inner root sheath formation.

In addition to the critical role of *Dlx3* in hair follicle differentiation, ablation of *E-cadherin* in keratinocytes has been shown to induce the progressive loss of hair follicle integrity (Young *et al.*, 2003; Tinkle *et al.*, 2004). We reason that CtBP1 contributes to hair follicle abnormality through multiple targets, including *Dlx3* and E-cadherin, and thus it may become more pronounced with aging. Future studies using an inducible model will help elucidate the impact of CtBP1 overexpression on epidermal and hair follicle differentiation at later stages. This model will also allow us to assess the role of CtBP1 in other pathological processes such as wound healing.

In summary, we report the CtBP1 transgenic mouse model to reveal *in vivo* CtBP1 transcriptional targets in keratinocytes

and functions associated with epidermal hyperproliferation, reduced differentiation, and E-cadherin downregulation. Further, CtBP1 has a role in hair morphogenesis, which had not been identified previously, and can only be appreciated with CtBP1 overexpression during differentiation of the hair follicle. Our data instigate future studies to determine whether pathologically induced CtBP1 overexpression has a role in skin pathogenesis in human diseases.

MATERIALS AND METHODS

Generation and identification of K5.CtBP1 mice

All animal experiments were performed with the approval of IACUC at the University of Colorado Denver. The 1.3 kb full-length wild-type human CtBP1 cDNA was inserted into the K5 expression vector (He *et al.*, 2002). K5.CtBP1 transgenic mice were generated with the B6D2 strain by microinjection of the transgene into the pronuclei of mouse embryos. Mice were genotyped by PCR analysis of tail DNA using primers specific for BK5 (tctgataggcagctgcacc) and CtBP1 (atcccagctgctgtggaagg). Throughout this study, all transgenic mice were heterozygous; all wild-type mice were littermates, and at least three independent analyses were performed for each assay, using three to five samples in each group.

Tissue histology, immunofluorescence, immunohistochemistry, western blotting, and *in situ* hybridization

Skin histology was visualized with hematoxylin and eosin (H&E) staining. Immunofluorescence, immunohistochemistry, and western blotting were performed on frozen and paraffin-embedded sections as previously described (Wang *et al.*, 1999). Immunofluorescence and western blotting were performed using antibodies against CtBP1 (Millipore, Billerica, MA), loricrin, filaggrin (Covance, San Diego, CA), Hoxc13 (Abnova, Walnut, CA), Gata3 (Santa Cruz, Santa Cruz, CA), E-cadherin (BD Biosciences, San Jose, CA), Dlx3 (Abcam, Cambridge, MA), and Keratin 14 (Fitzgerald, Acton, MA). The antibodies used in immunohistochemical analysis included CtBP1 (Millipore) and PCNA (Santa Cruz). For immunofluorescence, secondary antibodies to different species IgG were Alexa Fluor 594 (red) or 488 (green) conjugated (1:200 for all, Invitrogen, Grand Island, NY). For immunohistochemical analysis, we used secondary biotinylated antibodies to different species IgG (1:300, Vector Labs, Burlingame, CA) and developed using Vectastain ABC kit (Vector Labs). *In situ* hybridization with antisense digoxigenin-UTP-labeled RNA probes on sections of skin samples was performed as described (Han *et al.*, 2006).

Quantitative real-time reverse-transcriptase-PCR

Total RNA was isolated using TRIzol (Invitrogen) as previously described (Zhang *et al.*, 2006). One hundred nanograms of RNA from each sample were subjected to quantitative real-time reverse-transcriptase-PCR (ThermoFisher, Waltham, MA). An 18S probe was used as an internal control. Each sample was examined in triplicate. The relative RNA expression levels were determined by normalizing with internal controls, the values of which were calculated using the comparative Ct method.

Cell culture and transfections

Mouse keratinocytes were isolated from neonatal mouse skin as previously described (Han *et al.*, 2011), and cultured in PCT medium

(CELLnTEC, Bern, Switzerland). Fadu, a human HNSCC line, was purchased from ATCC and cultured in DMEM with 10% fetal bovine serum. To knock down CtBP1, cells were treated with siRNA against CtBP1 from Dharmacon (Lafayette, CO) using Lipofectamine 2000 (Invitrogen) for 48 hours and then collected. Keratinocytes were treated with BrdU; cells that incorporated BrdU were stained by FITC anti-BrdU.

ChIP and luciferase reporter assay

Keratinocytes were transfected with a CtBP1-expressing vector, either wild-type or the PLDLX-binding deficient mutant tagged with FLAG epitope. ChIP assay was performed with an anti-FLAG antibody as described previously (Zhang *et al.*, 2006). Primer sets spanning the *Dlx3* promoter were used to quantitative reverse transcriptase in real-time-amplify the ChIP sample. The pGL4.26 *DLX3* promoter luciferase reporter plasmid was generated by cloning a PCR-amplified – 286 to 0 bp fragment of the *Dlx3* promoter into the XhoI and NheI sites of the pGL4.26 vector (Promega, Madison, WI). *Dlx3* promoter-specific primers used were 5'-TATCTCGAGCCGCACAGCCAAC-3' (forward) and 5'-AATGCTAGCCAGCTCCGCC-3' (reverse). An empty renilla luciferase vector (pGL4.79) was used for normalization. Mouse keratinocytes were co-transfected with the reporters and CtBP1-expressing plasmids for 48 hours and luciferase activity was measured (Zhang *et al.*, 2002). Human Fadu cells were co-transfected with the reporters and siRNA to CtBP1 for 48 hours and luciferase activity was measured (Zhang *et al.*, 2002). Scrambled siRNA or empty plasmid was used for controls.

CONFLICT OF INTEREST

The authors state no conflict of interest.

ACKNOWLEDGMENTS

This work was supported by grants from the NIH, R01CA87949 (to XJW) and R01CA115468 and R03DA033982 (to QZ), pilot grants from P30CA046934 and P30 AR057212 (to QZ), and seed grants from Cancer League of Colorado and CCTSI (to QZ). HD was supported by grants from the Sixth People's Hospital of Shanghai. FL was supported by grants (no. 81102596) from the National Science Foundation of China (NSFC), The Shanghai Rising-Star Program (no. 12QA1403300), and The Innovation Program of Shanghai Municipal Education Commission (no. 12YZ067). We thank Laura Hoaglin and Pamela Fernandez for excellent technical support, Jiang Chen, Maranke Koster, Peter Koch, and Denis Roop for helpful comments, and Pamela J. Garl for editing the manuscript.

SUPPLEMENTARY MATERIAL

Supplementary material is linked to the online version of the paper at <http://www.nature.com/jid>

REFERENCES

- Berton TR, Matsumoto T, Page A *et al.* (2003) Tumor formation in mice with conditional inactivation of Brca1 in epithelial tissues. *Oncogene* 22:5415–26
- Boyd JM, Subramanian T, Schaeper U *et al.* (1993) A region in the C-terminus of adenovirus 2/5 E1a protein is required for association with a cellular phosphoprotein and important for the negative modulation of T24-ras mediated transformation, tumorigenesis and metastasis. *EMBO J* 12:469–78
- Candi E, Schmidt R, Melino G (2005) The cornified envelope: a model of cell death in the skin. *Nat Rev Mol Cell Biol* 6:328–40
- Chinnadurai G (2002) CtBP, an unconventional transcriptional corepressor in development and oncogenesis. *Mol Cell* 9:213–24
- Deng Y, Liu J, Han G *et al.* (2010) Redox-dependent Brca1 transcriptional regulation by an NADH-sensor CtBP1. *Oncogene* 29:6603–8

- Di LJ, Fernandez AG, De Siervi A *et al.* (2010) Transcriptional regulation of BRCA1 expression by a metabolic switch. *Nat Struct Mol Biol* 17:1406–13
- Feledy JA, Morasso MI, Jang SI *et al.* (1999) Transcriptional activation by the homeodomain protein distal-less 3. *Nucleic Acids Res* 27:764–70
- Godwin AR, Capecchi MR (1998) Hoxc13 mutant mice lack external hair. *Genes Dev* 12:11–20
- Grooteclaes M, Deveraux Q, Hildebrand J *et al.* (2003) C-terminal-binding protein corepresses epithelial and proapoptotic gene expression programs. *Proc Natl Acad Sci USA* 100:4568–73
- Grooteclaes ML, Frisch SM (2000) Evidence for a function of CtBP in epithelial gene regulation and anoikis. *Oncogene* 19:3823–8
- Han G, Li AG, Liang YY *et al.* (2006) Smad7-induced beta-catenin degradation alters epidermal appendage development. *Dev Cell* 11:301–12
- Han G, Li F, Ten Dijke P *et al.* (2011) Temporal smad7 transgene induction in mouse epidermis accelerates skin wound healing. *Am J Pathol* 179:1768–79
- Hassan MQ, Javed A, Morasso MI *et al.* (2004) Dlx3 transcriptional regulation of osteoblast differentiation: temporal recruitment of Msx2, Dlx3, and Dlx5 homeodomain proteins to chromatin of the osteocalcin gene. *Mol Cell Biol* 24:9248–61
- He W, Li AG, Wang D *et al.* (2002) Overexpression of Smad7 results in severe pathological alterations in multiple epithelial tissues. *EMBO J* 21:2580–90
- Hildebrand JD, Soriano P (2002) Overlapping and unique roles for C-terminal binding protein 1 (CtBP1) and CtBP2 during mouse development. *Mol Cell Biol* 22:5296–307
- Hwang J, Mehrani T, Millar SE *et al.* (2008) Dlx3 is a crucial regulator of hair follicle differentiation and cycling. *Development* 135:3149–59
- Jave-Suarez LF, Winter H, Langbein L *et al.* (2002) HOXC13 is involved in the regulation of human hair keratin gene expression. *J Biol Chem* 277:3718–26
- Kalinin A, Marekov LN, Steinert PM (2001) Assembly of the epidermal cornified cell envelope. *J Cell Sci* 114:3069–70
- Kim BK, Lee HY, Choi JH *et al.* (2012) Hairless plays a role in formation of inner root sheath via regulation of Dlx3 gene. *J Biol Chem* 287:16681–8
- Kitamura N, Motoi Y, Mori A *et al.* (2009) Suppressive role of C-terminal binding protein 1 in IL-4 synthesis in human T cells. *Biochem Biophys Res Commun* 382:326–30
- Kurek D, Garinis GA, van Doorninck JH *et al.* (2007) Transcriptome and phenotypic analysis reveals Gata3-dependent signalling pathways in murine hair follicles. *Development* 134:261–72
- Missero C, Di Cunto F, Kiyokawa H *et al.* (1996) The absence of p21Cip1/WAF1 alters keratinocyte growth and differentiation and promotes ras-tumor progression. *Genes Dev* 10:3065–75
- Morasso MI, Grinberg A, Robinson G *et al.* (1999) Placental failure in mice lacking the homeobox gene Dlx3. *Proc Natl Acad Sci USA* 96:162–7
- Morasso MI, Markova NG, Sargent TD (1996) Regulation of epidermal differentiation by a Distal-less homeodomain gene. *J Cell Biol* 135:1879–87
- Mroz EA, Baird AH, Michaud WA *et al.* (2008) COOH-terminal binding protein regulates expression of the p16INK4A tumor suppressor and senescence in primary human cells. *Cancer Res* 68:6049–53
- Nadauld LD, Phelps R, Moore BC *et al.* (2006) Adenomatous polyposis coli control of C-terminal binding protein-1 stability regulates expression of intestinal retinol dehydrogenases. *J Biol Chem* 281:37828–35
- Price JA, Bowden DW, Wright JT *et al.* (1998a) Identification of a mutation in DLX3 associated with tricho-dento-osseous (TDO) syndrome. *Hum Mol Genet* 7:563–9
- Price JA, Wright JT, Kula K *et al.* (1998b) A common DLX3 gene mutation is responsible for tricho-dento-osseous syndrome in Virginia and North Carolina families. *J Med Genet* 35:825–8
- Roukens MG, Alloul-Ramdhani M, Baan B *et al.* (2010) Control of endothelial sprouting by a Tel-CtBP complex. *Nat Cell Biol* 12:933–42
- Runkel F, Klawns M, Koch K *et al.* (2006) Morphologic and molecular characterization of two novel Krt71 (Krt2-fg) mutations: Krt71rco12 and Krt71rco13. *Mamm Genome* 17:1172–82
- Schaeper U, Boyd JM, Verma S *et al.* (1995) Molecular cloning and characterization of a cellular phosphoprotein that interacts with a conserved C-terminal domain of adenovirus E1A involved in negative modulation of oncogenic transformation. *Proc Natl Acad Sci USA* 92:10467–71
- Tanaka S, Miura I, Yoshiki A *et al.* (2007) Mutations in the helix termination motif of mouse type I IRS keratin genes impair the assembly of keratin intermediate filament. *Genomics* 90:703–11
- Tinkle CL, Lechler T, Pasolli HA *et al.* (2004) Conditional targeting of E-cadherin in skin: insights into hyperproliferative and degenerative responses. *Proc Natl Acad Sci USA* 101:552–7
- Tkatchenko AV, Visconti RP, Shang L *et al.* (2001) Overexpression of Hoxc13 in differentiating keratinocytes results in downregulation of a novel hair keratin gene cluster and alopecia. *Development* 128:1547–58
- Wang XJ, Liefer KM, Tsai S *et al.* (1999) Development of gene-switch transgenic mice that inducibly express transforming growth factor beta1 in the epidermis. *Proc Natl Acad Sci USA* 96:8483–8
- Young P, Boussadia O, Halfter H *et al.* (2003) E-cadherin controls adherens junctions in the epidermis and the renewal of hair follicles. *EMBO J* 22:5723–33
- Zhang Q, Piston DW, Goodman RH (2002) Regulation of corepressor function by nuclear NADH. *Science* 295:1895–7
- Zhang Q, Wang SY, Nottke AC *et al.* (2006) Redox sensor CtBP mediates hypoxia-induced tumor cell migration. *Proc Natl Acad Sci U S A* 103:9029–33