



ELSEVIER

Contents lists available at SciVerse ScienceDirect

Journal of Algebra

www.elsevier.com/locate/jalgebra

Quantum cohomology of the odd symplectic Grassmannian of lines

Clélia Pech*

Institut Fourier, 100 rue des Maths., St Martin d'Hères Cedex, France

ARTICLE INFO

Article history:

Received 6 January 2012

Available online 30 November 2012

Communicated by Shrawan Kumar

Keywords:

Quantum cohomology

Quasi-homogeneous spaces

Grassmannians

Pieri and Giambelli formulas

Exceptional collections in derived categories

ABSTRACT

Odd symplectic Grassmannians are a generalization of symplectic Grassmannians to odd-dimensional spaces. Here we compute the classical and quantum cohomology of the odd symplectic Grassmannian of lines. Although these varieties are not homogeneous, we obtain Pieri and Giambelli formulas that are very similar to the symplectic case. We notice that their quantum cohomology is semisimple, which enables us to check Dubrovin's conjecture for this case.

© 2012 Elsevier Inc. All rights reserved.

Introduction

The quantum cohomology of homogeneous varieties has been extensively studied (see [Tam] for references). Other well-known examples are toric varieties, yet apart from these settings, there are only few examples where the quantum cohomology has been explicitly determined. Quasi-homogeneous varieties provide interesting non-toric and non-homogeneous examples. Among these two Hilbert schemes have been studied, $\text{Hilb}(2, \mathbb{P}^2)$ [Gra01] and $\text{Hilb}(2, \mathbb{P}^1 \times \mathbb{P}^1)$ [Pon07].

In [Mih07] Mihai studied a family of varieties, the odd symplectic flag manifolds, which have many features in common with the symplectic flag manifolds. These varieties are interesting at least for two reasons; first, they are quasi-homogeneous, and secondly, since they have an action of the algebraic group Sp_{2n+1} (the odd symplectic group), whose properties are closely related to those of Sp_{2n} , they are expected to behave almost like homogeneous spaces and thus be relatively easy to deal with. The classical and quantum cohomology of symplectic Grassmannians has been described in [BKT09] and [BKT08], so one can ask whether it is possible to obtain similar results in the case of odd symplectic Grassmannians.

* Current address: Department of Mathematics, King's College London, Strand, WC2R 2LS London, UK.

E-mail addresses: clelia.pech@ujf-grenoble.fr, clelia.pech@kcl.ac.uk.

Here we deal with the case of the odd symplectic Grassmannian of lines $IG(2, 2n + 1)$, although some of the results about the classical cohomology hold in a more general setting. In 1.2 and 1.6 we use the natural embeddings of $IG(2, 2n + 1)$ in the usual Grassmannian and in the symplectic Grassmannian to compute classical Pieri (see 1.4) and Giambelli (see 1.7) formulas, as well as a presentation of the cohomology ring (see 1.8).

For the quantum cohomology the situation is more complicated. Since these varieties are not convex it is necessary to study the moduli spaces corresponding to invariants of degree one to show that they are smooth of the expected dimension. This is done in 2.1. Another difficulty is that since the group action is not transitive, an important transversality result, Kleiman’s lemma [Kle74, Thm. 2] no longer holds. So it will not be possible to force two Schubert varieties to meet transversely by an adequate choice of the defining flags as was done for instance in [Cos09]. Hence the Gromov–Witten invariants associated to Schubert varieties are not always enumerative. To solve this problem we replace Schubert varieties by another family of subvarieties and we use a transversality result of Graber [Gra01] suited for quasi-homogeneous spaces. In 2.5 we obtain a quantum Pieri formula and a presentation of the quantum cohomology ring. Finally, in 2.7, we check for odd symplectic Grassmannians of lines a conjecture of Dubrovin [Dub98, Conj. 4.2.2] relating semisimplicity of the quantum cohomology and the existence of a full exceptional collection in the derived category.

Our results show that there are many similarities with the symplectic case, since the classical and quantum Pieri formulas are almost the same in both cases. The Hasse diagrams are closely related as well (see 1.5). However, Poincaré duality is very different, since the Poincaré dual of a Schubert class is no longer always a single Schubert class (see 1.3). Moreover, contrary to what we prove here, the small quantum cohomology ring of the symplectic Grassmannian of lines is not semisimple (see [CP09]), and it is not known whether the Dubrovin conjecture holds in this case.

I wish to thank Laurent Manivel for his help on this subject.

1. Classical cohomology

Let $2 \leq m \leq n$ be integers, V be a \mathbb{C} -vector space of dimension $2n + 1$ and ω be an antisymmetric form of maximal rank on V . We denote its kernel by K . The *odd symplectic Grassmannian* is

$$IG_\omega(m, V) := \{ \Sigma \in \text{Gr}(m, V) \mid \Sigma \text{ is isotropic for } \omega \}.$$

It has an action of the *odd symplectic group*:

$$\text{Sp}(V) := \{ g \in \text{GL}(V) \mid \forall u, v \in V, \omega(gu, gv) = \omega(u, v) \}.$$

Up to isomorphism, $IG_\omega(m, V)$ does not depend on the $(2n + 1)$ -dimensional vector space V nor on the form ω , so we may denote it by $IG(m, 2n + 1)$. Similarly, from now on we denote $\text{Sp}(V)$ by Sp_{2n+1} . We recall some basic facts from [Mih07, Prop. 4.1 and 4.3]:

Proposition 1.

1. The odd symplectic Grassmannian $IG(m, 2n + 1)$ is a smooth subvariety of codimension $\frac{m(m-1)}{2}$ of the usual Grassmannian $\text{Gr}(m, 2n + 1)$.
2. Moreover, it has two orbits under the action of the odd symplectic group Sp_{2n+1} :
 - the closed orbit $\mathbb{O} := \{ \Sigma \in IG(m, 2n + 1) \mid \Sigma \supset K \}$, which is isomorphic to the symplectic Grassmannian $IG(m - 1, 2n)$;
 - the open orbit $\{ \Sigma \in IG(m, 2n + 1) \mid \Sigma \not\supset K \}$, which is isomorphic to the dual of the tautological bundle over the symplectic Grassmannian $IG(m, 2n)$.

For us, a *quasi-homogeneous space* will be an algebraic variety endowed with an action of an algebraic group with only finitely many orbits. Odd symplectic Grassmannians are examples of such spaces.

1.1. Schubert varieties

A \mathbb{C} -vector space V of dimension $2n + 1$ endowed with an antisymmetric form of maximal rank ω can be embedded in a symplectic space $(\bar{V}, \bar{\omega})$ of dimension $2n + 2$ such that $\bar{\omega}|_V = \omega$. This construction gives rise to a natural embedding $\mathbf{i}: \text{IG}(m, 2n + 1) \hookrightarrow \text{IG}(m, 2n + 2)$. It can be easily seen that \mathbf{i} identifies $\text{IG}(m, 2n + 1)$ with a Schubert subvariety of $\text{IG}(m, 2n + 2)$. Moreover this embedding enables us to obtain a description of the Schubert subvarieties of $\text{IG}(m, 2n + 1)$. In 1.1.1 we recall some facts about Schubert varieties in $\text{IG}(m, 2n)$, then in 1.1.2 we describe the Schubert varieties of $\text{IG}(m, 2n + 1)$ and introduce an indexation using partitions.

1.1.1. Schubert varieties in the symplectic Grassmannian

Here we recall the indexing conventions introduced in [BKT09, Def. 1.1]. Two kinds of combinatorial objects can be used to index Schubert varieties of the symplectic Grassmannian $\text{IG}(m, 2n)$, k -strict partitions (with $k := n - m$) and index sets:

Definition 1.

1. A k -strict partition is a weakly decreasing sequence of integers $\lambda = (\lambda_1 \geq \dots \geq \lambda_m \geq 0)$ such that $\lambda_j > k \Rightarrow \lambda_j > \lambda_{j+1}$.
2. An index set of length m for the symplectic Grassmannian is a subset $P = (p_1 < \dots < p_m) \subset [1, 2n]$ with m elements such that for all $1 \leq i, j \leq m$ we have $p_i + p_j \neq 2n + 1$.

Now if F_\bullet is an isotropic flag (i.e. a complete flag such that $F_{n-i}^\perp = F_{n+i}$ for all $0 \leq i \leq n$), to each admissible index set $P = (p_1, \dots, p_m)$ of length m we can associate the Schubert cell

$$X_P^\circ(F_\bullet) := \{ \Sigma \in \text{IG}(m, 2n) \mid \dim(\Sigma \cap F_{p_j}) = j, \forall 1 \leq j \leq m \}.$$

Moreover there is a bijection between k -strict partitions λ such that $\lambda_1 \leq 2n - m$ and index sets $P \subset [1, 2n]$ of length m , given by

$$\begin{aligned} \lambda &\mapsto P = (p_1, \dots, p_m) \quad \text{where } p_j = n + k + 1 - \lambda_j + \#\{i < j \mid \lambda_i + \lambda_j \leq 2k + j - i\}, \\ P &\mapsto \lambda = (\lambda_1, \dots, \lambda_m) \quad \text{where } \lambda_j = n + k + 1 - p_j + \#\{i < j \mid p_i + p_j > 2n + 1\}. \end{aligned}$$

The advantage of the representation by k -strict partitions is twofold: it mimics the indexation of Schubert classes of type A Grassmannians by partitions, and the codimension of the Schubert variety associated to a k -strict partition λ is easily computed as $|\lambda| = \sum_{j=1}^m \lambda_j$. In the next paragraph we will describe a similar indexation for the odd symplectic Grassmannian.

1.1.2. Schubert varieties in the odd symplectic Grassmannian

We now use Mihai’s description of the odd symplectic Grassmannian as a Schubert subvariety of $\text{IG}(m, 2n + 2)$ to define the Schubert varieties of the odd symplectic Grassmannian. We also introduce two indexations for them.

Schubert varieties of the odd symplectic Grassmannian will be defined with respect to an isotropic flag of \mathbb{C}^{2n+1} , i.e. a complete flag F_\bullet which is the restriction of an isotropic flag F_\bullet^+ of \mathbb{C}^{2n+2} . Denote by 1^m the partition λ^0 such that $\lambda_1^0 = \dots = \lambda_m^0 = 1$. It corresponds to the index set $P^0 = (2n + 2 - m, \dots, 2n + 1)$.

Proposition 2. *The embedding $\mathbf{i}: \text{IG}(m, 2n + 1) \rightarrow \text{IG}(m, 2n + 2)$ identifies $\text{IG}(m, 2n + 1)$ with the Schubert subvariety of $\text{IG}(m, 2n + 2)$ associated to the $(n + 1 - m)$ -strict partition λ^0 (or, equivalently, to the index set P^0).*

We define the Schubert varieties of $IG(m, 2n + 1)$ as the subvarieties of $IG(m, 2n + 1)$ of the form

$$X_P(F_\bullet) := \{ \Sigma \in IG(m, 2n + 1) \mid \dim(\Sigma \cap F_{p_j}) \geq j \text{ for all } j \},$$

where

- P is an index set of length m of $[1, 2n + 1]$, that is, an m -uple $P = (p_1 \leq \dots \leq p_m)$ with $1 \leq p_j \leq 2n + 1$ for all j and $p_i + p_j \neq 2n + 3$ for all i, j ;
- F_\bullet is an isotropic flag of \mathbb{C}^{2n+1} .

These varieties coincide with the Schubert varieties of $IG(m, 2n + 2)$ indexed by index sets P of $[1, 2n + 2]$ such that $P \leq P^0$ (for the lexicographical order), hence Proposition 2 implies that they define a cellular decomposition on $IG(m, 2n + 1)$.

Let us now describe another indexation of the Schubert varieties of $IG(m, 2n + 1)$ using partitions. If P is an index set of $[1, 2n + 1]$, we associate to it an $(n - m)$ -strict m -uple of weakly decreasing integers $\lambda = (\lambda_1 \geq \dots \geq \lambda_m \geq -1)$ defined by

$$\lambda_j = 2n + 2 - m - p_j + \#\{i < j \mid p_i + p_j > 2n + 3\} \quad \text{for all } 1 \leq j \leq m.$$

Conversely if $\lambda = (\lambda_1 \geq \dots \geq \lambda_m \geq -1)$ is any $(n - m)$ -strict m -uple of weakly decreasing integers such that $\lambda_1 \leq 2n + 1 - m$ and $(\lambda_m = -1 \Rightarrow \lambda_1 = 2n + 1 - m)$, then the assignment

$$p_j = 2n + 2 - m - \lambda_j + \#\{i < j \mid \lambda_i + \lambda_j \leq 2(n - m) + j - i\} \quad \text{for all } 1 \leq j \leq m$$

defines an index set of $[1, 2n + 1]$. It is easy to check that with respect to this indexation convention, the Schubert variety $X_\lambda(F_\bullet)$ has codimension $|\lambda|$ in $IG(m, 2n + 1)$.

Remark 1. For the case of the *odd symplectic Grassmannian of lines* $IG(2, 2n + 1)$, it follows that the indexing partitions can be either

- “usual” $(n - 2)$ -strict partitions $\lambda = (2n - 1 \geq \lambda_1 \geq \lambda_2 \geq 0)$;
- the “partition” $\lambda = (2n - 1, -1)$ corresponding to the class of the closed orbit \mathbb{O} .

1.2. Embedding in the symplectic Grassmannian

Now we draw some consequences of the embedding of $IG(2, 2n + 1)$ as a Schubert subvariety of a symplectic Grassmannian. Since we know the cohomology of $IG(2, 2n + 2)$, describing the restriction map \mathbf{i}^* will give us information on the cohomology of $IG(2, 2n + 1)$.

Let F_\bullet be an isotropic flag, $Y_{a,b}(F_\bullet)$ a Schubert subvariety of $IG(2, 2n + 2)$ and $\nu_{a,b}$ the associated Schubert class, where (a, b) is an $(n - 2)$ -strict partition. From Proposition 2, we know that $IG(2, 2n + 1)$ is isomorphic to the Schubert subvariety $Y_{1,1}(E_\bullet)$ of $IG(2, 2n + 2)$, where E_\bullet is an isotropic flag which we may assume to be in general position with respect to F_\bullet . Then it follows that $Y_{a,b}(F_\bullet)$ and $Y_{1,1}(E_\bullet)$ meet transversally, hence we can compute the restriction $\mathbf{i}^* \nu_{a,b}$ by computing the class of the intersection $Y_{a,b} \cap Y_{1,1}$ in $IG(2, 2n + 2)$ using the classical Pieri rules for $IG(2, 2n + 2)$ [BKT09, Thm. 1.1]:

$$\nu_{a,b} \cup \nu_{1,1} = \begin{cases} \nu_{a+1,b+1} & \text{if } a + b \neq 2n - 2, 2n - 1, \\ \nu_{a+1,b+1} + \nu_{a+2,b} & \text{if } a + b = 2n - 2 \text{ or } 2n - 1. \end{cases}$$

Remark 2. In the above formula, we should remove classes that are not indexed either by $(n - 2)$ -strict partitions $\lambda = (2n - 1 \geq \lambda_1 \geq \lambda_2 \geq 0)$ or by the special partition $\lambda = (2n - 1, -1)$. We will adopt this convention throughout the rest of the text to simplify formulas.

Denote by τ_λ the cohomology class associated to the Schubert variety $X_\lambda(F_\bullet) \subset IG(2, 2n + 1)$, where λ is an $(n - 2)$ -strict partition and F_\bullet is an isotropic flag of \mathbb{C}^{2n+1} . This class does not depend on the choice of the isotropic flag.

Looking at the incidence conditions for the corresponding Schubert varieties, we prove that for each $\tau_{c,d} \in H^*(IG(2, 2n + 1), \mathbb{Z})$, we have

$$\mathbf{i}_* \tau_{c,d} = \nu_{c+1,d+1}.$$

Using the projection formula $\mathbf{i}_*(\alpha \cup \mathbf{i}^*\beta) = \mathbf{i}_*\alpha \cup \beta$, we deduce

Lemma 1 (Restriction formula). *Let $\nu_{a,b} \in H^*(IG(2, 2n + 2), \mathbb{Z})$ be a Schubert class. Then its restriction to the odd symplectic Grassmannian $IG(2, 2n + 1)$ is given by*

$$\mathbf{i}^*\nu_{a,b} = \begin{cases} \tau_{a,b} & \text{if } a + b \neq 2n - 2, 2n - 1, \\ \tau_{a,b} + \tau_{a+1,b-1} & \text{if } a + b = 2n - 2 \text{ or } 2n - 1. \end{cases}$$

In particular we notice that \mathbf{i}^* is surjective and has kernel generated by the class ν_{2n} . So the classical cohomology of $IG(2, 2n + 1)$ is entirely determined by the classical cohomology of $IG(2, 2n + 2)$.

1.3. Poincaré duality

If X is a smooth algebraic variety and (γ_i) a homogeneous basis of its cohomology ring, we denote by (γ_i^\vee) the corresponding Poincaré dual basis. For homogeneous spaces and for odd symplectic Grassmannians, the basis (γ_i) we consider is the basis of Schubert classes. Here we compute Poincaré duality for $IG(2, 2n + 1)$.

If $\alpha = \nu_{a,b}$ is a Schubert class such that $b \geq 1$ or $(a, b) = (2n, 0)$, then there exists a unique class γ in $IG(2, 2n + 1)$ such that $\mathbf{i}_*\gamma = \alpha$. We denote it by α_- . We first prove

Lemma 2. *Let $\alpha = \nu_{a,b}$ be a cohomology class in $IG(2, 2n + 2)$ such that $b \geq 1$ or $(a, b) = (2n, 0)$. Then $\mathbf{i}^*\alpha^\vee = \alpha_-^\vee$.*

Proof. By definition of Poincaré duality, if α and β are two cohomology classes in $IG(2, 2n + 2)$, then

$$\int_{IG(2,2n+2)} \alpha \cup \beta^\vee = \delta_{\alpha,\beta},$$

where δ is the Kronecker symbol. So

$$\int_{IG(2,2n+2)} (\mathbf{i}_*\alpha_-) \cup \beta^\vee = \delta_{\alpha,\beta} = \int_{IG(2,2n+2)} \mathbf{i}_*(\alpha_- \cup \mathbf{i}^*\beta^\vee). \tag{1}$$

Expressing $\mathbf{i}^*\beta^\vee$ on the dual basis in $IG(2, 2n + 1)$, we get $\mathbf{i}^*\beta^\vee = \sum_\gamma x_{\beta,\gamma} \gamma^\vee$. Hence

$$\delta_{\alpha,\beta} = \sum_\gamma x_{\beta,\gamma} \int_{IG(2,2n+2)} \mathbf{i}_*(\alpha_- \cup \gamma^\vee) = \sum_\gamma x_{\beta,\gamma} \delta_{\alpha_-,\gamma}.$$

So $x_{\beta,\alpha_-} = \delta_{\alpha,\beta}$, and the result follows. \square

Finally, Poincaré duality in $IG(2, 2n + 1)$ takes the following form:




Fig. 1. Hasse diagram of IG(2, 7).




Fig. 2. Hasse diagram of IG(2, 6).

Proposition 3 (Poincaré duality).

$$\tau_{a,b}^\vee = \begin{cases} \tau_{2n-1-b, 2n-2-a} & \text{if } a + b < 2n - 2, \\ \tau_{2n-2-b, 2n-1-a} + \tau_{2n-1-b, 2n-2-a} & \text{if } a + b = 2n - 2 \text{ or } 2n - 1, \\ \tau_{2n-2-b, 2n-1-a} & \text{if } a + b > 2n - 1. \end{cases}$$

Proof. We will derive this result from Poincaré duality on IG(2, 2n + 2) using Lemmas 1 and 2. Indeed, we prove with the projection formula that if α is a class in IG(2, 2n + 2), then $\alpha^\vee = (\alpha^\vee \cup v_{1,1})_-$. Then using the Poincaré duality formula in IG(2, 2n + 2) proved in [BKT09, §4.1], an easy calculation gives the result. \square

Remark 3. This result is very different from what we get for the usual Grassmannians or even the symplectic or orthogonal ones. Indeed, the basis of Schubert classes is not self-dual. This fact will have many consequences; in particular, the Hasse diagram of IG(2, 2n + 1) (see Fig. 1) will be much less symmetric than the Hasse diagram of, say, IG(2, 2n + 2) (see Fig. 2).

1.4. Pieri formula

To compute the cup product of two cohomology classes in IG(2, 2n + 1), we need two ingredients: a *Pieri formula* describing the cup product of any Schubert class with a special class (that is, one of the classes τ_1 or $\tau_{1,1}$), and a *Giambelli formula* decomposing any Schubert class as a polynomial in τ_1 and $\tau_{1,1}$. In this paragraph we describe the Pieri formula as well as an alternative rule for multiplying Schubert classes and classes of the form τ_p with $0 \leq p \leq 2n - 1$ or $\tau_{2n-1,-1}$.

We start by expressing cohomology classes in IG(2, 2n + 1) in terms of cohomology classes in IG(2, 2n + 2) using Lemma 1:

$$\tau_{c,d} = \begin{cases} \mathbf{i}^* \mathcal{U}_{c,d} & \text{if } c + d \neq 2n - 2, 2n - 1, \\ \sum_{j=0}^{c-n} (-1)^{c-n-j} \mathbf{i}^* \mathcal{U}_{n-1+j, n-1-j} & \text{if } c + d = 2n - 2, \\ \sum_{j=c-n}^{n-1} (-1)^{j-c+n} \mathbf{i}^* \mathcal{U}_{n+j, n-1-j} & \text{if } c + d = 2n - 1. \end{cases}$$

Now combining this with the Pieri rule in $H^*(IG(2, 2n + 2), \mathbb{Z})$, we can prove a Pieri rule for IG(2, 2n + 1):

Proposition 4 (Pieri formula).

$$\tau_{a,b} \cup \tau_1 = \begin{cases} \tau_{a+1,b} + \tau_{a,b+1} & \text{if } a + b \neq 2n - 3, \\ \tau_{a,b+1} + 2\tau_{a+1,b} + \tau_{a+2,b-1} & \text{if } a + b = 2n - 3, \end{cases}$$

$$\tau_{a,b} \cup \tau_{1,1} = \begin{cases} \tau_{a+1,b+1} & \text{if } a + b \neq 2n - 4, 2n - 3, \\ \tau_{a+1,b+1} + \tau_{a+2,b} & \text{if } a + b = 2n - 4 \text{ or } 2n - 3. \end{cases}$$

We may also state a rule for multiplying by the Chern classes of the quotient bundle:

$$c_p(\mathcal{Q}) = \begin{cases} \tau_p & \text{if } 0 \leq p \leq 2n - 1 \text{ and } p \neq 2n - 2, \\ \tau_{2n-2} + \tau_{2n-1,-1} & \text{if } p = 2n - 2. \end{cases}$$

We prove in the same way as Proposition 4:

Proposition 5 (Another Pieri formula).

$$\tau_{a,b} \cup \tau_p = \begin{cases} \sum_{\substack{(a+1,b+1) \rightarrow (c+1,d+1) \\ c+d=a+b+p \\ d \geq 0 \text{ or } c=2n-1}} 2^{N((a+1,b+1),(c+1,d+1))} \tau_{c,d}, & \text{if } p \neq 2n - 2 \text{ or } (a + b \neq 2n - 1 \text{ and } (a, b) \neq (2n - 1, -1)), \\ (-1)^a \tau_{2n-1,2n-2} & \text{if } p = 2n - 2, a + b = 2n - 1 \text{ and } b \neq 0, \\ 0 & \text{if } p = 2n - 2 \text{ and } ((a, b) = (2n - 1, -1) \text{ or } (2n - 1, 0)), \end{cases}$$

where the relation $\lambda \rightarrow \mu$ and the integer $N(\lambda, \mu)$ are defined in [BKT09, Def. 1.3].

$$\tau_{a,b} \cup \tau_{2n-1,-1} = \begin{cases} (-1)^{a-1} \tau_{2n-1,2n-2} & \text{if } a + b = 2n - 1, \\ \tau_{2n-1,a-1} & \text{if } b = 0 \text{ and } a \neq 2n - 2, \\ \tau_{2n-1,2n-3} & \text{if } (a, b) = (2n - 1, -1), \\ 0 & \text{else.} \end{cases}$$

Notice that contrary to the symplectic case (and to the case of other homogeneous spaces) we sometimes get negative coefficients for the second Pieri rule. It is a consequence of the fact that we only have a quasi-homogeneous space, so it is not always possible to find representatives of the two Schubert varieties that intersect transversally. So even in degree 0 Gromov–Witten invariants associated to Schubert classes are not always enumerative, contrary to the case of homogeneous spaces. That is why we have to outline conditions in 2.2 to recover enumerativity for some invariants.

1.5. The Hasse diagram of $IG(2, 2n + 1)$

The Pieri rule from Proposition 4 enables us in particular to compute the multiplication by the hyperplane class τ_1 . The corresponding graph is called the *Hasse diagram* of $IG(2, 2n + 1)$. More precisely, the Hasse diagram of $IG(2, 2n + 1)$ is an oriented graph with multiplicity such that:

- its vertices are the Schubert classes of $IG(2, 2n + 1)$;
- two vertices $\tau_{a,b}$ and $\tau_{c,d}$ are related by an arrow of multiplicity r if $\tau_{c,d}$ appears with multiplicity r in the product $\tau_{a,b} \cup \tau_1$.

For instance see Fig. 1 for the Hasse diagram of $IG(2, 7)$. Arrows are going from left to right.

As a comparison, see also the Hasse diagram of the symplectic Grassmannian $IG(2, 6)$ in Fig. 2, and of $IG(2, 8)$ in Fig. 3.




Fig. 3. Hasse diagram of IG(2, 8).


Looking at these examples we notice that the Hasse diagram of IG(2, 7) contains the Hasse diagram of IG(2, 6) as a subgraph, the subgraph induced by the remaining vertices being isomorphic to the Hasse diagram of IG(1, 6). Moreover, the Hasse diagram of IG(2, 8) contains the Hasse diagram of IG(2, 7) as a subgraph, the subgraph induced by the remaining vertices being isomorphic to the Hasse diagram of IG(1, 6). This is a general fact. More precisely, we have the following decomposition of the Hasse diagrams of the even and odd symplectic Grassmannian:

Proposition 6.

- The Hasse diagram of IG(2, 2n + 1) is isomorphic to the disjoint union of:
 1. the Hasse diagram of IG(2, 2n), whose vertices are the classes in IG(2, 2n + 1) associated to the Schubert varieties not contained in the closed orbit;
 2. the Hasse diagram of the closed orbit $\mathbb{O} \cong \text{IG}(1, 2n)$;
 with parts 1 and 2 linked by the simple arrows joining τ_{2n-3} to $\tau_{2n-1,-1}$ and $\tau_{2n-2,a}$ to $\tau_{2n-1,a}$ for $0 \leq a \leq 2n - 3$.
- The Hasse diagram of IG(2, 2n) is isomorphic to the disjoint union of:
 1. the Hasse diagram of IG(2, 2n - 1), whose vertices are the classes in IG(2, 2n) associated to the Schubert varieties contained in $X_{1,1}$;
 2. the Hasse diagram of IG(1, 2n - 2), corresponding to the classes τ_\emptyset to τ_{2n-3} ;
 with parts 1 and 2 linked by the double arrow joining τ_{2n-3} to τ_{2n-2} and the simple arrows joining τ_p to $\tau_{p,1}$ for $1 \leq p \leq 2n - 3$.

Proof. We will denote by $\mathcal{H}_{\text{IG}(m,N)}$ the Hasse diagram of IG(m, N).

- Let G_1 be the subgraph of $\mathcal{H}_{\text{IG}(2,2n+1)}$ induced by the vertices τ_λ for λ such that $\lambda_1 < 2n - 1$. We need to prove that $G_1 = \mathcal{H}_{\text{IG}(2,2n)}$. First notice these graphs have the same set of vertices. Then use the diagram:



where i is the natural inclusion and $\phi(\Sigma) = \Sigma/K$ for each $\Sigma \in \mathbb{O}$. Looking at incidence conditions we notice that $\phi^*v_\lambda = i^*\tau_\lambda$ for each Schubert class v_λ of IG(2, 2n), and we get

$$\phi^*(v_1 \cup v_\lambda) = \phi^*v_1 \cup \phi^*v_\lambda = i^*(\tau_1 \cup \tau_\lambda),$$

hence G_1 and $\mathcal{H}_{\text{IG}(2,2n)}$ have the same arrows. Now the vertices of $\mathcal{H}_{\text{IG}(2,2n+1)}$ not contained in G_1 correspond to the classes τ_λ with $\lambda_1 = 2n - 1$, that is to the Schubert varieties contained in

the closed orbit $\mathbb{O} \cong \mathbb{P}^{2n-1}$. So the graph G_2 they induce is isomorphic to $IG(1, 2n)$. Finally, the arrows joining G_1 and G_2 are determined using the Pieri rule 4.

- For $IG(2, 2n)$ the result is simply a consequence of the isomorphism between $IG(2, 2n + 1)$ and the Schubert subvariety $X_{1,1}$ of $IG(2, 2n)$ stated in 1.2, and of the Pieri rule for $IG(2, 2n)$ proved in [BKT09, Thm. 1.1]. \square

This result can be easily generalized to all symplectic Grassmannians $IG(m, N)$:

Proposition 7.

- The Hasse diagram of $IG(m, 2n)$ is made of the union of:
 1. the Hasse diagram \mathcal{H}_1 of $IG(m, 2n - 1)$, whose vertices are the cohomology classes of $IG(m, 2n)$ associated to the Schubert varieties contained in X_{1^m} ;
 2. the Hasse diagram \mathcal{H}_2 of $IG(m - 1, 2n - 2)$.

The arrows from \mathcal{H}_2 to \mathcal{H}_1 are of two types:

 - simple arrows $\lambda \rightarrow \mu$ for λ, μ such that $\lambda_1 \leq 2n - 1 - m, \lambda_{m-1} \geq 1, \lambda_m = 0$ and $\mu = (\lambda_1, \dots, \lambda_{m-1}, 1)$;
 - double arrows $\lambda \rightarrow \mu$ for λ, μ such that $\lambda_1 = 2n - 1 - m, \lambda_m = 0$ and $\mu = (2n - m, \lambda_2, \dots, \lambda_m)$.

There is no arrow from \mathcal{H}_1 to \mathcal{H}_2 .
- The Hasse diagram of $IG(m, 2n + 1)$ is made of the union of:
 1. the Hasse diagram \mathcal{H}_1 of $IG(m, 2n)$, whose vertices are the cohomology classes associated to the Schubert varieties of $IG(m, 2n + 1)$ not contained in the closed orbit;
 2. the Hasse diagram \mathcal{H}_2 of the closed orbit $\mathbb{O} \cong IG(m - 1, 2n)$.

The arrows from \mathcal{H}_1 to \mathcal{H}_2 are simple and of two types:

 - $\lambda \rightarrow \mu$ for λ, μ such that $\lambda_1 = 2n - m$ and $\mu = (2n + 1 - m, \lambda_2, \dots, \lambda_m)$;
 - $\lambda \rightarrow \mu$ for λ, μ such that $2(n - m) + \#\{1 \leq i \leq m \mid \lambda_i \geq 1\} \leq \lambda_1 \leq 2n - 1 - m$ and $\mu = (2n + 1 - m, \lambda_2, \dots, \lambda_{\lambda_1 - 2(n - m)}, -1, \dots, -1)$.

There is no arrow from \mathcal{H}_2 to \mathcal{H}_1 .

The proof is very similar to the $m = 2$ case. However, the determination of the arrows between both parts of the Hasse diagram is a bit more complicated and uses a Pieri rule for the symplectic Grassmannian proved by Pragacz and Ratajski [PR96, Thm. 2.2], hence we will not give it here.

1.6. Embedding in the usual Grassmannian

The easiest way to find a Giambelli formula for $IG(2, 2n + 1)$ is to use the Giambelli formula on $Gr(2, 2n + 1)$ and to “pull it back” to $IG(2, 2n + 1)$. More precisely, we use the natural embedding:

$$j : IG(2, 2n + 1) \hookrightarrow Gr(2, 2n + 1).$$

We first prove that $j_*([IG(2, 2n + 1)]) = \sigma_1$, which means that j identifies $IG(2, 2n + 1)$ with a hyperplane section of $Gr(2, 2n + 1)$. It is enough to prove that if C is a line in $Gr(2, 2n + 1)$ not contained in $j(IG(2, 2n + 1))$, then C intersects $j(IG(2, 2n + 1))$ in only one point. Any line C not contained in $j(IG(2, 2n + 1))$ can be written as

$$C = \{ \Sigma \in Gr(2, 2n + 1) \mid V \subset \Sigma \subset W \},$$

where $\dim V = 1, \dim W = 3$ and $W \not\subset V^\perp$. Hence the only point in $C \cap j(IG(2, 2n + 1))$ is $\Sigma = W \cap V^\perp$.

Now using the same arguments as for Lemma 1 and the fact that $j_*([IG(2, 2n + 1)]) = \sigma_1$, we can prove

Lemma 3.

- If $a + b < 2n - 2$, then $\mathbf{j}^* \sigma_{a,b} = \tau_{a,b}$.
- If $a + b \geq 2n - 2$, then

$$\mathbf{j}^* \sigma_{a,b} = \tau_{a,b} + \tau_{a+1,b-1}.$$

This proves that the map \mathbf{j}^* is surjective and that its kernel is generated by the class

$$\sum_{i=0}^{n-1} (-1)^{n-i} \sigma_{n+i,n-i}.$$

1.7. Giambelli formula

With Lemma 3 and the Giambelli formula for $\text{Gr}(2, 2n + 1)$, we can prove a Giambelli formula with respect to τ_1 and $\tau_{1,1}$. First define $d_r := (\tau_{1^{1+j-i}})_{1 \leq i, j \leq r}$, with the convention that $\tau_{1^p} = 0$ if $p < 0$ or $p > 2$. We have

Proposition 8 (Giambelli formula).

$$\tau_{a,b} = \begin{cases} \tau_{1,1}^b d_{a-b} & \text{if } a + b \leq 2n - 3, \\ \sum_{q=0}^p (-1)^{p-q} \tau_{1,1}^{c-q} d_{2q} & \text{if } (a, b) = (c + 1 + p, c - 1 - p), \\ \sum_{q=p}^{2n-2-c} (-1)^{q-p} \tau_{1,1}^{c-q} d_{2q+1} & \text{if } (a, b) = (c + 1 + p, c - p), \end{cases}$$

where $n - 1 \leq c \leq 2n - 2$ and $0 \leq p \leq 2n - 2 - c$.

We can also state a Giambelli formula expressing classes in terms of the $e_p := c_p(\mathbb{Q})$:

Proposition 9 (Another Giambelli formula).

$$\tau_{a,b} = \begin{cases} e_a e_b - e_{a+1} e_{b-1} & \text{if } a + b \leq 2n - 3, \\ (-1)^{a-n} e_{n-1}^2 - e_a e_b + 2 \sum_{j=1}^{a-n} (-1)^{a-n-j} e_{n-1+j} e_{n-1-j} & \text{if } a + b = 2n - 2, \\ e_a e_b + 2 \sum_{j=1}^{2n-1-a} (-1)^j e_{a+j} e_{b-j} & \text{if } a + b \geq 2n - 1. \end{cases}$$

1.8. Two presentations for the classical cohomology ring

1.8.1. Presentation in terms of the classes e_p

Proposition 10 (Presentation of $H^*(\text{IG}(2, 2n + 1), \mathbb{Z})$). The ring $H^*(\text{IG}(2, 2n + 1), \mathbb{Z})$ is generated by the classes $(e_p)_{1 \leq p \leq 2n-1}$ and the relations are

$$\det(e_{1+j-i})_{1 \leq i, j \leq r} = 0 \quad \text{for } 3 \leq r \leq 2n, \tag{R1}$$

$$e_n^2 + 2 \sum_{i \geq 1} e_{n+i} e_{n-i} = 0. \tag{R2}$$

Proof. First of all, the quotient bundle \mathcal{Q} of $\text{IG}(2, 2n + 1)$ is the pullback by the restriction map \mathbf{i} of the quotient bundle \mathcal{Q}^+ on $\text{IG}(2, 2n + 2)$. So the $\mathbf{i}^*c_p(\mathcal{Q}^+) = c_p(\mathcal{Q}) = e_p$ for $1 \leq p \leq 2n$ generate $H^*(\text{IG}(2, 2n + 1), \mathbb{Z})$. But \mathcal{Q} having rank $2n - 1$, $\mathbf{i}^*c_{2n}(\mathcal{Q}^+) = 0$, hence the cohomology ring of $\text{IG}(2, 2n + 1)$ is generated by the $(e_p)_{1 \leq p \leq 2n-1}$. Then we follow the method from [BKT09, Thm. 1.2] to obtain presentations for the isotropic Grassmannians. Consider the graded ring $A := \mathbb{Z}[a_1, \dots, a_{2n-1}]$, where $\text{deg} a_i = i$. Set $a_0 = 1$, and $a_i = 0$ if $i < 0$ or $i > 2n - 1$. We also define $d_0 := 1$ and $d_r := \det(a_{1+j-i})_{1 \leq i, j \leq r}$ for $r > 0$. For all $r \geq 0$, set $b_r := a_r^2 + 2 \sum_{i \geq 1} (-1)^i a_{r+i} a_{r-i}$. Now let $\phi : A \rightarrow H^*(\text{IG}(2, 2n + 1), \mathbb{Z})$ be the degree-preserving morphism of graded rings sending a_i to e_i for all $1 \leq i \leq 2n - 1$. Since the e_p generate $H^*(\text{IG}(2, 2n + 1), \mathbb{Z})$, this morphism is surjective. To prove that relations (R1) and (R2) are satisfied, we must check that $\phi(d_r) = 0$ for all $r > 2$ and $\phi(b_n) = 0$.

(R1) Expanding the determinant d_r with respect to the first column, we get the identity

$$d_r = \sum_{i=1}^r (-1)^{i-1} a_i d_{r-i}.$$

Hence the identity on formal series:

$$\left(\sum_{i=0}^{2n-1} a_i t^i \right) \left(\sum_{i \geq 0} (-1)^i d_i t^i \right) = 1. \tag{2}$$

On $\text{IG}(2, 2n + 1)$ we have the following short exact sequence of vector bundles

$$0 \rightarrow \mathcal{S} \rightarrow \mathcal{O}_{\text{IG}(2, 2n+1)} \rightarrow \mathcal{Q} \rightarrow 0,$$

so $c(\mathcal{S})c(\mathcal{Q}) = 1$, where c denotes the total Chern class. But

$$c(\mathcal{Q}) = \sum_{i=0}^{2n-2} \tau_i t^i,$$

so (2) implies

$$c(\mathcal{S}) = \sum_{i \geq 0} (-1)^i \phi(d_i) t^i.$$

Since \mathcal{S} has rank 2, it follows that $\phi(d_r) = 0$ for all $r > 2$, hence the relations (R1).

(R2) From the presentation of $\text{IG}(2, 2n + 2)$ in [BKT09, Thm. 1.2], we know that

$$v_n^2 + 2 \sum_{i \geq 1} (-1)^i v_{n+i} v_{n-i} = 0$$

in $\text{IG}(2, 2n + 2)$. Pulling back by \mathbf{i} we get (R2).

Now consider the Poincaré polynomial of $\text{IG}(2, 2n + 1)$ computed in [Mih05, §2.2.3]:

$$P(\text{IG}(m, 2n + 1), q) = \frac{\prod_{i=1}^l (q^{2n+2-2i} - 1) \prod_{i=l+1}^m (q^{2n+4-2i} - 1)}{(q^m - 1)(q^{m-1} - 1) \dots (q - 1)}$$

for $m = 2l$. Evaluating this polynomial at $q = 1$, we get that the rank of $H^*(\text{IG}(2, 2n + 1))$ is $2n^2$.

As in the proof of [BKT09, Thm. 1.2], we will need the following lemma:

Lemma 4. *The quotient of the graded ring $\mathbb{Z}[a_1, \dots, a_d]$ with $\deg a_i = i$ modulo the relations*

$$\det(a_{1+j-i})_{1 \leq i, j \leq r} = 0, \quad m + 1 \leq r \leq m + d$$

is a free \mathbb{Z} -module of rank $\binom{m+d}{d}$.

To prove the previous lemma notice that the above presentation is nothing but the presentation of the cohomology ring of the usual Grassmannian $\text{Gr}(m, m + d)$. Now to conclude the proof of the proposition we use

Lemma 5. *Let $A = \mathbb{Z}[a_1, \dots, a_d]$ be a free polynomial ring generated by homogeneous elements a_i such that $\deg a_i = i$. Let I be an ideal in A generated by homogeneous elements c_1, \dots, c_d in A and $\phi : A/I \rightarrow H$ be a surjective ring homomorphism. Assume:*

C1. *H is a free \mathbb{Z} -module of rank $\prod_i \binom{\deg c_i}{\deg a_i}$.*

C2. *For every field K , the K -vector space $(A/I) \otimes_{\mathbb{Z}} K$ has finite dimension.*

Then ϕ is an isomorphism.

This result was proven in [BKT09, Lem. 1.1]. Apply it for

$$H = H^*(\text{IG}(2, 2n + 1), \mathbb{Z}), \quad I = (d_3, \dots, d_{2n}, b_n), \quad \text{and } A, \phi \text{ as above.}$$

Condition C1 is an immediate consequence of the rank calculation. For condition C2 it is enough to prove that A/I is a quotient of $A/(d_3, \dots, d_{2n+1})$. Indeed, by Lemma 4, the last module is a free \mathbb{Z} -module of finite rank. So we are left with proving that d_{2n+1} belongs to the ideal I . But the following identities of formal series hold:

$$\begin{aligned} \left(\sum_{i=0}^{2n-1} a_i t^i \right) \left(\sum_{i=0}^{2n-1} (-1)^i a_i t^i \right) &= \sum_{i=0}^{2n-1} (-1)^i b_i t^{2i}, \\ \left(\sum_{i=0}^{2n-1} (-1)^i a_i t^i \right) \left(\sum_{i \geq 0} d_i t^i \right) &= 1. \end{aligned}$$

Hence we get:

$$\sum_{i=0}^{2n-1} a_i t^i = \left(\sum_{i=0}^{2n-1} (-1)^i b_i t^{2i} \right) \left(\sum_{i \geq 0} d_i t^i \right).$$

Modding out by the ideal I , it yields:

$$\sum_{i=0}^{2n-1} a_i t^i \equiv \left(\sum_{i=0}^{n-1} (-1)^i b_i t^{2i} + \sum_{i=n+1}^{2n-1} (-1)^i b_i t^{2i} \right) \left(\sum_{i=0}^2 d_i t^i + \sum_{i \geq 2n+1} d_i t^i \right).$$

In degree $2n + 1$, we get $0 \equiv d_{2n+1}$, which ends the proof of the proposition. \square

1.8.2. Presentation in terms of τ_1 and $\tau_{1,1}$

First we will need a presentation for the symplectic Grassmannian $IG(2, 2n)$ in terms of ν_1 and $\nu_{1,1}$:

Proposition 11. *The ring $H^*(IG(2, 2n), \mathbb{Z})$ is generated by the classes $\nu_1, \nu_{1,1}$ and the relations are*

$$\frac{1}{\nu_1} \det(\nu_{1^{1+j-i}})_{1 \leq i, j \leq 2n-1} = 0,$$

$$\det(\nu_{1^{1+j-i}})_{1 \leq i, j \leq 2n} = 0.$$

Proof. We will use Lemma 5. Set $R := \mathbb{Z}[a_1, a_2]$, where $\deg a_i = i$. We denote by $\phi : R \rightarrow H^*(IG(2, 2n), \mathbb{Z})$ the surjective ring homomorphism given by $a_i \mapsto \tau_i$. We also use the convention that $a_0 = 1$ and $a_i = 0$ for $i \notin \{0, 1, 2\}$. For $r \geq 1$, set $\delta_r := \det(a_{1+j-i})_{1 \leq i, j \leq r}$. We have the recurrence relation

$$\delta_r = a_1 \delta_{r-1} - a_2 \delta_{r-2}, \tag{3}$$

which is equivalent to the identity of formal series

$$\left(\sum a_i t^i\right) \left(\sum (-1)^i \delta_i t^i\right) = 1.$$

But $\phi(a_i) = \tau_i = c_i(S^*)$. Moreover, as

$$0 \rightarrow S^\perp \rightarrow \mathcal{O}_{IG} \rightarrow S^* \rightarrow 0,$$

where we denote by S the tautological bundle on $IG(2, 2n)$, we have $c(S^\perp)c(S^*) = 1$, hence $\delta_r = c_r((S^\perp)^*) = c_r(Q)$ (Q being the quotient bundle on $IG(2, 2n)$). Since Q has rank $2n - 2$, we have $\phi(\delta_r) = 0$ for all $r > 2n - 2$, and in particular we get $\phi(\delta_{2n-1}) = \phi(\delta_{2n}) = 0$. We can write δ_{2q+1} as

$$\delta_{2q+1} = a_1 P_q(a_1, a_2),$$

where $P_q(a_1, a_2)$ is a homogeneous polynomial of degree $2q$. Now set $\delta'_{2q+1} := P_q(a_1, a_2)$. We want to prove that $\phi(\delta'_{2n-1}) = 0$. For this, since $IG(2, 2n)$ is a hyperplane section of the usual Grassmannian $Gr(2, 2n)$, we use Lefschetz's theorem. In particular, we obtain that the multiplication by the hyperplane class ν_1 is surjective from $H^{2n-2}(IG(2, 2n), \mathbb{Z})$ to $H^{2n-1}(IG(2, 2n), \mathbb{Z})$. But these vector spaces have the same dimension $n - 1$, so it is bijective. As we already know that $\phi(\delta_{2n-1}) = 0$ it implies that $\phi(\delta'_{2n-1}) = 0$. Now let $I := (\delta'_{2n-1}, \delta_{2n})$. We proved that $\phi(I) = 0$ so we may define $\bar{\phi} : R/I \rightarrow H^*(IG(2, 2n), \mathbb{Z})$. Now check that conditions C1 and C2 are satisfied:

- (C1) $H^*(IG(2, 2n), \mathbb{Z})$ is a free \mathbb{Z} -module of rank $2n(n - 1) = \frac{\deg(d_{2n-1})' \deg(d_{2n})}{\deg a_1 \deg a_2}$.
- (C2) For every field K , $(R/I) \otimes_{\mathbb{Z}} K$ is finite-dimensional. Indeed R/I is a quotient of $R/(d_{2n-1}, d_{2n})$, which is isomorphic with $H^*(Gr(2, 2n), \mathbb{Z})$, hence a free \mathbb{Z} -module of finite rank.

Finally Lemma 5 yields that $\bar{\phi}$ is an isomorphism, hence the result. \square

Now we deduce a presentation of $H^*(IG(2, 2n + 1), \mathbb{Z})$ using classes τ_1 and $\tau_{1,1}$:

Proposition 12 (Another presentation of $H^*(IG(2, 2n + 1), \mathbb{Z})$). The ring $H^*(IG(2, 2n + 1), \mathbb{Z})$ is generated by the classes $\tau_1, \tau_{1,1}$ and the relations are

$$\det(\tau_{1^{1+j-i}})_{1 \leq i, j \leq 2n} = 0,$$

$$\frac{1}{\tau_1} \det(\tau_{1^{1+j-i}})_{1 \leq i, j \leq 2n+1} = 0.$$

Proof. First notice that τ_1 and $\tau_{1,1}$ generate the cohomology ring of $IG(2, 2n + 1)$ since they are the pullbacks of the Chern classes of the dual tautological bundle over $Gr(2, 2n + 1)$ by the surjective restriction map \mathbf{j} . Then define $R := \mathbb{Z}[a_1, a_2]$, where $\deg a_i = i$. We denote by $\phi : R \rightarrow H^*(IG(2, 2n + 1), \mathbb{Z})$ the surjective ring homomorphism given by $a_i \mapsto \tau_{1^i}$. We also use the convention that $a_0 = 1$ and $a_i = 0$ for $i \notin \{0, 1, 2\}$. For $r \geq 1$, set $\delta_r := \det(a_{1+j-i})_{1 \leq i, j \leq r}$. On $Gr(2, 2n + 1)$ we know by the usual presentation (see for instance [ST97, §3]) that

$$\det(\sigma_{1^{1+j-i}})_{1 \leq i, j \leq 2n} = 0.$$

Now define δ'_{2q+1} as in the proof of Proposition 11. Using the embedding in the symplectic Grassmannian $IG(2, 2n + 2)$, we get that $\phi(\delta'_{2n+1}) = 0$. Indeed, we only have to pull back the relation $\frac{1}{v_1} \det(v_{1^{1+j-i}})_{1 \leq i, j \leq 2n+1} = 0$ proven in Proposition 11. Finally, set $l = (d_{2n}, d'_{2n+1})$ and apply Lemma 5. \square

2. Quantum cohomology

Our main goal in this section is to prove a quantum Pieri formula for $IG(2, 2n + 1)$. We denote the quantum product of two classes τ_λ and τ_μ as $\tau_\lambda \star \tau_\mu$. The degree of the quantum parameter q is equal to the index of $IG(2, 2n + 1)$, so $\deg q = 2n$.

Theorem 1 (Quantum Pieri rule for $IG(2, 2n + 1)$).

$$\tau_1 \star \tau_{a,b} = \begin{cases} \tau_{a+1,b} + \tau_{a,b+1} & \text{if } a + b \neq 2n - 3 \text{ and } a \neq 2n - 1, \\ \tau_{a,b+1} + 2\tau_{a+1,b} + \tau_{a+2,b-1} & \text{if } a + b = 2n - 3, \\ \tau_{2n-1,b+1} + q\tau_b & \text{if } a = 2n - 1 \text{ and } 0 \leq b \leq 2n - 3, \\ q(\tau_{2n-1,-1} + \tau_{2n-2}) & \text{if } a = 2n - 1 \text{ and } b = 2n - 2, \end{cases}$$

$$\tau_{1,1} \star \tau_{a,b} = \begin{cases} \tau_{a+1,b+1} & \text{if } a + b \neq 2n - 4, 2n - 3 \text{ and } a \neq 2n - 1, \\ \tau_{a+1,b+1} + \tau_{a+2,b} & \text{if } a + b = 2n - 4 \text{ or } 2n - 3, \\ q\tau_{b+1} & \text{if } a = 2n - 1 \text{ and } b \neq 2n - 3, \\ q(\tau_{2n-1,-1} + \tau_{2n-2}) & \text{if } a = 2n - 1 \text{ and } b = 2n - 3. \end{cases}$$

The previous theorem is proved in 2.5, and from this a quantum presentation is deduced in 2.6. To prove the quantum Pieri formula, we first study in 2.1 the moduli spaces of stable maps of degree 1 to $IG(2, 2n + 1)$. Then in 2.2 we describe conditions for the Gromov–Witten invariants to have enumerative meaning. Finally, in 2.3 and 2.4 we compute the invariants we need. From now on, we denote $IG(2, 2n + 1)$ by IG .

2.1. The moduli spaces $\overline{\mathcal{M}}_{0,r}(IG, 1)$

If X is a smooth projective variety we denote by $\overline{\mathcal{M}}_{g,n}(X, \beta)$ the moduli space of stable n -pointed maps f in genus g to X with degree $\beta \in H_2(X, \mathbb{Z})$. This moduli space is endowed with n evaluation maps $(ev_i)_{1 \leq i \leq n}$ mapping a stable map f to its value at the i -th marked point. We refer to [FP97] for more details. If X has Picard rank 1, which is the case when $X = IG(2, 2n + 1)$, then $\beta = dH$ for some

$d \geq 0$, H being the positive generator of the Picard group. In this situation, we will simply denote the degree as the integer d . In this section we prove

Proposition 13. *For every $r \in \mathbb{N}$, the moduli space $\overline{\mathcal{M}}_{0,r}(\text{IG}, 1)$ is a smooth projective variety. Moreover, it has the expected dimension $6n - 6 + r$.*

Proof. To prove this, we use a remark of Fulton and Pandharipande in [FP97, §0.4], which states that for all $r \geq 1$, the moduli space $\overline{\mathcal{M}}_{0,r}(\mathbb{P}^m, 1)$ is a locally trivial fibration over the variety $\text{Gr}(\mathbb{P}^1, \mathbb{P}^m)$ of lines in \mathbb{P}^m , having $\overline{\mathcal{M}}_{0,r}(\mathbb{P}^1, 1)$ as a fiber. Moreover, this last moduli space is isomorphic to the configuration space $\mathbb{P}^1[r]$ of Fulton–MacPherson. The fibration is simply the map

$$\begin{aligned} \overline{\mathcal{M}}_{0,r}(\mathbb{P}^m, 1) &\rightarrow \text{Gr}(\mathbb{P}^1, \mathbb{P}^m), \\ [f : (C; p_1, \dots, p_r) \rightarrow \mathbb{P}^m] &\mapsto f(C) \end{aligned}$$

which to any stable map of degree one associates its image line.

The Plücker embedding embeds IG as a closed subvariety of a projective space \mathbb{P}^m (with $m = (2n - 1)(n + 1)$). Under this embedding, lines in IG are lines in \mathbb{P}^m . From [FP97, §5.1], we know that this yields an embedding $\overline{\mathcal{M}}_{0,r}(\text{IG}, 1) \hookrightarrow \overline{\mathcal{M}}_{0,r}(\mathbb{P}^m, 1)$. If we denote by Y_1 the variety of lines on IG , we see that

$$[f, (C; p_1, \dots, p_r)] \in \overline{\mathcal{M}}_{0,r}(\text{IG}, 1) \iff ([f, (C; p_1, \dots, p_r)] \in \overline{\mathcal{M}}_{0,r}(\mathbb{P}^m, 1) \text{ and } f(C) \in Y_1).$$

Hence the following diagram:

$$\begin{array}{ccc} \overline{\mathcal{M}}_{0,r}(\mathbb{P}^m, 1) & \longrightarrow & \text{Gr}(\mathbb{P}^1, \mathbb{P}^m) \\ \uparrow & & \uparrow \\ \overline{\mathcal{M}}_{0,r}(\text{IG}, 1) & \longrightarrow & Y_1 \end{array}$$

is a fiber square, which means that the map $\overline{\mathcal{M}}_{0,r}(\text{IG}, 1) \rightarrow Y_1$ is locally trivial. Since the fiber $\mathbb{P}^1[r]$ is known to be smooth, we only need to prove that the variety of lines Y_1 is also smooth.

First notice that lines in IG are of the form

$$\mathcal{D}(U_1, U_3) := \{ \Sigma \in \text{IG}(2, 2n + 1) \mid U_1 \subset \Sigma \subset U_3 \}$$

where $\dim U_i = i$ and $U_3 \subset U_1^\perp$. Hence Y_1 is a subvariety of the (type A) flag variety $F(1, 3; 2n + 1)$. Let us denote by \mathcal{S}_1 and \mathcal{S}_3 the tautological bundle on $F(1, 3; 2n + 1)$ and consider the homogeneous vector bundle $\mathcal{E} := \mathcal{S}_1^* \otimes (\mathcal{S}_3/\mathcal{S}_1)^*$ on $F(1, 3; 2n + 1)$. Let also π be the projection map from the complete flag variety $F(\mathbb{C}^{2n+1})$ to the two-step flag variety $F(1, 3; 2n + 1)$. If we denote by $\mathcal{U}_1, \dots, \mathcal{U}_{2n+1}$ the tautological bundles on $F(\mathbb{C}^{2n+1})$, we see that $\mathcal{E} = \pi_*(\mathcal{U}_1^{-1} \otimes (\mathcal{U}_2/\mathcal{U}_1)^{-1})$. Hence

$$H^0(F(1, 3; 2n + 1), \mathcal{E}) = H^0(F(\mathbb{C}^{2n+1}), \mathcal{U}_1^{-1} \otimes (\mathcal{U}_2/\mathcal{U}_1)^{-1}) = \bigwedge^2 (\mathbb{C}^{2n+1})^*,$$

the last equality being a consequence of the Borel–Weil theorem. This implies that the form ω is a generic section of the vector bundle $\mathcal{S}_1^* \otimes (\mathcal{S}_3/\mathcal{S}_1)^*$ on $F(1, 3; 2n + 1)$. From the condition $U_3 \subset U_1^\perp$, it follows that the zero locus of this section is exactly the variety Y_1 . Moreover, the vector bundle \mathcal{E} is generated by its global sections. Hence Y_1 is smooth, and so is $\overline{\mathcal{M}}_{0,r}(\text{IG}, 1)$.

Finally, Y_1 having codimension 2 in $F(1, 3; 2n + 1)$, we have $\dim \overline{\mathcal{M}}_{0,r}(\text{IG}, 1) = \dim Y_1 + \dim \mathbb{P}^1[r] = 6n - 6 + r$. \square

2.2. Enumerativity of the invariants in $\overline{\mathcal{M}}_{0,2}(\mathbb{G}, 1)$ and $\overline{\mathcal{M}}_{0,3}(\mathbb{G}, 1)$

In this section we will use a Kleiman-type lemma for quasi-homogeneous spaces, due to Graber in [Gra01, Lem. 2.5]:

Lemma 6. *Let X be a variety endowed with an action of a connected algebraic group G with only finitely many orbits and Z an irreducible scheme with a morphism $f : Z \rightarrow X$. Let Y be a subvariety of X that intersects the orbit stratification properly. Then there exists a dense open subset U of G such that $\forall g \in U, f^{-1}(gY)$ is either empty or has pure dimension $\dim Y + \dim Z - \dim X$. Moreover, if X, Y and Z are smooth and we denote by Y_{reg} the subset of Y along which the intersection with the stratification is transverse, then the (possibly empty) open subset $f^{-1}(gY_{\text{reg}})$ is smooth.*

This enables us to prove the following enumerativity result for degree one Gromov–Witten invariants on $\mathbb{G}(2, 2n + 1)$.

Theorem 2 (Enumerativity of the Gromov–Witten invariants). *Let r be a positive integer and Y_1, \dots, Y_r be subvarieties of \mathbb{G} of codimension at least 2 intersecting the closed orbit generically transversely and representing cohomology classes $\gamma_1, \dots, \gamma_r$ such that*

$$\sum_{i=1}^r \text{codim } \gamma_i = \dim \overline{\mathcal{M}}_{0,r}(\mathbb{G}, 1).$$

Then there exists a dense open subset $U \subset \text{Sp}_{2n+1}^r$ such that for all $g_1, \dots, g_r \in U$, the Gromov–Witten invariant $I_1(\gamma_1, \dots, \gamma_r)$ is equal to the number of lines of \mathbb{G} incident to the translates $g_1 Y_1, \dots, g_r Y_r$.

Proof. The result is proven by successively applying the Transversality Lemma 6. First we prove that stable maps with reducible source do not contribute to the Gromov–Witten invariant by applying the lemma to the following diagram

$$\begin{array}{ccc} & & \overline{\mathcal{M}} \setminus \mathcal{M}^* \\ & & \downarrow \text{ev} \\ \underline{Y} \subset & \xrightarrow{i} & \mathbb{G}^r \end{array}$$

where $\underline{Y} = (Y_1, \dots, Y_r)$, $\text{ev} = \text{ev}_r \times \dots \times \text{ev}_1$, $\overline{\mathcal{M}} = \overline{\mathcal{M}}_{0,r}(\mathbb{G}, 1)$ and \mathcal{M}^* is the locus of map with irreducible source, which is a dense open subset by Proposition 13.

We should also prove that it is not possible for a line to be incident to one of the subvarieties Y_i in more than one point, since such a line would contribute several times to the invariant. Suppose for example that there exists a line L that intersects Y_1 in at least two points. Then any stable map f whose image curve is L corresponds to a map \tilde{f} in $\overline{\mathcal{M}}_{0,r+1}(\mathbb{G}, 1)$ (in fact in $\mathcal{M}_{0,r+1}^*(\mathbb{G}, 1)$) that contributes to the invariant $I_1(\gamma_1, \gamma_1, \dots, \gamma_r)$. By Proposition 13, $\overline{\mathcal{M}}_{0,r+1}(\mathbb{G}, 1)$ has dimension $6n - 5 + r$. Moreover, since each Y_i intersects the closed orbit generically transversely, $Y_1 \times Y_1 \times \dots \times Y_r$ intersects all $2^{r+1} \text{Sp}_{2n+1}^{r+1}$ -orbits of \mathbb{G}^{r+1} generically transversely. Hence applying Lemma 6 to the following diagram

$$\begin{array}{ccc} & & \mathcal{M}_{0,r+1}^*(\mathbb{G}, 1) \\ & & \downarrow \text{ev} \\ Y_1 \times Y_1 \times \dots \times Y_r \subset & \xrightarrow{i} & \mathbb{G}^{r+1} \end{array}$$

and using the fact that $\text{codim } \gamma_1 \geq 2$ we conclude that such a line cannot exist.

Now using

$$\begin{array}{ccc} & & \mathcal{M}^* \\ & & \downarrow \text{ev} \\ \text{Sing } \underline{Y} \subset \xrightarrow{i} & & \mathbb{G}^r \end{array}$$

where $\text{Sing } \underline{Y}$ denotes the singular locus of \underline{Y} , we may assume that \underline{Y} is smooth. Moreover, since Y_1, \dots, Y_r intersect the closed orbit generically transversely, a third application of Lemma 6 allows us to assume that this intersection is transverse everywhere. Finally, applying the lemma to

$$\begin{array}{ccc} & & \mathcal{M}^* \\ & & \downarrow \text{ev} \\ \underline{Y} \subset \xrightarrow{i} & & \mathbb{G}^r \end{array}$$

we conclude that there exists a dense open subset $U \subset \text{Sp}_{2n+1}^r$ such that for all $(g_1, \dots, g_r) \in U$, $\bigcap_{i=1}^r \text{ev}_i^{-1}(g_i Y_i)$ is a finite number of reduced points, which equals the number of lines incident to all the $g_i Y_i$. \square

Remark 4. Theorem 2 enables us to compute the Gromov–Witten invariants by geometric means. However, Schubert varieties will not be appropriate to perform this calculation. Indeed, the intersection of any Schubert variety and the closed orbit is not even proper. So we will instead use the restrictions of the Schubert varieties of the usual Grassmannian.

2.3. Computation of the invariants in $\overline{\mathcal{M}}_{0,2}(\mathbb{G}, 1)$

In this paragraph, we use Theorem 2 to compute all invariants of \mathbb{G} of the form $I_1(\alpha, \beta)$, where α and β are the classes of the restriction to \mathbb{G} of some Schubert varieties Y_1 and Y_2 of the usual Grassmannian, defined with respect to complete flags F_\bullet and G_\bullet . In order for the varieties Y_1 and Y_2 to verify the conditions of the theorem, we will need some technical conditions to hold for the defining flags F_\bullet and G_\bullet . We state these conditions in Lemma 7 and prove that they hold for generic flags: this is quite straightforward, and the list of conditions is in fact longer to state than to check. Then we compute the invariants in Proposition 14.

Notation 1. Denote by

- \mathbb{F}_n the variety of complete flags in \mathbb{C}^{2n+1} ;
- Λ_n the variety of antisymmetric 2-forms with maximal rank on \mathbb{C}^{2n+1} .

Lemma 7. Assume $n \geq 2$. Then the set of triples $(F_\bullet, G_\bullet, \omega) \in \mathbb{F}_n \times \mathbb{F}_n \times \Lambda_n$ such that the following hold

- (C1) $\forall 0 \leq p \leq 2n + 1, \omega|_{F_p}$ has maximal rank;
- (C2) $\forall 0 \leq p \leq 2n + 1, \omega|_{G_p}$ has maximal rank;
- (C3) $\forall 0 \leq p, q \leq 2n + 1, F_p \cap G_q$ has the expected dimension;
- (C4)_i $\dim(F_{2n+1-i} \cap G_{i+3} \cap F_1^\perp \cap G_1^\perp) = 1; (0 \leq i \leq 2n - 2);$
- (C5)_i $\dim F_{2n-i} \cap G_{i+3} \cap G_1^\perp = 1$ and $\dim(F_{2n-i} \cap G_{i+3} \cap G_1^\perp)^\perp \cap F_2 = 1; (0 \leq i \leq 2n - 2);$
- (C6)_i $\dim F_{2n+1-i} \cap G_{i+2} \cap F_1^\perp = 1$ and $\dim(F_{2n+1-i} \cap G_{i+2} \cap F_1^\perp)^\perp \cap G_2 = 1; (2 \leq i \leq 2n - 4);$
- (C7)_i $\dim(F_{2n-i} \cap G_{i+2})^\perp \cap F_2 = 1; (2 \leq i \leq 2n - 4);$

- (C8)_i $\dim(F_{2n-i} \cap G_{i+2})^\perp \cap G_2 = 1; (2 \leq i \leq 2n - 4);$
- (C9) $F_1 \not\subset G_1^\perp;$
- (C10) $G_1 \not\subset F_1^\perp;$
- (C11)_i $F_{2n-1-i} \cap G_{i+3} \cap G_1^\perp = 0; (0 \leq i \leq 2n - 6);$
- (C12)_i $F_{2n+1-i} \cap G_{i+1} \cap F_1^\perp = 0; (4 \leq i \leq 2n - 2);$

is a dense open subset in $\mathbb{F}_n \times \mathbb{F}_n \times \Lambda_n$.

Proof. $\mathbb{F}_n \times \mathbb{F}_n \times \Lambda_n$ is a (quasi-projective) irreducible variety. Moreover all conditions are clearly open. So it is enough to show that each of them is non-empty.

(C1), (C2) and (C3) Obvious.

(C4)_i Since $n \geq 2$, we may choose the flags F_\bullet and G_\bullet such that the subspace $A := F_{2n+1-i} \cap G_{i+3}$ has dimension 3 and A together with the lines $L := F_1$ and $L' := G_1$ are in direct sum. Then there exists a form $\omega \in \Lambda_n$ such that $A \cap L^\perp \cap L'^\perp$ has dimension 1.

(C5)_i As before we may choose F_\bullet and G_\bullet such that $A := F_{2n-i} \cap G_{i+3}$ has dimension 2 and $A, L := G_1$ and $B := F_2$ are complementary. So we may construct $\omega \in \Lambda_n$ such that $(A \cap L^\perp)^\perp \cap B$ has dimension 1. First construct ω_0 on $A \oplus B \oplus L$. Let $a \in A \setminus 0$ and $b \in B \setminus 0$. There exists ω_0 a symplectic form on $A \oplus B$ such that $\omega_0(a, b) \neq 0$. Then we extend ω_0 to ω defined on $A \oplus B \oplus L$ by setting $\omega(a, l) = 0, \omega(a', l) \neq 0$ and for instance $\omega(\beta, l) = 0$ for all $\beta \in B$, where l generates L and a, a' generate A .

(C6)_i As in (C5)_i.

(C7)_i We may choose F_\bullet and G_\bullet such that $L := F_{2n-i} \cap G_{i+2}$ has dimension 1 and is in direct sum with $A := F_2$. But then there exists $\omega \in \Lambda_n$ such that $A \not\subset L^\perp$.

(C8)_i As in (C7)_i.

(C9) G_1^\perp is a general hyperplane, so it does not contain F_1 .

(C10) As in (C9).

(C11)_i $F_{2n-1-i} \cap G_{i+3}$ is a line and G_1^\perp is a general hyperplane, so their intersection is zero.

(C12)_i As in (C11)_i. \square

We can now define the varieties we will use to compute the invariants, which will be restrictions of the Schubert varieties of the usual Grassmannian:

Lemma 8. Let $0 \leq j \leq n - 1$ and $0 \leq i \leq 2n - 1 - 2j$ be integers. Let

$$X_{i,j} := \{\Sigma \in \text{Gr} \mid \Sigma \cap F_{j+1} \neq 0, \Sigma \subset F_{2n+1-i-j}\},$$

be a subvariety of $\text{Gr} := \text{Gr}(2, 2n + 1)$, where F_\bullet is a complete flag satisfying condition (C1). Then:

1. $X_{i,j}$ and IG intersect generically transversely.
2. Let $Y_{i,j} := X_{i,j} \cap \text{IG}$. We have

$$[Y_{i,j}]^{\text{IG}} = \begin{cases} \tau_{2n-1-j,i+j} + \tau_{2n-j,i+j-1} & \text{if } j \neq 0 \text{ and } i \neq 2n - 1 - 2j, \\ \tau_{2n-j,2n-2-j} & \text{if } j \neq 0 \text{ and } i = 2n - 1 - 2j, \\ \tau_{2n-1,i} & \text{if } j = 0 \text{ and } i \neq 2n - 1, \\ 0 & \text{if } j = 0 \text{ and } i = 2n - 1, \end{cases}$$

where we denote by $[V]^{\text{IG}}$ (respectively by $[V]^{\text{Gr}}$) the class of the subvariety V in IG (respectively in Gr).

Proof. 1. In the Schubert cell $C_{i,j} \subset X_{i,j}$, a direct computation shows that $T_p X_{i,j} \not\subset T_p \text{IG}$ as soon as $F_{j+1} \not\subset F_{2n+1-i-j}^\perp$, which is true by condition (C1). So $C_{i,j} \cap \text{IG}$ is transverse. Applying again (C1), we notice that $C_{i,j} \cap \text{IG}$ is an open subset of $X_{i,j} \cap \text{IG}$.

2. We have $[X_{i,j}]^{\text{Gr}} = \sigma_{2n-1-j,i+j}$. Moreover, the previous item implies that $[Y_{i,j}]^{\text{Gr}} = \sigma_1[X_{i,j}]^{\text{Gr}}$. So

$$[Y_{i,j}]^{\text{Gr}} = \begin{cases} \sigma_{2n-1-j,i+j+1} + \sigma_{2n-j,i+j} & \text{if } j \neq 0 \text{ and } i \neq 2n-1-2j, \\ \sigma_{2n-j,2n-1-j} & \text{if } j \neq 0 \text{ and } i = 2n-1-2j, \\ \sigma_{2n-1,i+1} & \text{if } j = 0 \text{ and } i \neq 2n-1, \\ 0 & \text{if } j = 0 \text{ and } i = 2n-1. \end{cases}$$

Moreover, $[Y_{i,j}]^{\text{Gr}} = \mathbf{j}_* [Y_{i,j}]^{\text{IG}}$, $[Y_{i,j}]^{\text{IG}} = \sum_{p=0}^{\lfloor n-1-\frac{i}{2} \rfloor} \alpha_p \tau_{2n-1-p,i+p}$ and $\mathbf{j}_* \tau_{a,b} = \sigma_{a,b+1}$ for $a+b \geq 2n-1$, so we can determine the α_p by identifying both expressions. \square

We now assume all genericity conditions (C1)–(C12) are satisfied and prove

Proposition 14. *Let $0 \leq i \leq 2n-2$, $0 \leq 2j \leq 2n-2-i$ and $0 \leq 2l \leq i$ be integers. Set $Y_1 := Y_{i,j}(F_\bullet)$ and $Y_2 := Y_{2n-2-i,l}(G_\bullet)$, where the complete flags F_\bullet and G_\bullet as well as the form ω verify the transversality conditions of Lemma 7. Then*

1. *The intersections $Y_1 \cap \mathbb{O}$ and $Y_2 \cap \mathbb{O}$ are transverse. Moreover*

$$Y_1 \cap \mathbb{O} = \begin{cases} \emptyset & \text{if } i \text{ or } j \neq 0, \\ \{F_1 \oplus K\} & \text{if } i = j = 0, \end{cases}$$

$$Y_2 \cap \mathbb{O} = \begin{cases} \emptyset & \text{if } i \neq 2n-2 \text{ or } l \neq 0, \\ \{G_1 \oplus K\} & \text{if } i = 2n-2 \text{ and } l = 0. \end{cases}$$

2. *If j or $l \geq 2$, there exists no line passing through Y_1 and Y_2 . Else there exists a unique line passing through Y_1 and Y_2 . Therefore the Gromov–Witten invariant $I_1(\tau_{a,b}, \tau_{c,d})$ equals 1 if $a = c = 2n-1$ and $b + c = 2n-2$, and 0 in the other cases.*

Proof. 1. $Y_1 \cap \mathbb{O} = \{\Sigma \in \text{IG} \mid \Sigma \cap F_{j+1} \neq \emptyset, K \subset \Sigma \subset F_{2n+1-i-j}\}$, so if $i + j \neq 0$, then $K \subset F_{2n+1-i-j}$, which, according to (C1), implies that $Y_1 \cap \mathbb{O} = \emptyset$, so the intersection is transverse. Moreover if $i + j = 0$ we get $Y_1 \cap \mathbb{O} = \{F_1 \oplus K\}$. Denote by Σ_0 the point $K \oplus F_1$. To prove transversality at Σ_0 we use the embedding in the usual Grassmannian $\text{Gr} := \text{Gr}(2, 2n+1)$. It is well known that $T_{\Sigma_0} \text{Gr} = \text{Hom}(\Sigma_0, \mathbb{C}^{2n+1}/\Sigma_0)$. Now express $T_{\Sigma_0} Y_1$ and $T_{\Sigma_0} \mathbb{O}$ as subspaces of $T_{\Sigma_0} \text{Gr}$:

$$T_{\Sigma_0} Y_1 = \{\phi \in T_{\Sigma_0} \text{Gr} \mid \phi(f_1) = 0\},$$

$$T_{\Sigma_0} \mathbb{O} = \{\phi \in T_{\Sigma_0} \text{Gr} \mid \phi(k) = 0\},$$

where f_1 and k generate F_1 and K . We see that these subspaces are complementary in $T_{\Sigma_0} \text{Gr}$. Computing $\dim Y_1 = 2n-2$ and $\dim \mathbb{O} = 2n-1$ we conclude that they generate $T_{\Sigma_0} \text{IG}$. We can proceed in a similar fashion for $Y_2 \cap \mathbb{O}$.

2. We first study the case where j or $l \geq 2$. Let $\mathcal{D} := \mathcal{D}(V, W)$ be a line meeting Y_1 and Y_2 . Then we must have $V \subset F_{2n+1-i-j} \cap G_{i+3-l}$. But according to (C3), this subspace is either zero or it has codimension $2n+4-j-l$. So for $j+l \geq 3$, it is zero and there is no line. If $j=2$ and $l=0$ (and symmetrically if $j=0$ and $l=2$), we must have $V \subset F_{2n-1-i} \cap G_{i+3} \cap G_1^\perp = 0$, which is impossible by $(\text{C11})_i$ (respectively by $(\text{C12})_i$). So for a line to exist we must have j and $l \leq 1$.

Now assume $j, l \leq 1$. There are four cases to study:

- a) $j = l = 0$;
- b) $j = 1, l = 0$;
- c) $j = 0, l = 1$;
- d) $j = l = 1$.

a) Let $A = F_{2n+1-i} \cap G_{i+3}$. We have $\dim A = 3$ by (C3). But $V \subset A$ and $V \subset F_1^\perp \cap G_1^\perp$ since $F_1, G_1 \subset W$ and $W \subset V^\perp$. By (C4)_i, we have $\dim A \cap F_1^\perp \cap G_1^\perp = 1$, hence $V = A \cap F_1^\perp \cap G_1^\perp$. So $W \supset V + (F_1 \oplus G_1)$ (F_1 and G_1 are in direct sum by (C3)). To show equality, it is enough to prove that the sum is direct. If not then there exists a non-zero vector of the form $af_1 + bg_1$ in V , where f_1 and g_1 generate F_1 and G_1 . So $af_1 + bg_1 \in A \subset F_{2n+1-i}$, which implies $bg_1 \in F_{2n+1-i}$, hence $b = 0$ or $i = 0$. If $b = 0$, then $V = F_1$, and consequently $F_1 \subset G_1^\perp$, which is impossible by (C9). So $i = 0$. But then $af_1 + bg_1 \in G_3$, so $af_1 \in G_3$ and also $a = 0$. Hence $V = G_1 \subset F_1^\perp$, which is excluded by (C9).

b) Let $A = F_{2n-i} \cap G_{i+3}$. By (C3), $\dim A = 2$. By (C5)_i, $\dim A \cap G_1^\perp = 1$, so $V = A \cap G_1^\perp$. Moreover $\dim V^\perp \cap F_2 = 1$. We have $W \supset V + G_1 + V^\perp \cap F_2$. To determine W , it is enough to show that the sum is direct. First, $V + G_1$ is direct, because if it was not we would have $V = G_1$, so $G_1 \subset F_{2n-i}$, which is impossible by (C3). Finally the sum $V \oplus G_1 + V^\perp \cap F_2$ is direct, or we would have $V^\perp \cap F_2 \subset G_{i+3}$. But $\dim F_2 \cap G_{i+3} = 0$ by (C3) since $i \leq 2n - 4$.

c) This case is similar to 2b); the proof uses (C3) and (C6)_i.

d) By (C3), we get $\dim F_{2n-1} \cap G_{i+2} = 1$, so $V = F_{2n-1} \cap G_{i+2}$. We must have $\dim W \cap F_2 \neq 0$. But $V \not\subset F_2$, or else we would get $G_{i+2} \cap F_2 \neq 0$, which is impossible by (C3) since $i \leq 2n - 4$. Now $W \subset V^\perp$ implies $W \cap F_2 \subset V^\perp \cap F_2$, which has dimension 1 by (C7)_i. So $W \subset V^\perp \cap F_2 \oplus V$. Similarly, using (C8)_i, we get $W \cap G_2 = V^\perp \cap G_2$, so $W \supset V \oplus V^\perp \cap F_2 + V^\perp \cap G_2$. Now we only have to show that this sum is direct. If not, then there exists a non-zero vector of the form $av + bf_2$ in $V^\perp \cap G_2$, where v and f_2 generate V and $V^\perp \cap F_2$. As $v \in G_{i+2}$, we obtain $bf_2 \in G_{i+2}$, so $b = 0$ because $i \leq 2n - 4$. Hence $V^\perp \cap G_2 = V$ and consequently $V \subset G_2$ and $\dim F_{2n-i} \cap G_2 \leq 1$, which is impossible since $i \geq 2$.

The final formula for $I_1(\tau_{a,b}, \tau_{c,d})$ follows from a straightforward calculation. \square

2.4. Computation of some invariants in $\overline{\mathcal{M}}_{0,3}(\text{IG}, 1)$

In the previous section we computed the two-pointed invariants in IG, which is equivalent to computing the quantum terms of the product by the hyperplane class τ_1 . Indeed, the divisor axiom [KM94, §2.2.4] yields

$$I_1(\gamma_1, \gamma_2, \tau_1) = I_1(\gamma_1, \gamma_2),$$

where γ_1 and γ_2 are any cohomology classes. Hence to obtain a quantum Pieri rule for $\text{IG}(2, 2n + 1)$, we are left to compute the quantum product by $\tau_{1,1}$. So we have to determine all invariants of the form $I_1(\tau_{1,1}, \tau_\lambda, \tau_\mu)$ with $|\lambda| + |\mu| = 6n - 5$, that is to compute the number of lines through the following subvarieties:

$$Y_1 = \{\Sigma \in \text{IG} \mid \Sigma \cap F_{j+1} \neq \emptyset, \Sigma \subset F_{2n+2-i-j}\},$$

$$Y_2 = \{\Sigma \in \text{IG} \mid \Sigma \cap G_{l+1} \neq \emptyset, \Sigma \subset G_{i+3-l}\},$$

$$Y_3 = \{\Sigma \in \text{IG} \mid \Sigma \subset H\},$$

where $0 \leq i \leq 2n - 1, 0 \leq 2j \leq 2n - 1 - i$ and $0 \leq 2l \leq i$ are integers, F_\bullet and G_\bullet are isotropic flags and H is a hyperplane.

As before we use a genericity result which is proved in a similar way as Lemma 7:

Lemma 9. Assume $n \geq 2$. Then the set of 4-uples $(F_\bullet, G_\bullet, H, \omega) \in \mathbb{F}_n \times \mathbb{F}_n \times \mathbb{P}^{2n} \times \Lambda_n$ satisfying the following conditions

(C1) $\forall 0 \leq p \leq 2n + 1, \omega|_{F_p}$ has maximal rank;

(C2) $\forall 0 \leq p \leq 2n + 1, \omega|_{G_p}$ has maximal rank;

(C3) $\omega|_H$ is symplectic;

(C4) $\forall 0 \leq p, q \leq 2n + 1, F_p \cap G_q$ has the expected dimension;

(C5) $\forall 0 \leq p, q \leq 2n + 1, F_p \cap G_q \cap H$ has the expected dimension;

- (C6)_i $\dim(F_{2n+2-i} \cap G_{i+3} \cap H \cap F_1^\perp \cap G_1^\perp) = 1; (1 \leq i \leq 2n - 2);$
- (C7)_i $\dim F_{2n+1-i} \cap G_{i+3} \cap H \cap G_1^\perp = 1$ and $\dim(F_{2n+1-i} \cap G_{i+3} \cap H \cap G_1^\perp)^\perp \cap F_2 = 1; (0 \leq i \leq 2n - 3);$
- (C8)_i $\dim F_{2n+2-i} \cap G_{i+2} \cap H \cap F_1^\perp = 1$ and $\dim(F_{2n+2-i} \cap G_{i+2} \cap H \cap F_1^\perp)^\perp \cap G_2 = 1; (2 \leq i \leq 2n - 1);$
- (C9)_i $\dim(F_{2n+1-i} \cap G_{i+2} \cap H)^\perp \cap F_2 = 1; (2 \leq i \leq 2n - 3);$
- (C10)_i $\dim(F_{2n+1-i} \cap G_{i+2} \cap H)^\perp \cap G_2 = 1; (2 \leq i \leq 2n - 3);$
- (C11) $F_1 \not\subset G_1^\perp;$
- (C12) $G_1 \not\subset F_1^\perp;$
- (C13)_i $F_{2n-i} \cap G_{i+3} \cap H \cap G_1^\perp = 0; (0 \leq i \leq 2n - 5);$
- (C14)_i $F_{2n+2-i} \cap G_{i+1} \cap H \cap F_1^\perp = 0; (4 \leq i \leq 2n - 1);$
- (C15)_i $F_2 \cap G_{i+3} \cap G_1^\perp = 0; 0 \leq i \leq 2n - 3;$
- (C16)_i $G_2 \cap F_{2n+2-i} \cap F_1^\perp = 0; 2 \leq i \leq 2n - 1;$

is a dense open subset of $\mathbb{F}_n \times \mathbb{F}_n \times \mathbb{P}^{2n} \times \Lambda_n$.

Under these assumptions we can prove

Proposition 15.

1. The intersections $Y_i \cap \mathbb{O}$ are transverse. Moreover

$$Y_1 \cap \mathbb{O} = \begin{cases} \emptyset & \text{if } i + j \geq 2, \\ \{F_1 \oplus K\} & \text{if } i = 1 \text{ and } j = 0, \\ \{K \oplus L \mid L \subset F_2\} & \text{if } i = 0 \text{ and } j = 1, \end{cases}$$

$$Y_2 \cap \mathbb{O} = \begin{cases} \emptyset & \text{and } i \neq 2n - 2 \text{ or } l \neq 0, \\ \{G_1 \oplus K\} & \text{if } i = 2n - 2 \text{ and } l = 0, \end{cases}$$

$$Y_3 \cap \mathbb{O} = \emptyset.$$

- 2. If j or $l \geq 2$, there is no line meeting Y_1, Y_2 and Y_3 .
- 3. If j and $l \leq 1$, there is a unique line meeting Y_1, Y_2 and Y_3 . Therefore the Gromov–Witten invariant $I_1(\tau_{1,1}, \tau_{a,b}, \tau_{c,d})$ equals 1 if $a = c = 2n - 1$ and $b + c = 2n - 3$, and 0 in the other cases.

Proof. 1. The case of $Y_2 \cap \mathbb{O}$ has already been treated in the proof of Proposition 14. If $\Sigma \in Y_1 \cap \mathbb{O}$, we must have $K \subset F_{2n+2-i-j}$, so $i + j = 1$. If $i = 1$ and $j = 0$, then $Y_1 \cap \mathbb{O} = \{K \oplus F_1\}$, and transversality is proven as in Proposition 14. If $i = 0$ and $j = 1$, then $Y_1 \cap \mathbb{O} = \{K \oplus L \mid L \subset F_2\}$. Take $\Sigma_0 = K \oplus \langle f_2 \rangle$ where f_2 is a non-zero element in F_2 . Again we express $T_{\Sigma_0} Y_1$ and $T_{\Sigma_0} \mathbb{O}$ as subspaces of $T_{\Sigma_0} \text{Gr}$, where Gr is the usual Grassmannian:

$$T_{\Sigma_0} Y_1 = \{ \phi \in T_{\Sigma_0} \text{Gr} \mid \phi(f_2) \in F_2 / \langle f_2 \rangle, \phi(k) \perp f_2 \},$$

$$T_{\Sigma_0} \mathbb{O} = \{ \phi \in T_{\Sigma_0} \text{Gr} \mid \phi(k) = 0 \},$$

with k a generator of K . We see that the intersection of $T_{\Sigma_0} Y_1$ and $T_{\Sigma_0} \mathbb{O}$ has dimension 1. Computing $\dim Y_1 = 2n - 1$ and $\dim \mathbb{O} = 2n - 1$ we conclude that they generate $T_{\Sigma_0} \text{IG}$. Finally, $Y_3 \cap \mathbb{O} = \emptyset$ since $K \not\subset H$ by (C3).

2. By (C5), $F_{2n+2-i-j} \cap G_{i+3-l} \cap H = 0$ as soon as $j + l \geq 3$. Moreover if $j = 2$ and $l = 0$ then we get $W \supset G_1$, hence $V \subset F_{2n-i} \cap G_{i+3} \cap H \cap G_1^\perp$. But this space is zero by (C13)_i, so there is no line. By (C13)_i, we get the same result when $j = 0$ and $l = 2$.

3. There are four cases:

- a) $j = l = 0$;
- b) $j = 1, l = 0$;
- c) $j = 0, l = 1$;
- d) $j = l = 1$.

a) We have $V = F_{2n+2-i} \cap G_{i+3} \cap H \cap F_1^\perp \cap G_1^\perp$ by (C6)_i. Moreover $W \supset V + F_1 + G_1$. To obtain equality we only have to show that the sum is direct. First $V \neq F_1$ since $F_1 \not\subset G_1^\perp$ by (C11). Finally if $G_1 \subset V \oplus F_1$, as $V \subset F_1^\perp$, we would have $G_1 \subset F_1^\perp$, which is impossible by (C12).

b) We have $V = F_{2n+1-i} \cap G_{i+3} \cap H \cap G_1^\perp$ by (C7)_i. Moreover $W \subset V + G_1 + F_2 \cap V^\perp$. We prove now that this sum is direct. First $V \neq G_1$, or we would have $G_1 \subset H$, which is excluded by (C5). Now $F_2 \cap V^\perp \not\subset V \oplus G_1$ since $F_2 \cap G_{i+3} \cap G_1^\perp = 0$ for $i \leq 2n - 3$ by (C15)_i.

c) $V = F_{2n+2-i} \cap G_{i+2} \cap H \cap F_1^\perp$ by (C8)_i. Moreover $W \supset V + F_1 + G_2 \cap V^\perp$ (by (C9)_i and (C10)_i), and this sum is direct (same argument than in the previous case, using condition (C16)_i).

d) $V = F_{2n+1-i} \cap G_{i+2} \cap H, W \supset V + W \cap F_2 + W \cap G_2 = V + F_2 \cap V^\perp + G_2 \cap V^\perp$. This sum is direct; indeed, $F_2 \cap V^\perp \neq G_2 \cap V^\perp$ car $F_2 \cap G_2 = 0$ by (C4); in addition $V \not\subset F_2 \cap V^\perp \oplus G_2 \cap V^\perp$, or we would get $G_2 \cap F_{2n+1-i} \neq 0$, which is impossible by $i \geq 2$.

The final formula for $I_1(\tau_{1,1}, \tau_{a,b}, \tau_{c,d})$ follows from a straightforward calculation. \square

2.5. Quantum Pieri rule

We can now prove Theorem 1:

Proof of Theorem 1. We start with the invariants $I_1(\tau_1, \tau_{a,b}, \tau_{c,d})$, which are equal to the two-pointed invariants $I_1(\tau_{a,b}, \tau_{c,d})$ because of the divisor axiom. The first item of Proposition 14 enables us to apply the Enumerativity Theorem 2. Then we use the second item of Proposition 14. For $j = l = 0$ we get that for all $0 \leq i \leq 2n - 2$, we have $I_1(\tau_{2n-1,i}, \tau_{2n-1,2n-2-i}) = 1$. Then setting $j = 0$ and $l > 0$ we recursively get $I_1(\tau_{2n-1,i}, \tau_{2n-1-l,2n-2-i+l}) = 0$ (for all i and $l > 0$). Finally, setting j and $l > 0$ we get $I_1(\tau_{2n-1-j,i+j}, \tau_{2n-1-l,2n-2-i+l}) = 0$ (for all i and $j, l > 0$). Hence:

$$I_1(\tau_1, \tau_{a,b}, \tau_{c,d}) = \begin{cases} 1 & \text{if } a = c = 2n - 1, \\ 0 & \text{if } a \text{ or } c < 2n - 1. \end{cases}$$

Similarly, Proposition 15 and Theorem 2 imply

$$I_1(\tau_{1,1}, \tau_{a,b}, \tau_{c,d}) = \begin{cases} 1 & \text{if } a = c = 2n - 1, \\ 0 & \text{if } a \text{ or } c < 2n - 1. \end{cases}$$

Using the classical Pieri rule and Poincaré duality, we get our result. \square

Using the quantum Pieri formula we can fill out the Hasse diagram from Fig. 1 to obtain the quantum Hasse diagram of $IG(2, 7)$ in Fig. 4. As a comparison see the quantum Hasse diagram of $IG(2, 6)$ in Fig. 5.

2.6. Quantum presentation

Proposition 16 (Presentation of $QH^*(IG(2, 2n + 1), \mathbb{Z})$). *The ring $QH^*(IG(2, 2n + 1), \mathbb{Z})$ is generated by the classes $\tau_1, \tau_{1,1}$ and the quantum parameter q . The relations are*

$$\det(\tau_{1+j-i})_{1 \leq i, j \leq 2n} = 0,$$

$$\frac{1}{\tau_1} \det(\tau_{1+j-i})_{1 \leq i, j \leq 2n+1} + q = 0.$$




Fig. 4. Quantum Hasse diagram of IG(2, 7).




Fig. 5. Quantum Hasse diagram of IG(2, 6).

Proof. Siebert and Tian proved in [ST97, Prop. 2.2] that the quantum relations are obtained by evaluating the classical relations using the quantum product. Define δ_{2n} and δ'_{2n+1} as in the proof of Proposition 11 and denote by $\overline{\delta_{2n}}$ and $\overline{\delta'_{2n+1}}$ the same expressions with the cup product replaced by the quantum product.

Now we consider the quantum products $\Pi_a := (\tau_1)^{2(n-a)} \star (\tau_{1,1})^a$ for $0 \leq a \leq n$. For reasons of degree it has no q -term of degree greater than 1. First we prove that Π_a has no q -term if $a \neq 0, 1$. To prove this, we decompose Π_a for $a > 0$ as

$$\Pi_a = \tau_{1,1} \star ((\tau_1)^{2(n-a)} (\tau_{1,1})^{a-1}).$$

Notice that for degree reasons, $(\tau_1)^{2(n-a)} (\tau_{1,1})^{a-1}$ has no q -term. Moreover, if $a \geq 2$, the classical Pieri formula 4 implies that this product contains only classes $\tau_{c,d}$ with $c < 2n - 1$. Then we use the quantum Pieri formula 1 to conclude that there is no q -term in Π_a . We are now left with computing the q -term of Π_0 and Π_1 . Set $\alpha_p := (\tau_1)^p$ for $p \leq 2n - 1$. α_p has no q -term. We have $\Pi_0 = \tau_1 \star \alpha_{2n-1}$ and $\Pi_1 = \tau_{1,1} \star \alpha_{2n-2}$. We compute recursively the coefficients of τ_p and $\tau_{p-1,1}$ for $p \leq 2n - 3$ in α_p using the classical Pieri rule. We find

$$\alpha_p = \tau_p + (p - 1)\tau_{p-1,1} + \text{terms with lower first part.}$$

Then

$$\alpha_{2n-2} = \tau_{2n-1,-1} + (2n - 2)\tau_{2n-2} + \text{terms with lower first part}$$

and

$$\alpha_{2n-1} = (2n - 1)\tau_{2n-1} + \text{terms with lower first part.}$$

Finally we use the quantum Pieri rule to deduce that

$$\Pi_0 = \text{classical terms} + (2n - 1)q,$$

$$\Pi_1 = \text{classical terms} + q.$$

But

$$\begin{aligned} \overline{\delta_{2n}} &= \Pi_0 - (2n - 1)\Pi_1 + \text{linear combination of } \Pi_a \text{'s with } a \geq 2, \\ \overline{\delta'_{2n+1}} &= \Pi_0 - 2n\Pi_1 + \text{linear combination of } \Pi_a \text{'s with } a \geq 2, \end{aligned}$$

hence $\overline{\delta_{2n}} = \delta_{2n}$ and $\overline{\delta'_{2n+1}} = \delta'_{2n+1} - q$. \square

2.7. Around a conjecture of Dubrovin

In 1994, Dubrovin stated a conjecture relating properties of the quantum cohomology of Fano varieties and properties of their derived category:

Conjecture 1. (See Dubrovin [Dub98, Conj. 4.2.2].) *Let X be a Fano variety. The big quantum cohomology of X is generically semisimple if and only if its derived category of coherent sheaves $\mathcal{D}^b(X)$ admits a full exceptional collection.*

Let us recall some definitions involved in the statement of this conjecture:

Definition 2.

1. An (ordered) collection (E_1, \dots, E_m) of objects in $\mathcal{D}^b(X)$ is *exceptional* if $\text{Hom}_{\mathcal{D}^b(X)}(E_i, E_j) = 0$ for all $i > j$ and $\text{Hom}_{\mathcal{D}^b(X)}(E_i, E_i) = \mathbb{C}$ for all i .
2. A collection (E_1, \dots, E_m) of objects in $\mathcal{D}^b(X)$ is *full* if it generates the derived category, i.e. if the smallest triangulated subcategory of $\mathcal{D}^b(X)$ containing the collection is $\mathcal{D}^b(X)$ itself.
3. A finite-dimensional associative algebra is *semisimple* if it has no non-trivial nilpotent ideals. Moreover, the (big or small) quantum cohomology ring $\text{QH}^*(X)$ of a smooth projective variety X is said to be *generically semisimple* if it is semisimple as an algebra for generic values of the quantum parameters.

In this section, we check the conjecture for odd symplectic Grassmannians of lines. We first show that the (small) quantum cohomology ring of $\text{IG}(2, 2n + 1)$, localized at $q \neq 0$, is semisimple. To do this we adapt the presentation of Proposition 16 to make the symmetries more apparent:

Theorem 3.

1. The ring $\text{QH}^*(\text{IG}(2, 2n + 1), \mathbb{Z})$ is isomorphic to $R^{\mathbb{C}^2}$, where

$$R = \mathbb{Z}[x_1, x_2, q]/(h_{2n}(x_1, x_2), h_n(x_1^2, x_2^2) + q)$$

and x_1 and x_2 are the Chern roots of the tautological bundle S and $h_r(y_1, y_2)$ is the r -th complete symmetric function of the variables y_1, y_2 .

2. $\text{QH}^*(\text{IG}(2, 2n + 1), \mathbb{Z})_{q \neq 0}$ is semisimple.

Proof. 1. We use the recurrence relation (3) from Proposition 11 to prove that $\delta_r = h_r(x_1, x_2)$ for all r . Then

$$\delta'_{2n+1} = \frac{h_{2n+1}(x_1, x_2)}{x_1 + x_2} = h_n(x_1^2, x_2^2).$$

2. It is enough to prove the semisimplicity of R localized at $q \neq 0$. We may assume $q = -1$. Using $(x_1 - x_2)h_{2n}(x_1, x_2) = x_1^{2n+1} - x_2^{2n+1}$ and noticing that we must have $x_2 \neq 0$, the first relation implies that $x_1 = \zeta x_2$, where $\zeta \neq 1$ is a $(2n + 1)$ -th root of unity. Replacing in the second relation $h_n(x_1^2, x_2^2) - 1 = 0$, we get $x_1^{2n} = 1 + \zeta$. Since $\zeta \neq -1$, this equation has $2n$ distinct solutions. So we have $2n$ distinct solutions for x_1 , and for each x_1 we have $2n$ distinct solutions for x_2 , which gives us (at least) $4n^2$ distinct solutions for the pair (x_1, x_2) . But the number of solutions, counted with their multiplicity, should be equal to twice the rank of $H^*(\text{IG}(2, 2n + 1), \mathbb{Z})$, which is equal to $2n^2$. So there are no other solutions, and all solutions are simple. Hence the semisimplicity. \square

Since semisimplicity of the small quantum cohomology implies generic semisimplicity of the big one (see [Cio05, Lem. 3]), Theorem 3 proves that the big quantum cohomology of $\text{IG}(2, 2n + 1)$ is generically semisimple. So to confirm Dubrovin’s conjecture in this case it is enough to find a full exceptional collection.

In [Kuz08], Kuznetsov computed full exceptional collections for the symplectic Grassmannian of lines. His method can easily be adapted to the odd symplectic case. Here we denote the tautological bundle by \mathcal{U} instead of \mathcal{S} to avoid confusion with symmetric powers. We denote by \mathcal{Y}_{2n+1}^l the following collection of integer pairs, ordered lexicographically:

$$\mathcal{Y}_{2n+1}^l := \{(k, l) \in \mathbb{Z}^2 \mid 0 \leq k \leq 2n - 1 \text{ and } 0 \leq l \leq n - 1\}.$$

Theorem 4. The collection $\mathcal{C}^l := \{S^l \mathcal{U}^*(k) \mid (k, l) \in \mathcal{Y}_{2n+1}^l\}$ is a full exceptional collection in $\mathcal{D}^b(\text{IG}(2, 2n + 1))$.

Proof. We use the Lefschetz full exceptional collection for the Grassmannian $\text{Gr}(2, 2n + 1)$ introduced by Kuznetsov. This collection is different from the collection previously obtained by Kapranov in [Kap88] and more suitable for restriction to a hyperplane section.

Let

$$\mathcal{Y}_{2n+1} := \{(k, l) \in \mathbb{Z}^2 \mid 0 \leq k \leq 2n \text{ and } 0 \leq l \leq n - 1\}.$$

Then $\mathcal{C} := \{S^l \mathcal{U}^*(k) \mid (k, l) \in \mathcal{Y}_{2n+1}\}$ is a full exceptional collection in $\mathcal{D}^b(\text{Gr}(2, 2n + 1))$ (cf. [Kuz08, Thm. 4.1]). Since \mathcal{C} is a Lefschetz exceptional collection, it follows from [Kuz08, Prop. 2.4] that \mathcal{C}^l is an exceptional collection for $\text{IG}(2, 2n + 1)$. As in the even case, it remains to show by induction that it is full. We first introduce

$$\tilde{\mathcal{Y}}_{2n-1}^l = \{(k, l) \in \mathbb{Z}^2 \mid 0 \leq k \leq 2n - 1 \text{ and } 0 \leq l \leq n\}$$

and prove

Lemma 10. For all $(k, l) \in \tilde{\mathcal{Y}}_{2n-1}^l$, the vector bundle $S^l \mathcal{U}^*(k)$ lies in the subcategory of the derived category $\mathcal{D}^b(\text{IG}(2, 2n + 1))$ of $\text{IG}(2, 2n + 1)$ generated by the Lefschetz collection \mathcal{C}^l .

Proof. We have $\tilde{\mathcal{Y}}_{2n-1}^l \setminus \mathcal{Y}_{2n-1} = \{(0, n), \dots, (2n - 1, n)\}$. Moreover, we have the following exact sequences on $\text{Gr}(2, 2n + 1)$ for $0 \leq p \leq n$ from [Kuz08, Eq. (11)]:

$$\begin{aligned} 0 \rightarrow S^{2n-1-p} \mathcal{U}^* \rightarrow W \otimes S^{2n-2-p} \mathcal{U}^*(1) \rightarrow \dots \rightarrow \bigwedge^{2n-2-p} W \otimes \mathcal{U}^*(2n - 2 - p) \\ \rightarrow \bigwedge^{2n-1-p} W \otimes \mathcal{U}^*(2n - 1 - p) \rightarrow \bigwedge^p W^* \otimes \mathcal{O}(2n - p) \rightarrow \bigwedge^{p-1} W^* \otimes \mathcal{U}^*(2n - p) \\ \rightarrow \dots \rightarrow S^p \mathcal{U}^*(2n - p) \rightarrow 0. \end{aligned}$$

These exact sequences can be restricted to $\text{IG}(2, 2n + 1)$.

For $p = n$, we get a resolution of $S^n \mathcal{U}^*(n)$ by objects of the subcategory generated by \mathcal{C} . Tensoring by $\mathcal{O}(i)$ for $1 \leq i \leq n - 1$, we also get resolutions for $S^n \mathcal{U}^*(n + 1), \dots, S^n \mathcal{U}^*(2n - 1)$. Then for $p = n - 1$, we get a resolution of $S^n \mathcal{U}^*$, and again, after tensoring by $\mathcal{O}(i)$ for $1 \leq i \leq n - 2$, we obtain resolutions of $S^n \mathcal{U}^*(1), \dots, S^n \mathcal{U}^*(n - 2)$. To conclude the proof, we are left with finding a resolution of $S^n \mathcal{U}^*(n - 1)$ by objects of the subcategory generated by \mathcal{C} . Let $\bar{W} \supset W$ be a $(2n + 2)$ -dimensional vector space endowed with a symplectic form $\bar{\omega}$ extending ω . On $\text{Gr}_{\bar{\omega}}(2, \bar{W})$, we have the following bicomplex from [Kuz08, Prop. 5.3]:

$$\begin{array}{ccccccc} S^n \mathcal{U}^* & & & & & & \\ \downarrow & & & & & & \\ \bar{W}^* \otimes S^{n-1} \mathcal{U}^*(1) & \longrightarrow & S^n \mathcal{U}^*(1) & & & & \\ \downarrow & & \downarrow & & & & \\ \bigwedge^2 \bar{W}^* \otimes S^{n-2} \mathcal{U}^*(2) & \longrightarrow & \bar{W}^* \otimes S^{n-1} \mathcal{U}^*(2) & \longrightarrow & S^n \mathcal{U}^*(2) & & \\ \downarrow & & \downarrow & & \downarrow & & \\ \dots & & \dots & & \dots & & \\ \downarrow & & \downarrow & & \downarrow & & \\ \bigwedge^{n-1} \bar{W}^* \otimes \mathcal{U}^*(n - 1) & \longrightarrow & \bigwedge^{n-2} \bar{W}^* \otimes S^2 \mathcal{U}^*(n - 1) & \longrightarrow & \dots & \longrightarrow & S^n \mathcal{U}^*(n - 1) \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ \bigwedge^n \bar{W}^* \otimes \mathcal{O}(n) & \longrightarrow & \bigwedge^{n-1} \bar{W}^* \otimes \mathcal{U}^*(n) & \longrightarrow & \dots & \longrightarrow & \bar{W}^* \otimes S^{n-1} \mathcal{U}^*(n) \longrightarrow S^n \mathcal{U}^*(n) \end{array}$$

whose associated total complex is exact. Restricting this complex to

$$\text{Gr}_{\omega}(2, W) \subset \text{Gr}_{\bar{\omega}}(2, \bar{W}),$$

we get a complex such that every entry except $S^n \mathcal{U}^*(n - 1)$ is an object of the subcategory generated by \mathcal{C} . Hence $S^n \mathcal{U}^*(n - 1)$ also belongs to the latter category. \square

We will need a further lemma for the inductive step:

Lemma 11. *Let $V \subset W$ be a $(2n - 1)$ -dimensional vector space such that $\omega|_V$ has maximal rank. Let $X := \text{IG}(2, W)$, $X_V := \text{IG}(2, V)$ and $i_V : X_V \hookrightarrow X$ be the natural embedding. We get the following Koszul resolution for $i_{V*}\mathcal{O}_{X_V}$:*

$$\begin{aligned} 0 \rightarrow \mathcal{O}_X(-2) \rightarrow \mathcal{U}_X(-1) \oplus \mathcal{U}_X(-1) \rightarrow \mathcal{O}_X(-1)^{\oplus 3} \oplus S^2\mathcal{U}_X \rightarrow \mathcal{U}_X \oplus \mathcal{U}_X \\ \rightarrow \mathcal{O}_X \rightarrow i_{V*}\mathcal{O}_{X_V} \rightarrow 0. \end{aligned}$$

Proof. To each vector space V as above corresponds a section ϕ_V of $\mathcal{U}^* \oplus \mathcal{U}^*$ on $\text{IG}(2, W)$; moreover, the zero locus of ϕ_V is $\text{IG}(2, V) \subset \text{IG}(2, W)$. Since

$$\dim X_V = 4n - 7 = \dim X - 4 = \text{rk}(\mathcal{U}_X^* \oplus \mathcal{U}_X^*),$$

any such section ϕ_V is regular, hence the sheaf $i_*\mathcal{O}_{X_V}$ admits a Koszul resolution of the above form. \square

We may now use induction on n to prove the theorem. For $n = 1$, the result is obvious. Now assume that $n \geq 2$, that the result is proved for $n - 1$, and that the Lefschetz collection for n is not full. Then there exists an object $F \in \mathcal{D}^b(\text{IG}(2, 2n + 1))$ which is right orthogonal to all bundles in the collection, i.e. such that:

$$0 = \text{RHom}(S^l\mathcal{U}^*(k), F) = H^\bullet(X, S^l\mathcal{U}^*(-k) \otimes F)$$

for all $(k, l) \in \tilde{\mathcal{Y}}_{2n-1}^l$. Let V be such as in Lemma 11 and $i_V : X_V \hookrightarrow X$ be the embedding. We will prove $i_V^*F = 0$. Let $(k, l) \in \tilde{\mathcal{Y}}_{2n-1}^l$. Tensoring the resolution of Lemma 11 by $S^l\mathcal{U}^*(-k) \otimes F$, we get

$$\begin{aligned} 0 \rightarrow S^l\mathcal{U}(-2 - k) \otimes F \rightarrow (S^{l+1}\mathcal{U}(-1 - k) \otimes F \oplus S^{l-1}\mathcal{U}(-2 - k) \otimes F)^{\oplus 2} \\ \rightarrow (S^l\mathcal{U}(-1 - k) \otimes F)^{\oplus 4} \oplus S^{l+2}\mathcal{U}(-k) \otimes F \oplus S^{l-2}\mathcal{U}(-2 - k) \otimes F \\ \rightarrow (S^{l+1}\mathcal{U}(-k) \otimes F \oplus S^{l-1}\mathcal{U}(-1 - k) \otimes F)^{\oplus 2} \rightarrow S^l\mathcal{U}(-k) \otimes F \\ \rightarrow S^l\mathcal{U}(-k) \otimes F \otimes i_{V*}\mathcal{O}_{X_V} \rightarrow 0. \end{aligned}$$

Moreover

$$S^l\mathcal{U}(-k) \otimes F \otimes i_{V*}\mathcal{O}_{X_V} \cong i_{V*}(i_V^*(S^l\mathcal{U}(-k) \otimes F)) \cong i_{V*}(S^l\mathcal{U}(-k) \otimes i_V^*(F)).$$

If $(k, l) \in \mathcal{Y}_{2n-1}^l$, then $(k + 2, l)$, $(k + 2, l - 1)$, $(k + 1, l + 1)$, $(k + 1, l)$, $(k + 2, l - 2)$, $(k, l + 2)$, $(k, l + 1)$, $(k + 1, l - 1)$ and (k, l) are in $\tilde{\mathcal{Y}}_{2n-1}^l$. Hence the cohomology of the five first terms of the above complex vanishes, and

$$\text{RHom}_V(S^l\mathcal{U}^*(k), i_V^*F) = H^\bullet(X_V, S^l\mathcal{U}^*(-k) \otimes i_V^*F) = 0$$

for all $(k, l) \in \mathcal{Y}_{2n-1}^l$. By the induction hypothesis, we get $i_V^*F = 0$.

Lemma 12. *If, for some $F \in \mathcal{D}^b(\text{IG}(2, 2n + 1))$, we have $i_V^*F = 0$ for every $(2n - 1)$ -dimensional vector space V for which the conditions of Lemma 11 hold, then $F = 0$.*

Proof. If $F \neq 0$, let q be the maximal integer such that the cohomology sheaf $\mathcal{H}^q(F)$ is non-zero. Consider $P \in \text{Supp}(\mathcal{H}^q(F))$. As long as $n \geq 2$, there exists a $(2n - 1)$ -dimensional subspace V such that $P \subset V$ and $\omega|_V$ has maximal rank. It follows that $P \in \text{Supp}(i_V^* \mathcal{H}^q(F))$, hence $i_V^* \mathcal{H}^q(F) \neq 0$. The functor i_V^* being right exact, by choice of q we obtain that $\mathcal{H}^q(i_V^* F)$ is also non-zero, hence $i_V^* F \neq 0$. \square

This concludes the proof of the theorem. \square

It should be mentioned that it is not known whether the Dubrovin conjecture holds for the symplectic Grassmannian of lines. Indeed, although Kuznetsov has found a full exceptional collection for these varieties, Chaput and Perrin proved in [CP09, Thm. 4] that their small quantum cohomology is not semisimple. What happens for the big quantum cohomology is still unknown.

References

- [BKT08] A.S. Buch, A. Kresch, H. Tamvakis, A Giambelli formula for isotropic Grassmannians, preprint, arXiv:0811.2781 [math], 2008.
- [BKT09] A.S. Buch, A. Kresch, H. Tamvakis, Quantum Pieri rules for isotropic Grassmannians, *Invent. Math.* 178 (2) (2009) 345–405.
- [Cio05] Gianni Ciolli, On the quantum cohomology of some Fano threefolds and a conjecture of Dubrovin, *Internat. J. Math.* 16 (8) (2005) 823–839.
- [Cos09] Izzet Coskun, A Littlewood–Richardson rule for two-step flag varieties, *Invent. Math.* 176 (2) (2009) 325–395.
- [CP09] P.E. Chaput, N. Perrin, On the quantum cohomology of adjoint varieties, preprint, arXiv:0904.4824, 2009.
- [Dub98] Boris Dubrovin, Geometry and analytic theory of Frobenius manifolds, in: *Proceedings of the International Congress of Mathematicians*, Berlin, 1998, Doc. Math. Extra vol. II (1998) 315–326 (electronic).
- [FP97] W. Fulton, R. Pandharipande, Notes on stable maps and quantum cohomology, in: *Algebraic Geometry*, Santa Cruz, 1995, in: *Proc. Sympos. Pure Math.*, vol. 62, 1997, pp. 45–96.
- [Gra01] Tom Graber, Enumerative geometry of hyperelliptic plane curves, *J. Algebraic Geom.* 10 (4) (2001) 725–755.
- [Kap88] M.M. Kapranov, On the derived categories of coherent sheaves on some homogeneous spaces, *Invent. Math.* 92 (3) (1988) 479–508.
- [Kle74] S. Kleiman, The transversality of a general translate, *Compos. Math.* 28 (1974) 287–297.
- [KM94] M. Kontsevich, Y. Manin, Gromov–Witten classes, quantum cohomology, and enumerative geometry, *Comm. Math. Phys.* 164 (3) (1994) 525–562.
- [Kuz08] Alexander Kuznetsov, Exceptional collections for Grassmannians of isotropic lines, *Proc. Lond. Math. Soc.* (3) 97 (1) (2008) 155–182.
- [Mih05] I.A. Mihai, Variétés de drapeaux symplectiques impaires, PhD thesis, Institut Fourier, 2005.
- [Mih07] I.A. Mihai, Odd symplectic flag manifolds, *Transform. Groups* 12 (3) (2007) 573–599.
- [Pon07] Dalide Pontoni, Quantum cohomology of $\text{Hilb}^2(\mathbb{P}^1 \times \mathbb{P}^1)$ and enumerative applications, *Trans. Amer. Math. Soc.* 359 (11) (2007) 5419–5448.
- [PR96] Piotr Pragacz, Jan Ratajski, A Pieri-type theorem for Lagrangian and odd orthogonal Grassmannians, *J. Reine Angew. Math.* 476 (1996) 143–189.
- [ST97] Bernd Siebert, Gang Tian, On quantum cohomology rings of Fano manifolds and a formula of Vafa and Intriligator, *Asian J. Math.* 1 (4) (1997) 679–695.
- [Tam] H. Tamvakis, Quantum cohomology of homogeneous varieties: a survey, <http://www.math.umd.edu/~harryt/papers/report.07.pdf>.