

Letter to the Editor

Am. J. Hum. Genet. 78:1092, 2006

Response from Maraganore et al.

To the Editor:

In this issue, four independent research teams present new genetic association data for 13 SNPs previously reported by us to be potentially associated with Parkinson disease (PD [MIM 168600]).¹ Two groups^{2,3} report statistically significant association between one or more of these SNPs and PD, whereas two groups^{4,5} find no statistically significant association between PD and any of the SNPs investigated. In an accompanying letter,⁶ Dr. Richard H. Myers provides his qualitative assessment of the implications of these new results.

We have performed a Mantel-Haenszel analysis, using 10 of the 13 SNPs not displaying linkage disequilibrium (LD) with each other—combining the data of Li et al.,³ Farrer et al.,⁴ and Goris et al.⁵—to provide an overall quantitative assessment of the new results. The odds ratios (ORs) are reported for the SNP alleles that increase the risk of PD¹ (table 1). The X-linked SNP *rs7878232* was not included in this analysis, since subgroup-level data for males and females were not reported by all groups. The results of Clarimon et al.² were also not included, given the significant difference in SNP allele frequency observed between the European and Taiwanese control samples. This analysis reveals that none of the 10 SNPs shows statistically significant association with PD (i.e., $P < .05$). As pointed out in many of the accompanying letters, this failure to replicate may be due, in part, to differences in sample ascertainment and demographics.

A Mantel-Haenszel analysis combining these new results with those from tier 2 of Maraganore et al. reveal five SNPs with $P < .05$ and smaller effect sizes than were originally reported¹ (table 1). Although we are aware that these low P values may, at least in part, be explained by multiple testing, additional data are required to determine if these SNPs truly confer PD susceptibility or if they represent false-positive associations. Despite the small ORs, the point estimates of attributable risk for PD in the total data is still quite large for two of these SNPs (*rs10200894* population-attributable risk 0.27, 95% CI 0.04–0.77; *rs7520966* population-attributable

risk 0.21, 95% CI 0.1–0.39). If these are true associations, they may have substantial practical impact on PD.

We do not agree with Dr. Myers⁶ that our failure to identify an association between the *LRRK2* gene and PD in our original study is evidence of a false-negative result. Farrer et al. have reported elsewhere that only a very small number of the individuals with PD studied in our original whole-genome scan have a mutation in the *LRRK2* (MIM 609007) gene.⁷

We also do not consider the positive association findings between SNP *rs7702187* and PD in a Taiwanese population by Clarimon et al.² to be a replication of our original study results, since the SNP allele associated with PD susceptibility is not the same in the two studies. However, further work to follow up these results in the Taiwanese population seems warranted.

It is gratifying that our hypotheses have been tested rapidly by many groups. The Michael J. Fox Foundation, which funded our original research, also has a large-scale replication study under way. Given the low heritability estimates for PD,⁸ our initial study may have been underpowered for the detection of significant genetic associations, in part, because of the large number of genetic markers tested. Therefore, it may be prudent not to limit replication of our study to the 13 SNPs that we initially highlighted but to also consider additional SNPs and genes that had suggestive findings (as in the text files published in the online-only version of our original article).¹

DEMETRIUS M. MARAGANORE,¹
MARIZA DE ANDRADE,² TIMOTHY G. LESNICK,²
P. V. KRISHNA PANT,³ DAVID R. COX,³ AND
DENNIS G. BALLINGER³

Departments of ¹Neurology and ²Health Sciences Research, Mayo Clinic College of Medicine, Rochester, MN; and ³Perlegen Sciences, Mountain View, CA

Web Resources

The URLs for data presented herein are as follows:

dbSNP, <http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?CMD=search&DB=snp>
Online Mendelian Inheritance in Man (OMIM), <http://www.ncbi.nlm.nih.gov/Omim/> (for PD and *LRRK2*)

Table 1

Meta-Analysis of Genetic Association for 10 SNPs

dbSNP ACCESSION NUMBER	ALLELES (HIGH- RISK ALLELE)	MARAGANORE ET AL. ¹ TIERS 1 AND 2					META-ANALYSIS (REPLICATION STUDIES)					META-ANALYSIS (REPLICATION STUDIES AND MARAGANORE ET AL. ¹ TIER 2)				
		Cases	Allele Frequency (Controls)		OR (95% CI)	P	Cases	Allele Frequency (Controls)		OR (95% CI)	P	Cases	Allele Frequency (Controls)		OR (95% CI)	P
			Controls	Frequency				Controls	Frequency				Controls	Frequency		
rs10200894	C/G (C)	772	.88	1.84 (1.38–2.45)	1.70 × 10 ⁻⁵	1,566	1,546	.89	1.14 (.96–1.35)	.125	1,926	1,955	.89	1.25 (1.07–1.45)	.004	
rs11737074	G/A (A)	764	.19	1.50 (1.21–1.86)	1.55 × 10 ⁻⁴	1,563	1,542	.21	1.02 (.9–1.15)	.770	1,925	1,952	.21	1.09 (.97–1.21)	.142	
rs16851009	C/T (T)	741	.08	1.84 (1.36–2.49)	4.17 × 10 ⁻⁵	1,539	1,544	.1	.98 (.83–1.16)	.853	1,899	1,953	.1	1.08 (.93–1.26)	.312	
rs17329669	A/G (G)	768	.12	1.71 (1.33–2.21)	2.30 × 10 ⁻⁵	1,554	1,525	.12	1.13 (.97–1.32)	.102	1,914	1,933	.12	1.22 (1.06–1.39)	.004	
rs2245218	A/G (G)	770	.12	1.67 (1.29–2.14)	4.61 × 10 ⁻⁵	1,571	1,563	.16	.94 (.82–1.08)	.369	1,933	1,971	.16	1.02 (.9–1.16)	.752	
rs2313982	C/T (T)	740	.07	2.01 (1.44–2.79)	1.79 × 10 ⁻⁵	1,562	1,554	.09	.88 (.73–1.04)	.138	1,924	1,964	.09	1.01 (.86–1.18)	.935	
rs7320966	C/T (C)	769	.7	.67 (.55–.81)	2.96 × 10 ⁻⁵	1,563	1,550	.72	1.07 (.96–1.2)	.242	1,923	1,956	.72	1.15 (1.04–1.27)	.007	
rs7702187	T/A (T)	761	.81	1.74 (1.36–2.24)	7.62 × 10 ⁻⁶	1,541	1,541	.83	1.07 (.93–1.22)	.334	1,900	1,950	.82	1.14 (1.01–1.29)	.030	
rs7723605	T/C (C)	773	.11	1.78 (1.35–2.35)	3.30 × 10 ⁻⁵	1,567	1,571	.13	1.03 (.89–1.19)	.684	1,927	1,981	.13	1.12 (.98–1.28)	.105	
ss46548856	G/C (G)	765	.9	1.88 (1.38–2.57)	3.65 × 10 ⁻⁵	1,551	1,528	.9	1.12 (.94–1.33)	.196	1,913	1,933	.9	1.21 (1.03–1.42)	.016	

- Maraganore DM, de Andrade M, Lesnick TG, Strain KJ, Farrer MJ, Rocca WA, Pant PVK, Frazer KA, Cox DR, Ballinger DG (2005) High-resolution whole-genome association study of Parkinson disease. *Am J Hum Genet* 77:685–693
- Clarimon J, Scholz S, Fung H-C, Hardy J, Eerola J, Hellström O, Chen C-M, Wu Y-R, Tienari PJ, Singleton A (2006) Conflicting results regarding the semaphorin gene (*SEMA5A*) and the risk for Parkinson disease. *Am J Hum Genet* 78:1082–1084 (in this issue)
- Li Y, Rowland C, Schrodi S, Laird W, Tacey K, Ross D, Leong D, Catanese J, Sninsky J, Grupe A (2006) A case-control association study of the 12 single-nucleotide polymorphisms implicated in Parkinson disease by a recent genome scan. *Am J Hum Genet* 78:1090–1092 (in this issue)
- Farrer MJ, Haugarvoll K, Ross OA, Stone JT, Milkovic NM, Cobb SA, Whittle AJ, Lincoln SJ, Hulihan MM, Heckman MG, White LR, Aasly JO, Gibson JM, Gosal D, Lynch T, Wszolek ZK, Uitti RJ, Toft M (2006) Genomewide association, Parkinson disease, and *PARK10*. *Am J Hum Genet* 78:1084–1088 (in this issue)
- Goris A, Williams-Gray CH, Foltynie T, Compston DAS, Barker RA, Sawcer SJ (2006) No evidence for association with Parkinson disease for 13 single-nucleotide polymorphisms identified by whole-genome association screening. *Am J Hum Genet* 78:1088–1090 (in this issue)
- Myers RH (2006) Considerations for genomewide association studies in Parkinson disease. *Am J Hum Genet* 78:1081–1082 (in this issue)
- Farrer M, Stone J, Mata IF, Lincoln S, Kachergus J, Hulihan M, Strain KJ, Maraganore MD (2005) LRRK2 mutations in Parkinson disease. *Neurology* 65:738–740
- Rocca WA, McDonnell SK, Strain KJ, Bower JH, Ahlskog JE, Elbaz A, Schaid DJ, Maraganore DM (2004) Familial aggregation of Parkinson's disease: the Mayo Clinic Family Study. *Ann Neurol* 56:495–502

Address for correspondence and reprints: Dr. Demetrius Maraganore, Department of Neurology, Mayo Clinic College of Medicine, 200 First Street SW, Rochester, MN 55905. E-mail: dmaraganore@mayo.edu

© 2006 by The American Society of Human Genetics. All rights reserved.
0002-9297/2006/7806-0024\$15.00

References

- Maraganore DM, de Andrade M, Lesnick TG, Strain KJ, Farrer MJ, Rocca WA, Pant PVK, Frazer KA, Cox DR, Ballinger DG (2005) High-resolution whole-genome association study of Parkinson disease. *Am J Hum Genet* 77:685–693

