



ELSEVIER

Contents lists available at ScienceDirect

# Journal of Combinatorial Theory, Series B

[www.elsevier.com/locate/jctb](http://www.elsevier.com/locate/jctb)


## Excluding a group-labelled graph

 Jim Geelen<sup>a,1</sup>, Bert Gerards<sup>b,c</sup>
<sup>a</sup> Department of Combinatorics and Optimization, University of Waterloo, Waterloo, Canada N2L 3G1

<sup>b</sup> CWI, Postbus 94079, 1090 GB Amsterdam, The Netherlands

<sup>c</sup> Department of Mathematics and Computer Science, Eindhoven University of Technology, Postbus 513, 5600 MB Eindhoven, The Netherlands

### ARTICLE INFO

#### Article history:

Received 7 June 2006

Available online 9 August 2008

#### Keywords:

Group-labelled graphs

Gain graphs

Signed graphs

Graph minors

### ABSTRACT

This paper contains a first step towards extending the Graph Minors Project of Robertson and Seymour to group-labelled graphs. For a finite abelian group  $\Gamma$  and  $\Gamma$ -labelled graph  $G$ , we describe the class of  $\Gamma$ -labelled graphs that do not contain a minor isomorphic to  $G$ .

© 2008 Elsevier Inc. All rights reserved.

## 1. Introduction

Group-labelled graphs are a generalization of signed graphs. For a group  $\Gamma$ , a  $\Gamma$ -labelled graph is an oriented graph with edges labelled by elements of  $\Gamma$ . Here we are primarily interested in abelian groups, so we will use additive notation. A *minor* of a group-labelled graph  $G$  is any group-labelled graph obtained from  $G$  by any sequence of the following operations: vertex deletion, edge deletion, contracting zero-labelled edges, and *shifting at a vertex*, which, for a given vertex  $v$  and group element  $\gamma$ , amounts to adding  $\gamma$  to the label of each edge entering  $v$  and subtracting  $\gamma$  from the label of each edge leaving  $v$ .

We hope that the main results of the Graph Minors Project of Robertson and Seymour will extend to group-labelled graphs over any fixed finite abelian group. In particular, the following two conjectures, if true, would generalize the two main results of the Graph Minors Project; see [5] and [7].

**Conjecture 1.1.** *For any finite abelian group  $\Gamma$  and any infinite sequence  $G_1, G_2, \dots$  of  $\Gamma$ -labelled graphs, there exist integers  $i < j$  such that  $G_i$  is isomorphic to a minor of  $G_j$ .*

<sup>1</sup> This research was partially supported by the Natural Sciences and Engineering Research Council of Canada.

**Conjecture 1.2.** *For any finite abelian group  $\Gamma$  and any  $\Gamma$ -labelled graph  $H$ , there is a polynomial-time algorithm to determine whether or not a  $\Gamma$ -labelled graph  $G$  contains an  $H$ -minor.*

To avoid algorithmic complications in Conjecture 1.2, we assume that the group  $\Gamma$  is given by its addition table. In the above conjectures the finiteness of the group  $\Gamma$  is certainly necessary, however, there may be extensions to non-abelian groups.

Group-labelled graphs (also known as *gain graphs*) are closely related with several interesting classes of matroids; see Zaslavsky [8,9]. Proving Conjecture 1.1 would prove that two interesting classes of matroids are well-quasi-ordered with respect to taking minors and would go a long way towards well-quasi-ordering binary matroids.

The workhorse of the Graph Minors Project is the Graph Minors Structure Theorem [6], and the main result of this paper is an extension of that result to group-labelled graphs over finite abelian groups. The *complete*  $\Gamma$ -labelled graph on  $n$  vertices and  $2|\Gamma|\binom{n}{2}$  edges is denoted  $K(\Gamma, n)$  (a precise definition is given at the end of Section 2). For a subgroup  $\Gamma'$  of an abelian group  $\Gamma$ , we say that  $G$  is  $\Gamma'$ -balanced if it is shifting-equivalent to a  $\Gamma'$ -labelled graph. The following theorem is the main result of this paper.

**Theorem 1.3.** *Let  $\Gamma$  be a finite abelian group, let  $\Gamma'$  be a subgroup of  $\Gamma$ , let  $n \in \mathbb{N}$ , and let  $t = 8n|\Gamma|^2$ . If  $G$  is a  $\Gamma$ -labelled graph and  $H$  is a minor of  $G$  isomorphic to  $K(\Gamma', 4t)$ , then either*

- *there is a set  $X \subseteq V(G)$  with  $|X| < t$  such that the unique block of  $G - X$  that contains most of  $E(H)$  is  $\Gamma'$ -balanced, or*
- *there is a subgroup  $\Gamma''$  of  $\Gamma$  properly containing  $\Gamma'$  and a minor  $H'$  of  $G$  with  $E(H') \subseteq E(H)$  such that  $H'$  is isomorphic to  $K(\Gamma'', n)$ .*

The specialization of Theorem 1.3 to signed graphs is implicit in [3] and the proof of Theorem 1.3 is a routine extension of the proof given in that paper. The proof of Theorem 1.3 is constructive; considering  $n$  as a constant, in  $O(|V(G)|^5)$  time we can find either the set  $X$  or the minor  $H'$ .

We also prove the following easy result that is complementary to Theorem 1.3. The graph that is obtained from a group-labelled graph  $G$  by ignoring the orientation and the group-labels is denoted  $\tilde{G}$ .

**Theorem 1.4.** *Let  $\Gamma$  be a finite group and let  $n \in \mathbb{N}$ . Then there exists  $l \in \mathbb{N}$  such that if  $G$  is a  $\Gamma$ -labelled graph such that  $\tilde{G}$  has a  $K_l$ -minor, then  $G$  has a  $K(\{0\}, n)$ -minor.*

Theorems 1.3 and 1.4 are particularly useful when applied in conjunction with “tangles.” For the rest of the introduction we assume that the reader is familiar with the definitions in Graph minors, X [4].

Suppose that  $\mathcal{T}$  is a tangle of order  $k$  in a graph (or group-labelled graph)  $G$ . If  $X \subseteq V(G)$  with  $|X| \leq k - 2$ , then it is straightforward to show that there is a unique block  $B$  of  $G - X$  such that  $V(B) \cup X$  is not contained in the  $\mathcal{T}$ -small side of any separation of order  $< k$ ; we call  $B$  the  $\mathcal{T}$ -large block of  $G - X$ .

Let  $\Gamma$  be a finite abelian group and let  $n \in \mathbb{N}$ . Theorems 1.3 and 1.4 imply that there exist  $l, t \in \mathbb{N}$  such that if  $G$  is a  $\Gamma$ -labelled graph and  $\mathcal{T}$  is a tangle in  $G$  of order  $\geq t + 2$ , then either:

- (1)  $\mathcal{T}$  does not control a  $K_l$ -minor in  $\tilde{G}$ ,
- (2) there exists  $X \subseteq V(G)$  with  $|X| \leq t$  such that the  $\mathcal{T}$ -large block of  $G - X$  is  $\Gamma'$ -balanced for some proper subgroup  $\Gamma'$  of  $\Gamma$ , or
- (3)  $\mathcal{T}$  controls a  $K(\Gamma, n)$ -minor in  $G$ .

In the first case, we can use the Graph Minors Structure Theorem to describe the structure of  $\tilde{G}$ .

## 2. Group-labelled graphs

Let  $\Gamma$  be an abelian group. A  $\Gamma$ -labelled graph is an oriented graph with edges labelled by elements of  $\Gamma$ . More formally, if  $G$  is a  $\Gamma$ -labelled graph, then  $G$  has a vertex set  $V(G)$  and an edge set  $E(G)$ , and each edge  $e \in E(G)$  is assigned a head, denoted  $\text{head}_G(e)$ , in  $V(G)$ , a tail, denoted  $\text{tail}_G(e)$ , in  $V(G)$ , and a group-label, denoted  $\gamma_G(e)$ , in  $\Gamma$ . The head and tail of an edge are referred to as its ends.

Let  $G$  be a  $\Gamma$ -labelled graph. The graph obtained from  $G$  by ignoring the orientation and the group labels is denoted by  $\tilde{G}$ . By a walk in  $G$ , we mean a walk in  $\tilde{G}$ . If  $e \in E(G)$  and  $v$  is an end of  $e$ , then we let  $\gamma_G(e, v) = \gamma_G(e)$  if  $v = \text{head}_G(e)$  and  $\gamma_G(e, v) = -\gamma_G(e)$  if  $v = \text{tail}_G(e)$ . Let  $W = (v_0, e_1, v_1, e_2, v_2, \dots, e_k, v_k)$  be a walk of  $G$ . We let  $\gamma_G(W) = \gamma_G(e_1, v_1) + \dots + \gamma_G(e_k, v_k)$ . The length of  $W$  is  $k$ ; the ends of  $W$  are  $v_0$  and  $v_k$ ;  $W$  is closed if  $v_0 = v_k$ ;  $W$  is a path if  $v_0, v_1, \dots, v_k$  are distinct vertices; and  $W$  is a circuit if  $v_0, \dots, v_{k-1}$  are distinct,  $e_1, \dots, e_k$  are distinct, and  $v_0 = v_k$ . We let  $E(W) = \{e_1, \dots, e_k\}$  and  $V(W) = \{v_0, \dots, v_k\}$ .

### 2.1. Shifting

Let  $v \in V(G)$  and let  $\delta \in \Gamma$ . We obtain a new  $\Gamma$ -labelled graph  $G'$  from  $G$  by adding  $\delta$  to the label of each edge with head  $v$  and subtracting  $\delta$  from the label of each edge with tail  $v$ . We say that  $G'$  is obtained from  $G$  by shifting at  $v$ . Any  $\Gamma$ -labelled graph that is obtained from  $G$  by a sequence of shifting operations is said to be shifting-equivalent to  $G$ . Note that, if  $G'$  is shifting-equivalent to  $G$  and  $W$  is a closed walk of  $G$ , then  $\gamma_{G'}(W) = \gamma_G(W)$ .

We omit the elementary proof of the following result.

**Lemma 2.1.** *If  $G$  is a  $\Gamma$ -labelled graph, for some abelian group  $\Gamma$ , and  $T$  is a spanning tree of  $\tilde{G}$ , then  $G$  is switching-equivalent to some  $\Gamma$ -labelled graph  $G'$  with  $\gamma_{G'}(e) = 0$  for each  $e \in E(T)$ .*

### 2.2. Balanced labellings

Let  $\Gamma'$  be a subgroup of  $\Gamma$ . We say that  $G$  is  $\Gamma'$ -balanced if  $\gamma_G(C) \in \Gamma'$  for each circuit  $C$  of  $G$ . Note that, if  $\Gamma$  is abelian and  $C_1$  and  $C_2$  are circuits of  $G$  with  $E(C_1) = E(C_2)$ , then  $\gamma_G(C_1) = \pm \gamma_G(C_2)$ .

**Lemma 2.2.** *Let  $\Gamma$  be an abelian group and let  $G$  be a  $\Gamma$ -labelled graph. If  $G$  is  $\Gamma'$ -balanced for some subgroup  $\Gamma'$  of  $\Gamma$ , then  $G$  is switching-equivalent to a  $\Gamma'$ -labelled graph.*

**Proof.** By treating each component separately, we may assume that  $\tilde{G}$  is connected; let  $T$  be a spanning tree of  $\tilde{G}$ . By Lemma 2.1,  $G$  is switching-equivalent to some  $\Gamma$ -labelled graph  $G'$  with  $\gamma_{G'}(e) = 0$  for each  $e \in E(T)$ . Since  $\Gamma$  is abelian,  $G'$  is  $\Gamma'$ -balanced. Consider  $e \in E(G') - E(T)$  and a circuit  $C$  with  $E(C) \subseteq E(T) \cup \{e\}$ . We have  $\gamma_{G'}(e) = \pm \gamma_{G'}(C) \in \Gamma'$ . Hence  $G'$  is  $\Gamma'$ -labelled.  $\square$

**Lemma 2.3.** *Let  $\Gamma$  be an abelian group and let  $G$  be a  $\Gamma$ -labelled graph. If  $G$  is  $\Gamma'$ -balanced for some subgroup  $\Gamma'$  of  $\Gamma$ , then for any closed walk  $W$  of  $G$  we have  $\gamma_G(W) \in \Gamma'$ .*

**Proof.** By Lemma 2.2,  $G$  is switching-equivalent to a  $\Gamma'$ -labelled graph  $G'$ . Now, for any closed walk  $W$ , we have  $\gamma_G(W) = \gamma_{G'}(W) \in \Gamma'$ .  $\square$

We call  $G$  balanced if it is  $\{0\}$ -balanced. A set  $F \subseteq E(G)$  is balanced if the subgraph of  $G$  with edge-set  $F$  is balanced; that is,  $\gamma_G(C) = 0$  for each circuit  $C$  of  $G$  with  $E(C) \subseteq F$ .

### 2.3. Minors

A  $\Gamma$ -labelled graph  $H$  is a minor of  $G$  if it can be obtained from  $G$  via any sequence of the following operations: edge deletion, contraction of a zero-labelled edge, shifting at a vertex, and vertex

deletion. It is straightforward to see that if  $F$  is the set of edges that are contracted in obtaining a minor  $H$  of  $G$ , then  $F$  is balanced in  $G$ . Conversely, if  $F' \subseteq E(G)$  is balanced, then, by Lemma 2.2, we can shift so that each edge in  $F'$  is zero-labelled and then contract these edges. For any set  $A$  of zero-labelled edges of  $G$  we let  $G/A$  denote the minor of  $G$  obtained by contracting  $A$ . For a set  $S \subseteq V(G)$ , we let  $G[S]$  denote the  $\Gamma$ -labelled subgraph of  $G$  induced by  $S$  and let  $G - S = G[V(G) - S]$ .

The following result, whose easy proof we omit, provides a more tangible way to exhibit a minor in a signed graph.

**Lemma 2.4.** *Let  $G$  be a  $\Gamma$ -labelled graph, for some abelian group  $\Gamma$ , and let  $H$  be a minor of  $G$ . Then there is a  $\Gamma$ -labelled graph  $G'$  that is switching-equivalent to  $G$  and there exist vertex-disjoint trees  $(T(v) : v \in V(H))$  in  $G$  such that:*

- $\gamma_{G'}(e) = 0$  for each  $v \in V(H)$  and  $e \in E(T(v))$ ,
- $\gamma_{G'}(e) = \gamma_H(e)$  for each  $e \in E(H)$ , and
- $\text{head}_{G'}(e) \in V(T(\text{head}_H(e)))$  and  $\text{tail}_{G'}(e) \in V(T(\text{tail}_H(e)))$  for each  $e \in E(H)$ .

### 2.4. A-paths

Let  $A \subseteq V(G)$ . An  $A$ -path in  $G$  is a path of length at least one whose ends are both in  $A$ . The following result was proved by Chudnovsky et al. [2].

**Theorem 2.5.** *Let  $\Gamma$  be an abelian group, let  $G$  be a  $\Gamma$ -labelled graph, and let  $A \subseteq V(G)$ . Then for any  $k \in \mathbb{N}$  either*

- there exist vertex-disjoint  $A$ -paths  $P_1, \dots, P_k$  with  $\gamma_G(P_i) \neq 0$  for each  $i \in \{1, \dots, k\}$ , or
- there exists  $X \subseteq V(G)$  with  $|X| \leq 2(k - 1)$  such that  $\gamma_G(P) = 0$  for each  $A$ -path  $P$  in  $G$  disjoint from  $X$ .

For constant  $k$ , in  $O(|V(G)|^5)$  time we can find either the paths  $P_1, \dots, P_k$  or the set  $X$  guaranteed by Theorem 2.5; see Chudnovsky et al. [1].

We require the following elementary corollary.

**Corollary 2.6.** *Let  $\Gamma'$  be a subgroup of an abelian group  $\Gamma$ , let  $G$  be a  $\Gamma$ -labelled graph, and let  $A \subseteq V(G)$ . Then for any  $k \in \mathbb{N}$  either*

- there exist vertex-disjoint  $A$ -paths  $P_1, \dots, P_k$  with  $\gamma_G(P_i) \notin \Gamma'$  for each  $i \in \{1, \dots, k\}$ , or
- there exists  $X \subseteq V(G)$  with  $|X| \leq 2(k - 1)$  such that  $\gamma_G(P) \in \Gamma'$  for each  $A$ -path  $P$  in  $G$  disjoint from  $X$ .

**Proof.** Apply Theorem 2.5 to the quotient group  $\Gamma/\Gamma'$ .  $\square$

### 2.5. Blocks

A separation of  $G$  is a pair  $(G_1, G_2)$  of subgraphs of  $G$  such that  $E(G) = E(G_1) \cup E(G_2)$  and  $V(G) = V(G_1) \cup V(G_2)$ ; the order of the separation is  $|V(G_1) \cap V(G_2)|$ . We say that  $G$  is 2-connected if  $G$  is connected and for each separation  $(G_1, G_2)$  of  $G$  of order 1 either  $E(G_1) = \emptyset$  or  $E(G_2) = \emptyset$ . Note that if  $G$  is 2-connected and  $|V(G)| \geq 2$ , then  $G$  has no loops. A block of  $G$  is a maximal 2-connected subgraph of  $G$ .

**Lemma 2.7.** *Let  $G$  be a 2-connected  $\Gamma$ -labelled graph, for some abelian group  $\Gamma$ , and let  $\Gamma'$  be a subgroup of  $\Gamma$ . If  $u$  and  $v$  are distinct vertices of  $G$  such that  $\gamma_G(P) \in \Gamma'$  for each  $(u, v)$ -path  $P$  in  $G$ , then  $G$  is  $\Gamma'$ -balanced.*

**Proof.** Let  $C$  be a circuit of  $G$ . Since  $G$  is 2-connected, there exist two vertex-disjoint paths  $P$  and  $P'$  from  $\{u, v\}$  to  $V(C)$ . We may assume that  $P$  connects  $u$  to  $a \in V(C)$  and  $P'$  connects  $v$  to  $b \in V(C)$ . Furthermore, we may assume that  $P$  and  $P'$  each only meet  $C$  in one vertex. Let  $C'$  be a circuit of  $G$

starting at  $a$  with  $E(C') = E(C)$ . Let  $Q$  be the  $(u, v)$ -path obtained by following  $P$  from  $u$  to  $a$  then following  $C'$  to  $b$ , and then following  $P'$  backward to  $v$ . Now let  $Q'$  be the  $(v, u)$ -path obtained by following  $P'$  from  $v$  to  $b$  then following  $C'$  to  $a$ , and then following  $P$  backward to  $u$ . Note that

$$\gamma_G(C) = \pm \gamma_G(C') = \pm(\gamma_G(Q) + \gamma_G(Q')) \in \Gamma'.$$

Hence  $G$  is  $\Gamma'$ -balanced.  $\square$

### 2.6. Complete graphs

The complete  $\Gamma$ -labelled graph on  $n$  vertices, denoted  $K(\Gamma, n)$ , has vertex set  $\{1, \dots, n\}$  and edge set  $\{e(i, j, \sigma) : i, j \in V(K(\Gamma, n)), i \neq j, \sigma \in \Gamma\}$  where each edge  $e(i, j, \sigma)$  has tail  $i$ , head  $j$ , and label  $\sigma$ .

We can now prove Theorem 1.4, which we restate here for convenience.

**Theorem 2.8.** *Let  $\Gamma$  be a finite abelian group and let  $n \in \mathbb{N}$ . Then there exists  $l \in \mathbb{N}$  such that if  $G$  is a  $\Gamma$ -labelled graph such that  $\tilde{G}$  has a  $K_l$ -minor, then  $G$  has a  $K(\{0\}, n)$ -minor.*

**Proof.** By Ramsey's Theorem there exists an  $l$  such that, if we colour the edges of a clique on  $l$  vertices with  $2|\Gamma|$  colours, then there is a monochromatic subclique on  $2n$  vertices. We may assume that  $\tilde{G}$  is isomorphic to  $K_l$  and that  $V(G) = \{1, \dots, l\}$ . We partition  $E(G)$  into  $2|\Gamma|$  sets according to the label  $\gamma_G(e)$  of the edge  $e$  and to the sign of  $\text{head}_G(e) - \text{tail}_G(e)$ . By our choice of  $l$ , there exists a set  $X \subseteq \{1, \dots, l\}$  with  $|X| = 2n$ , an element  $\gamma \in \Gamma$ , and a sign  $\sigma \in \{-1, 1\}$  such that for each edge  $e$  of  $G[X]$  we have  $\gamma_G(e) = \gamma$  and  $\text{head}_G(e) - \text{tail}_G(e) = \sigma$ . By symmetry, we may assume that  $X = \{1, \dots, 2n\}$  and that  $\sigma = 1$ . Let  $H$  be the subgraph of  $G[X]$  with all edges of  $G$  having tail in  $\{1, \dots, n\}$  and head in  $\{n + 1, \dots, 2n\}$ . Now we obtain a  $K(\{0\}, n)$ -minor of  $H$  by shifting at each of  $\{n + 1, \dots, 2n\}$  so that all labels in  $H$  become zero, and then contracting a perfect matching.  $\square$

### 3. The main theorem

We need one more preliminary result.

**Lemma 3.1.** *Let  $\Gamma$  be a finite abelian group, let  $\Gamma'$  be a subgroup of  $\Gamma$ , let  $n \in \mathbb{N}$ , and let  $l = n|\Gamma|^2$ . If  $G$  is a  $\Gamma$ -labelled graph and  $M$  is a matching of size  $l$  in  $\tilde{G}$  such that  $G - M$  is isomorphic to  $K(\Gamma', 2l)$  and for each  $e \in M$  we have  $\gamma_G(e) \notin \Gamma'$ , then there is a subgroup  $\Gamma''$  of  $\Gamma$  properly containing  $\Gamma'$  and a minor  $H'$  of  $G$  with  $E(H') \subseteq E(G - M)$  such that  $H'$  is isomorphic to  $K(\Gamma'', n)$ .*

**Proof.** There exists  $M' \subseteq M$  with  $|M'| = n|\Gamma|$  and an element  $\sigma \in \Gamma - \Gamma'$  such that  $\gamma_G(e) = \sigma$  for each  $e \in M'$ . Let  $\Gamma''$  be the subgroup of  $\Gamma$  generated by  $\Gamma'$  and  $\sigma$  and let  $g$  be the order of  $\sigma$ .

By contracting some zero-labelled edges, we can find a minor with  $n$  vertex-disjoint directed paths of length  $g$  such that each edge in these paths has label  $\sigma$ . Then, by deleting edges, we obtain a minor  $G'$  of  $G$  with  $V(G') = \{v_k^i : 1 \leq i \leq n, 0 \leq k \leq g\}$  and with the following edges:

- For each  $i \in \{1, \dots, n\}$  and  $k \in \{1, \dots, g\}$ , we have an edge  $e_k^i \in M'$  with head  $v_k^i$ , tail  $v_{k-1}^i$ , and label  $\sigma$ .
- For each  $i, j \in \{1, \dots, n\}$  with  $i \neq j$ ,  $k \in \{1, \dots, g\}$ , and  $\gamma' \in \Gamma'$ , we have an edge  $e \in E(G) - M$  with tail  $v_0^j$ , head  $v_k^i$ , and label  $\gamma'$ .

Now we construct a minor  $H'$  of  $G'$  by shifting each vertex  $v_k^i$  by  $-k\sigma$  (so that each  $e_k^i$  is zero-labelled) and then contracting the edges  $\{e_k^i : 1 \leq i \leq n, 1 \leq k \leq g\}$ . It is straightforward to verify the  $H'$  is isomorphic to  $K(\Gamma'', n)$  and that  $E(H') \subseteq E(H)$ .  $\square$

We are now ready to prove the main result which we restate here for convenience.

**Theorem 3.2.** *Let  $\Gamma$  be a finite abelian group, let  $\Gamma'$  be a subgroup of  $\Gamma$ , let  $n \in \mathbb{N}$ , and let  $t = 8n|\Gamma|^2$ . Then if  $G$  is a  $\Gamma$ -labelled graph and  $H$  is a minor of  $G$  isomorphic to  $K(\Gamma', 4t)$ , then either*

- *there is a set  $X \subseteq V(G)$  with  $|X| < t$  such that the unique block of  $G - X$  that contains most of  $E(H)$  is  $\Gamma'$ -balanced, or*
- *there is a subgroup  $\Gamma''$  of  $\Gamma$  properly containing  $\Gamma'$  and a minor  $H'$  of  $G$  with  $E(H') \subseteq E(H)$  such that  $H'$  is isomorphic to  $K(\Gamma'', n)$ .*

**Proof.** Let  $l = n|\Gamma|^2$  and  $m = 4t$ .

We assume that:

**3.2.1.** *There is no set  $X \subseteq V(G)$  with  $|X| < t$  such that the block of  $G - X$  that contains most of  $E(H)$  is  $\Gamma'$ -balanced.*

By possibly shifting we may assume that there exist vertex-disjoint trees  $(T(v) : v \in V(H))$  in  $G$  such that:

- $\gamma_G(e) = 0$  for each  $v \in V(H)$  and  $e \in E(T(v))$ ,
- $\gamma_G(e) = \gamma_H(e)$  for each  $e \in E(H)$ , and
- $\text{head}_G(e) \in V(T(\text{head}_H(e)))$  and  $\text{tail}_G(e) \in V(T(\text{tail}_H(e)))$  for each  $e \in E(H)$ .

Consider any  $v \in V(H)$ . For each  $u \in V(H) - \{v\}$  we choose an edge  $e \in E(H)$  with  $u = \text{tail}_H(e)$  and  $v = \text{head}_H(e)$  and we let  $f_v(u) = \text{head}_G(e)$ ; thus  $f_v(u) \in V(T(v))$ . For each  $X \subseteq V(T(v))$  we let  $f_v^{-1}(X) = \{u \in V(H) - \{v\} : f_v(u) \in X\}$ .

We leave it to the reader to verify that we can choose a vertex  $a_v \in V(T(v))$  satisfying:

**3.2.2.** *For each edge  $e \in E(T(v))$  we have  $|f_v^{-1}(X)| \leq \frac{m-1}{2}$  where  $X$  is the vertex set of the component of  $T(v) - e$  that does not contain  $a_v$ .*

Now we let  $A = \{a_v : v \in V(H)\}$ . For notational convenience, for  $X \subseteq V(H)$ , we let  $T(X)$  denote the subgraph of  $G$  with components  $(T(v) : v \in X)$ . Our choice of  $A$  gives rise to the following result.

**3.2.3.** *Let  $S \subseteq V(H)$  with  $|S| > \frac{m+1}{2}$ , let  $L \subseteq S$ , let  $G'$  is the  $\Gamma$ -labelled subgraph of  $G$  induced by  $V(T(S))$ , and let  $B$  be the block of  $G'/E(T(L))$  that contains  $E(H[S])$ . If  $v \in S - L$ , then  $a_v \in V(B)$ .*

**Subproof.** Note that, since the edges of  $H[S]$  form a clique in  $G/E(T(V(H)))$ , there is a block of  $G'/E(T(L))$  that contains  $E(H[S])$ . Let  $G'' = G'/E(T(L))$ . If  $a_v \notin V(B)$ , then there is a separation  $(G_1, G_2)$  of  $G''$  of order 1 with  $E(B) \subseteq E(G_2)$  and  $a_v \in V(G_1) - V(G_2)$ . Let  $w$  be the vertex in  $V(G_1) \cap V(G_2)$ . Since  $E(H[S]) \subseteq E(B) \subseteq E(G_2)$  and since some edge in  $E(H[S])$  has an end in  $V(T(v))$ , we have  $w \in V(T(v))$ . Let  $e$  be the edge on the  $(a_v, w)$ -path in  $T(v)$  that is incident with  $w$  and let  $X$  be the vertex set of the component of  $T(v) - e$  that contains  $w$ . Note that, for each  $u \in S - \{v\}$ , we have  $f_v(u) \in X$ . Thus  $|f_v^{-1}(X)| \geq |S| - 1 > \frac{m-1}{2}$ , contradicting our choice of  $a_v$ .  $\square$

**3.2.4.** *There exist vertex-disjoint  $A$ -paths  $P_1, \dots, P_{4l}$  such that  $\gamma_G(P_i) \notin \Gamma'$  for each  $i \in \{1, \dots, 4l\}$ .*

**Subproof.** Suppose otherwise; then, by Corollary 2.6, there is a set  $X \subseteq V(G)$  with  $|X| < 8l$  such that  $\gamma_G(P) \in \Gamma'$  for each  $A$ -path  $P$  in  $G$  that is disjoint from  $X$ . Let  $S = \{v \in V(H) : V(T(v)) \cap X = \emptyset\}$  and let  $H' = H[S]$ . Now let  $B$  be the block of  $G - X$  that contains  $E(H')$  and let  $x$  and  $y$  be distinct vertices in  $\{a_v : v \in S\}$ . Note that  $|S| \geq \frac{m+1}{2}$ . By applying 3.2.3 with  $L = \emptyset$ , we have  $x, y \in V(B)$ , and, by our choice of  $X$ ,  $\gamma_B(P) \in \Gamma'$  for each  $(x, y)$ -path  $P$  in  $B$ . Then, by Lemma 2.7,  $B$  is  $\Gamma'$ -balanced. However  $|S| > \frac{3}{4}|V(H)|$  so  $|E(H')| > \frac{1}{2}|E(H)|$  and, hence,  $B$  is the unique block of  $G - X$  that contains most of  $E(H)$ , which contradicts 3.2.1.  $\square$

Let  $F = E(T(V(H)))$ . We choose vertex-disjoint  $A$ -paths  $P_1, \dots, P_{4l}$

- minimizing  $|E(P_1) - F| + \dots + |E(P_{4l}) - F|$ ,
- subject to  $\gamma_G(P_i) \notin \Gamma'$  for each  $i \in \{1, \dots, 4l\}$ .

Let  $A_0 \subseteq A$  be the set of ends of the paths  $P_1, \dots, P_{4l}$ , let  $W = \{v \in V(H) : A_0 \cap V(T(v)) \neq \emptyset\}$ , and let  $A_1 = V(H) - W$ .

**3.2.5.** For each  $v \in A_1$  and  $i \in \{1, \dots, 4l\}$ , we have  $V(T(v)) \cap V(P_i) = \emptyset$ .

**Subproof.** Suppose that  $T(v)$  meets one or more of  $P_1, \dots, P_{4l}$ . Then, for some path  $P_i$  and  $w \in V(P_i) \cap V(T(v))$ , the  $(a_v, w)$ -path  $Q$  in  $T(v)$  is internally disjoint from each of  $P_1, \dots, P_{4l}$ . Suppose that  $P_i$  is an  $(a_x, a_y)$ -path. Let  $Q_x$  denote the  $(a_v, a_x)$ -path with  $E(Q_x) \subseteq E(Q) \cup E(P_i)$  and let  $Q_y$  denote the  $(a_v, a_y)$ -path with  $E(Q_y) \subseteq E(Q) \cup E(P_i)$ . Note that  $\gamma_G(P_i) - \gamma_G(Q_y) + \gamma_G(Q_x) = 0$ . Then, since  $\gamma_G(P_i) \notin \Gamma'$ , either  $\gamma_G(Q_x) \notin \Gamma'$  or  $\gamma_G(Q_y) \notin \Gamma'$ . Moreover,  $|E(Q_x) - F| < |E(P) - F|$  and  $|E(Q_y) - F| < |E(P) - F|$ ; this contradicts our choice of  $P_1, \dots, P_{4l}$ .  $\square$

Let  $G_1 = G/E(T(A_1))$ . We may assume that the vertices of  $G_1$  are labelled so that  $H[A_1]$  is a subgraph of  $G_1$ ; thus  $G_1[A_1]$  is isomorphic to  $K(\Gamma', 16l)$ .

**3.2.6.** If  $X \subseteq V(G_1)$  and  $\gamma_G(P) \in \Gamma'$  for each  $A_1$ -path  $P$  in  $G_1 - X$ , then  $|X| \geq 2l$ .

**Subproof.** Suppose that  $|X| < 2l$ . For  $i \in \{1, \dots, 4l\}$  let  $W_i = V(T(x)) \cup V(T(y))$  where  $T(x)$  and  $T(y)$  are the trees that contain the ends of  $P_i$ . Since  $|X| < 2l$ , there is a path  $P_i$  such that  $X \cap (V(P_i) \cup W_i) = \emptyset$ . Suppose that  $W_i = V(T(x)) \cup V(T(y))$  and that  $P_i$  is an  $(a_x, a_y)$ -path. There is an  $(a_y, a_x)$ -path  $P'$  in  $G_1$  such that  $E(P') \subseteq E(T(x)) \cup E(T(y)) \cup E(H)$ ; thus  $\gamma_{G_1}(P') \in \Gamma'$ . Let  $B$  be the block of  $G_1 - X$  that contains  $G_1[(A_1 \cup \{x, y\}) - X]$ . By Lemma 2.7,  $B$  is  $\Gamma'$ -balanced. By 3.2.3,  $B$  contains  $a_x$  and  $a_y$ . Hence,  $P_i$  and  $P'$  are both contained in  $B$ . Let  $W'$  be the closed walk obtained by appending  $P'$  to  $P_i$ . Thus  $\gamma_B(W) = \gamma_B(P') + \gamma_B(P_i) \notin \Gamma'$ , contradicting Lemma 2.3.  $\square$

By the above claim and Corollary 2.6, there exist vertex-disjoint  $A_1$ -paths  $Q_1, \dots, Q_l$  in  $G_1$  such that  $\gamma_{G_1}(Q_i) \notin \Gamma'$  for each  $i \in \{1, \dots, l\}$ . Now the result follows immediately from Lemma 3.1.  $\square$

### Acknowledgments

These results were obtained during the *Workshop on Structural Graph Theory* at the Pacific Institute for the Mathematical Sciences (PIMS), Vancouver, Canada, 2003. We thank the organizers, Luis Goddyn and Bruce Reed, and the other participants, Maria Chudnovsky, Mike Lohman, Paul Seymour, Bruce Shepherd, and Adrian Vetta, for useful discussions. We also thank the referees for their helpful comments.

### References

- [1] M. Chudnovsky, W.H. Cunningham, J. Geelen, An algorithm for packing non-zero  $A$ -paths in group-labelled graphs, *Combinatorica* 28 (2008) 145–161.
- [2] M. Chudnovsky, J. Geelen, B. Gerards, L. Goddyn, M. Lohman, P. Seymour, Non-zero  $A$ -paths in group-labelled graphs, *Combinatorica* 26 (2006) 521–532.
- [3] J. Geelen, B. Gerards, B. Reed, P. Seymour, A. Vetta, On the odd-minor variant of Hadwiger’s conjecture, *J. Combin. Theory Ser. B* (2009), doi: 10.1016/j.jctb.2008.03.006, in press.
- [4] N. Robertson, P.D. Seymour, Graph minors. X. Obstructions to tree-decomposition, *J. Combin. Theory Ser. B* 52 (1991) 153–190.
- [5] N. Robertson, P.D. Seymour, Graph minors. XIII. The disjoint paths problem, *J. Combin. Theory Ser. B* 63 (1995) 65–110.
- [6] N. Robertson, P.D. Seymour, Graph minors. XVI. Excluding a non-planar graph, *J. Combin. Theory Ser. B* 89 (2003) 43–76.
- [7] N. Robertson, P.D. Seymour, Graph minors. XX. Wagner’s conjecture, *J. Combin. Theory Ser. B* 92 (2004) 325–357.
- [8] T. Zaslavsky, Biased graphs. I. Bias, balance, and gains, *J. Combin. Theory Ser. B* 47 (1989) 32–52.
- [9] T. Zaslavsky, Biased graphs. II. The three matroids, *J. Combin. Theory Ser. B* 51 (1991) 46–72.