


Europ. J. Combinatorics (2000) 21, 701–714

Article No. 10.1006/eujc.1999.0377

Available online at <http://www.idealibrary.com> on 



Decomposition and l_1 -Embedding of Weakly Median Graphs

HANS-JÜRGEN BANDELT AND VICTOR CHEPOI[†]

Weakly median graphs, being defined by interval conditions and forbidden induced subgraphs, generalize quasi-median graphs as well as pseudo-median graphs. It is shown that finite weakly median graphs can be decomposed with respect to gated amalgamation and Cartesian multiplication into 5-wheels, induced subgraphs of hyperoctahedra (alias cocktail party graphs), and 2-connected bridged graphs not containing K_4 or $K_{1,1,3}$ as an induced subgraph. As a consequence one obtains that every finite weakly median graph is l_1 -embeddable, that is, it embeds as a metric subspace into some \mathbb{R}^n equipped with the 1-norm.

© 2000 Academic Press

In this paper we continue to elaborate on a structure theory of graphs based on two fundamental operations, viz., Cartesian multiplication and gated amalgamation. While Cartesian multiplication is a standard operation, gated amalgamation seems to appear only in the context of median graphs and their generalizations; cf. [4, 6, 8, 23, 27]. An induced subgraph H of a graph G is called *gated* if for every vertex x outside H there exists a vertex x' (the *gate* of x) in H such that each vertex y of H is connected with x by a shortest path passing through the gate x' ; cf. [18]. G is a *gated amalgam* of two graphs G_1 and G_2 if G_1 and G_2 are (isomorphic to) two intersecting gated subgraphs of G whose union is all of G . A graph with at least two vertices is said to be *prime* if it is neither a proper Cartesian product nor a gated amalgam of smaller graphs. For instance, the only prime median graph is the two-vertex complete graph K_2 ; see Isbell [21] and van de Vel [26]. More generally, the prime quasi-median graphs are exactly the complete graphs; quasi-median graphs were introduced by Mulder [23] and further studied in [8, 14, 28]. The pseudo-median graphs form yet another class of graphs for which the prime members are known; see Bandelt and Mulder [6]. Unfortunately, the latter class is not closed under Cartesian multiplication and does not include all quasi-median graphs. In order to overcome these deficiencies we consider here the somewhat larger class of weakly median graphs, previously studied by Chepoi [10, 11] under the name ‘locally median’ graphs. First, we say that a graph G is *weakly modular* if its shortest-path metric $d = d_G$ satisfies the following two conditions:

- for any three vertices u, v, w with $1 = d(v, w) < d(u, v) = d(u, w)$ there exists a common neighbour x of v and w such that $d(u, x) = d(u, v) - 1$;
- for any four vertices u, v, w, z with $d(v, z) = d(w, z) = 1$ and

$$2 = d(v, w) \leq d(u, v) = d(u, w) = d(u, z) - 1,$$

there exists a common neighbour x of v and w such that $d(u, x) = d(u, v) - 1$.

For an illustration of these conditions see [9, Figure 2]. A *weakly median* graph is a weakly modular graph that does not contain any two vertices with an unconnected triple of common neighbours; see Figure 1.

Note that all bridged graphs are weakly modular; cf. [7, 10]. A graph is called *bridged* if G does not contain any isometric cycle of length greater than 3, that is, each cycle of length greater than 3 has a shortcut in G ; see Soltan and Chepoi [25] and Farber and Jamison [20].

[†]Current address: Laboratoire d’Informatique de Marseille, Université d’Aix Marseille II, Faculté des Sciences de Luminy, F-13288 Marseille Cedex 8, France.




FIGURE 1. Forbidden induced subgraphs.

Bridged graphs can easily be constructed since, according to [1, Corollaries 2.4 and 2.6], they admit certain vertex elimination schemes (relaxing simplicial elimination for chordal graphs). Now, a weakly median bridged graph G (i.e., a bridged graph in which the first two graphs of Figure 1 are forbidden induced subgraphs) is prime exactly when it has at least two vertices and does not have any cut vertex, that is, it is either K_2 or two-connected. Indeed, since G contains no induced 4-cycles, G cannot decompose as a nontrivial Cartesian product; G cannot be a gated amalgam because two-connected bridged graphs have no proper gated subgraphs other than singletons. This is an immediate consequence of the following two elementary facts: first, a gated subgraph cannot intersect a triangle in just a single edge; and second, the neighbourhood of any vertex induces a connected subgraph in a two-connected bridged graph. Furthermore, all wheels (whether bridged or not) are prime weakly median graphs; an n -wheel ($n \geq 4$) consists of a cycle of length n and a ‘central’ vertex adjacent to all vertices of the cycle. Finally, all multipartite graphs of the form $K_{i_1, i_2, i_3, \dots}$ (with $1 \leq i_j \leq 2$) different from K_1 , $K_{1,2}$, and $K_{2,2}$ are prime weakly median graphs. A particular instance is the α -octahedron $K_{2,2,2, \dots}$ (or *hyperoctahedron*, for short), which is the complement of the disjoint union of $\alpha \geq 3$ copies of K_2 . For convenience, we refer to induced subgraphs of hyperoctahedra as to *subhyperoctahedra* when they contain either K_4 or an induced 4-wheel $K_{1,2,2}$ (that is, whenever they constitute 1-skeletons of at least three-dimensional polyhedra).

THEOREM 1. *Every finite weakly median graph G (with more than one vertex) is obtained by successive applications of gated amalgamations from Cartesian products of the following prime graphs: two-vertex complete graphs, 5-wheels, subhyperoctahedra, and two-connected, K_4 - and $K_{1,1,3}$ -free bridged graphs. The latter bridged graphs are exactly the graphs which can be realized as plane graphs such that all inner faces are triangles and all inner vertices have degrees larger than 5. A weakly median graph is prime if and only if it does not have any proper gated subgraphs other than singletons.*

Particular instances of this result are the decomposition theorems for quasi-median graphs (with prime graphs being complete) [8, 23] and pseudo-median graphs [6].

An important feature of weakly median graphs is that they embed in rectilinear space. A finite graph G with shortest-path metric d_G is said to be l_1 -embeddable if there exists a distance-preserving embedding φ into some \mathbb{R}^n equipped with the 1-norm $\|\cdot\|_1$, that is,

$$d_G(x, y) = \|\varphi(x) - \varphi(y)\|_1$$

for all vertices x, y of G . Assouad and Deza [3] have shown that a graph G is l_1 -embeddable if and only if for some integer $\eta \geq 1$ it admits a *scale η embedding* ψ in some hypercube Q (being a Cartesian power of K_2), that is,

$$\eta \cdot d_G(x, y) = d_Q(\psi(x), \psi(y)) \quad \text{for all } x, y;$$




FIGURE 2. A weakly median bridged graph H and a fragment of a hypercube indicating the scale 2 embedding of H .

see Figure 2 for an instance of a scale 2 embedding. The graphs with scale 1 embeddings in hypercubes are thus the isometric subgraphs of hypercubes (characterized in [17]). Shpectorov [24] has proved that a finite graph is l_1 -embeddable if and only if it is an isometric subgraph of the Cartesian product of hyperoctahedra and ‘half-cubes’ (which are obtained from one parity half of a hypercube, with two vertices being adjacent exactly when their distance in the hypercube equals 2). Our following result shows that in order to decide whether a given graph G is l_1 -embeddable it suffices to check its *prime components*, i.e., those prime (gated) subgraphs from which G can be built up by successive Cartesian multiplications and gated amalgamations:

PROPOSITION 1. *A finite graph G is l_1 -embeddable if and only if every prime component of G is such. G has a scale η embedding in a hypercube if and only if every prime component does. G is l_1 -rigid if and only if every prime component is such.*

The particular instance of amalgamations along single vertices has already been dealt with in [16, Proposition 7.6.1].

It is well known that a graph G is l_1 -embeddable if and only if its metric d can be expressed in the form

$$d = \sum_{i=1}^m \lambda_i \cdot \delta_i \quad (\lambda_i > 0 \text{ for } i = 1, \dots, m)$$

as a positive linear combination of ‘split’ (alias ‘cut’) metrics δ_i that are associated with *splits* $\{A_i, B_i\}$ of G , i.e., partitions of the vertex-set into two parts, according to

$$\delta_i(x, y) = \begin{cases} 0 & \text{if either } x, y \in A_i \text{ or } x, y \in B_i, \\ 1 & \text{otherwise,} \end{cases}$$

for $i = 1, \dots, m$ (cf. [5, 16]). If, in addition, this decomposition of d is unique, then G is called *l_1 -rigid*. Necessarily, for each i the sets A_i and B_i occurring in the above decomposition constitute complementary *half-spaces* of G , that is, either set includes all shortest paths of G between any two of its vertices, and both sets together cover the vertex-set. Thus, G has a scale η embedding in a hypercube if and only if there is a collection \mathcal{Z} of splits such that any two adjacent vertices of G are separated by (i.e., in different parts of) exactly η splits of \mathcal{Z} . In particular, for $\eta = 1$ one obtains the well-known characterization of isometric subgraphs of hypercubes [17]: G is isometrically embeddable in a hypercube if and only if G is bipartite and $W(u, v) = \{x : d(u, x) < d(v, x)\}$ is a half-space for each pair u, v of adjacent vertices.

The specific information on half-spaces of weakly median graphs obtained in [11, 12] together with Theorem 1 and Proposition 1 enables us to prove the concluding result:

THEOREM 2. *Every finite weakly median graph G is l_1 -embeddable. G has a scale 2 embedding in a hypercube if and only if it does not contain $K_{1,1,1,1,2}$ (that is, K_6 minus an edge) as an induced subgraph. Furthermore, G is l_1 -rigid if and only if it is K_4 -free.*

A finite weakly median graph G which contains some induced $K_{1,1,1,1,2}$ has scale η embeddings only for $\eta \geq 4$. The minimum scale η then depends solely on the maximal induced subhyperoctahedra. To determine this number η for a particular subhyperoctahedron is not a trivial task; see [15] or [16, Chapter 7.4].

PROOF OF THEOREM 1

We commence by establishing a number of auxiliary results. Unless stated otherwise, G is always a weakly median graph (not necessarily finite). Recall that the *interval* $I(u, v)$ between two vertices u and v in G consists of all vertices x on shortest paths from u to v , that is, $d(u, x) + d(x, v) = d(u, v)$. An induced subgraph (or a subset of vertices) H is called *convex* if H includes every interval $I(u, v)$ of G between two vertices u, v from H . Half-spaces are thus the nonempty convex sets with nonempty convex complement. The smallest convex (or gated, respectively) subgraph containing a given subgraph S is the *convex hull* (or *gated hull*, respectively) of S . A subgraph H is said to be Δ -closed if, for every triangle having two vertices in H , the third vertex belongs to H as well; then the smallest Δ -closed subgraph containing S is the Δ -closure of S . In order to check whether a given subgraph of G is convex or gated the following lemma is useful, which essentially coincides with Theorem 7 of Chepoi [10] and can be proved quite easily by induction.

LEMMA 1. *A connected subgraph H of G is convex if and only if for every pair of vertices at distance 2 in H all their common neighbours belong to H . Moreover, a convex subgraph is gated if and only if it is Δ -closed.*

The next four lemmas provide the necessary information on gated hulls and isometric cycles in G . A *prism* is the Cartesian product $K_2 \square K_3$ of K_2 and K_3 , and a *house* is obtained from a prism by deleting one vertex.

LEMMA 2. *The convex hull of an induced (i) house, (ii) 5-cycle, (iii) 4-wheel, respectively, in G is a (i) prism, (ii) 5-wheel, (iii) (maximal) subhyperoctahedron, respectively.*

PROOF. Assertion (i) is obtained from [6, Lemma 4] and its proof.

Let C be an induced 5-cycle. Since G is weakly median there exists a common neighbour z of three vertices of C , two of which are adjacent with the third one being opposite to them. Then C and z constitute a 5-wheel, for otherwise, an induced house arises whose convex hull would not be a prism. If there is yet another vertex y in G adjacent to two non-adjacent vertices on C , then we would obtain either one of the forbidden induced subgraphs (see Figure 1) or an induced house whose convex hull is not a prism.

An induced 4-wheel can be extended to a maximal induced subhyperoctahedron H in G . Clearly H is included in the convex hull of this wheel. Suppose that H is not convex: then there exist two non-adjacent vertices u and v in H with a common neighbour y outside H . Since the third and fourth graphs of Figure 1 are forbidden, y is in fact adjacent to all pairs of non-adjacent vertices from H . Hence, as H together with y cannot induce a subhyperoctahedron, there exist two adjacent common neighbours w and x of u and v in H such that u, v, w, x, y induce the fourth graph of Figure 1, giving a contradiction. \square

LEMMA 3. *Induced 5-wheels and convex subhyperoctahedra containing an induced 4-wheel are gated in G and do not contain any proper gated subgraphs other than singletons.*

PROOF. In view of the preceding lemmas it suffices to show that induced 5-wheels and convex subhyperoctahedra containing an induced 4-wheel are Δ -closed. Clearly they do not have any proper gated subgraphs other than singletons.

Suppose that W is an induced 5-wheel which is not Δ -closed. Then two adjacent vertices u and v of W have a common neighbour y outside W . If, say, u is the central vertex of W , then y must also be adjacent to the two common neighbours of u and v in W in order to avoid forbidden subgraphs. This, however, contradicts the fact that induced 5-wheels are convex. Therefore u and v are peripheral vertices of the wheel. Let z be the central vertex of W , and let t be the vertex opposite to the edge uv on the cycle. If $d(t, y) = 3$, then as G is weakly median the two vertices of W different from t, u, v, z would have a common neighbour with y , which is necessarily outside W , contrary to convexity. Hence $d(t, y) = 2$, and so (again as G is weakly median) there exists a common neighbour x of t, y, z , which is impossible by what has just been shown.

Let H be a convex subhyperoctahedron which is not Δ -closed. Then there exists a vertex y outside H such that the neighbours of y in H form a complete subgraph K of size at least 2. Consider any induced 4-cycle C in H . If two vertices of K belong to C , then we would obtain an induced house the convex hull of which includes an induced 4-wheel, which is impossible. Therefore y and any vertex pair from K together with two (suitably chosen) non-adjacent vertices on C induce the first graph of Figure 1, a contradiction. \square

LEMMA 4. *There are no isometric odd cycles in G of length greater than 5.*

PROOF. Suppose the contrary, and choose a cycle C having minimal length among all isometric odd cycles of length at least 7. Let C consist of the vertices x_0, \dots, x_{2n} and edges $x_i x_{i+1}$ ($i = 0, \dots, 2n$, indices modulo $2n + 1$). Since x_0 and x_1 are at distance n from x_{n+1} , they have a common neighbour y_1 with $d(y_1, x_{n+1}) = n - 1$ (because G is weakly median). Further, x_2 and x_1 have a common neighbour y_2 with $d(y_2, x_{n+1}) = n - 2$. Continuing this way, we eventually obtain a shortest path $x_0, y_1, \dots, y_{n-1}, x_{n+1}$ such that each y_i is adjacent to x_i ($i = 1, \dots, n - 1$). Observe that each y_i is actually different from x_{i+1} because $d(x_0, x_{i+1}) = i + 1$ but $d(x_0, y_i) = i$. Now, a shortest path y_n, \dots, y_{2n-3} is constructed as follows: let y_i be a common neighbour of y_{i-1} and x_{i+2} with $d(x_0, y_i) = 2n - i - 2$ for $i = n, \dots, 2n - 3$. Again, each y_i ($i \geq n - 1$) must be different from x_{i+3} . We claim that the cycle induced by $x_0, x_1, \dots, x_{n-1}, y_{n-1}, y_n, \dots, y_{2n-3}$ is isometric. Indeed, suppose that for some $i = n, \dots, 2n - 3$ the distance from y_i to one of x_{i-n+1}, x_{i-n+2} is smaller than $n - 1$. Then $d(x_{i+2}, x_{i-n+1}) < n$ or $d(x_{i+2}, x_{i-n+2}) < n$ would follow, conflicting with the isometry of C . This proves the claim. Since the new isometric cycle has length $2n - 1$, we must have $n = 3$ by virtue of the initial minimality assumption. Thus, x_0, x_1, x_2, y_2, y_3 induce a 5-cycle with y_1 in its convex hull. Hence, by Lemma 2, x_2 and y_1 are adjacent. Now, by interchanging the roles of x_i and x_{8-i} for $i = 1, 2, 3$ we obtain yet another 5-wheel with central vertex y_1 , so that x_6 and y_1 must be adjacent as well. This, however, implies $d(x_2, x_6) = 2$, a final contradiction. \square

LEMMA 5. *If G does not contain any induced 4-wheel or 5-wheel, then the gated hull of any triangle is a two-connected bridged graph H , which does not have any proper gated subgraph other than singletons.*

PROOF. First suppose that some two-connected, weakly median bridged graph F contains a proper gated subgraph S which is not a singleton. Pick a vertex x outside S having a neighbour

w in S . Any neighbour v of w in S is connected with x by a path within the neighbourhood of w because F is bridged and two-connected. Then, however, as S is Δ -closed, all vertices on this path (including x) must belong to S , yielding a contradiction.

Now, there exists a maximal isometric two-connected bridged subgraph H of G that contains a given triangle. In the case where G is infinite, this follows from Zorn's lemma because directed unions preserve isometry, two-connectedness as well as the property of being bridged. To prove the lemma it thus suffices to show that H is gated (by what has just been proven).

First suppose that H is not convex. Then by Lemma 1, we can find non-adjacent vertices x, y in H having a common neighbour z in H and another one, v , outside H . Since H is two-connected and bridged, x and y are connected by a path P in H which is fully included in the neighbourhood of z . We may assume that v, x, y, z and P are chosen (in regard to the stated properties) so that P has minimal length. We claim that v and z must be adjacent. Suppose the contrary: then P has length at least 3, for otherwise, we would obtain the fourth graph of Figure 1 or the 4-wheel as an induced subgraph, both of which are forbidden here. By minimality, P has no neighbours of v other than x and y . Thus v, x, y, z together with the neighbour t of x on P induce a house. This house extends to an induced prism (according to Lemma 2); let w denote the common neighbour of t, v, y . Then w does not belong to H because H is bridged. Now, the vertices w, t, y, z and the subpath of P connecting t and y violate the minimality assumption. This proves the claim.

Next we show that the larger subgraph H' induced by H together with v is also isometric in G . Suppose the contrary: let u be a vertex of H such that the distance of u and v in H' exceeds the distance $d(u, v) = k \geq 2$ in G . As no shortest path from u to v in G can pass through one of the three neighbours x, y, z of v (because H is isometric), the distances $d(u, x), d(u, y), d(u, z)$ are necessarily between k and $k + 1$. So, we distinguish two cases.

Case 1. $d(u, z) = k + 1$.

If $d(u, x) = d(u, y) = k$, then x and y have a common neighbour t in the isometric bridged subgraph H such that $d(u, t) = k - 1$ (since H is weakly modular), thus yielding a forbidden 4-cycle (induced by t, x, z, y) in H . Therefore $d(u, y) = k + 1$, say. Then H contains a common neighbour t of y and z at distance k to u . In the weakly modular graph G we find a common neighbour w of t and v with $d(u, w) = k - 1$, whence t, v, w, y, z induce the fourth graph of Figure 1, which is impossible.

Case 2. $d(u, z) = k$.

Since now the vertices v and z are equidistant to u , they have a common neighbour w in G with $d(u, w) = k - 1$. As the distance between u and v in H' is larger than k , the vertex w lies outside H . In order to avoid forbidden induced subgraphs, w must be adjacent to both x and y . Now, replacing v by w and k by $k - 1$ we are back in Case 1, thus leading to a contradiction.

We conclude that the extended subgraph H' is indeed isometric. From what has been shown above we know that H' cannot include any induced 4-cycle. Therefore H' does not have any isometric even cycle at all: for otherwise, such a cycle C would contain v , and the two neighbours x', y' of the vertex z' opposite to v on C would admit a second common neighbour v' in G (because of weak modularity) which satisfies $d(v, v') = d(v, z') - 2$, so that a forbidden 4-cycle arises (whether v' belongs to H or not). By the hypothesis of the lemma (together with Lemma 2), H' is without induced 5-cycles, and hence by Lemma 4, it must be bridged. Clearly H' is two-connected, and therefore we arrive at a final contradiction to the maximality of H . This proves that H is in fact convex.

It remains to verify that H is Δ -closed as well. Suppose there exists a vertex v outside H having at least two neighbours x, y in H . Then, as H is convex, the neighbours of v in H form

a complete subgraph. Thus, v is a simplicial vertex of the extended subgraph H' induced by H and v . Hence H' is a bridged graph, which is evidently two-connected. As to isometry, consider any vertex u of H . If $d(u, y) < d(u, v)$, then u and v are at distance $d(u, v)$ in H' , too. We may therefore assume that $d(u, y) = d(u, v)$ because H is convex. Since G is weakly modular, we can find a common neighbour w of v and y at distance $d(u, v) - 1$ to u . As H is convex, w belongs to H . This shows that there is a shortest path in H' between u and v of length $d(u, v)$. It follows that H' is also a two-connected, isometric bridged subgraph, thus conflicting with the choice of H . Therefore H is Δ -closed, concluding the proof (by Lemma 1). \square

The next lemma ensures that the prime graphs listed in Theorem 1 actually encompass all two-connected, weakly median bridged graphs.

LEMMA 6. *A two-connected bridged graph H is weakly median if and only if either (1) H is a complete graph K_n ($n \geq 4$), or (2) H equals $K_{1,1,\dots,1,2}$ (i.e., a complete graph minus an edge, having more than four vertices), or (3) H does not contain K_4 or $K_{1,1,3}$ as an induced subgraph.*

PROOF. If H is of type (1) or (2), it is a subhyperoctahedron; if H satisfies (3), then H does not contain any forbidden induced subgraph of Figure 1 and hence is weakly median.

Conversely, suppose that H is weakly median and contains some K_4 but is not a subhyperoctahedron of type (1) or (2). Extend this K_4 to a maximal induced subhyperoctahedron H' , which is necessarily convex, being a complete graph or a complete graph minus an edge (since H has no induced 4-cycles). By the hypothesis, we can find a vertex z outside H' which forms a triangle together with two vertices from H' . Now, however, we arrive at a contradiction in that either $H' \cup \{z\}$ would induce a subhyperoctahedron or a forbidden induced subgraph (from Figure 1) would arise. \square

To characterize the two-connected, K_4 - and $K_{1,1,3}$ -free bridged graphs via their planar embeddings, we will make use of the following counting argument.

LEMMA 7. *Let G be a finite two-connected plane graph in which all inner faces are triangles and all inner vertices (i.e., the vertices not incident with the outer face) have degrees larger than 5. Then the numbers n_2 and n_3 of vertices with degrees 2 and 3 satisfy the inequality $2n_2 + n_3 \geq 6$.*

PROOF. Let f denote the number of inner faces of G , m the number of edges, n the number of vertices, and b the number of vertices incident with the outer face. Then

$$f - m + n = 1 \quad \text{and} \quad 3f + b = 2m$$

hold according to Euler's theorem and the hypothesis that all inner faces are triangles. Eliminating f yields

$$3n - b - m = 3.$$

The information on the vertex degrees is turned into the inequality

$$\begin{aligned} 2m &\geq 6(n - b) + 4(b - n_2 - n_3) + 3n_3 + 2n_2 \\ &= 6n - 2b - 2n_2 - n_3, \end{aligned}$$

whence

$$2n_2 + n_3 \geq 6n - 2b - 2m = 6,$$

as required. \square

We are now in position to identify the finite two-connected, K_4 - and $K_{1,1,3}$ -free bridged graphs with the plane graphs described in the preceding lemma, when choosing a planar embedding such that the outer face is bounded by the edges contained in exactly one triangle.

For the bridged graphs with the additional properties we construct a planar embedding recursively by employing the dismantling scheme of Anstee and Farber [1] (for a short proof, see [13]): there exists a vertex z dominated by some neighbour y in the sense that every vertex adjacent to z is also adjacent or identical to y . If the degree of z was larger than 3, then y and z together with three common neighbours would either induce $K_{1,1,3}$ or include some K_4 , contrary to the hypothesis. Therefore z has degree 2 or 3. We assume that G has at least four vertices and that the desired planar embeddings can be realized for all proper induced subgraphs which are two-connected.

Case 1. z has exactly two neighbours x and y (which are adjacent).

Then $G - z$ is a graph of the same kind, to which the induction hypothesis applies. Note that the edge xy belongs to exactly one triangle of $G - z$ because G is K_4 - and $K_{1,1,3}$ -free. Therefore we have chosen a planar embedding of $G - z$ with xy on the boundary of the outer face. Attaching the triangle x, y, z to $G - z$ so that z lies in the outer face of $G - z$, we obtain a plane graph with the required properties.

Case 2. z has exactly three common neighbours w, x , and y (such that y is adjacent to w and x).

Then w and x are not adjacent. If $G - z$ is not two-connected, then y is the unique cut vertex. Moreover, $G - \{y, z\}$ comprises exactly two components, which together with y induce either K_2 or two-connected subgraphs of G . In any case we can transform and combine the planar embeddings of these subgraphs so that wy and xy lie on one line for which one of the associated closed half-planes includes $G - z$. Placing z onto the complementary open half-plane and linking it with w, x, y produces the desired embedding. If $G - z$ is two-connected, then we could choose the planar embedding of $G - z$ right away, with wy and xy lying on the boundary of the outer face (since both edges belong to exactly one triangle of $G - z$). Locating z in this outer face we can extend the planar embedding to G , thereby creating two new triangles and turning y into an inner vertex. Take a minimal path P from $G - z$ in the neighbourhood of y which connects w and x . Then P together with y and z induce a k -wheel with $k \geq 6$, whence y satisfies the degree constraint.

As to the converse, let G be a plane graph satisfying the hypothesis of Lemma 7. We may assume that G has at least four vertices. Consider any triangle of G : together with its interior in the plane it constitutes a plane graph H to which Lemma 7 equally applies. We infer that each vertex of the boundary triangle must have degree 2 in H , that is, H includes no inner vertex. Hence all triangles of G constitute inner faces (and vice versa). In particular, G does not include any K_4 or $K_{1,1,3}$ as an induced subgraph. To show that G is bridged, we proceed by induction.

Case 1. There exist two adjacent vertices u and v separating G .

Then necessarily u and v both lie on the boundary of the outer face. We can thus decompose G into two plane subgraphs G_1 and G_2 whose boundaries intersect in the edge uv and cover the boundary of G . Certainly, G_1 and G_2 fulfil the hypothesis of Lemma 7 and hence are bridged by the induction hypothesis. Then G is bridged as well.

Case 2. G does not have any separating edge.

Then, by Lemma 7, the boundary contains some vertex v of degree 3 in G . Let w, x, y be the neighbours of v . One of the edges vw, vx, vy does not lie on the boundary, say vx . Since

vx is thus contained in two triangles, w and y must be adjacent to x . Further, x cannot be a boundary vertex, for otherwise, the edge vx would separate G . Therefore the plane subgraph $G - v$ is two-connected and inherits its inner faces and inner vertices from G . It follows from the induction hypothesis that $G - v$ is bridged. Suppose that G contains some isometric cycle C of length $2k$ or $2k + 1$ with $k \geq 2$. Then C includes v, w, y but not x . Substituting v by x creates a cycle in $G - v$ of the same length. This cycle must have a short cut, so that some vertex z on C is at distance k to v but $k - 1$ to x in G (as $G - v$ is clearly isometric in G). If w and y are at distance $k - 1$ to z , then (as $G - v$ is weakly modular) there exist common neighbours w' of w and x and y' of x and y , both at distance $k - 2$ to z . Recall that, for any two vertices y and z in a bridged graph, the neighbours of y on shortest paths between y and z form a complete subgraph [20, 25]. In particular, here either $w' = y'$ or w' and y' are adjacent. Then v, w, x, y together with w', y' induce a 4- or 5-wheel, so that x would become an inner vertex of G with degree smaller than 6, a contradiction. Therefore C must be an odd cycle such that exactly one of w, y is at distance k to z , say w . Then the neighbour $u \neq v$ of w on C must be adjacent to x because $G - v$ is bridged. Hence, as C is isometric, we infer that C is a 5-cycle comprising v, w, u, z, y , which together with x induces a 5-wheel, again a contradiction. We conclude that G is bridged.

This completes the proof of the second statement in Theorem 1, characterizing the specific bridged graphs.

The subsequent Lemmas 8 and 10 are needed to detect amalgams or products within G . Any gated subset S of G gives rise to a partition W_a ($a \in S$) of the vertex-set of G ; viz., the fibre W_a of a relative to S consists of all vertices x (including a itself) having a as their gate in S . For adjacent vertices a, b of S , let U_{ab} be the set of vertices from W_a which are adjacent to vertices from W_b .

LEMMA 8. *Let S be a gated subgraph of G . Then each fibre W_a relative to S is gated. There exists an edge between two distinct fibres W_a and W_b if and only if a and b are adjacent. Moreover, for any two adjacent vertices a, b of S , the sets U_{ab} and U_{ba} constitute isomorphic gated subgraphs of G under the canonical isomorphism $f_{ab} : U_{ab} \rightarrow U_{ba}$ that maps each vertex in U_{ab} to its unique neighbour in U_{ba} .*

PROOF. We adapt some arguments from [6, proof of Theorem 12]. If $x \in W_a$ and $y \in W_b$ are adjacent, then as $d(b, x) \leq d(b, y) + 1$ we obtain $d(a, x) \leq d(b, y)$ and by symmetry, $d(a, x) = d(b, y)$; therefore, since a and b are the gates of x and y , respectively, in S , a and b must be adjacent.

We claim that any vertex $v \in W_b$ has at most one neighbour in W_a for $a \neq b$. Suppose the contrary: let v be adjacent to two distinct vertices x, y from W_a . Then a and b are adjacent, and $d(a, v) = d(a, x) + 1 = d(a, y) + 1$, by what has just been shown. By weak modularity, x and y have a common neighbour z (necessarily in W_a) at distance $d(a, x) - 1$ from a . Further, as $d(b, x) = d(b, v) + 1 = d(b, z) + 1$, there exists a common neighbour w (necessarily in W_b) of v and z at distance $d(b, v) - 1$ from b . The five vertices v, w, x, y, z now induce the third or fourth graph of Figure 1, which is impossible. This proves the claim.

Each fibre W_a is connected because $I(a, x) \subseteq W_a$ for all $x \in W_a$. Then, by the above claim and Lemma 1, W_a is convex as well as Δ -closed and hence gated.

Let $x \in U_{ab}$ be adjacent to $x' \in U_{ba}$ (for some edge ab). Every neighbour w of x in $I(a, x) \subseteq W_a$ has the same distance to b as x' . Hence, by weak modularity, w and x' have a common neighbour w' , which necessarily belongs to W_b . Therefore $w \in U_{ab}$, and it follows by a straightforward induction that $I(a, x) \subseteq U_{ab}$. In particular, U_{ab} is connected. To prove that U_{ab} is gated, apply Lemma 1: let z be a common neighbour of $x, y \in U_{ab}$, which nec-

essarily belongs to W_a and is at distance 2 to the (respective) neighbours of x, y in U_{ba} ; then the gate of z in W_b is adjacent to z , showing that $z \in U_{ab}$, as desired.

Finally, let x and y be adjacent vertices in U_{ab} , with neighbours x' and y' , respectively, in U_{ba} . Since W_a and W_b are gated, x' and y' must be adjacent. We conclude that the neighbours map f_{ab} is an isomorphism from U_{ab} onto U_{ba} . \square

The *cycle space* of a graph with edge-set E is the subspace of $(\mathbf{GF}(2))^E$ comprising all unions of closed walks. The isometric cycles clearly generate this space. In the presence of weak modularity the triangles and induced 4-cycles generate all isometric cycles, as is easily seen by induction. Recall from Duchet *et al.* [19] or Jamison [22] that a graph is *null-homotopic* if its cycle space admits a basis constituted solely of triangles. We then record the following elementary fact.

LEMMA 9. *A weakly modular graph is null-homotopic whenever every induced 4-cycle extends to a 4-wheel.*

In the case that a proper gated subgraph S of G is two-connected and null-homotopic we can say more about the associated sets U_{ab} : the following lemma constitutes the tool for detecting proper decompositions of non-bipartite weakly median graphs.

LEMMA 10. *Let S be a gated two-connected and null-homotopic subgraph of G . Then the gated subgraphs U_{ab} (with a, b adjacent in S) are all isomorphic, and their union induces a gated subgraph H isomorphic to a Cartesian product $S \square U$ (where U may be any U_{ab}). If W_a and U_{ab} ($b \in S$) do not coincide for some $a \in S$, then G is the gated amalgam of W_a and $G - (W_a - U_{ab})$.*

PROOF. First we show that $U_{ab} = U_{ac}$ whenever a, b, c form a triangle in S . Let xy be an edge of G with $x \in U_{ab}$ and $y \in U_{ba}$. According to Lemma 8 and its proof, c is equidistant to x and y , whence there is a common neighbour z of x and y on shortest paths to c . Then $d(c, z) = d(c, x) - 1 = d(a, x)$, and hence as a is the gate of x in S we infer that z belongs to W_c . Therefore $z \in U_{ca}$ as well as $x \in U_{ac}$. Interchanging the roles of a, b, c , this proves

$$U_{ab} = U_{ac}, \quad U_{ba} = U_{bc}, \quad U_{ca} = U_{cb}.$$

Now assume that q and r are any two non-adjacent neighbours of a in S . Then, as S is two-connected, there exists a path P from q to r not passing through a . By C denote the closed walk from q to r along P and then back to q via the vertex a . To prove that $U_{aq} = U_{ar}$ we proceed by induction on the minimal number k of triangles whose (modulo 2) sum gives C (thereby using the null-homotopy of S). Since P does not include a , there must be some common neighbour s of a and r such that the closed walk obtained from C by substituting the pair a, r by the triplet a, s, r is the (modulo 2) sum of $k - 1$ triangles. Then $U_{aq} = U_{as} = U_{ar}$ by virtue of the induction hypothesis. This justifies the shorthand U_a for the sets U_{ab} .

We can verify that all subgraphs U_a ($a \in S$) are actually isomorphic using the same kind of argument: we claim that for every closed walk $a_0, a_1, \dots, a_n, a_0$ the composition $f_{a_n a_0} \circ f_{a_{n-1} a_n} \circ f_{a_{n-2} a_{n-1}} \circ \dots \circ f_{a_0 a_1}$ is the identity map. Indeed, this is evidently true for triangles ($n = 2$), by the first part of the proof. The general case is settled again by induction on the minimal number of triangles adding up to C . In particular, we get a unique isomorphism f_{rs} from U_r to U_s for any two (not necessarily adjacent) vertices $r, s \in S$, obtained by composing the isomorphisms f_{ab} along the edges ab of any path from r to s .

The product representation of H (the subgraph induced by the union of all sets U_a) is now immediate: pick any vertex a in S and consider the mapping

$$f : S \square U_a \rightarrow H$$

$$(s, x) \mapsto f_{as}(x).$$

This constitutes the desired isomorphism since (i) each mapping f_{as} is an isomorphism from U_a to U_s , (ii) the sets U_s ($s \in S$) partition H , (iii) there is an edge between U_r and U_s only if r and s are adjacent, and (iv) the isomorphism f_{rs} maps each vertex x onto a neighbour whenever r and s are adjacent.

Finally, assume $W_a \neq U_a$ for some $a \in S$. Since the subgraph $G - (W_a - U_a)$ and the gated fibre W_a together cover G and intersect in a gated subgraph (viz., U_a), G is the gated amalgam of $G - (W_a - U_a)$ and W_a . \square

Now, we have collected all the information that is necessary to conclude the proof of Theorem 1. Assume that G is neither a singleton nor any of the prime graphs listed in Theorem 1. Then, by Lemma 6, G is not a two-connected bridged graph. We have to show that G has a proper gated subgraph S with at least two vertices and G decomposes as a gated amalgam or Cartesian product. If G includes an induced 4- or 5-wheel, then by Lemmas 2 and 3 it has a proper gated subgraph S , which is a subhyperoctahedron or a 5-wheel. Since these graphs are null-homotopic and two-connected, Lemma 10 provides us with the required decomposition. So, we can assume that G is without induced 4- or 5-wheels. If G still contains some triangle, then by Lemma 5 we obtain a proper gated subgraph S which is bridged and two-connected. Since bridged graphs are null-homotopic (cf. Lemma 9), Lemma 10 applies again, yielding a proper decomposition of G . It remains to consider the case where G is triangle-free. Then, by Lemmas 2 and 4, there are no odd cycles at all, whence G is a median graph, in which any edge ab serves as a proper gated subgraph S . This subgraph S leads to a decomposition as stated in Lemma 10; cf. [21, 23].

In conclusion, note that Cartesian multiplication distributes overgated amalgamation, viz., the Cartesian product of a graph H with a gated amalgam of two graphs G_1 and G_2 equals the gated amalgam of $H \square G_1$ and $H \square G_2$. This completes the proof of Theorem 1.

PROOF OF PROPOSITION 1

In the case where $G = G_1 \square G_2$ is the Cartesian product of two nontrivial graphs G_1 and G_2 the assertions of Proposition 1 are evident; see [16, Proposition 7.5.2]. In fact, we may regard G_1 and G_2 as gated subgraphs of G intersecting in a single vertex. Then the half-spaces of G correspond to the pairs H_1, G_2 and G_1, H_2 where each H_i is a half-space of G_i .

As to gated amalgamation, the following observation is instrumental.

LEMMA 11. *Let G be a graph having a scale η embedding φ in a hypercube Q . Then every scale η embedding ψ of a gated subgraph S of G in some hypercube R extends to a scale η embedding of G in some hypercube containing R .*

PROOF. Let T be the convex hull of the image $\varphi(S)$ in the hypercube Q . For each vertex x of G let x' be the gate of x in S . We claim that $\varphi(x')$ is the gate of $\varphi(x)$ in the subhypercube T . Suppose it is not: then some vertex z from $T - \varphi(S)$ is this gate. Choose any half-space H of Q with $\varphi(x') \in H$ but $z \notin H$. Since $\varphi(x')$ is in the interval between $\varphi(x)$ and $\varphi(y)$ for each $y \in S$ (as φ is a scale embedding), it follows that H includes $\varphi(S)$ and hence T , yielding a contradiction. This proves the claim. In particular, the distance between the gates in T of two

vertices $\varphi(w)$, $\varphi(x)$ from the image of G equals $\eta d_G(w', x')$. Let U be a subhypercube of Q intersecting T in a single vertex such that the convex hull of T and U is all of Q . Letting φ_U denote the scale embedding φ of G in Q followed by the gate map onto U , we thus have

$$d_U(\varphi_U(w), \varphi_U(x)) = \eta(d_G(w, x) - d_G(w', x')).$$

Now, the required scale η extension of ψ is given by

$$x \mapsto (\psi(x'), \varphi_U(x)) \in R \square U.$$

Indeed, for vertices w, x of G ,

$$\begin{aligned} \eta d(w, x) &= \eta d_G(w', x') + d_U(\varphi_U(w), \varphi_U(x)) \\ &= d_R(\psi(w'), \psi(x')) + d_U(\varphi_U(w), \varphi_U(x)). \end{aligned}$$

□

Assume that G is the gated amalgam of two graphs G_1 and G_2 , which admit scale η embeddings φ_1 and φ_2 in hypercubes Q_1 and Q_2 , respectively. Let T be the convex hull of $\varphi_1(G_1 \cap G_2)$ in Q_1 . By virtue of Lemma 11 we can extend the restriction $\varphi_1|_{G_1 \cap G_2}$ to a scale η embedding ψ of G_2 in a hypercube R such that R intersects Q_1 only in T . The median graph $Q_1 \cup R$ extends isometrically to a hypercube Q . We can then regard the union $\varphi_1 \cup \psi$ as a mapping from G to Q , yielding the required scale η embedding.

As to l_1 -rigidity, observe that H is a half-space of G exactly when either H or its complement is a half-space of G_1 or G_2 not intersecting $G_1 \cap G_2$, or H is a gated amalgam of half-spaces H_i of G_i ($i = 1, 2$). This obviously implies that G is l_1 -rigid whenever G_1 and G_2 are such. Conversely assume that the l_1 -embeddable graph G contains a gated subgraph S which is not l_1 -rigid. Then G has some scale η embedding φ in a hypercube, while S admits yet another scale ξ embedding ψ in a hypercube such that $\varphi|_S$ and ψ induce different weighted systems of pairs of complementary half-spaces on S . Without loss of generality assume that the scales ξ and η are the same (since scales can be enlarged to arbitrary multiples). Then the extension of ψ to G guaranteed by Lemma 11 is essentially different from φ , showing that G is not l_1 -rigid.

PROOF OF THEOREM 2

In view of Proposition 1 it suffices to verify the assertions of the theorem only for the prime components of the given graph G . We may therefore assume that G is prime. First note that the 1-cube K_2 is trivially l_1 -rigid.

LEMMA 12. *In a prime K_4 -free weakly median graph G other than K_2 , any two adjacent vertices u and v are separated by exactly two distinct pairs of complementary half-spaces.*

PROOF. Let V denote the vertex-set of G . Since G is two-connected, null-homotopic, and K_4 -free, every edge belongs to exactly one or two triangles. Consider any triangle u, v, w in G . Recall from [12, Lemmas 9 and 10] that then the sets $W(u, v) = \{x \in V : d(u, x) < d(v, x)\}$ and $W(v, u) \cup W(w, u)$ are convex.

Case 1. w is the unique common neighbour of u and v .

Then every vertex x equidistant to u and v is closer to w than u and v (by weak modularity), whence the convex sets $W(u, v)$, $W(v, u)$, and $W(w, u) \cap W(w, v)$ partition V . It follows that

$W(u, v)$ and $W(v, u)$ are non-complementary half-spaces. If H is any half-space with $u \in H$ and $v, w \in V - H$, then necessarily $W(u, v) \subseteq H$ and $W(v, u) \cup W(w, u) \subseteq V - H$, thus yielding $W(u, v) = H$.

Case 2. u and v have exactly two common neighbours, w and w' .

Then every vertex x equidistant to u and v belongs to $W(w, u) \cap W(w, v)$ or $W(w', u) \cap W(w', v)$. Therefore $W(u, v) \cup W(w, v)$, $W(v, u) \cup W(w', u)$ and $W(u, v) \cup W(w', v)$, $W(v, u) \cup W(w, u)$ constitute two distinct pairs of complementary half-spaces. Let H be any half-space with $u \in H$ and $v \in V - H$. Since $u, v \in I(w, w')$, the vertices w and w' are separated by $H, V - H$, say $w \in H$ and $w' \in V - H$. Then necessarily $W(w, v) \subseteq H$ and $W(w', u) \subseteq V - H$, whence $H = W(u, v) \cup W(w, v)$. \square

From Lemma 12 and the observations preceding Theorem 2, we immediately infer that the graphs G of Lemma 12 have scale 2 embeddings in hypercubes. The proof further shows that the three splits of any triangle u, v, w uniquely extend to pairs of complementary half-spaces of G . This implies that the associated split metrics are linearly independent, thus establishing l_1 -rigidity.

To conclude the proof of Theorem 2, we can assume (by Theorem 1) that G is a subhyperoctahedron containing K_4 . The l_1 -embeddability of hyperoctahedra has been established by Assouad [2]. It is easy to see that the 4-octahedron $K_{2,2,2,2}$ has a scale 2 embedding in a 4-cube, but the minimum scale for $K_{1,1,1,1,2}$ equals 4; see [16, Lemma 7.4.6]. The scale 2 embeddable subhyperoctahedra containing K_4 are thus the subhyperoctahedra K_{i_1, i_2, i_3, i_4} with $1 \leq i_j \leq 2$ ($j = 1, 2, 3, 4$), all of which fail to be l_1 -rigid; cf. [16, Proposition 7.4.3]. This completes the proof of Theorem 2.

ACKNOWLEDGEMENTS

We thank an anonymous referee for her/his careful reading of the manuscript, which helped to improve the presentation. Part of this research was done when the second author was supported by the Alexander von Humboldt Stiftung.

REFERENCES

1. R. P. Anstee and M. Farber, On bridged graphs and cop-win graphs, *J. Comb. Theory, Ser. B*, **44** (1988), 22–28.
2. P. Assouad, Embeddability of regular polytopes and honeycombs in hypercubes, in: *The Geometric Vein., The Coxeter Festschrift*, Springer, Berlin, 1981, pp. 141–147.
3. P. Assouad and M. Deza, Espaces métriques plongeables dans un hypercube: aspects combinatoires, *Ann. Discrete Math.*, **8** (1980), 197–210.
4. H.-J. Bandelt and V. Chepoi, Cellular bipartite graphs, *Europ. J. Combinatorics*, **17** (1996), 121–134.
5. H.-J. Bandelt and A. W.M. Dress, A canonical decomposition theory for metrics on a finite set, *Adv. Math.*, **92** (1992), 47–105.
6. H.-J. Bandelt and H. M. Mulder, Pseudo-median graphs: decomposition via amalgamation and Cartesian multiplication, *Discrete Math.*, **94** (1991), 161–180.
7. H.-J. Bandelt and H. M. Mulder, Cartesian factorization of interval-regular graphs having no long isometric odd cycles, in: *Graph Theory, Combinatorics, and Applications*, volume 1, Y. Alavi, G. Chartrand, O. R. Oellermann and A. J. Schwenk (eds), Wiley, New York, 1991, pp. 55–75.

8. H.-J. Bandelt, H. M. Mulder and E. Wilkeit, Quasi-median graphs and algebras, *J. Graph Theory*, **18** (1994), 681–703.
9. H.-J. Bandelt and E. Pesch, Dismantling absolute retracts of reflexive graphs, *Europ. J. Combinatorics*, **10** (1989), 211–220.
10. V. Chepoi, Classifying graphs by means of metric triangles (in Russian), *Metody Diskret. Analiza*, **49** (1989), 75–93.
11. V. Chepoi, d -Convexity and local conditions on graphs (in Russian), *Issledovania po Prikladnoj Matematike i Informatike*, Chişinău, Stiinţa, (1990), 184–191.
12. V. Chepoi, Separation of two convex sets in convexity structures, *J. Geom.*, **50** (1994), 30–51.
13. V. Chepoi, Bridged graphs are cop-win graphs: an algorithmic proof, *J. Comb. Theory, Ser. B*, **69** (1997), 97–100.
14. F. R. K. Chung, R. L. Graham and M. E. Saks, A dynamic location problem for graphs, *Combinatorica*, **9** (1989), 111–131.
15. M. Deza and M. Laurent, l_1 -Rigid graphs, *J. Algebr. Comb.*, **3** (1994), 153–175.
16. M. Deza and M. Laurent, *Geometry of Cuts and Metrics*, Springer-Verlag, Berlin, 1997.
17. D. Ž. Djoković, Distance-preserving subgraphs of hypercubes, *J. Comb. Theory, Ser. B*, **14** (1973), 263–267.
18. A. W. M. Dress and R. Scharlau, Gated sets in metric spaces, *Aeq. Math.*, **34** (1987), 112–120.
19. P. Duchet, M. Las Vergnas and H. Meyniel, Connected cutsets of a graph and triangle bases of the cycle space, *Discrete Math.*, **62** (1986), 145–154.
20. M. Farber and R. E. Jamison, On local convexity in graphs, *Discrete Math.*, **66** (1987), 231–247.
21. J. R. Isbell, Median algebra, *Trans. Am. Math. Soc.*, **260** (1980), 319–362.
22. R. E. Jamison, On the null-homotopy of bridged graphs, *Europ. J. Combinatorics*, **8** (1987), 421–428.
23. H. M. Mulder, The interval function of a graph, *Math. Centre Tracts*, **132**, Amsterdam, 1980.
24. S. V. Shpectorov, On scale embeddings of graphs into hypercubes, *Europ. J. Combinatorics*, **14** (1993), 117–130.
25. V. P. Soltan and V. D. Chepoi, Conditions for invariance of set diameters under d -convexification in a graph, *Cybernetics*, **19** (1983), 750–756.
26. M. van de Vel, Matching binary convexities, *Topology Appl.*, **16** (1983), 207–235.
27. M. L.J. van de Vel, *Theory of Convex Structures*, North-Holland, Amsterdam, 1993.
28. E. Wilkeit, The retracts of Hamming graphs, *Discrete Math.*, **102** (1992), 197–218.

Received 22 March 1999 and accepted in revised form 29 November 1999

HANS-JÜRGEN BANDELT
*Mathematisches Seminar,
 Universität Hamburg,
 Bundesstr. 55,
 D-20146 Hamburg,
 Germany*
 AND

VICTOR CHEPOI
*SFB 343, Diskrete Strukturen in der Mathematik,
 Universität Bielefeld,
 D-33615 Bielefeld,
 Germany*