



ELSEVIER

Contents lists available at ScienceDirect

Redox Biology

journal homepage: www.elsevier.com/locate/redox

Research Paper

Possible involvement of membrane lipids peroxidation and oxidation of catalytically essential thiols of the cerebral transmembrane sodium pump as component mechanisms of iron-mediated oxidative stress-linked dysfunction of the pump's activity



T.I. Omotayo^a, G.S. Akinyemi^a, P.A. Omololu^a, B.O. Ajayi^a, A.A. Akindahunsi^a, J.B.T. Rocha^b, I.J. Kade^{a,*}

^a Department of Biochemistry, Federal University of Technology, Akure, Ondo State, Nigeria

^b Centro de Ciencias Naturais e Exatas, Programa Posgraduacao em Bioquimica Toxicologica, Universidade Federal de Santa Maria, RS, Brazil

ARTICLE INFO

Article history:

Received 18 November 2014

Received in revised form

21 December 2014

Accepted 22 December 2014

Available online 24 December 2014

Keywords:

Oxidative stress

Iron

Electrogenic sodium pump

Vitamin E

Dithiothreitol

Thiol oxidation

ABSTRACT

The precise molecular events defining the complex role of oxidative stress in the inactivation of the cerebral sodium pump in radical-induced neurodegenerative diseases is yet to be fully clarified and thus still open. Herein we investigated the modulation of the activity of the cerebral transmembrane electrogenic enzyme in Fe²⁺-mediated in vitro oxidative stress model. The results show that Fe²⁺ inhibited the transmembrane enzyme in a concentration dependent manner and this effect was accompanied by a biphasic generation of aldehydic product of lipid peroxidation. While dithiothreitol prevented both Fe²⁺ inhibitory effect on the pump and lipid peroxidation, vitamin E prevented only lipid peroxidation but not inhibition of the pump. Besides, malondialdehyde (MDA) inhibited the pump by a mechanism not related to oxidation of its critical thiols. Apparently, the low activity of the pump in degenerative diseases mediated by Fe²⁺ may involve complex multi-component mechanisms which may partly involve an initial oxidation of the critical thiols of the enzyme directly mediated by Fe²⁺ and during severe progression of such diseases; aldehydic products of lipid peroxidation such as MDA may further exacerbate this inhibitory effect by a mechanism that is likely not related to the oxidation of the catalytically essential thiols of the ouabain-sensitive cerebral electrogenic pump.

© 2014 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Reports have shown that the activity of Na⁺/K⁺-ATPase in many tissues and organs of mammalian systems can be profoundly inhibited under conditions of oxidative stress related diseases. Thus the pathophysiology, molecular and biochemical mechanisms underlying the reduced activity of the electrogenic enzyme in several free-radical linked degenerative and metabolic pathologies have been an area of intense study [1–5]. Although Na⁺/K⁺-ATPase is widely expressed in quite a number of tissues, the transmembrane protein is mainly expressed in brain and neuronal cells. Consequently, dysfunction of the pump has been characterized by neuronal hyperexcitability, depolarization and swelling [6]. Hence, the malfunctioning of the cerebral sodium pump has been associated with some specific neurological

disorders such as diabetes, Alzheimer's disease, epilepsy and bipolar disorder among others. In these neurological pathologies, constant depolarization across the cell membrane induces an imbalance in the amount of neurotransmitters that are released within the cell [3,4,7–10].

More importantly, it has been observed that neurological disorders associated with low activity of the cerebral sodium pump have also been characterized by increased oxidative stress indices in the brain of such human or disease models. Thus there is an inverse relationship between extent of oxidative stress mediated damage and the activity of the Na⁺/K⁺-ATPase in these neurological dysfunctions [11]. However, earlier investigators have attempted to unravel the underlying mechanisms that participate in free radicals-induced dysfunction of the pump. In this regard, emerging data led to some speculative conclusions.

Firstly, it was suggested that reactive oxygen species (ROS) may evoke an inhibition on the enzyme indirectly. Herein, the ROS may attack lipids in the membrane that anchor the transmembrane

* Corresponding author.

E-mail address: ijkade@yahoo.com (I.J. Kade).

enzyme and consequently leading to an altered membrane microviscosity, which ultimately result in loss of the enzyme function [12,13]. On the other hand, ROS have been shown to directly attack and eventually modify amino acid residues of the enzyme thus leading to loss of activity of the transmembrane protein [14]. Yet some authors have speculated that aldehydic products which arise from the lipid peroxidation process may also interact with the transmembrane enzyme ultimately leading to its inactivation [15,16].

Despite these findings and observations however, the precise biochemical and molecular events that characterize free radical-induced dysfunction of the cerebral enzyme appear complex and far from being completely understood. Partly, the complexity arises from the multifactorial agents and factors that mediate oxidative stress. Therefore, in order to better understand the molecular events associated with oxidative stress-induced inactivation of the cerebral pump, the study of the participation of these agents and factors must be holistic, all inclusive and total. Evidently, unravelling these complex mechanism(s) is thus still open.

One of the most potent neurotoxic agent that exhibit prooxidant effect under *in vitro* and *in vivo* conditions is iron especially Fe^{2+} . In fact, reports have shown that impaired iron metabolism is an initial cause of some neurodegeneration mediated by ROS [17]. Moreso, several common genetic and sporadic neurodegenerative disorders have been associated with dysregulated iron homeostasis in the central nervous system (CNS) [17]. In general, iron accumulation causes neurodegeneration chiefly by inducing the formation of free radicals which eventually damage cellular macromolecules such as lipids, DNA and proteins. Furthermore, iron-induced oxidative stress is particularly dangerous because it can cause further iron release from iron-containing proteins such as ferritin, heme-proteins, and Fe-S clusters, forming a destructive intracellular positive-feedback loop that exacerbates the toxic effects of brain iron overload [17].

The harmful and deleterious effect of iron especially in the brain has been the prime interest of researchers. Primarily, this is because the mammalian brain is highly susceptible to free radical attack due to high oxygen tension and a high content of polyunsaturated fatty acids in cell membrane phospholipids [18]. Evidences suggest that iron invasion is an initial cause of neuronal cell death and axonal degeneration [19–21]. Moreso, a cardinal pathology associated with several common sporadic neurodegenerative disorders such as Alzheimer's disease, Parkinson's disease and Huntington's disease has been associated with iron accumulation. Findings have shown that under *in vitro* models relating to these pathologies, there is a strong coincidence between localized disturbed iron homeostasis and neuronal cell death in the brain [19,20,22–26]. Furthermore, several *in vivo* studies of iron regulatory proteins using transgenic rodents [27,28] have indeed suggested that iron may be a pathogenic factor for neurodegeneration. Indeed experimental evidences have suggested that iron chelators [20,29,30] and genetic manipulation resulting in low iron levels [29,31,32] are neuroprotective strategies against these iron-induced neurodegenerative diseases. Other central nervous system (CNS) disorders that have been proposed to be associated with disturbed iron homeostasis include the Freiderich's Ataxia [26], Restless Leg Syndrome [33], Ischaemic/Haemorrhagic Stroke [34] and Multiple Sclerosis [25].

From the foregoing, it is apparent that iron-mediated ROS production may be a suitable model in unravelling the precise mechanisms of ROS-mediated dysfunction of the ouabain-sensitive electrogenic sodium pump. Really, some authors have studied iron-induced oxidative stress models to study the inactivation of the cerebral pump's activity. In this regard, Rauchova and colleagues observed that iron-induced altered membrane fluidity thus

leading to the inhibition of the Na^+/K^+ -ATPase activity at the initial process of lipid peroxidation. However, as lipid peroxidation progresses, the inhibition of the transmembrane enzyme strongly correlates only with the production of thiobarbituric acid reactive species (TBARS) and conjugated dienes (CD) [11]. However, their study is largely inconclusive and thus still open. Primarily, their study did not take into consideration the fact that the transmembrane enzyme is a sulphhydryl protein and thus the possibility of a direct or indirect interaction between iron and the enzyme was not clarified and hence the need for the present study.

Materials and methods

Chemicals

Adenosine triphosphate (ATP), α -tocopherol, cysteine, reduced glutathione, dithiothreitol, thiobarbituric acid (TBA), were obtained from Sigma (St. Louis, MO). All other chemicals which are of analytical grade were obtained from standard commercial suppliers.

Animals

Male adult Wistar rats (200–250 g) from our own breeding colony were used. Animals were kept in separate animal cages, on a 12-h light: 12-h dark cycle, at a room temperature of 22–24 °C, and with free access to food and water. The animals were used according to standard guidelines on the Care and Use of Experimental Animal Resources.

Preparation of tissue homogenate

Rats were decapitated under mild ether anaesthesia and the cerebral tissue (whole brain) was rapidly removed, placed on ice and weighed. The brain was immediately homogenized in cold 10 mM Tris-HCl, pH 7.4 (1/10, w/v) with 10 up-and-down strokes at approximately 1200 rev/min in a Teflon-glass homogenizer. The homogenate was centrifuged for 10 min at 4000g to yield a pellet that was discarded and a low-speed supernatant (S1).

Incubation systems for TBARS and sodium pump assays

Aliquot of S1 was used for the assays of thiobarbituric acid reactive substances (TBARS) as well as Na^+/K^+ -ATPase activity. For both assays the reaction mixture contained 3 mM MgCl₂, 125 mM NaCl, 20 mM KCl and 50 mM Tris-HCl, pH 7.4, FeSO_4 (final concentrations range of 1–100 μM), with and without α -tocopherol (final concentration, 1–100 μM) and with and without dithiothreitol (final concentration, 2 mM), with or without malondialdehyde (MDA) (1–100 μM) and 100–180 μg protein in a final volume of 500 μl . The reaction was initiated by addition of ATP to a final concentration of 3.0 mM. Controls were carried out under the same conditions with the addition of 0.1 mM ouabain. The reaction mixture was incubated at 37 °C for 30 min. At the end of the incubation time period, tubes were assayed for sodium pump activity and TBARS production.

Assay of sodium pump

The reaction system for the assay of the activity of cerebral Na^+/K^+ -ATPase was essentially the same as described above under the section "incubation systems for TBARS and sodium pump". However, at the end of the incubation time period (30–60 min), the reaction was stopped by addition of 5% trichloroacetic acid. Released inorganic phosphorous (P_i) was measured by the method

of Fiske and Subbarow [35]. Na^+/K^+ -ATPase activity was calculated by the difference between two assays (with and without ouabain). All the experiments were conducted at least three times and similar results were obtained. Protein was measured by the method of Lowry et al. [36], using bovine serum albumin as standard.

For all enzyme assays, incubation times and protein concentration were chosen to ensure the linearity of the reactions. All samples were run in duplicate. Controls with the addition of the enzyme preparation after mixing with trichloroacetic acid (TCA) were used to correct for non-enzymatic hydrolysis of substrates. Enzyme activity was expressed as nmol of phosphate (P_i) released $\text{min}^{-1} \text{mg protein}^{-1}$.

Assay of TBARS

The reaction system for the assay of production of the cerebral TBARS was essentially the same as described above under the section "incubation systems for TBARS and sodium pump". However, at the end of the incubation time period (30–60 min) production of TBARS was determined as described by the method of Ohkawa et al. [37] except that the buffer of colour reaction has a pH of 3.4. The colour reaction was developed by adding 200 μl 8.1% SDS to S1, followed by sequential addition of 500 μl acetic acid/HCl (pH 3.4) and 500 μl 0.8% of thiobarbituric acid (TBA). This mixture was incubated at 100 °C for 1 h. TBARS produced were measured at 532 nm and the absorbance was compared to that of a standard curve obtained using malondialdehyde (MDA).

Oxidation of thiols

The rate of dithiothreitol (DTT) oxidation was determined in the presence of 50 mM Tris-HCl, pH 7.4, and varying concentration of Fe^{2+} (10–100 μM). The rate of thiol oxidation was evaluated by measuring the disappearance of -SH groups. Free -SH groups were determined according to Ellman [38]. Incubation at 37 °C was initiated by the addition of DTT (final concentration, 2 mM). Aliquots of the reaction mixture (100 μl) were checked (at 1 h interval for 3 h) for the amount of -SH groups at 412 nm by the addition of the colour reagent 5'5'-dithio-bis (2-nitrobenzoic) acid (DTNB).

Statistical analysis

Results were analysed by appropriate analysis of variance (ANOVA) and this is indicated in text of results. Duncan's Multiple Range Test was applied. Differences between groups were considered to be significant when $p < 0.05$.

Results

Effect of Fe^{2+} , DTT and vitamin E on cerebral MDA production

In order to better explore the experimental results, separate and combined analysis of variance were conducted on the experimental data. As depicted in Fig. 1 (panel a), one-way ANOVA analysis indicate that Fe^{2+} elicited a biphasic effect on the production of aldehydic product of lipid peroxidation with lower




Fig. 1. Biphasic effect of iron (II) on MDA production from cerebral tissue homogenate (panel a). The influence of vitamin E (100 μM) (panel b), DTT (2 mM) (panel c) and combination of both 100 μM vitamin E and 2 mM DTT (panel d) on the biphasic effect iron (II) are shown as indicated. Data are presented as mean \pm SEM of at least three independent experiments carried out in different days. Data were analysed using appropriate analysis of variance followed by Duncan test as indicated in the description of results. *Significantly lower from control; #significantly higher than control ($p < 0.05$).

concentrations evoking an increase in lipid peroxidation and higher concentrations of Fe^{2+} appear to be counteracting the production of these aldehydic products of lipid peroxidation. Conversely, in the presence of vitamin E, a potent antioxidant, two-way ANOVA [(with and without vitamin E) \times (5 concentrations of Fe^{2+})], of the results obtained revealed that the production of aldehydic products of lipid peroxidation elicited by Fe^{2+} was markedly hindered (Fig. 1, panel b). However, since earlier data has shown that DTT could protect the oxidation of thiols especially on the cerebral sodium pump, attempt was made to investigate the possible inhibitory effect of this dithiol on lipid peroxidation. Similar to the result presented in Fig. 1 (panel b), two-way ANOVA [(with and without DTT) \times (5 concentrations of Fe^{2+})] clearly revealed that DTT have marked inhibitory effect on the biphasic pattern of aldehydic productions elicited by Fe^{2+} (Fig. 1, panel c). The possible effect of the combination of both vitamin E and DTT on the lipid peroxidation process was also tested. In this regard, three-way ANOVA [(with and without DTT) \times (with and without vitamin E) \times (5 concentrations of Fe^{2+})] clearly revealed that the lipid peroxidation process was inhibited in the presence of vitamin E whether alone or in combination with DTT (Fig. 1, panel d). Suffice to mention that for all analysis, mean values are considered significant from each other when $p < 0.05$.

Effect of Fe^{2+} , vitamin E and DTT on cerebral Na^+/K^+ -ATPase activity

In the reaction system for the assay of the cerebral sodium pump, the effect of increasing concentration of Fe^{2+} on the activity of the pump was tested. Herein it was observed that one-way ANOVA distinctly revealed that Fe^{2+} evoked a concentration-dependent decrease in the activity of the cerebral transmembrane sodium pump (Fig. 2, panel a). Based on this observed effect of

Fe^{2+} on the activity of the cerebral pump and considering Fig. 1a, it is rational to speculate that lipid peroxidation process may be linked to the observed inhibitory effect of Fe^{2+} on the activity of the enzyme. Hence the possible protective effect of vitamin E, a potent inhibitor of lipid peroxidation on the inhibitory effect of Fe^{2+} on the activity of the cerebral pump was tested. Two-way ANOVA analysis [(with and without vitamin E) \times (5 concentrations of Fe^{2+})] of the data obtained revealed that vitamin E had no protective effect on Fe^{2+} -induced inhibition of the activity of the cerebral pump (Fig. 2, panel b). Furthermore, since DTT has already been documented to protect against oxidation of sulphhydryl groups of the enzyme, the possible involvement of oxidation of the critical thiols of the enzyme as a component of its inhibition by Fe^{2+} was also investigated. Two-way ANOVA [(with and without DTT) \times (5 concentrations of Fe^{2+})] showed that DTT exerted considerable protective effect on the activity of the enzyme (Fig. 2, panel c). In fact, three-way ANOVA (with and without vitamin E) \times (with and without DTT) \times (5 concentrations of Fe^{2+}) revealed that when both vitamin E and DTT were combined they exerted a marked protective effect on the activity of the enzyme (Fig. 2, panel d).

Effect of vitamin E on activity of Na^+/K^+ -ATPase

The effect of increasing concentration of vitamin E on the activity of the ouabain sensitive cerebral electrogenic pump is presented in Fig. 3. One-way ANOVA apparently revealed that the transport function of the electrogenic pump was not affected by the potent antioxidant.




Fig. 2. Effect of iron (II) on the activity of the cerebral sodium pump (a). The different panels shows the effect of 100 μM vitamin E (panel b), 2 mM DTT (panel c) and combination of both vitamin E and DTT (panel d) on the activity of the cerebral sodium pump subjected to iron (II) assault. Data are presented as mean \pm SEM of at least three independent experiments carried out in different days. Data were analysed using appropriate analysis of variance followed by Duncan test as indicated in the description of results. *Significantly lower from control; #significantly higher than control ($p < 0.05$).




Fig. 3. Effect of vitamin E on Na⁺/K⁺-ATPase activity. Data are presented as mean ± SEM of three independent experiments carried out in different days. Data were analysed using one-way ANOVA followed by Duncan test ($p < 0.05$).

Effect of Fe²⁺ on oxidation of dithiothreitol

As shown in Fig. 4, the rate of DTT oxidation was increased by Fe²⁺. Two-way ANOVA (control/Fe²⁺ × 4 sample times) or (control/4 concentrations of Fe²⁺ × 4 sample times) revealed a significant Fe²⁺ versus time interaction ($p < 0.001$). In fact, the oxidation of DTT was increased as a function of time in the presence of Fe²⁺.

Effect of MDA on activity of Na⁺/K⁺-ATPase

Fig. 5 (panel a) shows the effect of increasing concentration of MDA on the activity of Na⁺/K⁺-ATPase. One-way ANOVA indicates that MDA inhibits the pump in a concentration dependent manner. Furthermore the inhibition of the cerebral enzyme by MDA was not markedly ($p < 0.05$) modified in the presence of DTT (Fig. 5, panel b).




Fig. 4. Effects of iron on the rate of dithiothreitol oxidation. The rate of oxidation was evaluated at the indicated times and concentrations of iron. Data are the means of five to seven independent experiments carried out in different days. Data are expressed as mean ± SEM and post-hoc comparisons were done by Duncan's multiple range test. *Significant difference in relation to the control.

Discussion

Excellent reports and reviews have clearly shown that the transmembrane Na⁺/K⁺-ATPase is a target in oxidative stress-linked degenerative diseases (such as diabetes, Alzheimer diseases etc.) in which free radicals participate in their pathophysiology [39,40] and thus in human and animal models of these degenerative diseases, there is an inverse relationship between extent of oxidative stress and activity of this important transmembrane enzyme [41,42]. However, the precise molecular events that characterize the role of oxidative stress in the dysfunction of the enzyme's activity is yet to be fully clarified. In any case, a definitive answer to the molecular puzzle on the precise role of oxidative stress in the loss of transport function of the Na⁺/K⁺-ATPase in degenerative diseases may represent a key step in the development of the proper treatment of such diseases.

Several authors have made elegant attempts to unravel the precise mechanisms that are involved in the pathophysiological processes leading to the inactivation of the pump's activity under conditions of oxidative stress. Thus far, available data clearly indicate that oxidative stress mediated altered membrane dynamics strongly correlated with altered activity of the sodium pump [43,44]. In this regard also, the sequential molecular events that eventually led to pump's malfunction remain obscure and thus experimental data may be warranted to further clarify these molecular processes.

Based on earlier findings [19,20,22–28], we decided to use an in vitro model of Fe²⁺-mediated oxidative stress to simulate possible in vivo oxidative stress situation in the brain. In fact, iron has been shown to react with superoxide anion (O₂^{•-}) and hydrogen peroxide (H₂O₂) to produce the hydroxyl radical (•OH) via the Fenton reaction [57]. These radicals can also lead to the formation of other reactive oxygen species (ROS) [56]. Ultimately, the overproduction of ROS can directly attack the polyunsaturated fatty acids of the cell membranes and induce lipid peroxidation. Hence, this model may be ideal to study the possible correlation between oxidative stress and the transport function of the transmembrane enzyme under study.

Firstly, it is worthwhile mentioning that at low concentrations, Fe²⁺ markedly increased the formation of aldehydic products of lipid peroxidation which is indicated by increase in the calculated malondialdehyde content of the brain tissue. However, at high concentration of Fe²⁺ this transition metal appears to inhibit the lipid peroxidation process (Fig. 1, panel a). This observation is a paradox and quite intriguing. In any case, earlier authors have suggested that when in excess, Fe²⁺ could compete with lipids as electron donors for Fe³⁺O₂^{•-} (perferryl ion), thereby inhibiting hydrogen abstraction and thus lipid peroxidation [45–47]. Conversely, there is the likelihood that the quantification of lipid peroxidation products using thiobarbituric acid could also detect lipid hydroperoxides. Hence, at high iron concentrations, there is the strong possibility that excess iron could likely decompose lipid hydroperoxides and ultimately decrease the observed level of thiobarbituric acid reactive substances observed in this study. Notwithstanding, at low concentrations, Fe²⁺-induced cerebral lipid peroxidation may be an excellent model to simulate in vivo situations involving oxidative stress assault on plasma membrane lipids in the brain and nervous cells. The implication of the above argument is that Fe²⁺ could disorganize membrane lipid bilayer arrangement of the brain cells and leading to altered membrane integrity and ultimately perturbed membrane fluidity. Expectedly, these series of events may lead to a parallel disorganization and eventual loss of function of the membrane bound protein.

Consistent with the above reasoning, Fig. 2 (panel a) clearly shows that Fe²⁺ inhibited the activity of the ouabain-sensitive electrogenic pump in a concentration dependent manner. In sharp




Fig. 5. Effect of varying concentrations of MDA on the activity of Na⁺/K⁺-ATPase in the absence (a) and presence of DTT (b). Data are presented as mean ± SEM of three independent experiments carried out in different days. Data were analysed using one-way ANOVA followed by Duncan test. *Significant difference from control at $p < 0.05$.

contrast with what was observed in Fig. 1 (panel a), high concentrations of the transition metal exerted a more profound inhibitory effect on the activity of the cerebral pump. Interpretation of this intriguing data may be complex. However, one could speculate that at low doses, Fe²⁺ may ultimately generate the highly reactive radical •OH via Fenton reactions. Next the reactive radical may assault cerebral membrane bound lipids, causing their peroxidation and ultimately altering membrane fluidity. In this case, the structural integrity of the transmembrane pump is adversely altered leading to a loss of its activity. On the other hand, there is a strong line of evidence from the present result to suggest that Fe²⁺ may bind directly to essential groups on the enzyme thus evoking a loss of its activity. This reason is plausible considering the fact that at high concentration there was inhibition of lipid peroxidation and yet the activity of the enzyme (Fig. 2, panel a).

Based on the above findings, we developed some conditions to further clarify the possible inter-relationship between of oxidative stress and the activity of the pump. In this regard, attempt was made to inhibit lipid peroxidation with an antioxidant agent that would possibly not affect the activity of the pump. Herein, vitamin E, a classical antioxidant was evaluated for its possible inhibitory effect on the activity of the cerebral sodium pump. The result (Fig. 3) clearly shows that this potent antioxidant when acting alone has no effect on the activity of the transmembrane sodium pump. With this fascinating result, this classical antioxidant was then employed to counteract the oxidative assault evoked by Fe²⁺ on brain lipids. As demonstrated in Fig. 1 (panel b), vitamin E effectively inhibited the lipid peroxidation process mediated by Fe²⁺ assaults at all concentrations tested. Conversely, while this potent antioxidant inhibited lipid peroxidation, it did not protect against inhibitory effect of Fe²⁺ on the activity of the enzyme (Fig. 2, panel b). This fact strongly suggests that in addition to inducing a peroxidation of cerebral lipids, the transition metal may also be interacting directly with this transmembrane protein.

Therefore, based on the latter speculation above, a reaction system was also developed to investigate the possibility of a direct interaction between Fe²⁺ and the transmembrane sodium pump that ultimately led to loss of its function. The reaction system was developed based on previous knowledge on the structure and function of the cerebral pump. Earlier authors and emerging data in our laboratory have consistently demonstrated that the cerebral sodium pump has sulphhydryl groups that are critical to its activity and that these thiols can be inhibited by oxidizing agents such as selenium-containing compounds and inorganic mercury [39,40,48–53]. In these reports [48,51,52], the authors observed that these compounds oxidize critical thiol groups on the enzyme and that endogenous thiols could prevent or relieve the inhibition

imposed on the enzyme by these thiol oxidizing compounds.

Therefore in order to explore the possibility of Fe²⁺ oxidizing the catalytically important sulphhydryl groups of the cerebral sodium pump, an in vitro method was developed to investigate the ability of Fe²⁺ to oxidize dithiothreitol alone. Interestingly, as shown in Fig. 4, Fe²⁺ profoundly increased the oxidation of the dithiol in a time dependent fashion. This observation when translated may indicate that the toxic effect of this transition metal on the activity of the cerebral pump may be partly linked with its ability to oxidize essential thiols on the electrogenic pump that may be critical to its catalytic function.

In order to verify the possible participation of the oxidation of essential thiol on the electrogenic sodium pump by Fe²⁺ as a component of its inhibitory mechanism on the pump's activity, the incubation condition developed allowed for the pre-incubation of dithiothreitol (DTT) with the cerebral enzyme before the assault of Fe²⁺. The data obtained (Fig. 2, panel c) shows that when DTT was preincubated with the enzyme before the assault of Fe²⁺, the dithiol completely prevented the inhibition of the enzyme's activity earlier evoked by Fe²⁺ assault as previously observed in Fig. 2 (panel a). Intriguing as this finding may seem, it is noteworthy that DTT also exerted a marked inhibitory effect on the lipid peroxidation process initiated by Fe²⁺ (Fig. 1, panel c). As a strong reductant, the mechanism by which DTT inhibited lipid peroxidation may be partly related to its ability to reduce the transition metal.

With the above findings describing the individual effect of vitamin E and the dithiol on both the activity of the cerebral sodium pump as well as the inhibition of lipid peroxidation processes, another incubation condition was carried to evaluate the combined effect of both vitamin E and the dithiol on the lipid peroxidation process and the activity of the pump. The results obtained clearly show that combination of vitamin E and DTT prevented lipid peroxidation evoked by Fe²⁺ (Fig. 1, panel d) and also relieved the inhibitory effect of Fe²⁺ on the activity of the electrogenic sodium pump (Fig. 2, panel d). From the foregoing, it is rational to summarize that there may be a strong basis to partly conclude that Fe²⁺ evoke a deleterious effect on the activity of the pump through direct and indirect mechanisms as earlier reported [11].

Moreso, it is a common knowledge that Fe²⁺ can ultimately generate the highly reactive •OH, and that this reactive oxygen species can attack proteins and ultimately damage several amino acid residues, including histidine, tryptophan, cysteine, proline, methionine, arginine and lysine. Hence, oxidative damage to several of these amino acid residues and/or to the peptide backbone of proteins has been reported to generate carbonyl products [48,54]. Thus the activity of the enzyme can also be compromised under direct assault of •OH. Suffice to mention that the

quantification of these carbonyl products was not a component of this study and may be a subject of future research.

However, Esterbauer et al. [55] have observed that the degradation products of lipid peroxidation process can have direct effect on the activity of enzyme protein molecule. Basically, lipid peroxidation products are mainly aldehydic products and can be generally quantified by reacting them with thiobarbituric acid to give a purple/pink colour. Among the aldehydic products, malondialdehyde (MDA) have been widely reported as the key aldehyde. Therefore, the study sought to also investigate the possible effect of malondialdehyde on the activity of the cerebral pump. As presented in Fig. 5 (panel a), MDA significantly inhibited the activity of the cerebral sodium pump in a concentration dependent manner. Further study also reveals that the inhibitory effect of MDA on the activity of the pump was not protected by the dithiol (Fig. 5, panel b).

In conclusion, within the limit of our present data and following the arguments thus far, it is apparent that the low activity of the pump in degenerative diseases mediated by Fe^{2+} may involve complex multi-component mechanisms which may partly involve an initial oxidation of the critical thiols of the enzyme directly mediated by Fe^{2+} . To the best of our knowledge, this fact has not been presented in the literature. In addition, however, the transition metal may also assault membrane lipids and thus evoke an altered membrane fluidity which ultimately leads to altered structural arrangement of the cerebral sodium pump with a concomitant loss or reduced activity of the enzyme. However, during severe oxidative-stress mediated progression of such degenerative diseases; aldehydic products of lipid peroxidation such as MDA may further exacerbate deleterious effect of Fe^{2+} by a mechanism that is likely not related to the oxidation of the essential thiols of the ouabain-sensitive cerebral electrogenic pump.

Acknowledgements

This work is partly supported by TWAS/COMSTECHE research grant (Grant No. 12-268 RG/PHA/AF/AC_C - UNESCO FR: 3240270862) awarded to IJK. Also the authors acknowledged the financial support of CAPES, FINEP, FAPERGS, PRONEX and CNPq, FINEP research grant Rede Instituto Brasileiro de Neurociencia (IBN-Net) # 01.06.0842-00 and the INCT for excitotoxicity and Neuroprotection-CNPq and CNPq-ProAfrica grant awarded to IJK and JBTR.

References

- [1] H. Smith, T.G. Janz, M. Erker, Digoxin toxicity presenting as altered mental status in a patient with severe chronic obstructive lung disease, *Heart & Lung: The Journal of Acute and Critical Care* 21 (1992) 78–80.
- [2] J.R. Smith, M. Gheorghiane, S. Goldstein, The current role of digoxin in the treatment of heart failure, *Coronary Artery Disease* 4 (1) (1993) 16–26. <http://dx.doi.org/10.1097/00019501-199301000-00003> 8269181.
- [3] R.S. El-Mallakh, J.L. Barrett, R. Jed Wyatt, The Na, K-ATPase hypothesis for bipolar disorder: Implications for normal development, *Journal of Child and Adolescent Psychopharmacology* 3 (1) (1993) 37–52. <http://dx.doi.org/10.1089/cap.1993.3.37> 19630595.
- [4] P.J. Christo, R.S. El-Mallakh, Possible role of endogenous ouabain-like compounds in the pathophysiology of bipolar illness, *Medical Hypotheses* 41 (4) (1993) 378–383. [http://dx.doi.org/10.1016/0306-9877\(93\)90089-9](http://dx.doi.org/10.1016/0306-9877(93)90089-9) 8289709.
- [5] P. Biver, A. Clerico, A. Paci, S. Balzan, A. Boldrini, C. Cipolloni, Endogenous digitalis-like factors: their possible pathophysiological implications with particular regard to the perinatal period, *Child Nephrology and Urology* 10 (3) (1990) 164–180 1962738.
- [6] G.J. Lees, Inhibition of sodium-potassium-ATPase: a potentially ubiquitous mechanism contributing to central nervous system neuropathology, *Brain Research Brain Research Reviews* 16 (3) (1991) 283–300. [http://dx.doi.org/10.1016/0165-0173\(91\)90011-V](http://dx.doi.org/10.1016/0165-0173(91)90011-V) 1665097.
- [7] A.K. Nagy, C.R. Houser, A.V. Delgado-Escueta, Synaptosomal ATPase activities in temporal cortex and hippocampal formation of humans with focal epilepsy, *Brain Research* 529 (1–2) (1990) 192–201. [http://dx.doi.org/10.1016/0006-8993\(90\)90827-X](http://dx.doi.org/10.1016/0006-8993(90)90827-X) 2178027.
- [8] A.K. Nagy, Ecto-ATPases of the nervous system, in: L. Plesner, T.L. Kirly, A. F. Knowles (Eds.), *Ecto-ATPase: Recent Progress on Structure and Function*, Plenum Press, New York, 1997, pp. 1–13.
- [9] S.I. Harik, M.J. Mitchell, R.N. Kalaria, Ouabain binding in the human brain. Effects of Alzheimer's disease and aging, *Archives of Neurology* 46 (9) (1989) 951–954. <http://dx.doi.org/10.1001/archneur.1989.00520450021013> 2549926.
- [10] G. Liguri, N. Taddei, P. Nassi, S. Latorraca, C. Nediani, S. Sorbi, Changes in Na^+ , K^+ (+)-ATPase, Ca^{2+} -ATPase and some soluble enzymes related to energy metabolism in brains of patients with Alzheimer's disease, *Neuroscience Letters* 112 (2–3) (1990) 338–342. [http://dx.doi.org/10.1016/0304-3940\(90\)90227-Z](http://dx.doi.org/10.1016/0304-3940(90)90227-Z) 2163043.
- [11] H. Rauchová, Z. Drahota, J. Koudelová, The role of membrane fluidity changes and thiobarbituric acid-reactive substances production in the inhibition of cerebral cortex Na^+ , K^+ -ATPase activity, *Physiological Research* 48 (1) (1999) 73–78 10470869.
- [12] I. Jamme, E. Petit, D. Divoux, A. Gerbi, J.M. Maixent, A. Nouvelot, Modulation of mouse cerebral Na^+ , K^+ -ATPase activity by oxygen free radicals, *Neuroreport* 7 (1) (1995) 333–337 8742483.
- [13] D.B. Stanimirovic, J. Wong, R. Ball, J.P. Durkin, Free radical-induced endothelial membrane dysfunction at the site of blood-brain barrier: relationship between lipid peroxidation, Na^+ , K^+ -ATPase activity, and ^{51}Cr release, *Neurochemical Research* 20 (12) (1995) 1417–1427. <http://dx.doi.org/10.1007/BF00970589> 8789603.
- [14] A.A. Boldyrev, E.R. Bulygina, Na^+ , K^+ -ATPase and oxidative stress, *Annals of the New York Academy of Sciences* 834 (1997) 666–668. <http://dx.doi.org/10.1111/j.1749-6632.1997.tb52345.x> 9405887.
- [15] R.J. Mark, M.A. Lovell, W.R. Markesbery, K. Uchida, M.P. Mattson, A role for 4-hydroxynonenal, an aldehydic product of lipid peroxidation, in disruption of ion homeostasis and neuronal death induced by amyloid beta-peptide, *Journal of Neurochemistry* 68 (1) (1997) 255–264. <http://dx.doi.org/10.1046/j.1471-4159.1997.68010255.x> 8978733.
- [16] W.G. Siems, S.J. Hapner, F.J. Van Kuijk, 4-Hydroxynonenal inhibits Na^+ (+)- K^+ (+)-ATPase, *Free Radical Biology and Medicine* 20 (2) (1996) 215–223. [http://dx.doi.org/10.1016/0891-5849\(95\)02041-1](http://dx.doi.org/10.1016/0891-5849(95)02041-1) 8746442.
- [17] E.L. MacKenzie, K. Iwasaki, Y. Tsuji, Intracellular iron transport and storage: from molecular mechanisms to health implications, *Antioxidants & Redox Signaling* 10 (6) (2008) 997–1030. <http://dx.doi.org/10.1089/ars.2007.1893> 18327971.
- [18] B. Halliwell, Reactive oxygen species and the central nervous system, *Journal of Neurochemistry* 59 (5) (1992) 1609–1623. <http://dx.doi.org/10.1111/j.1471-4159.1992.tb10990.x> 1402908.
- [19] T. Moos, T. Rosengren Nielsen, T. Skjørringe, E.H. Morgan, Iron trafficking inside the brain, *Journal of Neurochemistry* 103 (5) (2007) 1730–1740. <http://dx.doi.org/10.1111/j.1471-4159.2007.04976.x> 17953660.
- [20] L. Zecca, M.B. Youdim, P. Riederer, J.R. Connor, R.R. Crichton, Iron, brain ageing and neurodegenerative disorders, *Nature Reviews Neuroscience* 5 (11) (2004) 863–873. <http://dx.doi.org/10.1038/nrn1537> 15496864.
- [21] T. Moos, E.H. Morgan, The metabolism of neuronal iron and its pathogenic role in neurological disease: review, *Annals of the New York Academy of Sciences* 1012 (2004) 14–26. <http://dx.doi.org/10.1196/annals.1306.002> 15105252.
- [22] Y. Ke, Z.M. Qian, Brain iron metabolism: neurobiology and neurochemistry, *Progress in Neurobiology* 83 (3) (2007) 149–173. <http://dx.doi.org/10.1016/j.pneurobio.2007.07.009> 17870230.
- [23] D. Berg, H. Hochstrasser, Iron metabolism in parkinsonian syndromes, *Movement Disorders* 21 (9) (2006) 1299–1310. <http://dx.doi.org/10.1002/mds.21020> 16817199.
- [24] D.W. Lee, J.K. Andersen, D. Kaur, Iron dysregulation and neurodegeneration: the molecular connection, *Molecular Interventions* 6 (2) (2006) 89–97. <http://dx.doi.org/10.1124/mi.6.2.6> 16565471.
- [25] N. Abo-Krysha, L. Rashed, The role of iron dysregulation in the pathogenesis of multiple sclerosis: an Egyptian study, *Multiple Sclerosis* 14 (5) (2008) 602–608. <http://dx.doi.org/10.1177/1352458507085550> 18408021.
- [26] K.J. Barnham, A.I. Bush, Metals in Alzheimer's and Parkinson's diseases, *Current Opinion in Chemical Biology* 12 (2) (2008) 222–228. <http://dx.doi.org/10.1016/j.cbpa.2008.02.019> 18342639.
- [27] T. LaVaute, S. Smith, S. Cooperman, K. Iwai, W. Land, E. Meyron-Holtz, S. K. Drake, G. Miller, M. Abu-Asab, M. Tsokos, R. Switzer, A. Grinberg, P. Love, N. Tresser, T.A. Rouault, Targeted deletion of the gene encoding iron regulatory protein-2 causes misregulation of iron metabolism and neurodegenerative disease in mice, *Nature Genetics* 27 (2) (2001) 209–214. <http://dx.doi.org/10.1038/84859> 11175792.
- [28] T.A. Rouault, The role of iron regulatory proteins in mammalian iron homeostasis and disease, *Nature Chemical Biology* 2 (8) (2006) 406–414. <http://dx.doi.org/10.1038/nchembio807> 16850017.
- [29] D.R. Crapper McLachlan, A.J. Dalton, T.P. Kruck, M.Y. Bell, W.L. Smith, W. Kalow, D.F. Andrews, Intramuscular desferrioxamine in patients with Alzheimer's disease, *Lancet* 337 (8753) (1991) 1304–1308. [http://dx.doi.org/10.1016/0140-6736\(91\)92978-B](http://dx.doi.org/10.1016/0140-6736(91)92978-B) 1674295.
- [30] L. Benvenisti-Zarom, J. Chen, R.F. Regan, The oxidative neurotoxicity of clioquinol, *Neuropharmacology* 49 (5) (2005) 687–694. <http://dx.doi.org/10.1016/j.neuropharm.2005.04.023> 15992834.
- [31] D. Kaur, F. Yantiri, S. Rajagopalan, J. Kumar, J.Q. Mo, R. Boonplueang, V. Viswanath, R. Jacobs, L. Yang, M.F. Beal, D. DiMonte, I. Volitaskis, L. Ellerby, R.A. Cherny, A.I. Bush, J.K. Andersen, Genetic or pharmacological iron chelation

- prevents MPTP-induced neurotoxicity in vivo: a novel therapy for Parkinson's disease, *Neuron* 37 (6) (2003) 899–909. [http://dx.doi.org/10.1016/S0896-6273\(03\)00126-0](http://dx.doi.org/10.1016/S0896-6273(03)00126-0) 12670420.
- [32] J. Salazar, N. Mena, S. Hunot, A. Prigent, D. Alvarez-Fischer, M. Arredondo, C. Duyckaerts, V. Sazdovitch, L. Zhao, L.M. Garrick, M.T. Nuñez, M.D. Garrick, R. Raisman-Vozari, E.C. Hirsch, Divalent metal transporter 1 (DMT1) contributes to neurodegeneration in animal models of Parkinson's disease, *Proceedings of the National Academy of Sciences of the United States of America* 105 (47) (2008) 18578–18583. <http://dx.doi.org/10.1073/pnas.0804373105> 19011085.
- [33] S.L. Clardy, X. Wang, P.J. Boyer, C.J. Earley, R.P. Allen, J.R. Connor, Is ferroportin–hepcidin signaling altered in restless legs syndrome? *Journal of the Neurological Sciences* 247 (2) (2006) 173–179. <http://dx.doi.org/10.1016/j.jns.2006.04.008> 16759669.
- [34] Y. Hua, R.F. Keep, J.T. Hoff, G. Xi, Brain injury after intracerebral hemorrhage: the role of thrombin and iron, *Stroke* 38 (2 Suppl) (2007) 759–762. <http://dx.doi.org/10.1161/01.STR.0000247868.97078.10> 17261733.
- [35] C.H. Fiske, Y.J. Subbarow, *The colorimetric determination of phosphorus*, *Journal of Biological Chemistry* 66 (1925) 375–381.
- [36] O.H. Lowry, N.J. Rosebrough, A.L. Farr, R.J. Randall, Protein measurement with the folin phenol reagent, *Journal of Biological Chemistry* 193 (1) (1951) 265–275. <http://dx.doi.org/10.1016/j.jbc.1951.03.031> 14907713.
- [37] H. Ohkawa, N. Ohishi, K. Yagi, Assay for lipid peroxides in animal tissues by thiobarbituric acid reaction, *Analytical Biochemistry* 95 (2) (1979) 351–358. [http://dx.doi.org/10.1016/0003-2697\(79\)90738-3](http://dx.doi.org/10.1016/0003-2697(79)90738-3) 36810.
- [38] G.L. Ellman, Tissue sulfhydryl groups, *Archives of Biochemistry and Biophysics* 82 (1) (1959) 70–77. [http://dx.doi.org/10.1016/0003-9861\(59\)90090-6](http://dx.doi.org/10.1016/0003-9861(59)90090-6) 13650640.
- [39] I.J. Kade, C.W. Nogueira, J.B. Rocha, Diphenyl diselenide and streptozotocin did not alter cerebral glutamatergic and cholinergic systems but modulate antioxidant status and sodium pump in diabetic rats, *Brain Research* 1284 (2009) 202–211. <http://dx.doi.org/10.1016/j.brainres.2009.04.003> 19362073.
- [40] I.J. Kade, J.B. Rocha, Gallic acid modulates cerebral oxidative stress conditions and activities of enzyme-dependent signaling systems in streptozotocin-treated rats, *Neurochemical Research* 38 (4) (2013) 761–771. <http://dx.doi.org/10.1007/s11064-013-0975-6> 23381106.
- [41] D.A. Greene, S.A. Lattimer, A.A. Sima, Sorbitol, phosphoinositides, and sodium-potassium-ATPase in the pathogenesis of diabetic complications, *New England Journal of Medicine* 316 (10) (1987) 599–606. <http://dx.doi.org/10.1056/NEJM198703053161007> 3027558.
- [42] A.I. Winegrad, Banting lecture 1986. Does a common mechanism induce the diverse complications of diabetes? *Diabetes* 36 (3) (1987) 396–406. <http://dx.doi.org/10.2337/diab.36.3.396> 3026877.
- [43] E. Amler, R. Jasińska, Z. Drahota, J. Zborowski, Membrane lateral pressure as a modulator of glycerol-3-phosphate dehydrogenase activity, *FEBS Letters* 271 (1–2) (1990) 165–168. [http://dx.doi.org/10.1016/0014-5793\(90\)80398-3](http://dx.doi.org/10.1016/0014-5793(90)80398-3) 2226800.
- [44] B.P. Yu, E.A. Suescun, S.Y. Yang, Effect of age-related lipid peroxidation on membrane fluidity and phospholipase A₂: modulation by dietary restriction, *Mechanisms of Ageing and Development* 65 (1) (1992) 17–33. [http://dx.doi.org/10.1016/0047-6374\(92\)90123-U](http://dx.doi.org/10.1016/0047-6374(92)90123-U) 1405788.
- [45] G. Minotti, S.D. Aust, Redox cycling of iron and lipid peroxidation, *Lipids* 27 (3) (1992) 219–226. <http://dx.doi.org/10.1007/BF02536182> 1326072.
- [46] J.G. Goddard, G.D. Sweeney, Delayed, ferrous iron-dependent peroxidation of rat liver microsomes, *Archives of Biochemistry and Biophysics* 259 (2) (1987) 372–381. [http://dx.doi.org/10.1016/0003-9861\(87\)90503-0](http://dx.doi.org/10.1016/0003-9861(87)90503-0) 3426233.
- [47] F. Ursini, M. Maiorino, P. Hochstein, L. Ernster, Microsomal lipid peroxidation: mechanisms of initiation. The role of iron and iron chelators, *Free Radical Biology and Medicine* 6 (1) (1989) 31–36. [http://dx.doi.org/10.1016/0891-5849\(89\)90156-1](http://dx.doi.org/10.1016/0891-5849(89)90156-1) 2492247.
- [48] I.J. Kade, M.W. Paixão, O.E. Rodrigues, N.B. Barbosa, A.L. Braga, D.S. Ávila, C. W. Nogueira, J.B. Rocha, Comparative studies on dicholesteroyl diselenide and diphenyl diselenide as antioxidant agents and their effect on the activities of Na⁺/K⁺ ATPase and delta-aminolevulinic acid dehydratase in the rat brain, *Neurochemical Research* 33 (1) (2008) 167–178. <http://dx.doi.org/10.1007/s11064-007-9432-8> 17710541.
- [49] I.J. Kade, J.B. Rocha, Comparative study on the influence of subcutaneous administration of diphenyl and dicholesteroyl diselenides on sulphhydryl proteins and antioxidant parameters in mice, *Journal of Applied Toxicology* 30 (7) (2010) 688–693. <http://dx.doi.org/10.1002/jat.1542> 20583318.
- [50] N.B. Barbosa, J.B. Rocha, D.C. Wondracek, J. Perottoni, G. Zeni, C.W. Nogueira, Diphenyl diselenide reduces temporarily hyperglycemia: possible relationship with oxidative stress, *Chemico-Biological Interactions* 163 (3) (2006) 230–238. <http://dx.doi.org/10.1016/j.cbi.2006.08.004> 16965767.
- [51] V.C. Borges, J.B. Rocha, C.W. Nogueira, Effect of diphenyl diselenide, diphenyl ditelluride and ebselen on cerebral Na⁺, K⁺ ATPase activity in rats, *Toxicology* 215 (3) (2005) 191–197. <http://dx.doi.org/10.1016/j.tox.2005.07.002> 16095793.
- [52] T.I. Omotayo, J.B. Rocha, E.O. Ibukun, I.J. Kade, Inorganic mercury interacts with thiols at the nucleotide and cationic binding sites of the ouabain-sensitive cerebral electrogenic sodium pump, *Neurochemistry International* 58 (7) (2011) 776–784. <http://dx.doi.org/10.1016/j.neuint.2011.03.002> 21397648.
- [53] I.J. Kade, T.I. Omotayo, J.B.T. Rocha, Threshold protective effect of glutathione on inorganic mercury inhibition of cerebral sodium pump, *Biochemistry: An Indian Journal* 6 (2) (2012) 65–70.
- [54] A. Amici, R.L. Levine, L. Tsai, E.R. Stadtman, Conversion of amino acid residues in proteins and amino acid homopolymers to carbonyl derivatives by metal-catalyzed oxidation reactions, *Journal of Biological Chemistry* 264 (6) (1989) 3341–3346. [http://dx.doi.org/10.1016/0891-5849\(91\)90192-6](http://dx.doi.org/10.1016/0891-5849(91)90192-6) 1937131.
- [55] H. Esterbauer, R.J. Schaur, H. Zollner, Chemistry and biochemistry of 4-hydroxynonenal, malonaldehyde and related aldehydes, *Free Radical Biology and Medicine* 11 (1) (1991) 81–128. [http://dx.doi.org/10.1016/0891-5849\(91\)90192-6](http://dx.doi.org/10.1016/0891-5849(91)90192-6) 1937131.
- [56] S.J. Klebanoff, J.I. Gally, I.M. Goldstein, R. Snyderman (Eds.), *Oxygen metabolites from phagocytes*, Raven Press, New York, NY, 1992, pp. 541–588.
- [57] E. Graf, J.R. Mahoney, R.G. Bryant, J.W. Eaton, Iron catalyzed hydroxyl radical formation: Stringent requirement for free iron coordination site, *Journal of Biological Chemistry* 259 (1984) 3620–3624.