

Original Article

Distribution patterns of flavonoids from three *Momordica* species by ultra-high performance liquid chromatography quadrupole time of flight mass spectrometry: a metabolomic profiling approach



Ntakadzeni Edwin Madala^{a,*}, Lizelle Piater^a, Ian Dubery^a, Paul Steenkamp^{a,b}

^a Department of Biochemistry, University of Johannesburg, Auckland Park, South Africa

^b CSIR Biosciences, Natural Products and Agroprocessing Group, Pretoria, South Africa

ARTICLE INFO

Article history:

Received 15 January 2016

Accepted 14 March 2016

Available online 20 May 2016

Keywords:

Momordica

UHPLC-qTOF-MS

Flavonoids

Principal component analysis

Chemotaxonomy

Hierarchical cluster analysis

ABSTRACT

Plants from the *Momordica* genus, Cucurbitaceae, are used for several purposes, especially for their nutritional and medicinal properties. Commonly known as bitter gourds, melon and cucumber, these plants are characterized by a bitter taste owing to the large content of cucurbitacin compounds. However, several reports have shown an undisputed correlation between the therapeutic activities and polyphenolic flavonoid content. Using ultra-high performance liquid chromatography quadrupole time of flight mass spectrometry in combination with multivariate data models such as principal component analysis and hierarchical cluster analysis, three *Momordica* species (*M. foetida* Schumach., *M. charantia* L. and *M. balsamina* L.) were chemo-taxonomically grouped based on their flavonoid content. Using a conventional mass spectrometric-based approach, thirteen flavonoids were tentatively identified and the three species were found to contain different isomers of the quercetin-, kaempferol- and isorhamnetin-O-glycosides. Our results indicate that *Momordica* species are overall very rich sources of flavonoids but do contain different forms thereof. Furthermore, to the best of our knowledge, this is a first report on the flavonoid content of *M. balsamina* L.

© 2016 Sociedade Brasileira de Farmacognosia. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Momordica species are versatile plants belonging to the family Cucurbitaceae and are often referred to by various common names such as bitter gourd, bitter cucumber and bitter melon (Nagarani et al., 2014a). In different parts of the world, *Momordica* plants are consumed as a vegetable and are known for their bitter taste due to the presence of phytochemicals such as alkaloids and cucurbitacins (Chen et al., 2005; Rios et al., 2005; Nagarani et al., 2014a). Within this genus there are several species widely distributed across the globe, mainly in the tropical and subtropical regions of Africa, Asia and Australia. In a recent review article by Nagarani et al. (2014a), undisputed scientific evidence on the origin of these plants was presented and believed to be endemic to India, while earlier biogeographical origins of these species can be found elsewhere (Dey et al., 2006; Singh et al., 2007; Gaikwad et al., 2008).

Apart from the nutritious value, *Momordica* species are also used for their medicinal properties. For example, *M. charantia* has

been used in Chinese folk medicine for the treatment of different chronic diseases (Zhang, 1992). In some parts of South Africa, these species are currently used as medication for sugar diabetes and chronic hypertension diseases, without any scientific backing. Due to the wide range of phytochemicals in *Momordica* (Nagarani et al., 2014a), it is very difficult to point out one active compound, even though numerous metabolites with known pharmacological activities have been identified (Singh et al., 2011; Kenny et al., 2013; Nagarani et al., 2014b). *Momordica* species are known to contain large quantities of polyphenolic compounds (Kubola and Siriamornpun, 2008) and amongst these are flavonoids which are known to possess several therapeutic activities. Further reports have suggested a possible link between the medicinal properties of *Momordica* species and their flavonoid content (Horax et al., 2005; Lin and Tang, 2007; Wu and Ng, 2008; Zhu et al., 2012). The distribution of flavonoids in these plants represents another interesting dimension. For instance, Kenny et al. (2013) reported the presence of flavonoids in the bitter melon (*M. charantia*) fruit, however, the levels vary across different sections of the fruit. In a separate study, Nagarani et al. (2014b) reported a very interesting distribution pattern across different species of *Momordica* (*M. tuberosa*, *M. charantia* and *M. cochinchinensis*). Here, the flavonoid rutin was detected in

* Corresponding author.

E-mail: emadala@uj.ac.za (N.E. Madala).

the former two species but absent in the later. As such, the aim of the current study was to investigate the flavonoid distribution patterns within the three *Momordica* species (namely *M. charantia* L., *M. foetida* Schumach. and *M. balsamina* L.) currently used for diabetic control and nutritional value in the Limpopo (Northern) regions of South Africa.

In addition to their nutritional and medicinal properties, plant metabolites can also be used to taxonomically classify plants to ensure that the correct specie(s) are used for medicinal purposes. In the past, flavonoids have been used for chemo-taxonomical classification of plants (Iwashina, 2000; El Shabrawy et al., 2014; Martucci et al., 2014). To achieve our objective, the current study was divided into two parts: the first aiming to establish a chemo-taxonomical relationship between the three species using a metabolomic profiling approach with the aid of UHPLC–qTOF–MS and multivariate data models, and in the second, the flavonoid composition of the three species was investigated using targeted MS-based flavonoid identification strategies presented elsewhere (Cuyckens and Claeys, 2004).

Materials and methods

Plant materials and chemicals

Momordica plants, Cucurbitaceae, were collected in and around the Venda region of South Africa with the help of the local farmers. Briefly, *M. charantia* L. was collected from a farm in the Nwanedi farming area, about 80 km south of the Zimbabwean border. The other two species, *M. balsamina* L. and *M. foetida* Schumach. were collected from various villages around Thohoyandou. The species were identified with the help of Mr Philip Ramela (Madzivhandila College of Agriculture, South Africa) and for further confirmation, the plant materials were also compared to the national herbarium specimens at the South African National Biodiversity Institute (SANBI) (Pretoria, South Africa). Voucher herbarium specimens (with voucher number NEM003 (*M. balsamina*), NEM004 (*M. charantia*) and NEM005 (*M. foetida*)) were prepared and deposited to the Department of Botany, University of Johannesburg. Unless stated otherwise, all the chemicals were of analytical grade and obtained from various internationally reputable suppliers. Both the methanol and acetonitrile (Romil, MicroSep, South Africa) were used for UHPLC–qTOF–MS analyses. Water was purified with a Milli-Q Gradient A10 system (Millipore, Billerica, MA, USA). Leucine encephalin, rutin and formic acid were purchased from Sigma Aldrich, Germany. Quercetin-3-glucose, quercetin-4'-glucose and quercetin-7-glucose were purchased from Phytolab (Vestenbergsgreuth, Germany).

Metabolite extraction

The leaves of the three *Momordica* species were air-dried at 37 °C for three consecutive days. Metabolites were extracted from the four independently crushed leaf samples (2 g), representing four independent biological replicates, using 80% aqueous methanol (20 ml). For maximum extraction, the homogenate was placed on an orbital shaker at room temperature (25 °C) for at least 30 min. After the extraction, the tissue debris was removed by centrifugation at 5000 × g for 10 min. The supernatant was dried to at least 1 ml using a rotary evaporator operating at 55 °C under negative pressure vacuum. The 1 ml extract was subsequently dried to completeness using a vacuum concentrator centrifuge (Vacufuge, Eppendorf, Germany) operating at 55 °C. Prior to UHPLC–MS analyses, the pelleted extract residues were re-constituted in 1 ml of 50% aqueous methanol and filtered through 0.22 μm nylon filters.

UHPLC–MS analyses

For UHPLC–MS analyses, a previously described method (Madala et al., 2014a) was used. Briefly, one (1) μl of the extracts was analysed on a Waters Acquity BEH C8 column (150 mm × 2.1 mm, 1.7 μm particle size) and the temperature controlled at 60 °C. Here, three technical replicates were analyzed and randomized during the UHPLC–MS analyses using online randomizing software (www.random.org/lists/) to avoid technical bias. A binary solvent mixture was used consisting of water containing 10 mM formic acid (pH 2.3) (eluent A) and acetonitrile containing 10 mM formic acid (eluent B). The initial conditions were 98% A at a flow rate of 0.4 ml/min and maintained for 1 min, followed by multiple gradients to 5% A at 26 min. The conditions were kept constant for 1 min and then changed to the initial conditions. The total chromatographic run time was 30 min. Chromatographic elution was monitored with the aid of a photo diode array (PDA) detector and MS.

For MS detection, a high resolution mass spectrometer (Waters SYNAPT G1 Q-TOF system), operating in V-optics and electrospray negative mode, was used. Leucine enkephalin (50 pg/ml) was used as reference lockmass calibrant to obtain typical mass accuracies between 1 and 5 mDa. The optimal conditions for analysis were as follows: capillary voltage of 2.5 kV, the sampling cone at 30 V and the extraction cone at 4 V. The scan time was 0.1 s covering the 100–1000 Da mass range. The source temperature was 120 °C and the desolvation temperature was set at 450 °C. Nitrogen was used as the nebulization gas at a flow rate of 700 l/h. To obtain better metabolite coverage and fragmentation patterns thereof, the MS was operated at different collision energy (CE) levels as reported elsewhere (Madala et al., 2012). For comparison purposes, authentic standards (quercetin-3-glucose, quercetin-4'-glucose and quercetin-7-glucose) were also analyzed using the same conditions. All the acquisition and analysis of data were controlled by Waters MassLynx™ v4.1 software (SCN 704).

Multivariate data analyses

Primary raw data was analyzed by data alignment, peak finding, peak integration and retention time (Rt) correction using a Markerlynx XS™ software (Waters Corporation, Milford, USA) with the following processing parameters: Rt range of 7–12 min, mass range of 100–1000 Da, mass tolerance of 0.02 Da, Rt window of 0.2 min. The resulting datasets were exported to the SIMCA-P software version 13.0 (Umetrics, Umea, Sweden) for Principal Component Analysis (PCA) and Hierarchical Cluster Analysis (HCA). Before the models were computed, all the data were mean centered and Pareto-scaled. For HCA analysis, the Ward distance algorithm was used to calculate the distance between the different generated clusters (Madala et al., 2014b).

Results and discussion

Classification of *Momordica* species based on their flavonoid content

Analyses of the crude aqueous-methanol extracts prepared from the leaves of the three *Momordica* species were conducted using an UHPLC–qTOF–MS operating in negative electrospray ionization (ESI) mode. The data obtained was automatically processed by MarkerLnx™ software targeting the flavonoid region (7–12 min) of the chromatograms (Fig. 1). The resulting files were further exported to the SIMCA-P version 13 software for multivariate data analyses. The resulting PCA and HCA are shown in Fig. 2. In the past it was noted that there seems to exist a tendency amongst




Fig. 1. Selected ion chromatograms (XIC) showing relative chromatographic elution rates of different flavonoid isomers in the three *Momordica* species. The XIC were extracted using the aglycone m/z of 301, 285 and 315 representing quercetin, kaempferol and isorhamnetin.

taxonomically-related plants to produce very similar phytochemicals and, as such, flavonoids have been used for chemo-taxonomical classification (El Shabrawy et al., 2014; Martucci et al., 2014). To evaluate the taxonomical connection/relationship between the three species, PCA was performed and the score plot showed that

the three species are distinct from one another (Fig. 2A). However, PCA models are known to be less superior in showing sample relationships (Madala et al., 2014b). To overcome this challenge, HCA results (Fig. 2B) were evaluated and it was seen that *M. foetida* forms a distinctive group whilst the other two species





Fig. 2. Principal component analysis (PCA) score plots (A) and hierarchical cluster analysis (HCA) showing separation of samples representing the three *Momordica* species based on their phytochemical content.



Aglycone	R ¹	R ²	R ³	[Y ₀ -H] ⁻ / Y ₀ ⁻
Quercetin	H	H	OH	300/301
Kaempferol	H	H	H	284/285
Isorhamnetin	H	H	OCH ₃	314/315

Scheme 1. The structure of common flavonoid aglycones and characteristic fragments ions ([Y₀-H]⁻ and Y₀⁻) formed during homolytic and heterolytic sugar cleavage, respectively.


form a very close clade. Overall, these results suggest a relatively close relationship between *M. balsamina* and *M. charantia*, but these two (as a group/clade) have a distance relationship with *M. foetida*.

General approach for identification of flavonoids

From UHPLC-qTOF-MS base ion peak (BIP) chromatograms, ions representing typical flavonoid aglycone fragments were selected and used to generate the single ion chromatograms (Fig. 1). From the MS spectra, molecular formulae of the pseudo-molecular ions ([M-H]⁻) representing quercetin (*m/z* 300.021/301.029), kaempferol (*m/z* 284.025/285.040) and isorhamnetin (*m/z* 315.045) were generated and selected based on the criterion that the mass difference between the measured and calculated mass is below 5 ppm. The generated molecular formulae were further used for compound identification searches using

the Dictionary of Natural Products (DNP) online database (<http://dnp.chemnetbase.com/>) and the KNapSack database (http://kanaya.naist.jp/KNAPSACK_jsp/top.html). Mass spectra generated at different CEs were also used to elucidate the position and number of sugar molecules attached to the aglycone skeleton. The proposed identities of the flavonoids and their respective MS fragmentation patterns were also compared to literature (Cuyckens and Claeys, 2004; Gobbo-Neto et al., 2008).

Structurally, flavonoids are polyphenolic compounds with a nuclear structure base of C₆-C₃-C₆ (Scheme 1, Cuyckens and Claeys, 2004; Tan et al., 2014). These compounds exist as either aglycones or remain glycosylated with different sugar moieties. It is also worth mentioning that sugar attachment on the flavonoid aglycone moiety can happen at different positions and, as such, deducing the sugar position becomes a difficult analytical task. This adds to the structural complexity which makes flavonoids difficult to identify. However, it has been reported that during



[M-H] ⁻	Rt (min)	R ¹	R ²	R ³	<i>M. balsamina</i>	<i>M. charantia</i>	<i>M. foetida</i>	Identity	Molecular formula (< 5 ppm)
625	7.2	gly	gly	OH	x	x	x	quercetin-3,7- <i>O</i> -diglucopyranoside	C ₂₇ H ₃₀ O ₁₇
595	7.46	di-gly	H	OH	x	x	x	quercetin 3- <i>O</i> -sambubioside	C ₂₆ H ₂₈ O ₁₆
609	7.83	di-gly	H	OH	x	x	x	quercetin 3- <i>O</i> -rutinoside	C ₂₇ H ₃₀ O ₁₆
579	8.12	di-gly	H	OH	x	x	x	quercetin-3- <i>O</i> -arabinoside-rhamnopyranoside	C ₂₆ H ₂₇ O ₁₅
609	8.3	gly	gly	OCH ₃	x			isorhamnetin-3- <i>O</i> -glucoside-7- <i>O</i> -arabinopyranoside	C ₂₇ H ₃₀ O ₁₆
593	8.48	di-gly	H	OH	x	x	x	quercetin-3,7-di-rhamnopyranoside	C ₂₇ H ₃₀ O ₁₅
593	8.56	di-gly	H	H	x	x		kaempferol-3- <i>O</i> -rutinoside	C ₂₇ H ₃₀ O ₁₅
447	8.85	gly	H	H	x	x	x	kaempferol-3- <i>O</i> -glucoside	C ₂₁ H ₂₀ O ₁₁
533	9.32	H	gly	H	x		x	kaempferol-7- <i>O</i> -malonylglucoside	C ₂₄ H ₂₂ O ₁₄
533	9.5	gly	H	H			x	kaempferol-3- <i>O</i> -malonylglucoside	C ₂₄ H ₂₂ O ₁₄
477	9.6	gly	H	OCH ₃	x	x		isorhamnetin 3- <i>O</i> -glucoside	C ₂₂ H ₂₂ O ₁₂
533	9.87	H	H	gly			x	kaempferol-4'- <i>O</i> -malonylglucoside	C ₂₄ H ₂₂ O ₁₄
489	10.01	gly	H	H			x	kaempferol-3- <i>O</i> -acetylglucoside	C ₂₃ H ₂₂ O ₁₂

Fig. 3. Identities of thirteen flavonoid isomers detected in the *Momordica* species.




Fig. 4. Mass spectra generated at different collision energy levels showing the respective fragmentation patterns of quercetin (A), kaempferol (B) and isorhamnetin (C) containing flavonoids.

negative mode ESI, deprotonated flavonoid-*O*-glycosides $[M-H]^-$ ions undergo a series of fragmentation stages during MS analyses. This fragmentation is typically characterized by the immediate loss of sugar which can happen in two different ways. As shown

in **Scheme 1**, removal of the sugar moiety can either happen through heterolytic cleavage accompanied by proton migration or homolytic sugar cleavage to produce an aglycone product ion $Y_0^{\bullet-}$ or an aglycone radical ion $[Y_0-H]^-$ respectively (Zhou et al.,

2014). The ratio of $[Y_0-H]^{\bullet-}/Y_0^{\bullet-}$ has been used with some success for the determination of the sugar position. For instance, Geng et al. (2009) could positively distinguish the different isomers of quercetin based on the sugar position using three different mass analyzers. However, these observations were found to be instrument (MS analyzer) specific. As such, in the current study, the proposal made by Geng et al. (2009) was re-affirmed with our Q-TOF-MS using three positional isomers of quercetin glycosides, namely quercetin-3-glucose, quercetin-4'-glucose and quercetin-7-glucose. Our results (supplementary file 1) suggest that our Q-TOF-MS instrument is also capable of showing the sugar position on the aglycone flavonoid moieties. Furthermore, it is known that flavonoids can be glycosylated at different positions with two or more sugars attaching in the same or different positions (Zhou et al., 2014). Here, at moderate CE levels, molecules showing an abundant $[Y_0-H]^{\bullet-}$ were identified as those glycosylated at 3' position and those with an intense $Y_0^{\bullet-}$ were regarded as those glycosylated at 7' or 4' position (Geng et al., 2009). Furthermore, molecules exhibiting equal intensities of both radical aglycones ($[Y_0-H]^{\bullet-}/Y_0^{\bullet-}$) were regarded as those glycosylated in the two possible glycosylation sites (Geng et al., 2009; Ablajan and Tuoheti, 2013; Zhou et al., 2014). Therefore, with the aid of the above information and thorough visual inspection of the mass spectra, the flavonoid composition of the three *Momordica* species could be precisely determined (Fig. 3).

Identification of quercetin flavonoids

For flavonoids containing the quercetin aglycone, m/z 300.021/301.029 was used to generate the extracted ion chromatograms representing possible quercetin-bearing molecules (Fig. 1). By comparing the MS chromatograms generated using a CE of 3 eV and that of 30 eV, at least five quercetin flavonoid isomers were positively identified from all three plant species (Fig. 3). However, not all three species were found to contain all the isomers. More interestingly, the higher CE (30 eV) MS chromatograms were capable of showing the position of the sugar more efficiently than the lower CE (3 eV) (Fig. 4A–C). This observation is consistent with previously published data (Geng et al., 2009) where it was shown that $[Y_0-H]^{\bullet-}/Y_0^{\bullet-}$ ratio increases concomitant with an increase in CE.

Identification of kaempferol flavonoids

Similarly, kaempferol-bearing flavonoids were also identified using the approach as described for quercetin flavonoids. However, m/z 284.025 ($[Y_0-H]^{\bullet-}$) and 285.040 ($Y_0^{\bullet-}$), representing the deprotonated forms of kaempferol aglycones, were used to generate the extracted ion chromatogram (Fig. 1) showing the different retention times of kaempferol flavonoids isomers. At least six kaempferol-containing flavonoids were positively identified between the three species, with *M. foetida* containing large quantities of these compounds, in comparison to the other two species (Fig. 3).

Identification of isorhamnetin flavonoids

In comparison to the former two types of flavonoids, the isorhamnetin class was found to be the least abundant in the three *Momordica* species investigated. Here, only two different isomers (Fig. 3) were identified, with the *M. balsamina* having three isomers and *M. foetida* only one. Surprisingly, the *M. charantia* species was found to not contain any isorhamnetin-bearing flavonoids. It is these differences in the flavonoid content which makes the current study important because it reveals interesting underlying biochemical differences between the these species which, in part, could

possibly be used for taxonomical and other biological classification systems.

Conclusion

While some reports on the flavonoids from *M. charantia* and *M. foetida* exist, to the best of our knowledge this is the first report on the flavonoid composition of *M. balsamina*. The results suggest that the three species are chemo-taxonomically related and contain very similar flavonoid compositions. From the results it can also be seen that *M. charantia* and *M. balsamina* are more closely related to each other and, as a group, are distantly related to *M. foetida*. Above all, the results of the current study confirm the *Momordica* species as a rich source of structurally diverse flavonoids. The results also reaffirm the use of LC-MS in combination with multivariate data models to be a feasible approach to study metabolite distribution patterns between closely related plant species.

Authors' contributions

NEM, IAD and LAP planned the study. NEM collected the plants and extracted the metabolites. NEM, LAP, IAD and PAS conducted the UHPLC-qTOF-MS, multivariate data models and executed metabolite identification. All authors read and approved the final manuscript.

Conflicts of interest

The authors declare no conflicts of interest.

Acknowledgements

The authors would like to thank the University of Johannesburg and the NRF for financial support. Mr Muade is thanked for the donation of *M. charantia* plants.

Appendix A. Supplementary data

Supplementary material related to this article can be found, in the online version, at doi:10.1016/j.bjp.2016.03.009.

References

- Ablajan, K., Tuoheti, A., 2013. Fragmentation characteristics and isomeric differentiation of flavonol O-rhamnosides using negative ion electrospray ionization tandem mass spectrometry. *Rapid Commun. Mass Spectrom.* 27, 451–460.
- Chen, J., Chiu, M., Nie, R., Cordell, G., Qiu, S., 2005. Cucurbitacins and cucurbitane glycosides: structures and biological activities. *Nat. Prod. Rep.* 22, 386–399.
- Cuyckens, F., Claeys, M., 2004. Mass spectrometry in the structural analysis of flavonoids. *J. Mass Spectrom.* 39, 1–15.
- Dey, S.S., Singh, A.K., Chandel, D., Behera, T.K., 2006. Genetic diversity of bitter gourd *Momordica charantia* genotypes revealed by RAPD markers and agronomic traits. *Sci. Hortic.* 109, 21–28.
- El Shabrawy, M.O., Hosni, H.A., El Garf, I.A., Marzouk, M.M., Kawashty, S.A., Saleh, N.A., 2014. Flavonoids from *Allium myrianthu* Boiss. *Biochem. Syst. Ecol.* 56, 125–128.
- Gaikwad, A.B., Behera, T.K., Singh, A.K., Chandel, D., Karihaloo, J.L., Staub, J.E., 2008. Amplified fragment length polymorphism analysis provides strategies for improvement of bitter gourd (*Momordica charantia* L.). *HortScience* 43, 127–133.
- Geng, P., Sun, J., Zhang, R., He, J., Abliz, Z., 2009. An investigation of the fragmentation differences of isomeric flavonol-O-glycosides under different collision-induced dissociation based mass spectrometry. *Rapid Commun. Mass Spectrom.* 23, 1519–1524.
- Gobbo-Neto, L., Gates, P.J., Lopes, N.P., 2008. Negative ion 'chip-based' nanospray tandem mass spectrometry for the analysis of flavonoids in glandular trichomes of *Lychnophora ericoides* Mart. (Asteraceae). *Rapid Commun. Mass Spectrom.* 22, 3802–3808.
- Horax, R., Hettiarachchy, N., Islam, S., 2005. Total phenolic contents and phenolic acid constituents in 4 varieties of bitter melons (*Momordica charantia*) and antioxidant activities of their extracts. *J. Food Sci.* 70, C275–C280.
- Iwashina, T., 2000. The structure and distribution of the flavonoids in plants. *J. Plant Res.* 113, 287–299.

- Kenny, O., Smyth, T.J., Hewage, C.M., Brunton, N.P., 2013. Antioxidant properties and quantitative UPLC–MS analysis of phenolic compounds from extracts of fenugreek (*Trigonella foenum-graecum*) seeds and bitter melon (*Momordica charantia*) fruit. *Food Chem.* 141, 4295–4302.
- Kubola, J., Siriamornpun, S., 2008. Phenolic contents and antioxidant activities of bitter melon (*Momordica charantia* L.) leaf, stem and fruit fraction extracts in vitro. *Food Chem.* 110, 881–890.
- Lin, J.Y., Tang, C.Y., 2007. Determination of total phenolic and flavonoid contents in selected fruits and vegetables, as well as their stimulatory effects on mouse splenocyte proliferation. *Food Chem.* 101, 140–147.
- Martucci, M.E.P., De Vos, R.C., Carollo, C.A., Gobbo-Neto, L., 2014. Metabolomics as a potential chemotaxonomical tool: application in the genus *Vernonia* Schreb. *PLoS ONE* 9, e93149.
- Madala, N.E., Steenkamp, P.A., Piater, L.A., Dubery, I.A., 2012. Collision energy alteration during mass spectrometric acquisition is essential to ensure unbiased metabolomic analysis. *Anal. Bioanal. Chem.* 404, 367–372.
- Madala, N.E., Tugizimana, F., Steenkamp, P.A., 2014a. Development and optimization of an UPLC–QTOF–MS/MS method based on an in-source collision induced dissociation approach for comprehensive discrimination of chlorogenic acids isomers from *Momordica* plant species. *J. Anal. Met. Chem.*, <http://dx.doi.org/10.1155/2014/650879>.
- Madala, N.E., Piater, L.A., Steenkamp, P.A., Dubery, I.A., 2014b. Multivariate statistical models of metabolomic data reveals different metabolite distribution patterns in isonitrosoacetophenone-elicited *Nicotiana tabacum* and *Sorghum bicolor* cells. SpringerPlus, <http://dx.doi.org/10.1186/2193-1801-3-254>.
- Nagarani, G., Abirami, A., Siddhuraju, P., 2014a. Food prospects and nutraceutical attributes of *Momordica* species: a potential tropical bioresources – a review. *Food Sci. Hum. Wellness* 3, 117–126.
- Nagarani, G., Abirami, A., Siddhuraju, P., 2014b. A comparative study on antioxidant potentials, inhibitory activities against key enzymes related to metabolic syndrome, and anti-inflammatory activity of leaf extract from different *Momordica* species. *Food Sci. Hum. Wellness* 3, 36–46.
- Rios, J., Escandell, J., Recio, M., 2005. New insights into bioactivity of cucurbitacins. In: Atta-ur-Rahman (Ed.), *Studies in Natural Products Chemistry*. Amsterdam, pp. 429–469.
- Singh, A.K., Behera, T.K., Chandel, D., Sharma, P., Singh, N.K., 2007. Assessing genetic relationships among bitter melon (*Momordica charantia* L.) accessions using inter-simple sequence repeat (ISSR) markers. *J. Hortic. Sci. Biotechnol.* 82, 217–222.
- Singh, J., Cumming, E., Manoharan, G., Kalasz, H., Adeghate, E., 2011. Suppl 2: Medicinal chemistry of the anti-diabetic effects of *Momordica charantia*: active constituents and modes of actions. *Open Med. Chem. J.* 5, 70.
- Tan, S.P., Parks, S.E., Stathopoulos, C.E., Roach, P.D., 2014. Extraction of flavonoids from bitter melon. *Food Nutr. Sci.* 5, 458–465.
- Wu, S.J., Ng, L.T., 2008. Antioxidant and free radical scavenging activities of wild bitter melon (*Momordica charantia* Linn. var. *abbreviate* Ser.) in Taiwan. *LWT – Food. Sci. Technol.* 41, 323–330.
- Zhou, H., Tang, W., Zeng, J., Tang, C., 2014. Screening of terpene lactones and flavonoid glycosides in *Ginkgo biloba* capsule by UPLC–Orbitrap High Resolution MS, with emphasis on isomer differentiation. *J. Food Nutr. Res.* 2, 369–376.
- Zhang, Q.C., 1992. Preliminary report on the use of *Momordica charantia* extract by HIV patients. *J. Naturpath. Med.* 3, 65–69.
- Zhu, Y., Dong, Y., Qian, X., Cui, F., Guo, Q., Zhou, X., Wang, Y., Zhang, Y., Xiong, Z., 2012. Effect of superfine grinding on antidiabetic activity of bitter melon powder. *Int. J. Mol. Sci.* 13, 14203–14218.