

Contents lists available at [ScienceDirect](http://www.sciencedirect.com)

Virology

journal homepage: www.elsevier.com/locate/yviro

Tyrosine-phosphorylation of AAV2 vectors and its consequences on viral intracellular trafficking and transgene expression

Li Zhong^{a,b,c,d}, Baozheng Li^a, Giridhararao Jayandharan^a, Cathryn S. Mah^{a,b,c,d},
Lakshmanan Govindasamy^{b,c,e}, Mavis Agbandje-McKenna^{b,c,e}, Roland W. Herzog^{a,b,c,f},
Kirsten A. Weigel-Van Aken^{a,b,c,d,f}, Jacqueline A. Hobbs^{c,f,g}, Sergei Zolotukhin^{a,b,c,d,f},
Nicholas Muzyczka^{b,c,d,f}, Arun Srivastava^{a,b,c,d,f,*}

^a Division of Cellular and Molecular Therapy, Department of Pediatrics, University of Florida College of Medicine, Gainesville, FL, USA

^b Powell Gene Therapy Center, University of Florida College of Medicine, Gainesville, FL, USA

^c Genetics Institute, University of Florida College of Medicine, Gainesville, FL, USA

^d Shands Cancer Center, University of Florida College of Medicine, Gainesville, FL, USA

^e Department of Biochemistry and Molecular Biology, University of Florida College of Medicine, Gainesville, FL, USA

^f Department of Molecular Genetics and Microbiology, University of Florida College of Medicine, Gainesville, FL, USA

^g Department of Psychiatry, University of Florida College of Medicine, Gainesville, FL, USA

ARTICLE INFO

Article history:

Received 6 May 2008

Returned to author for revision

7 July 2008

Accepted 7 August 2008

Available online 2 October 2008

Keywords:

AAV2 vectors

Capsid proteins

Tyrosine-phosphorylation

Ubiquitination

Proteasomes

Intracellular trafficking

Gene expression

Gene therapy

ABSTRACT

We have documented that epidermal growth factor receptor protein tyrosine kinase (EGFR-PTK) signaling negatively affects intracellular trafficking and transduction efficiency of recombinant adeno-associated virus 2 (AAV2) vectors. Specifically, inhibition of EGFR-PTK signaling leads to decreased ubiquitination of AAV2 capsid proteins, which in turn, facilitates viral nuclear transport by limiting proteasome-mediated degradation of AAV2 vectors. In the present studies, we observed that AAV capsids can indeed be phosphorylated at tyrosine residues by EGFR-PTK in *in vitro* phosphorylation assays and that phosphorylated AAV capsids retain their structural integrity. However, although phosphorylated AAV vectors enter cells as efficiently as their unphosphorylated counterparts, their transduction efficiency is significantly reduced. This reduction is not due to impaired viral second-strand DNA synthesis since transduction efficiency of both single-stranded AAV (ssAAV) and self-complementary AAV (scAAV) vectors is decreased by ~68% and ~74%, respectively. We also observed that intracellular trafficking of tyrosine-phosphorylated AAV vectors from cytoplasm to nucleus is significantly decreased, which results from ubiquitination of AAV capsids followed by proteasome-mediated degradation, although downstream consequences of capsid ubiquitination may also be affected by tyrosine-phosphorylation. These studies provide new insights into the role of tyrosine-phosphorylation of AAV capsids in various steps in the virus life cycle, which has implications in the optimal use of recombinant AAV vectors in human gene therapy.

© 2008 Elsevier Inc. All rights reserved.

Introduction

Recombinant adeno-associated virus 2 (AAV2) vectors have gained attention as an alternative to the more commonly used retrovirus- and adenovirus-based vectors for gene transfer and gene therapy (Berns and Giraud, 1996; Muzyczka, 1992). A number of Phase I/II clinical trials have been initiated, or are currently underway, that utilize recombinant AAV2 vectors for the potential gene therapy of human diseases such as cystic fibrosis, α -1 anti-trypsin deficiency, Parkinson's disease,

Batten's disease, and muscular dystrophy (Flotte et al., 1996, 2004; Kay et al., 2000; Snyder and Francis, 2005). Recombinant AAV2 vectors have also been reported to transduce a wide variety of cells and tissues *in vitro* and *in vivo* (Flotte et al., 1993; Muzyczka, 1992; Snyder et al., 1997; Xiao, Li, and Samulski, 1996). Several fundamental steps in the life cycle of AAV2 vectors, such as viral binding and entry (Kashiwakura et al., 2005; Qing et al., 1999; Summerford et al., 1999; Summerford and Samulski, 1998), intracellular trafficking (Douar et al., 2001; Hansen et al., 2000, 2001; Sanlioglu et al., 2000; Zhao et al., 2006), uncoating (Thomas et al., 2004; Zhong et al., 2004c), second-strand DNA synthesis and transgene expression (Ferrari et al., 1996; Fisher et al., 1996; Qing et al., 1997; Zhong et al., 2004a, 2004b, 2004d, 2008b), and viral genome integration into host cell chromosome (McCarty, Young, and Samulski, 2004; Tan et al., 2001; Zhong et al., 2006), have been studied extensively.

* Corresponding author. Division of Cellular and Molecular Therapy, Cancer and Genetics Research Complex, 1376 Mowry Road, Room 492-A, University of Florida College of Medicine, Gainesville, FL 32610, USA. Fax: +1 352 273 8342.

E-mail address: aruns@peds.ufl.edu (A. Srivastava).




Fig. 1. Autoradiographic image of SDS-polyacrylamide gel for *in vitro* phosphorylation of AAV2 capsids by CKII and EGFR-PTK. Approximately 4.8×10^{10} particles of intact (I) and denatured (D) ssAAV2-adiponectin (K9) vectors were incubated in the absence of a protein kinase (lanes 1 and 2). CKII was incubated in the absence (lane 3) or presence of intact (I) (lane 4) or denatured (D) (lane 5) K9 vectors. Similarly, EGFR-PTK was incubated in the absence (lane 6) or presence of intact (I) (lane 7) or denatured (D) (lane 8) K9 vectors as described under Materials and methods. The arrows indicate the phosphorylated forms of AAV2 capsid proteins, and the closed and open arrowheads denote the autophosphorylated CKII and EGFR-PTK, respectively. These results are representative of two independent experiments.

Previous studies have shown that the ubiquitin–proteasome pathway plays a critical role in intracellular trafficking of AAV2 vectors (Ding et al., 2005, 2006; Douar et al., 2001; Duan et al., 2000). We recently reported that perturbations in EGFR-PTK signaling affects AAV2 transduction efficiency by not only augmenting viral second-strand DNA synthesis, but also by facilitating intracellular trafficking

from the cytoplasm to nucleus (Zhong et al., 2007). These studies led to the hypothesis that prior to exiting the late endosomes, intact AAV2 particles become phosphorylated at tyrosine residues by EGFR-PTK, which leads to ubiquitination and subsequent degradation of a substantial fraction of the vectors by the cytoplasmic proteasomes, which negatively impacts the efficiency of their transport to the nucleus.

We report here that intact AAV2 capsids can be phosphorylated at tyrosine residues by EGFR-PTK, but not at serine/threonine residues by casein kinase II (CKII), under cell-free conditions *in vitro*. We also document that tyrosine-phosphorylation of AAV2 capsids negatively affects the viral intracellular trafficking and transgene expression in intact cells *in vivo*. These studies provide new insights into the role of tyrosine-phosphorylation of AAV2 capsid in various steps in the life cycle of AAV2, which have implications in the optimal use of recombinant AAV2 vectors in human gene therapy.

Results

Intact AAV2 capsids can be phosphorylated at tyrosine residues by EGFR-PTK, but not at serine/threonine residues by CKII, in in vitro phosphorylation assays

One of the predictions from our recently published model was that AAV2 vectors become phosphorylated at tyrosine residues by EGFR-PTK in the late endosome (Zhong et al., 2007). However, whether intact AAV2 capsid proteins can be phosphorylated at tyrosine and/or serine/threonine residues has not been reported. Using *in vitro* phosphorylation assays, we first examined whether intact or denatured AAV2 capsid could be phosphorylated by casein kinase II (CKII) or EGFR-PTK. As shown in Fig. 1, the results clearly indicate that denatured AAV2 capsid proteins (lanes 5 and 8) can be readily phosphorylated *in vitro* by CKII and EGFR-PTK at serine/threonine and tyrosine residues, respectively. However, intact AAV2 capsids could not be phosphorylated by CKII (lane 4), but phosphorylation at




Fig. 2. (A) Western blot analyses of *in vitro* phosphorylation of AAV2 capsids by EGFR-PTK from two different packaging systems. Approximately 3×10^{10} particles of AAV2 vectors were incubated in the absence of EGFR-PTK (lanes 3, 5, and 7) and EGFR-PTK was mock-incubated (lane 1) or incubated in the absence (lane 2) or presence of AAV2 vectors (lanes 4, 6, and 8) as described under Materials and methods. The reaction samples were analyzed by Western blotting using anti-p-Tyr antibody for detection of phospho-tyrosine containing capsid proteins. The arrows indicate the phosphorylated forms of AAV2 capsid proteins, and the open arrowhead denotes the autophosphorylated EGFR-PTK. K9: AAV2-adiponectin (baculovirus-based AAV2 packaging system); RFP: ssAAV2-RFP (293 cells-based AAV2 packaging system); scEGFP: scAAV2-CBAP-EGFP (293 cell-based AAV2 packaging system). These results are representative of two independent experiments. (B) Western blot analyses of *in vitro* phosphorylation of AAV2 capsids by EGFR-PTK followed by separation of intact virions and free capsid proteins. Approximately 3×10^{10} particles of AAV2 vectors were incubated in the absence of EGFR-PTK (lane 2) and EGFR-PTK was incubated in the absence (lane 1) or presence of AAV2 vectors (lane 3) followed by separation of >100 kDa fraction (intact virions) (lanes 4, 5, and 6) and 30–100 kDa fraction (free capsid proteins) (lane 7, 8, and 9) using centrifugal filter devices [(Ultracel YM-100 (KD) and YM-30 (KD)] as described under Materials and methods. The samples were analyzed by Western blotting using anti-p-Tyr antibody for detection of phospho-tyrosine containing capsid proteins (upper panel) and anti-AAV2 cap (B1) antibody for detection of total capsid proteins (lower panel). The arrows indicate the phosphorylated and total AAV2 capsid proteins, and the open arrowhead denotes the autophosphorylated EGFR-PTK.




Fig. 3. DNA slot blot analysis for AAV2 entry into HeLa cells after *in vitro* phosphorylation of AAV2 capsids by EGFR-PTK. HeLa cells were mock-infected (lane 1) or infected with AAV2-*lacZ* vectors, which were either mock-incubated (lane 2) or pre-incubated with ATP (lane 3), EGFR-TPK (lane 4) or both (lane 5). Following digestion with trypsin to degrade adsorbed and unadsorbed vectors, low- M_r DNA samples were isolated 2 h post-infection and analyzed on by slot blot hybridization using a ^{32}P -labeled *lacZ* DNA probe.

tyrosine residues by EGFR-PTK was readily observed (lane 7). Tyrosine-phosphorylation of AAV2 capsids by EGFR-PTK occurred regardless of the vector production method or the encapsidated transgene. For example, Western blot analyses using anti-phosphotyrosine (anti-p-Tyr) antibody revealed that phospho-tyrosines could not be detected in single-stranded AAV2 (ssAAV2) vectors containing the canine adiponectin gene (K9) produced by the baculovirus-based packaging system, or ssAAV2 vectors containing the red fluorescence protein (RFP), or self-complementary AAV2 (scAAV2) vectors containing the enhanced green fluorescent protein (EGFP) gene generated by the 293 cell-based packaging system (Fig. 2A, lanes 3, 5, and 7). However, all vector preparations could be phosphorylated by EGFR-PTK (Fig. 2A, lanes 4, 6, and 8). We also examined whether *in vitro* phosphorylation conditions *per se* lead to vector instability. To this end, following *in vitro* phosphorylation of AAV2 capsids by EGFR-PTK, intact virions (>100 kDa) were separated from free capsid proteins (30–100 kDa) using centrifugal filter devices [(Ultracel YM-100 (KD) and YM-30 (KD)] and analyzed on Western blots using anti-p-Tyr antibody for detection of phospho-tyrosine containing capsid proteins as well as anti-AAV2 capsid (B1) antibody for detection of total capsid proteins. As shown in Fig. 2B, both the tyrosine-phosphorylated (upper panel) and AAV2 capsid proteins (lower panel) were detected in the > 100 kDa fraction (lanes 5 and 6). No signals were detected in the 30–100 kDa fraction (lanes 8 and 9). We conclude, therefore, that tyrosine-phosphorylated AAV2 vectors retain their structural integrity.

Tyrosine-phosphorylated AAV2 vectors are infectious, but their transduction efficiency is significantly reduced

We next evaluated whether tyrosine-phosphorylation affected the biological activity of AAV2 vectors by examining their ability to bind and enter the target cell as well as their ability to express the encapsidated transgene. For the first set of experiments, recombinant AAV2-*lacZ* vectors were pre-incubated with ATP alone, EGFR-PTK alone, or ATP+EGFR-PTK, and used to infect HeLa cells at 37 °C

for 2 h under identical conditions. Following exhaustive digestion with trypsin to remove any free virions, low- M_r DNA samples were isolated at 2 h post-infection and analyzed on DNA slot-blot using a ^{32}P -labeled *lacZ* DNA probe. As can be seen in Fig. 3, *in vitro* phosphorylation of AAV2 capsids by EGFR-PTK had no effect on viral binding and entry into HeLa cells.









For the second set of experiments, both ssAAV2 and scAAV2 vectors were used since tyrosine-phosphorylation negatively affects viral second-strand DNA synthesis of ssAAV2 vectors (Mah et al., 1998; Qing et al., 2001, 1997; Zhong et al., 2004a, 2004b, 2004d), and scAAV2 vectors bypass the requirement for viral second-strand DNA synthesis (McCarty et al., 2003; Wang et al., 2003). HeLa cells were mock-infected or infected with ssAAV2-RFP or scAAV2-EGFP vectors, which were either mock-treated or incubated with ATP alone, EGFR-PTK alone, or ATP+EGFR-PTK, as described above. Transgene expression was detected by fluorescence microscopy 48 h post-infection. As is evident from the results shown in Fig. 4, *in vitro* phosphorylation of viral capsids by EGFR-PTK decreased transgene expression by ~68% for ssAAV2 vectors (panels A, B) and by ~74% for scAAV2 vectors (panels C, D). Treatment of viral capsids with CKII, on the other hand, had no effect on either ssAAV2 vectors (panels A, B) or scAAV2 vectors (panels C, D). Thus, we conclude that although tyrosine-phosphorylated AAV2 vectors enter cells as efficiently as their unphosphorylated counterparts, their transduction efficiency is significantly reduced. This reduction is not due to impaired viral second-strand DNA synthesis since transduction efficiency of both ssAAV2 and scAAV2 vectors is affected.

AAV2 capsids are phosphorylated at tyrosine residues following infection, and *in vitro* phosphorylation of viral capsids at tyrosine residues leads to ubiquitination of intact AAV2 vectors

Since AAV2 vectors could be phosphorylated by EGFR-PTK *in vitro*, we also wished to evaluate whether intact AAV2 particles also undergo tyrosine-phosphorylation following infection. This was carried out as follows. HeLa cells were either mock-treated or treated with sodium orthovanadate (NaOV), a specific inhibitor of protein phosphatases, and then were either mock-infected or infected with AAV2 vectors for 2 h at 37 °C. Whole cell lysates (WCLs) were prepared 4 h post-infection and equivalent amounts of proteins were immunoprecipitated first with anti-AAV2 capsid antibody (A20) followed by Western blot analyses using anti-p-Tyr antibody. These results are shown in Fig. 5A. As can be seen, tyrosine-phosphorylated AAV2 capsid proteins (P-Tyr-AAV2 Cap) were readily detectable in AAV2-infected cells (lanes 1 and 2). No signal was detectable in mock-infected cells (lane 3), and the extent of tyrosine-phosphorylation of AAV2 capsid proteins was increased following treatment with NaOV (lane 2). These results suggest that AAV2 vectors are indeed phosphorylated at tyrosine residues following infection of cells.

We next examined why tyrosine-phosphorylated AAV2 vectors failed to efficiently transduce the target cell. The ubiquitin–proteasome pathway plays an important role in the cell by specifically degrading both endogenous and foreign proteins (Pickart, 2001; Schwartz and Ciechanover, 1999). Direct evidence for ubiquitination of AAV2 capsid proteins in HeLa cells and *in vitro* ubiquitination assays

Fig. 4. (A) Comparative analyses of AAV2-mediated transduction efficiency in HeLa cells after *in vitro* phosphorylation of AAV2 capsids by EGFR-PTK. HeLa cells were infected by ssAAV2-RFP vectors, which were pre-incubated with ATP, EGFR-TPK, or both. Transgene expression was detected by fluorescence microscopy 48 h post-infection. Original magnification 100 \times . (B) Quantitative analyses of AAV2 transduction efficiency in HeLa cells. Images from five visual fields were analyzed quantitatively by ImageJ analysis software. Transgene expression was assessed as total area of red fluorescence (pixel 2) per visual field (mean \pm SD). Analysis of variance (ANOVA) was used to compare test results with the control and they were determined to be statistically significant. * $P < 0.05$ vs. ssAAV2-RFP. (C) Infection of HeLa cells with scAAV2-EGFP vectors pre-incubated with ATP, EGFR-TPK, or both. Transgene expression was detected by fluorescence microscopy at 48 h post-infection. Original magnification 100 \times . (D) Quantitative analyses of AAV2 transduction efficiency was assessed as described above, and were determined to be statistically significant. * $P < 0.05$ vs. scAAV2-EGFP. (E) Comparative analyses of AAV2-mediated transduction efficiency in HeLa cells after *in vitro* phosphorylation of AAV2 capsids by CKII. HeLa cells were infected by ssAAV2-RFP vectors, which were pre-incubated with ATP, CKII, or both. Transgene expression was detected by fluorescence microscopy 48 h post-infection. Original magnification 100 \times . (F) Quantitative analyses of AAV2 transduction efficiency was assessed as described above. (G) Infection of HeLa cells with scAAV2-EGFP vectors pre-incubated with ATP, CKII, or both. Transgene expression was detected by fluorescence microscopy 48 h post-infection. Original magnification 100 \times . (H) Quantitative analyses of AAV2 transduction efficiency were assessed as described above.






Fig. 5. (A) Detection of intracellular tyrosine-phosphorylation of intact AAV2 vectors. Whole cell lysates (WCLs) prepared from HeLa cells untreated (lanes 1 and 3) or treated with NaOV (lane 2) following infection with K9 vectors (lanes 1 and 2) or mock-infection (lane 3), were immuno-precipitated with anti-AAV2 capsid antibody A20 followed by Western blot analyses with anti-p-Tyr antibody. (B) Detection of intracellular ubiquitin-conjugation of intact AAV2 vectors after *in vitro* phosphorylation of viral capsids by EGFR-PTK. Western blot analyses of ubiquitinated AAV2 capsid proteins in HeLa cells following transduction with ssAAV2-K9 vectors after *in vitro* phosphorylation of viral capsids by EGFR-PTK. Whole cell lysates (WCLs) prepared from HeLa cells untreated or treated with MG132 following mock-infected (lanes 1 and 2), or infected with K9 vectors, which were mock-incubated (lanes 3 and 4), or incubated with ATP (lanes 7 and 8), EGFR-PTK (lanes 9 and 10) or both (lanes 5 and 6), were immuno-precipitated with anti-AAV2 capsid antibody A20 followed by Western blot analyses with anti-Ub monoclonal antibody P4D1. This result is representative of two independent experiments.

has been reported (Yan et al., 2002), where only denatured AAV2 capsids, but not intact AAV2, could be ubiquitinated *in vitro*, which suggested that either a conformational change or a modification, such as phosphorylation of intact AAV2 capsid is required before ubiquitination can ensue. A number of studies have reported that phosphorylation of cellular proteins at tyrosine or serine/threonine residues is required for efficient ubiquitination and degradation of these proteins (Brown et al., 1995; Cenciarelli et al., 1996; Paolini et al., 1999; Penrose et al., 2004).

Since in our previously published studies (Zhong et al., 2007), we reported that inhibition of EGFR-PTK signaling decreases ubiquitination of total cellular proteins as well as AAV2 capsid proteins, we examined whether *in vitro* phosphorylation of viral capsids by EGFR-PTK leads to ubiquitination of AAV2 capsids in intact cells. This was carried out as follows. HeLa cells were either mock-treated or treated with MG132, a specific inhibitor of proteasomes, and then were either mock-infected or infected with AAV2 vectors, with and without tyrosine-phosphorylation, for 2 h at 37 °C. Whole cell lysates (WCLs) were prepared 4 h post-infection and equivalent amounts of proteins were immuno-precipitated first with anti-AAV2 capsid antibody (A20) followed by Western blot analyses using anti-Ub monoclonal antibody. These results are shown in Fig. 5B. As can be seen, the ubiquitinated AAV2 capsid proteins (Ub-AAV2 Cap) were undetectable in mock-infected cells (lanes 1 and 2), the signal of ubiquitinated AAV2 capsid proteins was weaker in untreated cells (lanes 3, 5, 7, and 9), but a significant accumulation of ubiquitinated AAV2 capsid proteins occurred following treatment with MG132 (lanes 4, 6, 8, and 10). Most interestingly, *in vitro* phosphorylation of viral capsids by EGFR-PTK dramatically increased the extent of accumulation of MG132-induced ubiquitinated AAV2 capsid proteins (lane 6). These results strongly suggest that *in vitro* phosphorylation of capsids by EGFR-PTK leads to ubiquitination of AAV2 capsids in intact cells. However, since MG132 potentiates capsid ubiquitination as well as increases transduction, it is also possible that capsid ubiquitination serves as a positive sorting signal and not as a negative degradation signal. Thus, tyrosine-phosphorylation of AAV2 capsids may limit transduction by affecting other events during infection.

In vitro phosphorylation of viral capsids at tyrosine residues leads to impaired intracellular trafficking to the nucleus

The ubiquitin–proteasome pathway plays a critical role in AAV2 intracellular trafficking, and proteasome inhibitors can promote AAV2 nuclear transport, leading to augmentation of AAV2 transduction (Ding et al., 2005, 2006; Douar et al., 2001; Duan et al., 2000). We have also documented that inhibition of EGFR-PTK signaling facilitates nuclear transport of AAV2 vectors (Zhong et al., 2004c, 2007), which suggested that phosphorylation of AAV2 capsids by EGFR-PTK might also be involved in AAV2 trafficking. To further examine this hypothesis, we examined the cytoplasmic and nuclear distribution of AAV2 genomes following *in vitro* phosphorylation of AAV2 vectors. HeLa cells were infected with AAV2-*lacZ* vectors, with or without tyrosine-phosphorylation, as described above. Nuclear and cytoplasmic fractions were obtained 18 h post-infection, and low- M_r DNA samples were isolated and analyzed on Southern blots using a ^{32}P -labeled *lacZ* DNA probe. These results are shown in Fig. 6, panel A. A significant fraction of the input ssAAV2 DNA was present in the cytoplasm following infection with AAV2 vectors that were mock-treated or pre-incubated with ATP alone, or with EGFR-TPK alone (lanes 2, 3, and 4), and only a small fraction was detected in the nucleus (lanes 7, 8, and 9). Interestingly, however, in cells infected with AAV2 vectors that were phosphorylated by EGFR-PTK, the input ssAAV2 DNA in the cytoplasmic fraction was increased (lane 5) with a corresponding decrease in the nuclear fraction (lane 10). Densitometric scanning of autoradiographs (Fig. 6, panel B) indicated that ~35% of the input ssAAV2 DNA was present in the nuclear fraction in cells infected with AAV2 vectors in the absence of tyrosine-phosphorylation, but only ~16% following *in vitro* phosphorylation at tyrosine residues. These data are consistent with our previous studies (Zhong et al., 2007), and corroborate that intracellular trafficking of tyrosine-phosphorylated AAV2 vectors from the cytoplasm to the nucleus is significantly decreased, which is due most likely to increased ubiquitination of AAV2 capsids followed by proteasome-mediated degradation.

Discussion

Our original observation that specific inhibitors of cellular protein tyrosine kinases in general, and EGFR-PTK in particular, but not inhibitors of cellular serine/threonine kinases, dramatically increase the transduction efficiency of ssAAV vectors (Mah et al., 1998; Qing et al., 1997), prompted us to pursue a series of studies in which we identified that a cellular chaperone protein, FKBP52, is phosphorylated at tyrosine residues by EGFR-PTK, and that tyrosine-phosphorylated FKBP52 strongly inhibits the viral second-strand DNA synthesis, and consequently the transgene expression (Qing et al., 2001, 2003; Zhong et al., 2004a, 2004b, 2004d, 2008b). More recently, however, we observed that in addition to negatively impacting viral second-strand DNA synthesis and transgene expression in the nucleus, EGFR-PTK-signaling also negatively regulates viral intracellular trafficking in the cytoplasm (Zhong et al., 2007). Based on these studies, we hypothesized that either prior to or during viral egress from the late endosome, EGFR-PTK catalyzes the phosphorylation of intact AAV2 capsids, which is a putative signal for ubiquitination. Ubiquitinated AAV2 capsids are subsequently targeted for degradation by the cellular proteasome. That intact AAV2 vectors can indeed be phosphorylated at tyrosine residues by EGFR-PTK, both *in vitro* and *in vivo*, but not at serine/threonine residues by CKII, as described in the present studies, lends strong support to our hypothesis.

Although the Ub/proteasome pathway has previously been shown to play an essential role in AAV2 nuclear transport (Ding et al., 2005, 2006; Douar et al., 2001; Duan et al., 2000), our data also provide direct evidence that EGFR-PTK-mediated tyrosine-phosphorylation of AAV2 capsids leads to ubiquitination, and subsequent degradation by the proteasome, which impairs viral nuclear transport as well as transduction efficiency of AAV2 vectors. Whether AAV2 vectors can be phosphorylated at serine/threonine residues by other cellular serine/threonine kinase(s), and what consequences, if any, of serine/threonine phosphorylation of capsids have on the wild-type (WT) AAV2 biology, remains unclear. What has become clear from our current studies is that phosphorylation of the surface-exposed tyrosine residues on recombinant AAV2 vector capsids is not desirable since site-directed mutagenesis of these tyrosine residues resulted in the generation of AAV2 vectors that are capable of transducing a wide variety of cells and tissues *in vitro* and *in vivo* significantly more efficiently than those with the WT capsids (Zhong et al., 2008a).

Further studies on the precise role of tyrosine-phosphorylation of AAV2 capsids are likely to lead not only to a better understanding of

various steps in the virus life cycle, but also to aid in the development of novel recombinant AAV vectors especially in view of the fact that the surface-exposed tyrosine residues are conserved in AAV serotypes 1 through 10, which has implications in the potential use of AAV serotype vectors in human gene therapy.

Materials and methods

Cells, viruses, plasmids, antibodies, and chemicals

The human cervical carcinoma cell line, HeLa, was obtained from the American Type Culture Collection (ATCC, Rockville, MD). The cell line was maintained as monolayer cultures in Iscove's-modified Dulbecco's medium (IMDM) supplemented with 10% newborn calf serum (NCS) and 1% (by volume) of 100× stock solution of antibiotics (10,000 U penicillin + 10,000 µg streptomycin). Highly purified stocks of ssAAV2 vectors containing the β-galactosidase (*lacZ*) reporter gene, red fluorescence protein (RFP) gene, or scAAV2 vectors containing the enhanced green fluorescence protein (EGFP) gene driven by the chick β-actin (CBA) promoter (ssAAV2-*lacZ*, ssAAV2-RFP, or scAAV2-EGFP) were generated as described previously (Auricchio et al., 2001). ssAAV2 vectors containing the canine adiponectin gene (ssAAV2-K9) were generated by a baculovirus-based packaging system as described previously (Urabe et al., 2002). Physical particle titers of recombinant vector stocks were determined by quantitative DNA slot blot analysis. An AAV2-helper plasmid, pACG-2, containing the AAV2 *rep* gene with an ACG start codon, was generously provided by Dr. R. Jude Samulski (University of North Carolina at Chapel Hill, Chapel Hill, NC). A self-complementary AAV2 (scAAV2) cloning vector, pdsCBA-EGFP, was a kind gift from Dr. Xiao Xiao (University of North Carolina at Chapel Hill). Horseradish peroxidase (HRP)-conjugated antibody specific for ubiquitin (Ub) (mouse monoclonal immunoglobulin G₁ [IgG₁], clone P4D1), antibody specific for phosphorylated tyrosine residues (anti-p-Tyr) (mouse monoclonal IgG_{2b}, clone PY99), HRP-conjugated goat anti-mouse IgG_{2b} antibody, HRP-conjugated goat anti-mouse IgG₁ antibody, normal mouse IgG and protein G plus-agarose were purchased from Santa Cruz Biotechnology (Santa Cruz, CA). Antibody specific for intact AAV2 particles (mouse monoclonal IgG₃, clone A20) and antibody specific for AAV2 capsid proteins (mouse monoclonal IgG₁, clone B1) were obtained from Research Diagnostics, Inc. (Flanders, NJ). MG132 was purchased from Calbiochem (La Jolla, CA). All other chemicals were purchased from Sigma-Aldrich Co. (St. Louis, MO).




Fig. 6. Detection of intracellular trafficking of AAV2 vectors to the nucleus after *in vitro* phosphorylation of viral capsids by EGFR-PTK. (A) Southern blot analyses of cytoplasmic and nuclear distribution of AAV2 genomes in HeLa cells after *in vitro* phosphorylation of AAV2 capsids by EGFR-PTK. HeLa cells were mock-infected (lanes 1 and 6) or infected by AAV2-*lacZ* vectors, which were mock-incubated (lanes 2 and 7) or pre-incubated with ATP (lanes 3 and 8), EGFR-PTK (lanes 4 and 9) or both (lanes 5 and 10). Nuclear and cytoplasmic fractions were prepared 18 h post-infection, low- M_r DNA samples were isolated and electrophoresed on 1% agarose gels followed analyzed by Southern blot hybridization using a 32 P-labeled *lacZ* DNA probe. (B) Densitometric scanning of autoradiographs for the quantitation of relative amounts of viral genomes. These results are representative of two independent experiments.

In vitro phosphorylation assays

In vitro phosphorylation by CKII (Calbiochem, La Jolla, CA) was carried out as previously described (Qing et al., 2001; Zhong et al., 2004d). Briefly, the complete reaction mixture contained identical amount of intact or denatured AAV2 vectors ($\sim 3\text{--}4.8 \times 10^{10}$ viral particles), 20 mM Tris-HCl, 50 mM KCl, 10 mM MgCl₂, 50 mM Na₃VO₄, 10 μ Ci (0.37 mBq) γ -³²P-ATP and 500 U (500,000 U/ml) purified CKII. *In vitro* phosphorylation by EGFR-PTK was carried out as described previously (Qing et al., 2001; Weber et al., 1984; Zhong et al., 2004d), with the following modifications. The complete reaction mixture contained identical amounts of intact or denatured AAV2 vectors ($\sim 3\text{--}4.8 \times 10^{10}$ viral particles), 20 mM HEPES, 4 mM MgCl₂, 10 mM MnCl₂, 10 μ Ci (0.37 mBq) γ -³²P-ATP and 10 U (15,000 U/mg) purified EGFR-PTK (Sigma-Aldrich Co., St. Louis, MO) with the indicated appropriate controls. The reaction mixtures were incubated at 30 °C for 1 h. The phosphorylated AAV2 capsid proteins were separated from free γ -³²P-ATP on 10% SDS-polyacrylamide gels followed by autoradiography with Kodak 7 X-Omat film at -70 °C. In some experiments, *in vitro* phosphorylation assays by EGFR-PTK were also carried out by adding 200 μ M ATP instead of γ -³²P-ATP. The phosphorylated AAV2 capsid proteins were identified by Western blotting with anti-p-Tyr antibody as described below.

Separation of intact AAV2 virions and AAV2 capsid proteins using ultrafiltration assay

After *in vitro* phosphorylation assays, the reaction samples plus 450 μ l of phosphate buffered saline (PBS, pH7.4) were added into the upper part of the YM-100 Microcon[®] centrifugal filter device (cutoff value: 100 kDa) (Millipore Corporation, Bedford, MA) and centrifuged at 14,000 \times g for 24 min at 4 °C. Approximately 10–20 μ l of ultrafiltrates was collected. The flow through samples were added into the upper part of the YM-30 Microcon[®] centrifugal filter device (cutoff value: 30 kDa) (Millipore Corporation, Bedford, MA) and centrifuged at 14,000 \times g for 24 min at 4 °C. Approximately 10–20 μ l of ultrafiltrates was collected. The ultrafiltrate samples were analyzed by Western blotting using anti-p-Tyr and anti-AAV2 cap (B1) antibodies as described below.

Preparation of whole cell lysates (WCLs) and co-immunoprecipitations

WCLs were prepared as described previously (Zhao et al., 2006; Zhong et al., 2004d; Zhong and Su, 2002), with the following modifications. Briefly, 2×10^6 HeLa cells were either mock-treated, treated with, 1 mM NaOV for 2 h and then were infected with AAV2-adiponectin vectors at 10^4 particles/cell for 2 h at 37 °C. In some experiments cells were either mock-treated or treated with 4 μ M MG132 for 4 h and then were infected with AAV2-adiponectin vectors at 10^4 particles/cell for 2 h at 37 °C, which were mock-incubated or pre-incubated with ATP, EGFR-TPK, or both. For immunoprecipitations, cells were treated with 0.01% trypsin and washed extensively with PBS to remove any adsorbed and unadsorbed virus particles after treatment or at 4 h post-infection and then resuspended in 1 ml hypotonic buffer (20 mM HEPES pH 7.5, 5 mM KCl, 0.5 mM MgCl₂) containing 1 mM DTT, 10 mM NaF, 2 mM Na₃VO₄, 0.5 mM PMSF, 10 μ g/ml aprotinin, 10 μ g/ml leupeptin and 10 μ g/ml. WCLs were prepared by homogenization in a tight-fitting Duall tissue grinder until about 95% cell lysis was achieved as monitored by trypan blue dye exclusion assay. WCLs were cleared of non-specific binding by incubation with 0.25 mg of normal mouse IgG together with 20 μ l of protein G plus-agarose beads for 60 min at 4 °C in an orbital shaker. After preclearing, 2 μ g of capsid antibody against intact AAV2 particles (A20) (mouse IgG₃) or 2 μ g of normal mouse IgG (as a negative control) was added and incubated at 4 °C for 1 h, followed by precipitation with protein G-agarose

beads at 4 °C for 12 h in a shaker. Pellets were collected by centrifugation at 2500 rpm for 5 min at 4 °C and washed four times with PBS. After the final wash, supernatants were aspirated and discarded, and pellets were resuspended in equal volume of 2 \times SDS sample buffer. Twenty μ l of resuspended pellet solutions were used for Western blotting with HRP-conjugated anti-Ub antibody as described below.

Western blot analyses

Western blotting was performed as described previously (Zhao et al., 2006; Zhong et al., 2004d; Zhong and Su, 2002). For *in vitro* phosphorylation assays, the reaction mixtures were separated by 10% SDS-polyacrylamide gel electrophoresis (SDS-PAGE) and transferred to Immobilon-P membranes (Millipore, Bedford, MA). Membranes were blocked at 4 °C for 12 h with 5% nonfat milk in 1 \times Tris-buffered saline (TBS, 20 mM Tris-HCl, pH 7.5, 150 mM NaCl). Membranes were treated with monoclonal anti-p-Tyr antibody (1:300 dilution) or monoclonal anti-AAV2 capsid proteins antibody followed by horseradish peroxidase-conjugated anti-mouse IgG_{2b} or anti-mouse IgG₁ (1:10,000 dilution). For immunoprecipitations, resuspended pellet solutions were boiled for 2–3 min and 20 μ l of samples were used for SDS-PAGE. After blocking at 4 °C for 12 h with 5% nonfat milk in 1 \times Tris-buffered saline, membranes were treated with monoclonal HRP-conjugated anti-Ub antibody (1:2000 dilution). Immuno-reactive bands were visualized using chemiluminescence (ECL-plus, Amersham Pharmacia Biotech, Piscataway, NJ).

Isolation of nuclear and cytoplasmic fractions from HeLa cells

Nuclear and cytoplasmic fractions from HeLa cells were isolated as described previously (Zhong et al., 2004c, 2007). Briefly, mock-infected or recombinant ssAAV2-*lacZ* vector-infected cells were washed twice with PBS at 18 h post-infection, treated with 0.01% trypsin and washed extensively with PBS to remove any adsorbed and unadsorbed virus particles. Cell pellets were gently resuspended in 200 μ l hypotonic buffer (10 mM HEPES, pH 7.9, 1.5 mM MgCl₂, 10 mM KCl, 0.5 mM DTT, 0.5 mM PMSF) and incubated on ice for 5 min, after which 10 μ l 10% NP-40 was added, and observed under a light microscope. Samples were mixed gently and centrifuged for 5 min at 500 rpm at 4 °C. Supernatants (cytoplasmic fraction) were decanted and stored on ice. Pellets (nuclear fractions) were washed twice with 1 ml hypotonic buffer and stored on ice. The purity of each fraction was determined to be >95%, as measured by the absence of acid phosphatase activity (nuclear fractions) and absence of histone H3 (cytoplasmic fractions) as described previously (Hansen et al., 2000; Zhong et al., 2004c).

Southern blot analysis for AAV2 trafficking

Low-M_r DNA samples from nuclear and cytoplasmic fractions were isolated and electrophoresed on 1% agarose gels or 1% alkaline-agarose gels followed by Southern blot hybridization using a ³²P-labeled *lacZ*-specific DNA probe as described previously (Hansen et al., 2000; Zhong et al., 2004c). Densitometric scanning of autoradiographs for the quantitation was evaluated with ImageJ analysis software (NIH, Bethesda, MD).

Recombinant AAV2 vector transduction assays

Approximately 1×10^5 HeLa cells were plated in each well in 12-well plates and incubated at 37 °C for 12 h. Cells were washed once with IMDM and then infected at 37 °C for 2 h with recombinant AAV2 vectors as described previously (Qing et al., 2001; Zhong et al., 2004a, 2004d). Cells were incubated in complete IMDM containing 10% NCS and 1% antibiotics for 48 h. The transduction efficiency was measured

by EGFP imaging using a Leica DM IRB/E fluorescence microscope (Leica Microsystems Wetzlar GmbH, Wetzlar, Germany). Images from three to five visual fields of mock-infected and vector-infected HeLa cells at 48 h post-infection were analyzed quantitatively by ImageJ analysis software (NIH, Bethesda, MD, USA). Transgene expression was assessed as total area of green fluorescence (pixel²) per visual field (mean±SD). Analysis of variance (ANOVA) was used to compare between test results and the control and they were determined to be statistically significant.

Acknowledgments

We thank Drs. R. Jude Samulski and Xiao Xiao for their kind gifts of recombinant AAV plasmids, and Dr. Shangzhen Zhou for her help with recombinant AAV2 vectors. We also thank Drs. Kenneth I. Berns and Dennis Steindler for a critical review of this manuscript. This research was supported in part by a research grant 8187368876 from the Roche Foundation for Anemia Research (to LZ), a Scientist Development Grant from the American Heart Association (to CSM), and Public Health Service grants R01 GM-082946 and P01 HL-51811 (to MA-M and SZ), R01 GM-082946 and P01 HL-51811 (to MA-M, NM and SZ), R01 DK-062302 (to SZ), P01 HL-078810-Project 3 (to RWH), and R01 EB-002073, R01 HL-65570 and R01 HL-07691, and P01 DK 058327-Project 1 (to AS) from the National Institutes of Health.

References

- Auricchio, A., Hildinger, M., O'Connor, E., Gao, G.P., Wilson, J.M., 2001. Isolation of highly infectious and pure adeno-associated virus type 2 vectors with a single-step gravity-flow column. *Hum. Gene Ther.* 12 (1), 71–76.
- Berns, K.I., Giraud, C., 1996. Biology of adeno-associated virus. *Curr. Top. Microbiol. Immunol.* 218, 1–23.
- Brown, K., Gerstberger, S., Carlson, L., Franzoso, G., Siebenlist, U., 1995. Control of I kappa B-alpha proteolysis by site-specific, signal-induced phosphorylation. *Science* 267 (5203), 1485–1488.
- Cenciarelli, C., Wilhelm Jr., K.G., Guo, A., Weissman, A.M., 1996. T cell antigen receptor ubiquitination is a consequence of receptor-mediated tyrosine kinase activation. *J. Biol. Chem.* 271 (15), 8709–8713.
- Ding, W., Zhang, L., Yan, Z., Engelhardt, J.F., 2005. Intracellular trafficking of adeno-associated viral vectors. *Gene Ther.* 12 (11), 873–880.
- Ding, W., Zhang, L.N., Yeaman, C., Engelhardt, J.F., 2006. rAAV2 traffics through both the late and the recycling endosomes in a dose-dependent fashion. *Mol. Ther.* 13 (4), 671–682.
- Douar, A.M., Poulard, K., Stockholm, D., Danos, O., 2001. Intracellular trafficking of adeno-associated virus vectors: routing to the late endosomal compartment and proteasome degradation. *J. Virol.* 75 (4), 1824–1833.
- Duan, D., Yue, Y., Yan, Z., Yang, J., Engelhardt, J.F., 2000. Endosomal processing limits gene transfer to polarized airway epithelia by adeno-associated virus. *J. Clin. Invest.* 105 (11), 1573–1587.
- Ferrari, F.K., Samulski, T., Shenk, T., Samulski, R.J., 1996. Second-strand synthesis is a rate-limiting step for efficient transduction by recombinant adeno-associated virus vectors. *J. Virol.* 70 (5), 3227–3234.
- Fisher, K.J., Gao, G.P., Weitzman, M.D., DeMatteo, R., Burda, J.F., Wilson, J.M., 1996. Transduction with recombinant adeno-associated virus for gene therapy is limited by leading-strand synthesis. *J. Virol.* 70 (1), 520–532.
- Flotte, T.R., Afione, S.A., Conrad, C., McGrath, S.A., Solow, R., Oka, H., Zeitlin, P.L., Guggino, W.B., Carter, B.J., 1993. Stable in vivo expression of the cystic fibrosis transmembrane conductance regulator with an adeno-associated virus vector. *Proc. Natl. Acad. Sci. U. S. A.* 90 (22), 10613–10617.
- Flotte, T., Carter, B., Conrad, C., Guggino, W., Reynolds, T., Rosenstein, B., Taylor, G., Walden, S., Wetzel, R., 1996. A phase I study of an adeno-associated virus-CFTR gene vector in adult CF patients with mild lung disease. *Hum. Gene Ther.* 7 (9), 1145–1159.
- Flotte, T.R., Brantly, M.L., Spencer, L.T., Byrne, B.J., Spencer, C.T., Baker, D.J., Humphries, M., 2004. Phase I trial of intramuscular injection of a recombinant adeno-associated virus alpha 1-antitrypsin (rAAV2-CB-hAAT) gene vector to AAT-deficient adults. *Hum. Gene Ther.* 15 (1), 93–128.
- Hansen, J., Qing, K., Kwon, H.J., Mah, C., Srivastava, A., 2000. Impaired intracellular trafficking of adeno-associated virus type 2 vectors limits efficient transduction of murine fibroblasts. *J. Virol.* 74 (2), 992–996.
- Hansen, J., Qing, K., Srivastava, A., 2001. Adeno-associated virus type 2-mediated gene transfer: altered endocytic processing enhances transduction efficiency in murine fibroblasts. *J. Virol.* 75 (9), 4080–4090.
- Kashiwakura, Y., Tamayose, K., Iwabuchi, K., Hirai, Y., Shimada, T., Matsumoto, K., Nakamura, T., Watanabe, M., Oshimi, K., Daida, H., 2005. Hepatocyte growth factor receptor is a coreceptor for adeno-associated virus type 2 infection. *J. Virol.* 79 (1), 609–614.
- Kay, M.A., Manno, C.S., Ragni, M.V., Larson, P.J., Couto, L.B., McClelland, A., Glader, B., Chew, A.J., Tai, S.J., Herzog, R.W., Arruda, V., Johnson, F., Scallan, C., Skarsgard, E., Flake, A.W., High, K.A., 2000. Evidence for gene transfer and expression of factor IX in haemophilia B patients treated with an AAV vector. *Nat. Genet.* 24 (3), 257–261.
- Mah, C., Qing, K., Khuntirat, B., Ponnazhagan, S., Wang, X.S., Kube, D.M., Yoder, M.C., Srivastava, A., 1998. Adeno-associated virus type 2-mediated gene transfer: role of epidermal growth factor receptor protein tyrosine kinase in transgene expression. *J. Virol.* 72 (12), 9835–9843.
- McCarty, D.M., Fu, H., Monahan, P.E., Toulson, C.E., Naik, P., Samulski, R.J., 2003. Adeno-associated virus terminal repeat (TR) mutant generates self-complementary vectors to overcome the rate-limiting step to transduction in vivo. *Gene Ther.* 10 (26), 2112–2118.
- McCarty, D.M., Young Jr., S.M., Samulski, R.J., 2004. Integration of adeno-associated virus (AAV) and recombinant AAV vectors. *Annu. Rev. Genet.* 38, 819–845.
- Muzyczka, N., 1992. Use of adeno-associated virus as a general transduction vector for mammalian cells. *Curr. Top. Microbiol. Immunol.* 158, 97–129.
- Paolini, R., Serra, A., Molfetta, R., Piccoli, M., Frati, L., Santoni, A., 1999. Tyrosine kinase-dependent ubiquitination of CD16 zeta subunit in human NK cells following receptor engagement. *Eur. J. Immunol.* 29 (10), 3179–3187.
- Penrose, K.J., Garcia-Alai, M., de Prat-Gay, G., McBride, A.A., 2004. Casein Kinase II phosphorylation-induced conformational switch triggers degradation of the papillomavirus E2 protein. *J. Biol. Chem.* 279 (21), 22430–22439.
- Pickart, C.M., 2001. Mechanisms underlying ubiquitination. *Annu. Rev. Biochem.* 70, 503–533.
- Qing, K., Wang, X.S., Kube, D.M., Ponnazhagan, S., Bajpai, A., Srivastava, A., 1997. Role of tyrosine phosphorylation of a cellular protein in adeno-associated virus 2-mediated transgene expression. *Proc. Natl. Acad. Sci. U. S. A.* 94 (20), 10879–10884.
- Qing, K., Mah, C., Hansen, J., Zhou, S., Dwarki, V., Srivastava, A., 1999. Human fibroblast growth factor receptor 1 is a co-receptor for infection by adeno-associated virus 2. *Nat. Med.* 5 (1), 71–77.
- Qing, K., Hansen, J., Weigel-Kelley, K.A., Tan, M., Zhou, S., Srivastava, A., 2001. Adeno-associated virus type 2-mediated gene transfer: role of cellular FKBP52 protein in transgene expression. *J. Virol.* 75 (19), 8968–8976.
- Qing, K., Li, W., Zhong, L., Tan, M., Hansen, J., Weigel-Kelley, K.A., Chen, L., Yoder, M.C., Srivastava, A., 2003. Adeno-associated virus type 2-mediated gene transfer: role of cellular T-cell protein tyrosine phosphatase in transgene expression in established cell lines in vitro and transgenic mice in vivo. *J. Virol.* 77 (4), 2741–2746.
- Sanlioglu, S., Benson, P.K., Yang, J., Atkinson, E.M., Reynolds, T., Engelhardt, J.F., 2000. Endocytosis and nuclear trafficking of adeno-associated virus type 2 are controlled by rac1 and phosphatidylinositol-3 kinase activation. *J. Virol.* 74 (19), 9184–9196.
- Schwartz, A.L., Ciechanover, A., 1999. The ubiquitin-proteasome pathway and pathogenesis of human diseases. *Annu. Rev. Med.* 50, 57–74.
- Snyder, R.O., Francis, J., 2005. Adeno-associated viral vectors for clinical gene transfer studies. *Curr. Gene Ther.* 5 (3), 311–321.
- Snyder, R.O., Miao, C.H., Pattijn, G.A., Spratt, S.K., Danos, O., Nagy, D., Gown, A.M., Winther, B., Meuse, L., Cohen, L.K., Thompson, A.R., Kay, M.A., 1997. Persistent and therapeutic concentrations of human factor IX in mice after hepatic gene transfer of recombinant AAV vectors. *Nat. Genet.* 16 (3), 270–276.
- Summerford, C., Samulski, R.J., 1998. Membrane-associated heparan sulfate proteoglycan is a receptor for adeno-associated virus type 2 virions. *J. Virol.* 72 (2), 1438–1445.
- Summerford, C., Bartlett, J.S., Samulski, R.J., 1999. AlphaVbeta5 integrin: a co-receptor for adeno-associated virus type 2 infection. *Nat. Med.* 5 (1), 78–82.
- Tan, M., Qing, K., Zhou, S., Yoder, M.C., Srivastava, A., 2001. Adeno-associated virus 2-mediated transduction and erythroid lineage-restricted long-term expression of the human beta-globin gene in hematopoietic cells from homozygous beta-thalassemic mice. *Mol. Ther.* 3 (6), 940–946.
- Thomas, C.E., Storm, T.A., Huang, Z., Kay, M.A., 2004. Rapid uncoating of vector genomes is the key to efficient liver transduction with pseudotyped adeno-associated virus vectors. *J. Virol.* 78 (6), 3110–3122.
- Urabe, M., Ding, C., Kotin, R.M., 2002. Insect cells as a factory to produce adeno-associated virus type 2 vectors. *Hum. Gene Ther.* 13 (16), 1935–1943.
- Wang, Z., Ma, H.I., Li, J., Sun, L., Zhang, J., Xiao, X., 2003. Rapid and highly efficient transduction by double-stranded adeno-associated virus vectors in vitro and in vivo. *Gene Ther.* 10 (26), 2105–2111.
- Weber, W., Bertics, P.J., Gill, G.N., 1984. Immunoaffinity purification of the epidermal growth factor receptor. Stoichiometry of binding and kinetics of self-phosphorylation. *J. Biol. Chem.* 259 (23), 14631–14636.
- Xiao, X., Li, J., Samulski, R.J., 1996. Efficient long-term gene transfer into muscle tissue of immunocompetent mice by adeno-associated virus vector. *J. Virol.* 70 (11), 8098–8108.
- Yan, Z., Zak, R., Luxton, G.W., Ritchie, T.C., Bantel-Schaal, U., Engelhardt, J.F., 2002. Ubiquitination of both adeno-associated virus type 2 and 5 capsid proteins affects the transduction efficiency of recombinant vectors. *J. Virol.* 76 (5), 2043–2053.
- Zhao, W., Zhong, L., Wu, J., Chen, L., Qing, K., Weigel-Kelley, K.A., Larsen, S.H., Shou, W., Warrington Jr., K.H., Srivastava, A., 2006. Role of cellular FKBP52 protein in intracellular trafficking of recombinant adeno-associated virus 2 vectors. *Virology* 353 (2), 283–293.
- Zhong, L., Su, J.Y., 2002. Isoflurane activates PKC and Ca(2+)-calmodulin-dependent protein kinase II via MAP kinase signaling in cultured vascular smooth muscle cells. *Anesthesiology* 96 (1), 148–154.
- Zhong, L., Chen, L., Li, Y., Qing, K., Weigel-Kelley, K.A., Chan, R.J., Yoder, M.C., Srivastava, A., 2004a. Self-complementary adeno-associated virus 2 (AAV)-T cell protein tyrosine phosphatase vectors as helper viruses to improve transduction efficiency of

- conventional single-stranded AAV vectors in vitro and in vivo. *Mol. Ther.* 10 (5), 950–957.
- Zhong, L., Li, W., Yang, Z., Chen, L., Li, Y., Qing, K., Weigel-Kelley, K.A., Yoder, M.C., Shou, W., Srivastava, A., 2004b. Improved transduction of primary murine hepatocytes by recombinant adeno-associated virus 2 vectors in vivo. *Gene Ther.* 11 (14), 1165–1169.
- Zhong, L., Li, W., Yang, Z., Qing, K., Tan, M., Hansen, J., Li, Y., Chen, L., Chan, R.J., Bischof, D., Maina, N., Weigel-Kelley, K.A., Zhao, W., Larsen, S.H., Yoder, M.C., Shou, W., Srivastava, A., 2004c. Impaired nuclear transport and uncoating limit recombinant adeno-associated virus 2 vector-mediated transduction of primary murine hematopoietic cells. *Hum. Gene Ther.* 15 (12), 1207–1218.
- Zhong, L., Qing, K., Si, Y., Chen, L., Tan, M., Srivastava, A., 2004d. Heat-shock treatment-mediated increase in transduction by recombinant adeno-associated virus 2 vectors is independent of the cellular heat-shock protein 90. *J. Biol. Chem.* 279 (13), 12714–12723.
- Zhong, L., Li, W., Li, Y., Zhao, W., Wu, J., Li, B., Maina, N., Bischof, D., Qing, K., Weigel-Kelley, K.A., Zolotukhin, I., Warrington Jr., K.H., Li, X., Slayton, W.B., Yoder, M.C., Srivastava, A., 2006. Evaluation of primitive murine hematopoietic stem and progenitor cell transduction in vitro and in vivo by recombinant adeno-associated virus vector serotypes 1 through 5. *Hum. Gene Ther.* 17 (3), 321–333.
- Zhong, L., Zhao, W., Wu, J., Li, B., Zolotukhin, S., Govindasamy, L., Agbandje-McKenna, M., Srivastava, A., 2007. A dual role of EGFR protein tyrosine kinase signaling in ubiquitination of AAV2 capsids and viral second-strand DNA synthesis. *Mol. Ther.* 15 (7), 1323–1330.
- Zhong, L., Li, B., Mah, C.S., Govindasamy, L., Agbandje-McKenna, M., Cooper, M.C., Herzog, R.W., Weigel Van-Aken, K.A., Hobbs, J.A., Zolotukhin, S., Muzyczka, N., Srivastava, A., 2008a. Next generation of recombinant adeno-associated virus 2 vectors: Point mutations in tyrosines lead to high-efficiency transduction at lower doses. *Proc. Natl. Acad. Sci. U. S. A.* 105 (22), 7827–7832.
- Zhong, L., Zhou, X., Li, Y., Qing, K., Xiao, X., Samulski, R.J., Srivastava, A., 2008b. Single-polarity recombinant adeno-associated virus 2 vector-mediated transgene expression in vitro and in vivo: mechanism of transduction. *Mol. Ther.* 16 (2), 290–295.