

Available online at www.sciencedirect.com**ScienceDirect**

Nuclear Physics B 906 (2016) 289–320

www.elsevier.com/locate/nuclphysb

Flavour symmetries in a renormalizable $SO(10)$ model

P.M. Ferreira^{a,b}, W. Grimus^{c,*}, D. Jurčiukonis^d, L. Lavoura^e^a ISEL—Instituto Superior de Engenharia de Lisboa, Instituto Politécnico de Lisboa, Portugal^b Centro de Física Teórica e Computacional, Faculdade de Ciências, Universidade de Lisboa, Portugal^c University of Vienna, Faculty of Physics, Boltzmannngasse 5, A-1090 Wien, Austria^d University of Vilnius, Institute of Theoretical Physics and Astronomy, A. Goštauto st. 12, Vilnius 01108, Lithuania^e CFTP, Instituto Superior Técnico, Universidade de Lisboa, 1049-001 Lisboa, Portugal

Received 21 October 2015; received in revised form 22 February 2016; accepted 5 March 2016

Available online 9 March 2016

Editor: Tommy Ohlsson

Abstract

In the context of a renormalizable supersymmetric $SO(10)$ Grand Unified Theory, we consider the fermion mass matrices generated by the Yukawa couplings to a $\mathbf{10} \oplus \mathbf{120} \oplus \mathbf{126}$ representation of scalars. We perform a complete investigation of the possibilities of imposing flavour symmetries in this scenario; the purpose is to reduce the number of Yukawa coupling constants in order to identify potentially predictive models. We have found that there are only 14 inequivalent cases of Yukawa coupling matrices, out of which 13 cases are generated by \mathbb{Z}_n symmetries, with suitable n , and one case is generated by a $\mathbb{Z}_2 \times \mathbb{Z}_2$ symmetry. A numerical analysis of the 14 cases reveals that only two of them—dubbed A and B in the present paper—allow good fits to the experimentally known fermion masses and mixings.

© 2016 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP³.

* Corresponding author.

E-mail addresses: pmmferreira@fc.ul.pt (P.M. Ferreira), walter.grimus@univie.ac.at (W. Grimus), darius.jurciukonis@tfai.vu.lt (D. Jurčiukonis), balio@cftp.tecnico.ulisboa.pt (L. Lavoura).

<http://dx.doi.org/10.1016/j.nuclphysb.2016.03.011>

0550-3213/© 2016 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP³.

1. Introduction

$SO(10)$ is a popular gauge group for the construction of Grand Unified Theories (GUTs). The reason is that its 16-plet accommodates at once all the chiral fields of one fermion family. Now [1,2],

$$(\mathbf{16} \otimes \mathbf{16})_S = \mathbf{10} \oplus \mathbf{126}, \quad (1a)$$

$$(\mathbf{16} \otimes \mathbf{16})_{AS} = \mathbf{120}, \quad (1b)$$

where the subscripts “S” and “AS” stand for, respectively, the symmetric and the antisymmetric parts of the tensor product. Therefore, in a renormalizable theory the scalars occurring in the Yukawa couplings belong solely to the irreducible representations (irreps) $\mathbf{10}$, $\overline{\mathbf{126}}$, and $\mathbf{120}$.¹ Previously, in the so-called “minimal supersymmetric $SO(10)$ GUT” (for an incomplete list of references see Refs. [3–5]) the $\mathbf{120}$ was absent. However, inconsistencies in the fit of the experimental masses and mixings of the fermions—in particular, a tension between the seesaw and GUT scales [6]—led to the inclusion of the 120-plet; the resulting theory has been called [7] the “new minimal supersymmetric $SO(10)$ GUT” (NMSGUT)—see Ref. [8] and the references therein.²

It has turned out that the NMSGUT, which contains three 16-plets of fermionic fields and one multiplet of scalars for each of the irreps in the right-hand sides of equations (1), is quite a successful theory and is capable of accommodating all the available data on the fermion masses and mixings, including the recent neutrino oscillation data [11,12]; this has been demonstrated by numerical fits [13].³ However, adding a 120-plet to the 10-plet and the 126-plet of scalars leads to a proliferation of parameters in the Yukawa couplings; one might want to restrict the number of parameters in order to obtain potentially predictive scenarios. Attempts in this direction have been made: in Ref. [15], texture zeros were placed in the mass matrices; in Ref. [16], a \mathbb{Z}_2 flavour symmetry has been imposed together with a CP symmetry; in Ref. [17], real Yukawa couplings were assumed and CP was broken solely by the imaginary vacuum expectation values (VEVs) of the $\mathbf{120}$.

In the present paper we pursue the approach of Ref. [16] by investigating all the possible flavour symmetries acting on the Yukawa couplings in the NMSGUT. We firstly perform a complete discussion by using only minimal assumptions; we thereby identify all the possible cases and their symmetry groups. Thereafter, all the cases are subjected to a numerical analysis in order to identify the viable ones. Partially anticipating our results, no non-Abelian flavour symmetry groups are permitted and there are 14 inequivalent cases, out of which 13 pertain to one-generator Abelian groups and only one case has a two-generator symmetry group $\mathbb{Z}_2 \times \mathbb{Z}_2$. However, the numerical analysis rules out almost all the cases, leaving only two viable ones which are compatible with the data on the fermion masses and mixings.

In section 2 we fix the notation, display the basic formulas needed for our investigation, and set forth our assumptions. In section 3 we list all the 14 cases. The results of the numerical analysis are presented in section 4. The conclusions of our work are given in section 5. The analysis of two specific problems that arise in family symmetry-furnished GUTs is deferred to Appendix A. The

¹ The representations $\mathbf{10}$ and $\mathbf{120}$ are self-conjugate.

² A completely different approach is $SO(10)$ GUT models in extra dimensions—see for instance Ref. [9] and the references therein—or with a hidden sector [10].

³ Note that skipping the $\overline{\mathbf{126}}$ of scalars does not allow for a good fit of even the charged-fermion sector alone [14].

discussion of the possibility of one further group generator is left to [Appendix B](#). [Appendix C](#) focuses on the derivation of some inequalities among the VEVs of the various $SO(10)$ scalar representations.

2. Notation, framework, and assumptions

The relevant fermion mass matrices are given by (see for instance Refs. [\[2,18\]](#))

$$M_d = k_d H + \kappa_d G + v_d F, \tag{2a}$$

$$M_u = k_u H + \kappa_u G + v_u F, \tag{2b}$$

$$M_\ell = k_\ell H + \kappa_\ell G - 3v_d F, \tag{2c}$$

$$M_D = k_u H + \kappa_D G - 3v_u F, \tag{2d}$$

where M_d , M_u , and M_ℓ are the mass matrices of the down-type quarks, the up-type quarks, and the charged leptons, respectively, while M_D is the neutrino Dirac mass matrix. The Yukawa-coupling matrices H , G , and F are associated with the scalar irreps **10**, **120**, and $\overline{\mathbf{126}}$, respectively. Those matrices have the (anti)symmetry properties

$$H^T = H, \tag{3a}$$

$$G^T = -G, \tag{3b}$$

$$F^T = F. \tag{3c}$$

The coefficients k_d , v_d , κ_d , and κ_ℓ are the VEVs of the Higgs doublet components in the respective $SO(10)$ scalar irreps which contribute to the Higgs doublet H_d of the Minimal Supersymmetric Standard Model (MSSM). The remaining coefficients— k_u , v_u , κ_u , and κ_D —refer to H_u . The light-neutrino mass matrix is obtained as

$$\mathcal{M}_\nu = M_L - M_D M_R^{-1} M_D^T \tag{4}$$

with

$$M_L = w_L F, \tag{5a}$$

$$M_R = w_R F, \tag{5b}$$

where w_L and w_R are the VEVs of scalar triplets of the Pati–Salam [\[19\]](#) group $SU(4)_c \times SU(2)_L \times SU(2)_R$, which are part of the scalar 126-plet of $SO(10)$. The first term in the right-hand side of equation [\(4\)](#) corresponds to the contribution of the type II seesaw mechanism [\[20\]](#) and the second term to the contribution of the type I seesaw mechanism [\[21\]](#). Thus,

$$\frac{w_R}{v_d} \mathcal{M}_\nu = \frac{w_L w_R}{v_d^2} M_d^F - M_D (M_d^F)^{-1} M_D^T, \tag{6}$$

where $M_d^F \equiv v_d F$ is the component of the down-type-quark mass matrix arising from the Yukawa coupling to the $\overline{\mathbf{126}}$ of scalars. One sees that

- a complex factor $w_L w_R / v_d^2$ parameterizes the strength of the type II seesaw contribution relative to the strength of the type I seesaw contribution; and
- the overall magnitude of the neutrino masses relative to the charged-fermion masses is parameterized by a dimensionless factor $|w_R / v_d|$.

The mass Lagrangian of the “light” fermions reads

$$\mathcal{L}_{\text{mass}} = -\bar{d}_L M_d d_R - \bar{u}_L M_u u_R - \bar{\ell}_L M_\ell \ell_R - \frac{1}{2} \bar{\nu}_L \mathcal{M}_\nu (\nu_L)^c + \text{H.c.}, \quad (7)$$

with $(\nu_L)^c = C \bar{\nu}_L^T$ being the charge-conjugate of ν_L . One diagonalizes the “Hermitian mass matrices” as

$$U_d^\dagger (M_d M_d^\dagger) U_d = \text{diag} (m_d^2, m_s^2, m_b^2), \quad (8a)$$

$$U_u^\dagger (M_u M_u^\dagger) U_u = \text{diag} (m_u^2, m_c^2, m_t^2), \quad (8b)$$

$$U_\ell^\dagger (M_\ell M_\ell^\dagger) U_\ell = \text{diag} (m_e^2, m_\mu^2, m_\tau^2), \quad (8c)$$

$$U_\nu^\dagger (\mathcal{M}_\nu \mathcal{M}_\nu^\dagger) U_\nu = \text{diag} (m_1^2, m_2^2, m_3^2), \quad (8d)$$

where the matrices $U_{d,u,\ell,\nu}$ are unitary and $|m_3^2 - m_1^2| \gg m_2^2 - m_1^2 > 0$. The fermion mixing matrices are then

$$V \equiv U_{\text{CKM}} = U_u^\dagger U_d, \quad (9a)$$

$$U_{\text{PMNS}} = U_\ell^\dagger U_\nu. \quad (9b)$$

The neutrino mass spectrum is dubbed “normal” if $m_3^2 > m_1^2$ and “inverted” otherwise.

We make the following assumptions:

- All three matrices H , F , and G are non-zero.
- $\det F \neq 0$.
- No generation decouples.

The second assumption is necessary for the type I seesaw mechanism. The third assumption is an experimental fact.

If the Lagrangian is invariant under a flavour symmetry \mathcal{S}_0 , then, due to the $SO(10)$ structure of the Yukawa couplings we obtain the following relations:

$$\mathcal{S}_0 : \begin{cases} W^T H W e^{i\alpha} = H, \\ W^T G W e^{i\beta} = G, \\ W^T F W e^{i\gamma} = F, \end{cases} \quad (10)$$

where W is the 3×3 unitary matrix which acts on the three matter 16-plets under \mathcal{S}_0 . Without loss of generality we take W to be diagonal. The scalar multiplets **10**, **120**, and $\overline{\mathbf{126}}$ transform under \mathcal{S}_0 with the phase factors $e^{i\alpha}$, $e^{i\beta}$, and $e^{i\gamma}$, respectively. (One of the phase factors may be absorbed into W .)

3. The 14 cases

3.1. A single flavour symmetry

A single symmetry transformation \mathcal{S}_0 leads to 13 inequivalent cases. We refrain from going through the tedious arguments leading to these cases; we merely list them instead. In the following, generic non-zero entries in the Yukawa coupling matrices are denoted “ \times ”. For each case, we also give the Abelian group through which the Yukawa-coupling matrices can be enforced.

Case A

$$\mathbb{Z}_2 : W = \text{diag}(+1, +1, -1), \quad e^{i\alpha} = +1, \quad e^{i\beta} = -1, \quad e^{i\gamma} = +1, \quad (11a)$$

$$H \sim \begin{pmatrix} \times & \times & 0 \\ \times & \times & 0 \\ 0 & 0 & \times \end{pmatrix}, \quad G \sim \begin{pmatrix} 0 & 0 & \times \\ 0 & 0 & \times \\ \times & \times & 0 \end{pmatrix}, \quad F \sim \begin{pmatrix} \times & \times & 0 \\ \times & \times & 0 \\ 0 & 0 & \times \end{pmatrix}. \quad (11b)$$

Case B

$$\mathbb{Z}_2 : W = \text{diag}(+1, +1, -1), \quad e^{i\alpha} = -1, \quad e^{i\beta} = -1, \quad e^{i\gamma} = +1, \quad (12a)$$

$$H \sim \begin{pmatrix} 0 & 0 & \times \\ 0 & 0 & \times \\ \times & \times & 0 \end{pmatrix}, \quad G \sim \begin{pmatrix} 0 & 0 & \times \\ 0 & 0 & \times \\ \times & \times & 0 \end{pmatrix}, \quad F \sim \begin{pmatrix} \times & \times & 0 \\ \times & \times & 0 \\ 0 & 0 & \times \end{pmatrix}. \quad (12b)$$

Case C

$$\mathbb{Z}_2 : W = \text{diag}(+1, -1, +1), \quad e^{i\alpha} = -1, \quad e^{i\beta} = +1, \quad e^{i\gamma} = +1, \quad (13a)$$

$$H \sim \begin{pmatrix} 0 & \times & 0 \\ \times & 0 & \times \\ 0 & \times & 0 \end{pmatrix}, \quad G \sim \begin{pmatrix} 0 & 0 & \times \\ 0 & 0 & 0 \\ \times & 0 & 0 \end{pmatrix}, \quad F \sim \begin{pmatrix} \times & 0 & \times \\ 0 & \times & 0 \\ \times & 0 & \times \end{pmatrix}. \quad (13b)$$

Case A₁

$$\mathbb{Z}_4 : W = \text{diag}(+1, -1, \pm i), \quad e^{i\alpha} = +1, \quad e^{i\beta} = \mp i, \quad e^{i\gamma} = -1, \quad (14a)$$

$$H \sim \begin{pmatrix} \times & 0 & 0 \\ 0 & \times & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad G \sim \begin{pmatrix} 0 & 0 & \times \\ 0 & 0 & 0 \\ \times & 0 & 0 \end{pmatrix}, \quad F \sim \begin{pmatrix} 0 & \times & 0 \\ \times & 0 & 0 \\ 0 & 0 & \times \end{pmatrix}. \quad (14b)$$

Case A'₁

$$U(1) : W = \text{diag}(1, e^{2i\sigma}, e^{i\sigma}), \quad e^{i\alpha} = 1, \quad e^{i\beta} = e^{-i\sigma}, \quad e^{i\gamma} = e^{-2i\sigma}, \quad (15a)$$

$$H \sim \begin{pmatrix} \times & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad G \sim \begin{pmatrix} 0 & 0 & \times \\ 0 & 0 & 0 \\ \times & 0 & 0 \end{pmatrix}, \quad F \sim \begin{pmatrix} 0 & \times & 0 \\ \times & 0 & 0 \\ 0 & 0 & \times \end{pmatrix}. \quad (15b)$$

Case A''₁

$$U(1) : W = \text{diag}(e^{2i\sigma}, 1, e^{i\sigma}), \quad e^{i\alpha} = 1, \quad e^{i\beta} = e^{-3i\sigma}, \quad e^{i\gamma} = e^{-2i\sigma}, \quad (16a)$$

$$H \sim \begin{pmatrix} 0 & 0 & 0 \\ 0 & \times & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad G \sim \begin{pmatrix} 0 & 0 & \times \\ 0 & 0 & 0 \\ \times & 0 & 0 \end{pmatrix}, \quad F \sim \begin{pmatrix} 0 & \times & 0 \\ \times & 0 & 0 \\ 0 & 0 & \times \end{pmatrix}. \quad (16b)$$

Case A₂

$$U(1) : W = \text{diag}(e^{i\sigma}, e^{-i\sigma}, 1), \quad e^{i\alpha} = 1, \quad e^{i\beta} = e^{-i\sigma}, \quad e^{i\gamma} = 1, \quad (17a)$$

$$H \sim \begin{pmatrix} 0 & \times & 0 \\ \times & 0 & 0 \\ 0 & 0 & \times \end{pmatrix}, \quad G \sim \begin{pmatrix} 0 & 0 & \times \\ 0 & 0 & 0 \\ \times & 0 & 0 \end{pmatrix}, \quad F \sim \begin{pmatrix} 0 & \times & 0 \\ \times & 0 & 0 \\ 0 & 0 & \times \end{pmatrix}. \quad (17b)$$

Case D₁

$$\mathbb{Z}_3: W = \text{diag}(\omega^2, \omega, 1), \quad e^{i\alpha} = 1, \quad e^{i\beta} = \omega, \quad e^{i\gamma} = \omega, \quad (18a)$$

$$H \sim \begin{pmatrix} 0 & \times & 0 \\ \times & 0 & 0 \\ 0 & 0 & \times \end{pmatrix}, \quad G \sim \begin{pmatrix} 0 & 0 & \times \\ 0 & 0 & 0 \\ \times & 0 & 0 \end{pmatrix}, \quad F \sim \begin{pmatrix} 0 & 0 & \times \\ 0 & \times & 0 \\ \times & 0 & 0 \end{pmatrix}. \quad (18b)$$

Case D₂

$$\mathbb{Z}_3: W = \text{diag}(\omega, \omega^2, 1), \quad e^{i\alpha} = 1, \quad e^{i\beta} = \omega^2, \quad e^{i\gamma} = \omega, \quad (19a)$$

$$H \sim \begin{pmatrix} 0 & \times & 0 \\ \times & 0 & 0 \\ 0 & 0 & \times \end{pmatrix}, \quad G \sim \begin{pmatrix} 0 & 0 & \times \\ 0 & 0 & 0 \\ \times & 0 & 0 \end{pmatrix}, \quad F \sim \begin{pmatrix} \times & 0 & 0 \\ 0 & 0 & \times \\ 0 & \times & 0 \end{pmatrix}. \quad (19b)$$

Case D₃

$$\mathbb{Z}_3: W = \text{diag}(\omega, 1, \omega^2), \quad e^{i\alpha} = 1, \quad e^{i\beta} = 1, \quad e^{i\gamma} = \omega, \quad (20a)$$

$$H \sim \begin{pmatrix} 0 & 0 & \times \\ 0 & \times & 0 \\ \times & 0 & 0 \end{pmatrix}, \quad G \sim \begin{pmatrix} 0 & 0 & \times \\ 0 & 0 & 0 \\ \times & 0 & 0 \end{pmatrix}, \quad F \sim \begin{pmatrix} \times & 0 & 0 \\ 0 & 0 & \times \\ 0 & \times & 0 \end{pmatrix}. \quad (20b)$$

Case D'₁

$$U(1): W = \text{diag}(e^{-i\sigma}, e^{i\sigma}, e^{3i\sigma}), \quad e^{i\alpha} = 1, \quad e^{i\beta} = e^{-2i\sigma}, \quad e^{i\gamma} = e^{-2i\sigma}, \quad (21a)$$

$$H \sim \begin{pmatrix} 0 & \times & 0 \\ \times & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad G \sim \begin{pmatrix} 0 & 0 & \times \\ 0 & 0 & 0 \\ \times & 0 & 0 \end{pmatrix}, \quad F \sim \begin{pmatrix} 0 & 0 & \times \\ 0 & \times & 0 \\ \times & 0 & 0 \end{pmatrix}. \quad (21b)$$

Case D'₂

$$U(1): W = \text{diag}(e^{i\sigma}, e^{-i\sigma}, e^{3i\sigma}), \quad e^{i\alpha} = 1, \quad e^{i\beta} = e^{-4i\sigma}, \quad e^{i\gamma} = e^{-2i\sigma}, \quad (22a)$$

$$H \sim \begin{pmatrix} 0 & \times & 0 \\ \times & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad G \sim \begin{pmatrix} 0 & 0 & \times \\ 0 & 0 & 0 \\ \times & 0 & 0 \end{pmatrix}, \quad F \sim \begin{pmatrix} \times & 0 & 0 \\ 0 & 0 & \times \\ 0 & \times & 0 \end{pmatrix}. \quad (22b)$$

Case D'₃

$$U(1): W = \text{diag}(e^{i\sigma}, e^{3i\sigma}, e^{-i\sigma}), \quad e^{i\alpha} = 1, \quad e^{i\beta} = 1, \quad e^{i\gamma} = e^{-2i\sigma}, \quad (23a)$$

$$H \sim \begin{pmatrix} 0 & 0 & \times \\ 0 & 0 & 0 \\ \times & 0 & 0 \end{pmatrix}, \quad G \sim \begin{pmatrix} 0 & 0 & \times \\ 0 & 0 & 0 \\ \times & 0 & 0 \end{pmatrix}, \quad F \sim \begin{pmatrix} \times & 0 & 0 \\ 0 & 0 & \times \\ 0 & \times & 0 \end{pmatrix}. \quad (23b)$$

In equations (18a), (19a), and (20a) $\omega \equiv \exp(\pm i2\pi/3)$.

We note that only case A had been discussed earlier, in Ref. [16]. Cases A₁ and A₂ have Yukawa-coupling matrices which are restrictions (*i.e.* they contain extra zero matrix elements)

of those of case A; cases A'_1 and A''_1 have Yukawa-coupling matrices which are more restrictive than those of case A_1 .

We demonstrate in [Appendix A](#) that the scalar potential of the NMSGUT can consistently be modified in order to incorporate the \mathbb{Z}_2 symmetries present in cases A, B and C.

3.2. A second flavour symmetry

The list of 13 cases in the previous subsection does not necessarily comprise all the Yukawa-coupling matrices obtainable through flavour symmetries, because in each of those 13 cases either one or more further symmetry transformations might be operative and lead to more restrictive Yukawa-coupling matrices and thus to new cases. Let us denote a generic further symmetry transformation, different from \mathcal{S}_0 of equation (10), by \mathcal{S}_1 :

$$\mathcal{S}_1 : \begin{cases} X^T H X e^{i\alpha_1} = H, \\ X^T G X e^{i\beta_1} = G, \\ X^T F X e^{i\gamma_1} = F. \end{cases} \tag{24}$$

In principle, the symmetry \mathcal{S}_1 might either commute or not commute with \mathcal{S}_0 . However, as shown in [Appendix B](#), by using our assumptions of section 2 one may demonstrate that X always commutes with W , *i.e.* that \mathcal{S}_1 commutes with \mathcal{S}_0 . Even more surprisingly, only one new case ensues, which we denote by the letter E and is a subcase of both case A and case C^4 :

Case E

$$\mathbb{Z}_2^{(1)} : W = \text{diag}(+1, +1, -1), \quad e^{i\alpha} = +1, \quad e^{i\beta} = -1, \quad e^{i\gamma} = +1, \tag{25a}$$

$$\mathbb{Z}_2^{(2)} : X = \text{diag}(+1, -1, +1), \quad e^{i\alpha_1} = -1, \quad e^{i\beta_1} = +1, \quad e^{i\gamma_1} = +1, \tag{25b}$$

$$H \sim \begin{pmatrix} 0 & \times & 0 \\ \times & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad G \sim \begin{pmatrix} 0 & 0 & \times \\ 0 & 0 & 0 \\ \times & 0 & 0 \end{pmatrix}, \quad F \sim \begin{pmatrix} \times & 0 & 0 \\ 0 & \times & 0 \\ 0 & 0 & \times \end{pmatrix}. \tag{25c}$$

Note that $\mathbb{Z}_2^{(1)}$ is the symmetry (11a) of case A while $\mathbb{Z}_2^{(2)}$ is the symmetry (13a) of case C.

There are no possible cases for a flavour group with three or more generators.

3.3. Summary

From the assumptions stated in section 2 we have obtained the following results:

- There are 14 inequivalent cases.
- All the cases except E can be obtained from a single flavour symmetry transformation.
- The flavour groups with one generator are the cyclic groups \mathbb{Z}_2 (in the cases A, B, and C), \mathbb{Z}_3 (in the cases D_k with $k = 1, 2, 3$), and \mathbb{Z}_4 (in case A_1). The remaining cases have a $U(1)$ symmetry.⁵

⁴ It is also a subcase of case B, as can be seen when one interchanges the first and third generations in the matrices of equations (12).

⁵ This $U(1)$ must be broken explicitly by the scalar potential, which we did not consider here, lest a Goldstone boson arises. Therefore, a full model will have a suitable cyclic symmetry group instead of $U(1)$.

Table 1
The number of parameters in the Hermitian mass matrices for each case.

Cases	A	B	C	A_1, D_k
# parameters in the $M_x M_x^\dagger$ for $x = d, \ell, u$	13 moduli and 10 phases	11 moduli and 7 phases	10 moduli and 6 phases	9 moduli and 5 phases
# extra parameters in $\mathcal{M}_\nu \mathcal{M}_\nu^\dagger$	3 moduli and 2 phases	3 moduli and 2 phases	3 moduli and 2 phases	3 moduli and 2 phases

- In case E there are two symmetry transformations which commute with each other; the flavour group is $\mathbb{Z}_2 \times \mathbb{Z}_2$.
- Our scenario does not admit non-Abelian flavour groups.

4. Fitting the cases to the data

In this section we report on our numerical study of cases A, B, C, A_1 , and D_k ($k = 1, 2, 3$). We have not studied the cases A'_1 and A''_1 because they are restrictions of case A_1 and we have found that case is unable to fit the data well (details will be given later). Analogously, the cases D'_k are restrictions of the cases D_k ; since we have found that the cases D_k do not work well, we did not need to bother with the cases D'_k . Finally, case E is a restriction of case C (and also of cases A and B); since case C is unable even to correctly fit the charged-fermion masses, case E can be discarded outright.

We did not attempt to fit case A_2 because we knew beforehand that such an attempt would be unsuccessful. Indeed, case A_2 yields M_d and M_u of the Fritzsch form [22], which has long been known to be unable to simultaneously fit the quark masses and the CKM matrix.

4.1. Parameter counting

In order to get a feeling for the ability for fitting the data that each case ought to have, it is instructive to count the number of parameters in each of the cases—see Table 1. For instance, in case A_1 the charged-fermion mass matrices may be written, after adequate rephasings,

$$M_d = \begin{pmatrix} a & 0 & f e^{i\theta_2} \\ 0 & c e^{i\theta_1} & b \\ -f e^{i\theta_2} & b & d \end{pmatrix}, \quad (26a)$$

$$M_\ell = \begin{pmatrix} 3a & 0 & g e^{i\theta_5} \\ 0 & c e^{i\theta_1} & 3b \\ -g e^{i\theta_5} & 3b & d \end{pmatrix}, \quad (26b)$$

$$M_u = \begin{pmatrix} ta & 0 & l e^{i\theta_4} \\ 0 & r c e^{i(\theta_1+\theta_3)} & tb \\ -l e^{i\theta_4} & tb & r d e^{i\theta_3} \end{pmatrix}, \quad (26c)$$

with five phases $\theta_{1,2,3,4,5}$ and nine real and non-negative parameters (“moduli”) $a, b, c, d, f, g, l, t \equiv |v_u/v_d|$, and $r \equiv |k_u/k_d|$. Moreover, the neutrino mass matrix is

$$\mathcal{M}_\nu = \left| \frac{v_d}{w_R} \right| \begin{pmatrix} Ca & -(rch/b) e^{i(\theta_1+\theta_3)} & 0 \\ -(rch/b) e^{i(\theta_1+\theta_3)} & 6rct e^{i(\theta_1+\theta_3)} & Cb - (r^2 cd/b) e^{i(\theta_1+2\theta_3)} \\ 0 & Cb - (r^2 cd/b) e^{i(\theta_1+2\theta_3)} & 6rt d e^{i\theta_3} - h^2/a \end{pmatrix}, \quad (27)$$

viz. it contains two extra complex parameters C and h , plus the real parameter $|v_d/w_R|$, making an extra three moduli and two phases.

One sees in Table 1 that $\mathcal{M}_\nu \mathcal{M}_\nu^\dagger$ always contains three moduli and two phases beyond the parameters which appear in the charged-fermion Hermitian mass matrices. It is easy to understand the reasons for that: one extra complex parameter originates in κ_D of equation (2d); another complex parameter originates in $w_L w_R / v_d^2$ in the right-hand side of equation (6); and there is an extra modulus $|w_R/v_d|$ in the left-hand side of equation (6).⁶

Case A is the one that has most parameters, hence most degrees of freedom, in the mass matrices. In Ref. [16] that case has been numerically studied under some restrictive assumptions; we have repeated that study under the same restrictive assumptions, but using the updated values for the charged-fermion masses given in Ref. [23].

The restriction of case A analyzed in Ref. [16] contains 13 moduli and 6 phases in the $M_x M_x^\dagger$ ($x = d, \ell, u$), plus an extra two moduli and one phase in $\mathcal{M}_\nu \mathcal{M}_\nu^\dagger$. The original “minimal supersymmetric $SO(10)$ GUT” [3] has 11 moduli and 8 phases in the $M_x M_x^\dagger$, plus an extra two moduli and one phase in $\mathcal{M}_\nu \mathcal{M}_\nu^\dagger$. We see that both those models are comparable to our case B in their numbers of parameters.

The $M_x M_x^\dagger$ are supposed to be able to fit 13 observables: the nine charged-fermion masses and the four observables in the CKM matrix. One must take into account that phases usually do not help much in fitting observables; the moduli are most relevant. Additionally, if one also takes into account $\mathcal{M}_\nu \mathcal{M}_\nu^\dagger$, then we have to fit five parameters more—the three lepton mixing angles, the ratio $r_{\text{solar}}^2 \equiv (m_2^2 - m_1^2) / |m_3^2 - m_1^2|$, and $|m_3^2 - m_1^2|$ itself. We have used the fixed value $|m_3^2 - m_1^2| = 2.5 \times 10^{-15} \text{ MeV}^2$, which just allows us to determine the overall scale of \mathcal{M}_ν , *viz.* $|v_d/w_R|$.

4.2. χ^2 function

In order to test the viability of each case, and to find adequate numerical values for its parameters, we construct a χ^2 function

$$\chi^2(x) = \sum_{i=1}^n \left\{ H[f_i(x) - \bar{O}_i] \left(\frac{f_i(x) - \bar{O}_i}{\delta_+ O_i} \right)^2 + H[\bar{O}_i - f_i(x)] \left(\frac{\bar{O}_i - f_i(x)}{\delta_- O_i} \right)^2 \right\}, \tag{28}$$

where n is the total number of observables (masses and mixing parameters) to be fitted. In equation (28), H is the Heaviside step function, \bar{O}_i is the central value of each observable O_i , $\delta_\pm O_i$ are the upper and lower errors of that observable, and $f_i(x)$ is the value of that observable, in any given case, when the parameters of that case have the values $x = \{x_\alpha\}$. The data are fitted by minimizing $\chi^2(x)$ with respect to the x_α .

We have used the mean values \bar{O} and the errors $\delta_\pm O$ given in Tables 2–4. We have taken the charged-fermion masses in Table 2, which are renormalized at $M_{\text{GUT}} = 2 \times 10^{16} \text{ GeV}$, from the last column of Table V of Ref. [23].⁷ These are values computed using the renormalization-group equations of the MSSM with $\tan \beta = 10$; we leave it for some later, more detailed study the task

⁶ Note that the overall phase of \mathcal{M}_ν is unphysical.

⁷ For other determinations of the values of the running quark and lepton masses, evolved from the electroweak scale to the GUT scale through the renormalization group of the MSSM, see Ref. [24].

Table 2

The values of the charged-fermion masses used in our fits.

observable	m_d/MeV	m_s/MeV	m_b/MeV
$\bar{O}_{-\delta-O}^{+\delta+O}$	$0.70^{+0.31}_{-0.30}$	13^{+4}_{-4}	790^{+40}_{-40}
observable	m_e/MeV	m_μ/MeV	m_τ/MeV
$\bar{O}_{-\delta-O}^{+\delta+O}$	$0.283755495^{+2.4 \times 10^{-8}}_{-2.5 \times 10^{-8}}$	$59.9033617^{+5.4 \times 10^{-6}}_{-5.4 \times 10^{-6}}$	$1021.95^{+0.11}_{-0.12}$
observable	m_u/MeV	m_c/MeV	m_t/MeV
$\bar{O}_{-\delta-O}^{+\delta+O}$	$0.49^{+0.20}_{-0.17}$	236^{+37}_{-36}	92200^{+9600}_{-7800}

Table 3

The values of the CKM-matrix observables used in our fits.

observable	$ V_{12} $	$ V_{13} $	$ V_{23} $	$10^5 J$
$\bar{O}_{-\delta-O}^{+\delta+O}$	$0.22536^{+0.00183}_{-0.00183}$	$0.00355^{+0.00045}_{-0.00045}$	$0.0414^{+0.0036}_{-0.0036}$	$3.06^{+0.63}_{-0.60}$

Table 4

The values of the neutrino and lepton-mixing observables used in our fits. “NH” refers to a normal neutrino mass spectrum and “IH” to an inverted one.

observable	r_{solar}^2 (NH)	$\sin^2 \theta_{12}$ (NH)	$\sin^2 \theta_{13}$ (NH)	$\sin^2 \theta_{23}$ (NH)
$\bar{O}_{-\delta-O}^{+\delta+O}$	$0.0306^{+0.0050}_{-0.0038}$	$0.323^{+0.052}_{-0.045}$	$0.0234^{+0.0060}_{-0.0057}$	$0.567^{+0.076}_{-0.175}$
observable	r_{solar}^2 (IH)	$\sin^2 \theta_{12}$ (IH)	$\sin^2 \theta_{13}$ (IH)	$\sin^2 \theta_{23}$ (IH)
$\bar{O}_{-\delta-O}^{+\delta+O}$	$0.0319^{+0.0053}_{-0.0039}$	$0.323^{+0.052}_{-0.045}$	$0.0240^{+0.0057}_{-0.0057}$	$0.573^{+0.067}_{-0.172}$

of fitting the data for other values of $\tan \beta$. The values of the CKM mixing angles in Table 3 are low-energy values and were taken from equation (12.27) of Ref. [25]; we have multiplied the error bars given in that equation by a factor of three in order to obtain adequately large intervals. The values in Table 4 are the 3σ intervals given for each observable in Ref. [11].

In order to assess the fitting ability of each case, we have firstly attempted to fit only the charged-fermion masses (nine observables, given in Table 2), secondly the charged-fermion masses together with the CKM matrix (four more observables, given in Table 3), and, finally, all that together with the neutrino masses and the PMNS matrix (four observables more, given in Table 4). The total χ^2 function is thus the sum of three terms:

$$\chi_{\text{total}}^2 = \chi_{\text{masses}}^2 + \chi_{\text{CKM}}^2 + \chi_{\nu}^2. \quad (29)$$

For the neutrino masses, we have analysed both possibilities of a normal or inverted neutrino mass spectrum; indeed, for each set of values for the parameters x , we have computed the eigenvalues of $\mathcal{M}_\nu \mathcal{M}_\nu^*$ and thereby determined the type of neutrino mass spectrum; we have then chosen accordingly the input values in the computation of the function χ_{ν}^2 .

In some cases we have not been able to find a reasonably small value of χ_{masses}^2 alone; in those cases, further analysis by considering χ_{CKM}^2 and χ_{ν}^2 made no sense. Similarly, in some other cases a sufficiently low value of $\chi_{\text{masses}}^2 + \chi_{\text{CKM}}^2$ could not be achieved, so we did not have to consider χ_{ν}^2 . Finally, even when χ_{total}^2 could be correctly fitted, we still had to check whether $|w_R/v_d|$ turned out in the right range. Indeed, since v_d must be of order the Fermi scale 100 GeV

and w_R must be of order the grand-unification scale 10^{16} GeV, we must require $|w_R/v_d|$ to be 10^{14} or even larger. We had to check some other inequalities, the exposition of which we defer to section 4.4.

4.3. Numerical method

The minimization of $\chi^2(x)$ is a difficult task because the various parameters x_α may differ by several orders of magnitude and because there always is a large number of local minima. We have spent much time in the numerical analysis trying to find absolute minima; this has involved various fitting options and restrictions of the parameters for each particular case. Still, we cannot be 100% sure that we have found the absolute minimum for all cases—the possibility remains that a better solution exists somewhere in parameter space.

For the numerical minimization of the χ^2 functions we have employed the Differential Evolution (DE) algorithm. This is a stochastic algorithm that exploits a population of potential solutions in order to effectively probe the parameter space. It was first introduced in Ref. [26] and it has been modified several times since then.

The effectiveness of the DE algorithm strongly depends on control parameters. We have performed preliminary tests in order to hand-tune the appropriate ranges for the control parameters in each case. Also, in the χ^2 function of equation (28), we have modified the errors $\delta_\pm O_i$ randomly (within the range of magnitude of the true errors) according to the behaviour of the fits; we have thus been able to test, for each case, more local minima—defined as the points where the minimization algorithm converges—and to find the minima closer to the global minimum.

All the numerical calculations were implemented by using the programming language Fortran.

4.4. Case B

4.4.1. Theoretical treatment

We choose a weak basis in which the Yukawa-coupling matrix F is diagonal. After an interchange of the first and third generations,

$$k_d H = \begin{pmatrix} 0 & d & h \\ d & 0 & 0 \\ h & 0 & 0 \end{pmatrix}, \quad \kappa_d G = \begin{pmatrix} 0 & f & g \\ -f & 0 & 0 \\ -g & 0 & 0 \end{pmatrix}, \quad v_d F = \begin{pmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{pmatrix}. \quad (30)$$

Without loss of generality, we assume the parameters a , b , and c to be non-negative real. Then the mass matrices are given by

$$M_d = \begin{pmatrix} a & d + f & h + g \\ d - f & b & 0 \\ h - g & 0 & c \end{pmatrix}, \quad (31a)$$

$$M_\ell = \begin{pmatrix} -3a & d + (\kappa_\ell/\kappa_d) f & h + (\kappa_\ell/\kappa_d) g \\ d - (\kappa_\ell/\kappa_d) f & -3b & 0 \\ h - (\kappa_\ell/\kappa_d) g & 0 & -3c \end{pmatrix}, \quad (31b)$$

$$M_u = \begin{pmatrix} (v_u/v_d) a & (k_u/k_d) d + (\kappa_u/\kappa_d) f & (k_u/k_d) h + (\kappa_u/\kappa_d) g \\ (k_u/k_d) d - (\kappa_u/\kappa_d) f & (v_u/v_d) b & 0 \\ (k_u/k_d) h - (\kappa_u/\kappa_d) g & 0 & (v_u/v_d) c \end{pmatrix}, \quad (31c)$$

$$M_D = \begin{pmatrix} -3(v_u/v_d)a & (k_u/k_d)d + (\kappa_D/\kappa_d)f & (k_u/k_d)h + (\kappa_D/\kappa_d)g \\ (k_u/k_d)d - (\kappa_D/\kappa_d)f & -3(v_u/v_d)b & 0 \\ (k_u/k_d)h - (\kappa_D/\kappa_d)g & 0 & -3(v_u/v_d)c \end{pmatrix}. \quad (31d)$$

We rewrite the mass matrices (31) as

$$M_d = \begin{pmatrix} a & k_1 & k_3 \\ k_2 & b & 0 \\ k_4 & 0 & c \end{pmatrix}, \quad (32a)$$

$$M_\ell = \begin{pmatrix} -3a & k_5 & k_7 \\ k_6 & -3b & 0 \\ k_8 & 0 & -3c \end{pmatrix}, \quad (32b)$$

$$M_u = \begin{pmatrix} ta & k_9 & k_{11} \\ k_{10} & tb & 0 \\ k_{12} & 0 & tc \end{pmatrix}, \quad (32c)$$

$$M_D = \begin{pmatrix} -3ta & k_{13} & k_{15} \\ k_{14} & -3tb & 0 \\ k_{16} & 0 & -3tc \end{pmatrix}, \quad (32d)$$

where $t \equiv v_u/v_d$. The $k_{1,2,\dots,16}$ are not all independent. We choose $k_{1,2,3,4,5,9,10,13}$ as parameters, while

$$k_6 = k_1 + k_2 - k_5, \quad (33a)$$

$$k_7 = \frac{k_1k_4 + k_3k_5 - k_2k_3 - k_4k_5}{k_1 - k_2}, \quad (33b)$$

$$k_8 = \frac{k_1k_3 + k_4k_5 - k_2k_4 - k_3k_5}{k_1 - k_2}, \quad (33c)$$

$$k_{11} = \frac{(k_1k_3 - k_2k_4)k_9 + (k_1k_4 - k_2k_3)k_{10}}{k_1^2 - k_2^2}, \quad (33d)$$

$$k_{12} = \frac{(k_1k_3 - k_2k_4)k_{10} + (k_1k_4 - k_2k_3)k_9}{k_1^2 - k_2^2}, \quad (33e)$$

$$k_{14} = k_9 + k_{10} - k_{13}, \quad (33f)$$

$$k_{15} = \frac{k_{13}(k_3 - k_4)}{k_1 - k_2} + \frac{(k_9 + k_{10})(k_1k_4 - k_2k_3)}{k_1^2 - k_2^2}, \quad (33g)$$

$$k_{16} = \frac{k_{13}(k_4 - k_3)}{k_1 - k_2} + \frac{(k_9 + k_{10})(k_1k_3 - k_2k_4)}{k_1^2 - k_2^2}. \quad (33h)$$

From equations (6), (30), and (32d) it is easy to compute

$$\frac{w_R}{v_d} \mathcal{M}_\nu = \begin{pmatrix} w_L w_R \\ v_d^2 \end{pmatrix} - 9t^2 \begin{pmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{pmatrix} - \begin{pmatrix} k_{13}^2/b + k_{15}^2/c & -3t(k_{13} + k_{14}) & -3t(k_{15} + k_{16}) \\ -3t(k_{13} + k_{14}) & k_{14}^2/a & k_{14}k_{16}/a \\ -3t(k_{15} + k_{16}) & k_{14}k_{16}/a & k_{16}^2/a \end{pmatrix}. \quad (34)$$

Next, we multiply M_u and \mathcal{M}_ν by phase factors $\exp(-i \arg t)$, defining

$$M'_u \equiv \exp(-i \arg t) M_u, \tag{35a}$$

$$\mathcal{M}'_\nu \equiv \exp(-2i \arg t) \mathcal{M}_\nu. \tag{35b}$$

This phase change leads to the redefinitions

$$k'_p \equiv k_p \exp(-i \arg t) \quad \text{for } p = 9, \dots, 16. \tag{36}$$

Crucially, equations (33) remain valid when using the k'_p instead of the k_p for $p = 9, \dots, 16$. One obtains

$$M'_u = \begin{pmatrix} |t|a & k'_9 & k'_{11} \\ k'_{10} & |t|b & 0 \\ k'_{12} & 0 & |t|c \end{pmatrix}, \tag{37a}$$

$$\begin{aligned} \frac{w_R}{v_d} \mathcal{M}'_\nu &= (\hat{C} - 9|t|^2) \begin{pmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{pmatrix} \\ &- \begin{pmatrix} k'_{13}{}^2/|b| + k'_{15}{}^2/|c| & -3|t|(k'_{13} + k'_{14}) & -3|t|(k'_{15} + k'_{16}) \\ -3|t|(k'_{13} + k'_{14}) & k'_{14}{}^2/|a| & k'_{14}k'_{16}/|a| \\ -3|t|(k'_{15} + k'_{16}) & k'_{14}k'_{16}/|a| & k'_{16}{}^2/|a| \end{pmatrix}, \end{aligned} \tag{37b}$$

where $\hat{C} \equiv (w_L w_R / v_d^2) \exp(-2i \arg t)$.

In equations (32a), (32b), and (37) one observes that the mass matrices of case B may be parameterized through five real quantities $a, b, c, |t|$, and $|w_R/v_d|$, plus nine complex parameters $k_{1,2,3,4,5}, k'_{9,10,13}$, and \hat{C} . This justifies the third column of Table 1.

4.4.2. Inequalities

We fix the scale $|w_R/v_d|$ in the left-hand side of equation (37b) by requiring the difference of the squared neutrino masses $|m_3^2 - m_1^2|$ to be equal to the atmospheric mass scale $2.5 \times 10^{-3} \text{ eV}^2$. Afterwards, we compute $|v_d|$ by identifying $|w_R|$ with the unification scale $M_{\text{GUT}} = 2 \times 10^{16} \text{ GeV}$. Finally, we calculate $|v_u| = |t v_d|$ from the value of the parameter $|t|$ of the fit.

In the supersymmetric GUT that we envisage there is only one scalar doublet with hypercharge $+1/2$, viz. H_d , at the Fermi mass scale; there is also only one scalar doublet with hypercharge $-1/2$, viz. H_u , at that scale. Those two doublets have VEVs

$$\langle H_d^0 \rangle_0 = \frac{174 \text{ GeV}}{\sqrt{1 + \tan^2 \beta}} \quad \text{and} \quad \langle H_u^0 \rangle_0 = \frac{(174 \text{ GeV}) \tan \beta}{\sqrt{1 + \tan^2 \beta}}, \tag{38}$$

respectively, where $\tan \beta = 10$ in our fit. According to the inequalities (C.11),

$$(\langle H_d^0 \rangle_0)^2 \geq |v_d|^2 + |k_d|^2 + |\kappa_d|^2 + \frac{1}{3} |\kappa_\ell|^2, \tag{39a}$$

$$(\langle H_u^0 \rangle_0)^2 \geq |v_u|^2 + |k_u|^2 + |\kappa_u|^2 + \frac{1}{3} |\kappa_D|^2. \tag{39b}$$

Therefore, we have first of all to enforce the inequalities

$$\frac{(2 \times 10^{16} \text{ GeV})^2}{|w_R/v_d|^2} < \frac{(174 \text{ GeV})^2}{1 + \tan^2 \beta}, \tag{40a}$$

$$|t|^2 \frac{(2 \times 10^{16} \text{ GeV})^2}{|w_R/v_d|^2} < \frac{(174 \text{ GeV})^2 \tan^2 \beta}{1 + \tan^2 \beta}, \tag{40b}$$

on our fits. The inequality (40a) reads $|w_R/v_d| > 1.155 \times 10^{15}$, which is a quite useful lower bound.

Our fit fixes

$$\frac{\kappa_\ell}{\kappa_d} = \frac{k_5 - k_6}{k_1 - k_2}, \tag{41a}$$

$$\frac{\kappa_u}{\kappa_d} = \frac{k_9 - k_{10}}{k_1 - k_2}, \tag{41b}$$

$$\frac{\kappa_D}{\kappa_d} = \frac{k_{13} - k_{14}}{k_1 - k_2}, \tag{41c}$$

$$\frac{k_u}{k_d} = \frac{k_9 + k_{10}}{k_1 + k_2}. \tag{41d}$$

Therefore, from the inequalities (39),

$$|k_d|^2 + \left(1 + \frac{1}{3} \left| \frac{k_5 - k_6}{k_1 - k_2} \right|^2\right) |\kappa_d|^2 \leq \left(\langle H_d^0 \rangle_0 \right)^2 - |v_d|^2, \tag{42a}$$

$$\left| \frac{k'_9 + k'_{10}}{k_1 + k_2} \right|^2 |k_d|^2 + \frac{|k'_9 - k'_{10}|^2 + (1/3) |k'_{13} - k'_{14}|^2}{|k_1 - k_2|^2} |\kappa_d|^2 \leq \left(\langle H_u^0 \rangle_0 \right)^2 - |v_u|^2. \tag{42b}$$

We must now face the additional fact that the Yukawa couplings cannot be too large, lest the theory ceases to be perturbative and/or Landau poles arise in the Yukawa couplings. Let $y > 0$ denote the maximum value that we accept for the absolute value of the Yukawa couplings; we may take y somewhere between 1 and 10. From equations (30),

$$\max(|d|, |h|) < |k_d| y, \tag{43a}$$

$$\max(|f|, |g|) < |\kappa_d| y, \tag{43b}$$

$$\max(a, b, c) < |v_d| y. \tag{43c}$$

We have directly enforced the inequality (43c) on our fits. The other two inequalities (43) may be put together with the inequalities (42) to derive

$$\max\left(|k_1 + k_2|^2, |k_3 + k_4|^2\right) + \left(1 + \frac{1}{3} \left| \frac{k_5 - k_6}{k_1 - k_2} \right|^2\right) \max\left(|k_1 - k_2|^2, |k_3 - k_4|^2\right) \leq 4 \left[\left(\langle H_d^0 \rangle_0 \right)^2 - |v_d|^2 \right] y^2, \tag{44a}$$

$$\left| \frac{k'_9 + k'_{10}}{k_1 + k_2} \right|^2 \max\left(|k_1 + k_2|^2, |k_3 + k_4|^2\right) + \left(|k'_9 - k'_{10}|^2 + \frac{1}{3} |k'_{13} - k'_{14}|^2 \right) \times \frac{\max\left(|k_1 - k_2|^2, |k_3 - k_4|^2\right)}{|k_1 - k_2|^2} \leq 4 \left[\left(\langle H_u^0 \rangle_0 \right)^2 - |v_u|^2 \right] y^2. \tag{44b}$$

To summarize, we have enforced on our fits the inequalities (40), (43c), and (44).

Table 5
The values of the parameters for out best fit of case B.

Parameter	Value
a/MeV	219.850545793720272
b/MeV	0.561919252512016
c/MeV	28.64031991278612
$ t $	1.558686846443802
k_1/MeV	$106.613768172192835 \exp(i 2.726661945096518)$
k_2/MeV	$3.214360308597388 \exp(i 5.73665831290545)$
k_3/MeV	$750.563494049026872 \exp(i 4.735747016077402)$
k_4/MeV	$20.603622366818627 \exp(i 1.445003798120007)$
k_5/MeV	$10.49565466331894 \exp(i 4.813783368633092)$
k'_9/MeV	$12964.825027004273579 \exp(i 3.963802982387908)$
k'_{10}/MeV	$19.353350623796356 \exp(i 5.971066682817606)$
k'_{13}/MeV	$22682.297777225823666 \exp(i 5.075844560968867)$
\hat{C}	$3291007.008905897848 \exp(i 2.868704037387841)$
$ w_R/v_d $	1.67257×10^{15}

4.4.3. Fit

We have found that case B is able to fit *perfectly* all the observables. This is true irrespective of whether the neutrino mass spectrum is normal or inverted. However, when the neutrino mass spectrum is inverted some of the inequalities in the previous subsection always turn out to be violated; this happens because either $|w_R/v_d| < 10^{15}$ is too small or $|t| > 300$ is so large that the inequality (44b) ends up being violated.

For a normal neutrino mass spectrum, on the other hand, there are fits in which all the inequalities are observed. In Table 5 we give the values of the mass-matrix parameters that lead to the best fit which we have been able to achieve. The value of χ^2_{total} for this fit is smaller than 10^{-3} , *i.e.*, for all practical purposes, it is zero. The smallest neutrino mass for this fit is $m_1 \approx 0.006$ eV, while $m_1 + m_2 + m_3 \approx 0.07$ eV.

It is interesting to observe in Table 5 that the best fit is achieved for a very large value of $|\hat{C}| \sim 10^6$, meaning that the type-II seesaw mechanism dominates over the type-I. For this fit, the matrices U_d and U_u are almost diagonal, with U_d mostly identical with U_{CKM} . On the other hand, U_ν is almost completely a rotation between the first two generations, while U_ℓ is largely, but not exclusively, a rotation between the second and third generations; both rotations are almost maximal.

Since very perfect fits can be obtained in case B, we suspect that this case has too many degrees of freedom and has little or no predictive power. However, since such a study is very time-consuming, we leave it for later investigation.

4.5. Non-viable cases

We have found that all the cases except cases A and B either fail to fit the observables adequately or give a much too low value for $|w_R/v_d|$. (For us, an acceptable fit is one in which all the observables simultaneously are within their ranges in Tables 2–4.) Indeed, case C even fails to adequately fit the charged-fermion masses alone, while cases A₁, D₂, and D₃ are unable to

Table 6

Description of the minimization results for the cases that fail. The pull is defined as $H[f_i(x) - \bar{O}_i] [f_i(x) - \bar{O}_i] / \delta_+ O_i + H[\bar{O}_i - f_i(x)] [f_i(x) - \bar{O}_i] / \delta_- O_i$.

Case	χ^2 of best fit	Pulls larger than one in absolute value	Remarks
A ₁	$\chi_{\text{masses}}^2 \sim 10^{-6}$		
	$\chi_{\text{masses}}^2 + \chi_{\text{CKM}}^2 = 11.57$	$m_d : -1.83$	
		$m_s : -1.49$	
		$m_b : +2.17$	
	$\chi_{\text{total}}^2 = 19.26$	$m_d : -1.79$	normal hierarchy
		$m_s : -1.44$	
		$m_b : +2.27$	
		$\sin^2 \theta_{23} : -2.28$	
D ₂	$\chi_{\text{masses}}^2 = 3.21$	$m_d : -1.74$	
	$\chi_{\text{masses}}^2 + \chi_{\text{CKM}}^2 = 12.75$	$m_d : -2.09$	
		$m_b : +2.54$	
D ₃	$\chi_{\text{masses}}^2 \sim 10^{-6}$		
	$\chi_{\text{masses}}^2 + \chi_{\text{CKM}}^2 = 11.86$	$m_d : -1.42$	
		$m_s : -3.14$	
C	$\chi_{\text{masses}}^2 = 107.59$	$m_s : +1.03$	
		$m_b : -10.32$	

acceptably fit the charged-fermion masses together with the CKM matrix. The best results that we were able to find for all the cases are given in Table 6.

Only case D₁ is able to fit all the observables, but all those good fits yield $|w_R/v_d| < 3 \times 10^{13}$. This is unacceptable since, with $\tan \beta = 10$, $|v_d| = \langle H_d^0 \rangle_0 \approx 17.3$ GeV then leads to $|w_R| \lesssim 5 \times 10^{14}$ GeV, which is almost two orders of magnitude below the unification scale $M_{\text{GUT}} = 2 \times 10^{16}$ GeV. If we enforce a more realistic $|w_R/v_d| > 10^{15}$ on case D₁, then we are only able to obtain poor fits with $\chi_{\text{total}}^2 \gtrsim 60$.

4.6. Case A

Case A has much too many degrees of freedom, so it is adequate to try and constrain it somewhat. We follow Ref. [16], in which real Yukawa-coupling matrices (due to an additional CP symmetry) F , G , and H were enforced and, moreover, $w_L = 0$ has been assumed, thereby discarding the type-II seesaw mechanism. Under these assumptions, the authors of Ref. [16] have parameterized

$$M_d = \begin{pmatrix} x + e^{i\zeta_d} a & e^{i\xi_d} f & e^{i\xi_d} g \\ -e^{i\xi_d} f & y + e^{i\zeta_d} b & e^{i\zeta_d} d \\ -e^{i\xi_d} g & e^{i\zeta_d} d & z + e^{i\zeta_d} c \end{pmatrix}, \quad (45a)$$

$$M_\ell = \begin{pmatrix} x - 3e^{i\zeta_d} a & r_\ell e^{i\xi_\ell} f & r_\ell e^{i\xi_\ell} g \\ -r_\ell e^{i\xi_\ell} f & y - 3e^{i\zeta_d} b & -3e^{i\zeta_d} d \\ -r_\ell e^{i\xi_\ell} g & -3e^{i\zeta_d} d & z - 3e^{i\zeta_d} c \end{pmatrix}, \quad (45b)$$

$$M_u = \begin{pmatrix} r_H x + r_F e^{i\zeta_u} a & r_u e^{i\xi_u} f & r_u e^{i\xi_u} g \\ -r_u e^{i\xi_u} f & r_H y + r_F e^{i\zeta_u} b & r_F e^{i\zeta_u} d \\ -r_u e^{i\xi_u} g & r_F e^{i\zeta_u} d & r_H z + r_F e^{i\zeta_u} c \end{pmatrix}, \tag{45c}$$

$$M_D = \begin{pmatrix} r_H x - 3r_F e^{i\zeta_u} a & r_D e^{i\xi_D} f & r_D e^{i\xi_D} g \\ -r_D e^{i\xi_D} f & r_H y - 3r_F e^{i\zeta_u} b & -3r_F e^{i\zeta_u} d \\ -r_D e^{i\xi_D} g & -3r_F e^{i\zeta_u} d & r_H z - 3r_F e^{i\zeta_u} c \end{pmatrix}, \tag{45d}$$

$$\left| \frac{w_R}{v_d} \right| \mathcal{M}_\nu = \frac{1}{a(bc - d^2)} M_D \begin{pmatrix} bc - d^2 & 0 & 0 \\ 0 & ac & -ad \\ 0 & -ad & ab \end{pmatrix} M_D^T, \tag{45e}$$

where

$$r_\ell e^{i\xi_\ell} \equiv \frac{\kappa_\ell}{|\kappa_d|}, \tag{46a}$$

$$r_u e^{i\xi_u} \equiv \frac{\kappa_u}{|\kappa_d|}, \tag{46b}$$

$$r_D e^{i\xi_D} \equiv \frac{\kappa_D}{|\kappa_d|}. \tag{46c}$$

$$r_H \equiv \left| \frac{k_u}{k_d} \right|, \tag{46d}$$

$$r_F e^{i\zeta_u} \equiv \frac{v_u}{|v_d|}. \tag{46e}$$

In this parameterization, there are six phases (ξ_ℓ , ξ_u , ξ_D , ζ_u , ξ_d , and ζ_d) and 15 moduli (x , y , z , a , b , c , d , f , g , r_ℓ , r_H , r_u , r_F , r_D , and $|w_R/v_d|$).

As usual, we firstly fit the charged-fermion masses, the mixing angles, and r_{solar}^2 . Secondly we adjust the factor $|w_R/v_d|$ in the left-hand side of equation (45e) in such a way that $|m_3^2 - m_1^2| = 2.5 \times 10^{-3} \text{ eV}^2$. Thirdly we compute $|v_d| = |v_d/w_R| (2 \times 10^{16} \text{ GeV})$ and $|v_u| = r_F |v_d|$. Finally, we check that

$$|v_d|^2 < ((H_d^0)_0)^2 \approx (17.3 \text{ GeV})^2, \tag{47a}$$

$$|v_u|^2 < ((H_u^0)_0)^2 \approx (173 \text{ GeV})^2. \tag{47b}$$

We also check that

$$\max(a^2, b^2, c^2, d^2) < y^2 |v_d|^2 \tag{48}$$

for some $1 < y < 10$; we also require

$$\max(x^2, y^2, z^2) + \left(1 + \frac{r_\ell^2}{3}\right) \max(f^2, g^2) < y^2 [((H_d^0)_0)^2 - |v_d|^2], \tag{49a}$$

$$r_H^2 \max(x^2, y^2, z^2) + \left(r_u^2 + \frac{r_D^2}{3}\right) \max(f^2, g^2) < y^2 [((H_u^0)_0)^2 - |v_u|^2]. \tag{49b}$$

In Ref. [16] an explicit fit of case A—under the above restrictions $w_L = 0$ and real Yukawa-coupling matrices—to some data was presented. However, the authors of Ref. [16] have used the charged-fermion masses give in Ref. [24] and have used the upper bound on $\sin^2 \theta_{13}$ that existed at the time. We have attempted to fit case A both to the updated charged-fermion masses of Ref. [23] and to the now extant value of $\sin^2 \theta_{13}$. We could achieve an excellent fit when the

Table 7

The values of the parameters for out best fit of case A with a normal neutrino mass spectrum. The fit has $\chi_{\text{total}}^2 \approx 0.005$.

Parameter	Value
x/MeV	-0.476561625448
y/MeV	-63.004302166872
z/MeV	410.084319821441
a/MeV	-0.372645355981
b/MeV	-91.581208223942
c/MeV	-342.559232981562
d/MeV	-172.410208448655
f/MeV	-3.322328858814
g/MeV	-0.261790479555
$r_\ell e^{i\xi_\ell}$	$4.197350155392 \exp(i 3.160952468)$
$r_u e^{i\xi_u}$	$6.938241672636 \exp(i 2.800761399333)$
$r_D e^{i\xi_D}$	$5682.169770871835 \exp(i 4.151626745)$
$r_F e^{i\xi_u}$	$131.838888425156 \exp(i 3.114060300)$
r_H	100.325400021876
ξ_d/rad	1.736825028772
ζ_d/rad	2.935971894656
$ w_R/v_d $	$1.90053716 \times 10^{16}$

neutrino mass spectrum is normal and a passable one when the mass spectrum is inverted; those fits are presented in [Tables 7 and 8](#), respectively.

For the fit of [Table 7](#) one has $m_1 + m_2 + m_3 \approx 0.06$ eV. The fit of [Table 8](#) has $m_1 + m_2 + m_3 \approx 0.1$ eV.

5. Conclusions

In this paper we have considered a supersymmetric $SO(10)$ GUT in which the fermion masses are generated by renormalizable Yukawa couplings. Consequently, the scalar multiplets under consideration belong to the irreps **10**, $\overline{\mathbf{126}}$, and **120** of $SO(10)$. We have assumed that there is a single scalar multiplet belonging to each of these three irreps; some further mild assumptions are listed in section 2. We have analysed the prospects of imposing flavour symmetries in this scenario, potentially making it predictive. An exhaustive discussion has revealed 14 cases compatible with our scenario. For the numerical examination of those cases we have used the charged-fermion masses evaluated at the GUT scale through renormalization-group running in the context of the Minimal Supersymmetric Standard Model. Interestingly, the numerical analysis ruled out all 14 cases except case A—see equation (11)—and case B—see equation (12). We have demonstrated that both cases A and B allow excellent fits to the data when the neutrino mass spectrum is normal; when that spectrum is inverted, case A can still fit the data but we were unable to find a fit for case B.

Thus, we have come to the conclusion that within the NMSGUT [7], which has renormalizable Yukawa couplings just as the ones considered here, there are at most two possibilities to reduce

Table 8

The values of the parameters for out best fit of case A with an inverted neutrino mass spectrum. This fit has $\chi_{\text{total}}^2 \approx 0.8$.

Parameter	Value
x/MeV	0.980345675289
y/MeV	13.317045360098
z/MeV	834.100031282343
a/MeV	1.230521136698
b/MeV	18.085613337982
c/MeV	82.58146771928
d/MeV	36.879698307221
f/MeV	-2.572520483121
g/MeV	3.267046126672
$r_\ell e^{i\xi_\ell}$	5.389802407484 exp(i 5.291743244393)
$r_u e^{i\xi_u}$	9.506621363405 exp(i 6.456544563269)
$r_D e^{i\xi_D}$	19058.47748201563 exp(i 4.698412501341)
$r_F e^{i\xi_u}$	93.384741884164 exp(i 3.1289172068552067)
r_H	119.091394096965
ξ_d/rad	6.182757601569
ζ_d/rad	4.027845889022
$ w_R/v_d $	$9.0899658664 \times 10^{16}$

the number of Yukawa couplings through flavour symmetries, while remaining in agreement with the data.

Acknowledgements

We thank Ivo de Medeiros Varzielas for help in the initial stages of the fitting procedure. The work of W.G. was partly supported by a grant from the project ‘‘NUPHYS 2013’’ 20NIMLRD0287/CC 1140 of the *Associaao do Instituto Superior Tecnico para a Investigaao e Desenvolvimento* (IST-ID). D.J. thanks the Lithuanian Academy of Sciences for support through the project DaFi2015. The work of L.L. was supported through the projects PEst-OE-FIS-UI0777-2013, PTDC/FIS-NUC/0548-2012, and CERN-FP-123580-2011 of *Fundaao para a Ciencia e a Tecnologia* (FCT); those projects are partially funded through POCTI (FEDER), COMPETE, QREN, and the European Union.

Appendix A. The scalar potential with an additional 45

This appendix addresses two problems that may in general arise in a renormalizable supersymmetric $SO(10)$ GUT furnished with additional symmetries:

- How to promote the full mixing among the Higgs doublets residing in the **10**, **120**, and $\overline{\mathbf{126}}$ of $SO(10)$.
- How to achieve the full breaking of $SO(10)$ to the SM gauge group by using only renormalizable interactions.

Both these problems can be solved in the NMSGUT, but they are non-trivial in the context of our symmetry-furnished cases, especially when the symmetry is larger than \mathbb{Z}_2 .

According to Refs. [4,27,28], in the NMSGUT there are five scalar irreps: the **10**, the **120**, the **126**, the $\overline{\mathbf{126}}$, and the **210**. The **10**, the **120**, and the $\overline{\mathbf{126}}$ have Yukawa couplings; the **126** and the **210** do not. The **210** is needed, together with the **126** and $\overline{\mathbf{126}}$, in order to break $SO(10)$ down to the SM gauge group.

In our models, we propose to add to the NMSGUT one further scalar irrep—the **45**, which is the adjoint of $SO(10)$. The full superpotential is then⁸

$$\begin{aligned}
 V_{\text{super}} = & \lambda_1 \mathbf{10} \mathbf{10} + \lambda_2 \mathbf{45} \mathbf{45} + \lambda_3 \mathbf{120} \mathbf{120} + \lambda_4 \mathbf{210} \mathbf{210} + \lambda_5 \mathbf{126} \overline{\mathbf{126}} \\
 & + \lambda_6 \mathbf{210} \mathbf{210} \mathbf{210} + \lambda_7 \mathbf{45} \mathbf{45} \mathbf{210} + \lambda_8 \mathbf{126} \overline{\mathbf{126}} \mathbf{210} \\
 & + \lambda_9 \mathbf{10} \mathbf{126} \mathbf{210} + \lambda_{10} \mathbf{10} \overline{\mathbf{126}} \mathbf{210} + \lambda_{11} \mathbf{120} \mathbf{120} \mathbf{210} + \lambda_{12} \mathbf{10} \mathbf{120} \mathbf{210} \\
 & + \lambda_{13} \mathbf{120} \mathbf{126} \mathbf{210} + \lambda_{14} \mathbf{120} \overline{\mathbf{126}} \mathbf{210} + \lambda_{15} \mathbf{126} \overline{\mathbf{126}} \mathbf{45} \\
 & + \lambda_{16} \mathbf{10} \mathbf{120} \mathbf{45} + \lambda_{17} \mathbf{120} \mathbf{126} \mathbf{45} + \lambda_{18} \mathbf{120} \overline{\mathbf{126}} \mathbf{45}.
 \end{aligned} \tag{A.1}$$

In order to go from the superpotential to the scalar potential one must square the partial derivative relative to each superfield. Thus, the scalar potential is of the form

$$\begin{aligned}
 V = & \left| \lambda_1 \mathbf{10} + \lambda_9 \mathbf{126} \mathbf{210} + \lambda_{10} \overline{\mathbf{126}} \mathbf{210} + \lambda_{12} \mathbf{120} \mathbf{210} + \lambda_{16} \mathbf{120} \mathbf{45} \right|^2 \\
 & + \left| \lambda_2 \mathbf{45} + \lambda_7 \mathbf{45} \mathbf{210} + \lambda_{15} \mathbf{126} \overline{\mathbf{126}} \right. \\
 & \left. + \lambda_{16} \mathbf{10} \mathbf{120} + \lambda_{17} \mathbf{120} \mathbf{126} + \lambda_{18} \mathbf{120} \overline{\mathbf{126}} \right|^2 \\
 & + \left| \lambda_3 \mathbf{120} + \lambda_{11} \mathbf{120} \mathbf{210} + \lambda_{12} \mathbf{10} \mathbf{210} + \lambda_{13} \mathbf{126} \mathbf{210} + \lambda_{14} \overline{\mathbf{126}} \mathbf{210} \right. \\
 & \left. + \lambda_{16} \mathbf{10} \mathbf{45} + \lambda_{17} \mathbf{126} \mathbf{45} + \lambda_{18} \overline{\mathbf{126}} \mathbf{45} \right|^2 \\
 & + \left| \lambda_4 \mathbf{210} + \lambda_6 \mathbf{210} \mathbf{210} + \lambda_7 \mathbf{45} \mathbf{45} + \lambda_8 \mathbf{126} \overline{\mathbf{126}} + \lambda_9 \mathbf{10} \mathbf{126} \right. \\
 & \left. + \lambda_{10} \mathbf{10} \overline{\mathbf{126}} + \lambda_{11} \mathbf{120} \mathbf{120} + \lambda_{12} \mathbf{10} \mathbf{120} + \lambda_{13} \mathbf{120} \mathbf{126} + \lambda_{14} \mathbf{120} \overline{\mathbf{126}} \right|^2 \\
 & + \left| \lambda_5 \overline{\mathbf{126}} + \lambda_8 \overline{\mathbf{126}} \mathbf{210} + \lambda_9 \mathbf{10} \mathbf{210} + \lambda_{13} \mathbf{120} \mathbf{210} \right. \\
 & \left. + \lambda_{15} \overline{\mathbf{126}} \mathbf{45} + \lambda_{17} \mathbf{120} \mathbf{45} \right|^2 \\
 & + \left| \lambda_5 \mathbf{126} + \lambda_8 \mathbf{126} \mathbf{210} + \lambda_{10} \mathbf{10} \mathbf{210} + \lambda_{14} \mathbf{120} \mathbf{210} \right. \\
 & \left. + \lambda_{15} \mathbf{126} \mathbf{45} + \lambda_{18} \mathbf{120} \mathbf{45} \right|^2.
 \end{aligned} \tag{A.2}$$

The **10** and the **120** do not have any component which is invariant under the SM gauge group, therefore they are not allowed to acquire a VEV at the GUT scale. Thus, at the GUT scale the relevant potential is just

⁸ One may check that no term is missing in equation (A.1) by studying Table 820 of Ref. [29].

$$\begin{aligned}
 V_{\text{GUT}} = & \left| \lambda_9 \mathbf{126} \mathbf{210} + \lambda_{10} \overline{\mathbf{126}} \mathbf{210} \right|^2 + \left| \lambda_2 \mathbf{45} + \lambda_7 \mathbf{45} \mathbf{210} + \lambda_{15} \mathbf{126} \overline{\mathbf{126}} \right|^2 \\
 & + \left| \lambda_{13} \mathbf{126} \mathbf{210} + \lambda_{14} \overline{\mathbf{126}} \mathbf{210} + \lambda_{17} \mathbf{126} \mathbf{45} + \lambda_{18} \overline{\mathbf{126}} \mathbf{45} \right|^2 \\
 & + \left| \lambda_4 \mathbf{210} + \lambda_6 \mathbf{210} \mathbf{210} + \lambda_7 \mathbf{45} \mathbf{45} + \lambda_8 \mathbf{126} \overline{\mathbf{126}} \right|^2 \\
 & + \left| \lambda_5 \overline{\mathbf{126}} + \lambda_8 \overline{\mathbf{126}} \mathbf{210} + \lambda_{15} \overline{\mathbf{126}} \mathbf{45} \right|^2 \\
 & + \left| \lambda_5 \mathbf{126} + \lambda_8 \mathbf{126} \mathbf{210} + \lambda_{15} \mathbf{126} \mathbf{45} \right|^2.
 \end{aligned} \tag{A.3}$$

In our case A there is a symmetry $\mathbf{120} \rightarrow -\mathbf{120}$. We must extend it and make $\mathbf{45} \rightarrow -\mathbf{45}$ too. Then the symmetry implies $\lambda_{12} = \lambda_{13} = \lambda_{14} = \lambda_{15} = 0$. Equation (A.2) becomes

$$\begin{aligned}
 V^{(\text{case A})} = & \left| \lambda_1 \mathbf{10} + \lambda_9 \mathbf{126} \mathbf{210} + \lambda_{10} \overline{\mathbf{126}} \mathbf{210} + \lambda_{16} \mathbf{120} \mathbf{45} \right|^2 \\
 & + \left| \lambda_2 \mathbf{45} + \lambda_7 \mathbf{45} \mathbf{210} + \lambda_{16} \mathbf{10} \mathbf{120} + \lambda_{17} \mathbf{120} \mathbf{126} + \lambda_{18} \mathbf{120} \overline{\mathbf{126}} \right|^2 \\
 & + \left| \lambda_3 \mathbf{120} + \lambda_{11} \mathbf{120} \mathbf{210} + \lambda_{16} \mathbf{10} \mathbf{45} + \lambda_{17} \mathbf{126} \mathbf{45} + \lambda_{18} \overline{\mathbf{126}} \mathbf{45} \right|^2 \\
 & + \left| \lambda_4 \mathbf{210} + \lambda_6 \mathbf{210} \mathbf{210} + \lambda_7 \mathbf{45} \mathbf{45} + \lambda_8 \mathbf{126} \overline{\mathbf{126}} + \lambda_9 \mathbf{10} \mathbf{126} \right. \\
 & \left. + \lambda_{10} \mathbf{10} \overline{\mathbf{126}} + \lambda_{11} \mathbf{120} \mathbf{120} \right|^2 \\
 & + \left| \lambda_5 \overline{\mathbf{126}} + \lambda_8 \overline{\mathbf{126}} \mathbf{210} + \lambda_9 \mathbf{10} \mathbf{210} + \lambda_{17} \mathbf{120} \mathbf{45} \right|^2 \\
 & + \left| \lambda_5 \mathbf{126} + \lambda_8 \mathbf{126} \mathbf{210} + \lambda_{10} \mathbf{10} \mathbf{210} + \lambda_{18} \mathbf{120} \mathbf{45} \right|^2
 \end{aligned} \tag{A.4}$$

and equation (A.3) becomes

$$\begin{aligned}
 V_{\text{GUT}}^{(\text{case A})} = & \left| \lambda_9 \mathbf{126} \mathbf{210} + \lambda_{10} \overline{\mathbf{126}} \mathbf{210} \right|^2 + \left| \lambda_2 \mathbf{45} + \lambda_7 \mathbf{45} \mathbf{210} \right|^2 \\
 & + \left| \lambda_{17} \mathbf{126} \mathbf{45} + \lambda_{18} \overline{\mathbf{126}} \mathbf{45} \right|^2 \\
 & + \left| \lambda_4 \mathbf{210} + \lambda_6 \mathbf{210} \mathbf{210} + \lambda_7 \mathbf{45} \mathbf{45} + \lambda_8 \mathbf{126} \overline{\mathbf{126}} \right|^2 \\
 & + \left| \lambda_5 \overline{\mathbf{126}} + \lambda_8 \overline{\mathbf{126}} \mathbf{210} \right|^2 + \left| \lambda_5 \mathbf{126} + \lambda_8 \mathbf{126} \mathbf{210} \right|^2.
 \end{aligned} \tag{A.5}$$

The third line of equation (A.4) indicates that the $\mathbf{120}$ fully mixes with both the $\mathbf{10}$ and the $\overline{\mathbf{126}}$. The first and last lines of equation (A.5) indicate that the potential for the $\mathbf{45}$, $\mathbf{126}$, $\overline{\mathbf{126}}$, and $\mathbf{210}$ allows all of them to acquire VEVs.

In our case B there is a symmetry $\mathbf{10} \rightarrow -\mathbf{10}$, $\mathbf{120} \rightarrow -\mathbf{120}$, $\mathbf{45} \rightarrow -\mathbf{45}$. This implies $\lambda_9 = \lambda_{10} = \lambda_{13} = \lambda_{14} = \lambda_{15} = \lambda_{16} = 0$. Equation (A.2) becomes

$$\begin{aligned}
 V^{(\text{case B})} = & \left| \lambda_1 \mathbf{10} + \lambda_{12} \mathbf{120} \mathbf{210} \right|^2 \\
 & + \left| \lambda_2 \mathbf{45} + \lambda_7 \mathbf{45} \mathbf{210} + \lambda_{17} \mathbf{120} \mathbf{126} + \lambda_{18} \mathbf{120} \overline{\mathbf{126}} \right|^2 \\
 & + \left| \lambda_3 \mathbf{120} + \lambda_{11} \mathbf{120} \mathbf{210} + \lambda_{12} \mathbf{10} \mathbf{210} + \lambda_{17} \mathbf{126} \mathbf{45} + \lambda_{18} \overline{\mathbf{126}} \mathbf{45} \right|^2 \\
 & + \left| \lambda_4 \mathbf{210} + \lambda_6 \mathbf{210} \mathbf{210} + \lambda_7 \mathbf{45} \mathbf{45} \right|^2
 \end{aligned}$$

$$\begin{aligned}
 & + \lambda_8 \mathbf{126} \overline{\mathbf{126}} + \lambda_{11} \mathbf{120} \mathbf{120} + \lambda_{12} \mathbf{10} \mathbf{120} \Big|^2 \\
 & + \left| \lambda_5 \overline{\mathbf{126}} + \lambda_8 \overline{\mathbf{126}} \mathbf{210} + \lambda_{17} \mathbf{120} \mathbf{45} \right|^2 \\
 & + \left| \lambda_5 \mathbf{126} + \lambda_8 \mathbf{126} \mathbf{210} + \lambda_{18} \mathbf{120} \mathbf{45} \right|^2
 \end{aligned} \tag{A.6}$$

and equation (A.3) becomes

$$\begin{aligned}
 V_{\text{GUT}}^{(\text{case B})} & = \left| \lambda_2 \mathbf{45} + \lambda_7 \mathbf{45} \mathbf{210} \right|^2 + \left| \lambda_{17} \mathbf{126} \mathbf{45} + \lambda_{18} \overline{\mathbf{126}} \mathbf{45} \right|^2 \\
 & + \left| \lambda_4 \mathbf{210} + \lambda_6 \mathbf{210} \mathbf{210} + \lambda_7 \mathbf{45} \mathbf{45} + \lambda_8 \mathbf{126} \overline{\mathbf{126}} \right|^2 \\
 & + \left| \lambda_5 \overline{\mathbf{126}} + \lambda_8 \overline{\mathbf{126}} \mathbf{210} \right|^2 + \left| \lambda_5 \mathbf{126} + \lambda_8 \mathbf{126} \mathbf{210} \right|^2.
 \end{aligned} \tag{A.7}$$

The third line of equation (A.6) indicates that the **120** fully mixes with both the **10** and the $\overline{\mathbf{126}}$. Equation (A.7) demonstrates that the potential for the **45**, **126**, $\overline{\mathbf{126}}$, and **210** allows all of them to acquire VEVs.

Case C may be treated in a similar fashion. Our remaining cases have symmetries \mathbb{Z}_n with $n > 2$ and are much more problematic. Anyway, we do not need to worry about those cases since we already know that they are unable to fit the phenomenological data.

Appendix B. Investigation of a second symmetry

In this appendix we take all 13 cases of subsection 3.1 and consider, for each of them, the possibility of a second flavour symmetry defined in equation (24). Without loss of generality we set $e^{i\beta_1} = 1$ in that equation.

The conclusion of this appendix is that, beyond those 13 cases, only one new case arises which does not contradict our assumptions—case E in equation (25).

B.1. Cases A_1 , A'_1 , A''_1 , and A_2

In all these four cases,

$$G = \begin{pmatrix} 0 & 0 & d \\ 0 & 0 & 0 \\ -d & 0 & 0 \end{pmatrix} \tag{B.1}$$

with $d \neq 0$. Since

$$X^T G X = G \Leftrightarrow G X = X^* G, \tag{B.2}$$

we find

$$x_{12} = x_{21} = x_{23} = x_{32} = 0, \quad X = \begin{pmatrix} x_{11} & 0 & x_{13} \\ 0 & x_{22} & 0 \\ -x_{13}^* & 0 & x_{11}^* \end{pmatrix}. \tag{B.3}$$

In these four cases the matrix F has the form

$$F = F_1 \equiv \begin{pmatrix} 0 & a & 0 \\ a & 0 & 0 \\ 0 & 0 & b \end{pmatrix}. \tag{B.4}$$

We require $\det F \neq 0$, hence $a \neq 0$ and $b \neq 0$. Using

$$e^{i\gamma_1} F X = X^* F \tag{B.5}$$

with the matrix X of equation (B.3), we obtain that X must be diagonal:

$$X = \text{diag} \left(e^{i\gamma_1/2}, e^{-3i\gamma_1/2}, e^{-i\gamma_1/2} \right). \tag{B.6}$$

Now we look for the consequences of

$$e^{i\alpha_1} H X = X^* H. \tag{B.7}$$

With a diagonal X , equation (B.7) can only force either one or more matrix elements of H to be zero. In the case A_1 , if one sets one matrix element of H to zero then one simply recovers the cases A'_1 and A''_1 . In the cases A'_1 and A''_1 , the number of non-vanishing elements of H is already minimal. In the case A_2 we have $X^T H X = e^{-i\gamma_1} H$, therefore either $\alpha_1 = \gamma_1$ and H is not restricted by S_1 or $\alpha_1 \neq \gamma_1$ and $H = 0$, which is excluded by our assumptions.

In summary, departing from cases A_1, A'_1, A''_1 , or A_2 no new cases can ensue from a second symmetry.

B.2. Cases D_2, D_3, D'_2 , and D'_3

In these cases equation (B.1) is still valid, therefore equation (B.3) also holds. In all four cases

$$F = F_2 \equiv \begin{pmatrix} b & 0 & 0 \\ 0 & 0 & a \\ 0 & a & 0 \end{pmatrix}, \tag{B.8}$$

with $a \neq 0$ and $b \neq 0$. Using equation (B.5) then yields

$$X = \text{diag} \left(e^{-i\gamma_1/2}, e^{-3i\gamma_1/2}, e^{i\gamma_1/2} \right), \tag{B.9}$$

i.e. X is once again diagonal.

We next consider equation (B.7). In case D_3 we obtain

$$H = \begin{pmatrix} 0 & 0 & r \\ 0 & s & 0 \\ r & 0 & 0 \end{pmatrix} = e^{i\alpha_1} X^T H X = e^{i\alpha_1} \begin{pmatrix} 0 & 0 & r \\ 0 & e^{-3i\gamma_1} s & 0 \\ r & 0 & 0 \end{pmatrix}. \tag{B.10}$$

In case D_2 we have

$$H = \begin{pmatrix} 0 & r & 0 \\ r & 0 & 0 \\ 0 & 0 & s \end{pmatrix} = e^{i\alpha_1} X^T H X = e^{i\alpha_1} \begin{pmatrix} 0 & e^{-2i\gamma_1} r & 0 \\ e^{-2i\gamma_1} r & 0 & 0 \\ 0 & 0 & e^{i\gamma_1} s \end{pmatrix}. \tag{B.11}$$

Thus, equation (B.7) can at most set either $r = 0$ or $s = 0$. If $s = 0$ then one recovers case D'_2 from case D_2 and case D'_3 from case D_3 . If $r = 0$ then, through an interchange of the first and third generations, one recovers case A'_1 from case D_2 and case A''_1 from case D_3 . Therefore, no new cases arise from the enforcement of the symmetry S_1 on any of these four cases.

B.3. Cases D_1 and D'_1

Equations (B.1) and (B.3) once again hold. Now

$$F = F_3 \equiv \begin{pmatrix} 0 & 0 & a \\ 0 & b & 0 \\ a & 0 & 0 \end{pmatrix}, \tag{B.12}$$

with $a \neq 0$ and $b \neq 0$. Equation (B.5) then yields that either $e^{i\gamma_1} = +1$ and

$$X = \text{diag}(e^{i\psi}, \pm 1, e^{-i\psi}) \tag{B.13}$$

or $e^{i\gamma_1} = -1$ and

$$X = \begin{pmatrix} 0 & 0 & e^{i\varphi} \\ 0 & \pm i & 0 \\ -e^{-i\varphi} & 0 & 0 \end{pmatrix}. \tag{B.14}$$

In case D_1 and with equation (B.13) one obtains

$$H = \begin{pmatrix} 0 & r & 0 \\ r & 0 & 0 \\ 0 & 0 & s \end{pmatrix} = e^{i\alpha_1} X^T H X = e^{i\alpha_1} \begin{pmatrix} 0 & \pm e^{i\psi} r & 0 \\ \pm e^{i\psi} r & 0 & 0 \\ 0 & 0 & e^{-2i\psi} s \end{pmatrix}. \tag{B.15}$$

With equation (B.14) one arrives instead at

$$H = \begin{pmatrix} 0 & r & 0 \\ r & 0 & 0 \\ 0 & 0 & s \end{pmatrix} = e^{i\alpha_1} X^T H X = e^{i\alpha_1} \begin{pmatrix} e^{-2i\varphi} s & 0 & 0 \\ 0 & 0 & \pm i e^{i\varphi} r \\ 0 & \pm i e^{i\varphi} r & 0 \end{pmatrix}. \tag{B.16}$$

Thus, the possibility (B.14) implies $H = 0$, which contradicts our assumptions. With equation (B.15) then either $s = 0$ and one recovers case D'_1 or $r = 0$ and the second generation decouples. We conclude that the enforcement of the symmetry \mathcal{S}_1 on cases D_1 and D'_1 cannot lead to new cases.

B.4. Cases A and B, step 1: X may be chosen to be diagonal

In cases A and B we may perform a weak-basis transformation such that G acquires the form (B.1) while the forms of H and F are kept unchanged:

$$\text{case A: } H \sim \begin{pmatrix} \times & \times & 0 \\ \times & \times & 0 \\ 0 & 0 & \times \end{pmatrix}, \quad G \sim \begin{pmatrix} 0 & 0 & \times \\ 0 & 0 & 0 \\ \times & 0 & 0 \end{pmatrix}, \quad F \sim \begin{pmatrix} \times & \times & 0 \\ \times & \times & 0 \\ 0 & 0 & \times \end{pmatrix}; \tag{B.17a}$$

$$\text{case B: } H \sim \begin{pmatrix} 0 & 0 & \times \\ 0 & 0 & \times \\ \times & \times & 0 \end{pmatrix}, \quad G \sim \begin{pmatrix} 0 & 0 & \times \\ 0 & 0 & 0 \\ \times & 0 & 0 \end{pmatrix}, \quad F \sim \begin{pmatrix} \times & \times & 0 \\ \times & \times & 0 \\ 0 & 0 & \times \end{pmatrix}. \tag{B.17b}$$

This is achieved through a unitary rotation of the first and second generations, which does not alter the matrix $W = \text{diag}(+1, +1, -1)$ for these cases. In the new basis (B.17), equation (B.3) holds.

Next we consider equation (B.5). With $F \equiv (f_{ij})$, it reads

$$e^{i\gamma_1} \begin{pmatrix} f_{11}x_{11} & f_{12}x_{22} & f_{11}x_{13} \\ f_{12}x_{11} & f_{22}x_{22} & f_{12}x_{13} \\ -f_{33}x_{13}^* & 0 & f_{33}x_{11}^* \end{pmatrix} = \begin{pmatrix} f_{11}x_{11}^* & f_{12}x_{11}^* & f_{33}x_{13}^* \\ f_{12}x_{22}^* & f_{22}x_{22}^* & 0 \\ -f_{11}x_{13} & -f_{12}x_{13} & f_{33}x_{11} \end{pmatrix}. \tag{B.18}$$

Let us suppose that X is not diagonal, *i.e.* that x_{13} is non-zero. Then equation (B.18) tells us that $f_{12} = 0$, *i.e.* that F is diagonal. Now we invoke $e^{i\alpha_1} H X = X^* H$. In case A the matrix $H \equiv (h_{ij})$ has the same form as the matrix F , hence we may conclude, from the analogue of equation (B.18), that $h_{12} = 0$ just as $f_{12} = 0$, *i.e.* H is diagonal too. But then the second generation decouples, which runs against our assumptions. For case B the equation $e^{i\alpha_1} H X = X^* H$ reads

$$e^{i\alpha_1} \begin{pmatrix} -h_{13}x_{13}^* & 0 & h_{13}x_{11}^* \\ -h_{23}x_{13}^* & 0 & h_{23}x_{11}^* \\ h_{13}x_{11} & h_{23}x_{22} & h_{13}x_{13} \end{pmatrix} = \begin{pmatrix} h_{13}x_{13}^* & h_{23}x_{13}^* & h_{13}x_{11}^* \\ 0 & 0 & h_{23}x_{22}^* \\ h_{13}x_{11} & h_{23}x_{11} & -h_{13}x_{13} \end{pmatrix}, \tag{B.19}$$

hence $h_{23} = 0$ and the second generation decouples.

We conclude that the hypothesis $x_{13} \neq 0$ leads to a contradiction with our assumptions. Thus, cases A and B do not admit a non-diagonal X .

B.5. Cases A and B, step 2: the forms of F and X

With a diagonal matrix X , the equation $X^T F X e^{i\gamma_1} = F$ yields

$$x_{11}^2 f_{11} e^{i\gamma_1} = f_{11}, \tag{B.20a}$$

$$x_{22}^2 f_{22} e^{i\gamma_1} = f_{22}, \tag{B.20b}$$

$$x_{11} x_{22} f_{12} e^{i\gamma_1} = f_{12}, \tag{B.20c}$$

$$x_{33}^2 f_{33} e^{i\gamma_1} = f_{33}. \tag{B.20d}$$

Since $\det F \neq 0$, f_{33} cannot vanish. Therefore, equation (B.20d) gives $x_{33} = \varepsilon e^{-i\gamma_1/2}$, where $\varepsilon = \pm 1$.

Let us firstly suppose that $x_{11} = x_{22}$. In this case we must have $x_{11}^2 = e^{-i\gamma_1}$, else $f_{11} = f_{22} = f_{12} = 0$ and $\det F = 0$. Consequently, $x_{11} = \eta e^{-i\gamma_1/2}$, where $\eta = \pm 1$. In this case the matrix F cannot be restricted any further by S_1 .

Since $X = e^{-i\gamma_1/2} \text{diag}(\eta, \eta, \varepsilon)$, $X^T H X = e^{-i\gamma_1} H$ in case A and $X^T H X = \varepsilon \eta e^{-i\gamma_1} H$ in case B. This means that the equation $e^{i\alpha_1} X^T H X = H$ either does not restrict H any further, or it enforces $H = 0$ (depending on the choice for $e^{i\alpha_1}$). Since $H = 0$ runs against our assumptions, we conclude that, with $x_{11} = x_{22}$, the symmetry S_1 does not restrict the Yukawa-coupling matrices any further, *i.e.* it does not lead to any new cases.

So we are lead to consider $x_{11} \neq x_{22}$. Then, only two possibilities for X remain, which are compatible with $\det F \neq 0$: either

$$X = X_a \equiv e^{-i\gamma_1/2} \text{diag}(\eta, -\eta, \varepsilon), \tag{B.21a}$$

$$F = F_a \equiv \text{diag}(f_{11}, f_{22}, f_{33}), \tag{B.21b}$$

or

$$X = X_b \equiv e^{-i\gamma_1/2} \text{diag}(e^{i\rho}, e^{-i\rho}, \varepsilon), \tag{B.22a}$$

$$F = F_1, \tag{B.22b}$$

with F_1 given by equation (B.4) and $e^{2i\rho} \neq 1$.

We must remember that X must be of the form (B.3), viz. that $x_{33} = x_{11}^*$. Therefore,

$$e^{i\gamma_1} = e^{-i\gamma_1} = \eta\varepsilon \text{ if } X = X_a, \tag{B.23a}$$

$$e^{i(\gamma_1-\rho)} = e^{i(\rho-\gamma_1)} = \varepsilon \text{ if } X = X_b. \tag{B.23b}$$

B.6. Cases A and B, step 3: the form of H

Case A, $X = X_a, F = F_a$: In this case the equation $e^{i\alpha_1} X^T H X = H$ gives

$$e^{i(\alpha_1-\gamma_1)} \begin{pmatrix} h_{11} & -h_{12} & 0 \\ -h_{12} & h_{22} & 0 \\ 0 & 0 & h_{33} \end{pmatrix} = \begin{pmatrix} h_{11} & h_{12} & 0 \\ h_{12} & h_{22} & 0 \\ 0 & 0 & h_{33} \end{pmatrix}. \tag{B.24}$$

If $e^{i(\alpha_1-\gamma_1)} \neq \pm 1$, then $H = 0$ contradicts our assumptions. If $e^{i(\alpha_1-\gamma_1)} = 1$, then $h_{12} = 0$ and the second generation decouples. If $e^{i(\alpha_1-\gamma_1)} = -1$, then we get case E—see equations (25), because the choice $e^{i\gamma_1} = e^{i\gamma_1/2} = \eta = \varepsilon = 1$ indeed leads to the symmetry $\mathbb{Z}_2^{(2)}$ of that equation.

Case B, $X = X_a, F = F_a$: In this case the equation $e^{i\alpha_1} X^T H X = H$ gives

$$e^{i(\alpha_1-\gamma_1)} \eta\varepsilon \begin{pmatrix} 0 & 0 & h_{13} \\ 0 & 0 & -h_{23} \\ h_{13} & -h_{23} & 0 \end{pmatrix} = \begin{pmatrix} 0 & 0 & h_{13} \\ 0 & 0 & h_{23} \\ h_{13} & h_{23} & 0 \end{pmatrix}. \tag{B.25}$$

In order to avoid decoupling of the second generation, we must choose $e^{i(\alpha_1-\gamma_1)} \eta\varepsilon = -1$ and $h_{13} = 0$. We then obtain a case which is equivalent to case E after the interchange of the first and third generations.

Case A, $X = X_b, F = F_1$: In this case the equation $e^{i\alpha_1} X^T H X = H$ gives

$$e^{i(\alpha_1-\gamma_1)} \begin{pmatrix} e^{2i\rho} h_{11} & h_{12} & 0 \\ h_{12} & e^{-2i\rho} h_{22} & 0 \\ 0 & 0 & h_{33} \end{pmatrix} = \begin{pmatrix} h_{11} & h_{12} & 0 \\ h_{12} & h_{22} & 0 \\ 0 & 0 & h_{33} \end{pmatrix}. \tag{B.26}$$

Since $e^{2i\rho} \neq 1$, through a choice of the phases we may achieve either case A_1 or case A'_1 or case A''_1 or case A_2 ; no new case arises.

Case B, $X = X_b, F = F_1$: In this case the equation $e^{i\alpha_1} X^T H X = H$ gives

$$e^{i(\alpha_1-\gamma_1)} \varepsilon \begin{pmatrix} 0 & 0 & e^{i\rho} h_{13} \\ 0 & 0 & e^{-i\rho} h_{23} \\ e^{i\rho} h_{13} & e^{-i\rho} h_{23} & 0 \end{pmatrix} = \begin{pmatrix} 0 & 0 & h_{13} \\ 0 & 0 & h_{23} \\ h_{13} & h_{23} & 0 \end{pmatrix}. \tag{B.27}$$

In order to avoid decoupling of the second generation we must choose $e^{i(\alpha_1-\gamma_1-\rho)} \varepsilon = 1$ and $h_{13} = 0$; this case is equivalent to D'_2 through the interchange of the first and third generations.

B.7. Case C

In case C, it is convenient to choose a weak basis where

$$H \sim \begin{pmatrix} 0 & \times & 0 \\ \times & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad G \sim \begin{pmatrix} 0 & 0 & \times \\ 0 & 0 & 0 \\ \times & 0 & 0 \end{pmatrix}, \quad F \sim \begin{pmatrix} \times & 0 & \times \\ 0 & \times & 0 \\ \times & 0 & \times \end{pmatrix}. \tag{B.28}$$

This weak basis is achieved, starting from the form (13b) of the matrices H , G , and F , through a unitary rotation mixing the first and third generations; such a rotation does not alter the matrix W in equation (13a).

With G of equation (B.28) we know that X has to obey equation (B.3). It is then easy to see that $HXe^{i\alpha_1} = X^*H$ requires X to be diagonal with $e^{i\alpha_1}x_{11} = x_{22}^*$. Therefore, X can be parameterized as

$$X = \text{diag} \left(e^{i\psi}, e^{-i(\alpha_1+\psi)}, e^{-i\psi} \right). \tag{B.29}$$

With this X , the equation $X^T F X e^{i\gamma_1} = F$ can only force one or more matrix elements of F to be zero.

If $e^{i\gamma_1} \neq 1$, one obtains $f_{13} = f_{31} = 0$ and, therefore, case E. Then, because of $\det F \neq 0$, all f_{ii} must be non-zero and it is easy to show that this leads to $e^{i\gamma_1} = -1$, $e^{i\psi} = i\eta$, and $e^{i\alpha_1} = \varepsilon$ with $\eta^2 = \varepsilon^2 = 1$. Summarizing, we have

$$X = i\eta (1, -\varepsilon, -1), \quad e^{i\alpha_1} = \varepsilon, \quad e^{i\beta_1} = +1, \quad e^{i\gamma_1} = -1. \tag{B.30}$$

By choosing $\varepsilon = -1$ and absorbing $(i\eta)^2 = -1$ into the phase factors, we arrive at $\mathbb{Z}_2^{(1)}$ of equation (25). Note that in this subsection the symmetry S_0 is given by $\mathbb{Z}_2^{(2)}$ and S_1 by $\mathbb{Z}_2^{(1)}$, since we started from case C. Thus, in the present subsection, the notation for $\mathbb{Z}_2^{(1)}$ and $\mathbb{Z}_2^{(2)}$ is exchanged compared to equation (25).

Moving to $e^{i\gamma_1} = 1$ and taking again into account $\det F \neq 0$, we have $f_{22} \neq 0$. However, it is neither possible to enforce $f_{11} = 0$ while keeping $f_{33} \neq 0$ nor to enforce $f_{33} = 0$ while keeping $f_{11} \neq 0$; with $f_{11} = f_{33} = 0$ one recovers case D'_1 .

One thus concludes that enforcing an extra symmetry on case C can only lead to cases E or D'_1 , or else to a violation of our assumptions.

Appendix C. Precise definition of the matrices F , G , H

The aim of this appendix is to precisely define the matrices F , G , and H through equations (C.5) and thereby to extract the useful inequalities (C.11), which we employ in subsection 4.4.2.

The MSSM contains two Higgs doublets, H_d and H_u , with hypercharges $+1/2$ and $-1/2$, respectively. Their corresponding VEVs are $v \cos \beta$ and $v \sin \beta$, respectively, where $v = 174 \text{ GeV}$. When one neglects the effects of the electroweak scale, these two doublets are, by assumption, the only scalar zero-modes extant at the GUT scale; this requires a minimal finetuning condition [30,27]. Each of the scalar irreps **10**, $\overline{\mathbf{126}}$, **126**, and **210** contains one doublet with the quantum numbers of H_d ; the **120** contains two such doublets. The doublet H_d is a superposition of these six doublets with amplitudes $\bar{\alpha}_j$ ($j = 1, \dots, 6$). Let α_j denote the analogous coefficients for H_u . The normalization conditions are

$$\sum_{j=1}^6 |\bar{\alpha}_j|^2 = \sum_{j=1}^6 |\alpha_j|^2 = 1. \tag{C.1}$$

It follows from equations (C.1) that

$$|\bar{\alpha}_1|^2 + |\bar{\alpha}_2|^2 + |\bar{\alpha}_5|^2 + |\bar{\alpha}_6|^2 \leq 1, \tag{C.2a}$$

$$|\alpha_1|^2 + |\alpha_2|^2 + |\alpha_5|^2 + |\alpha_6|^2 \leq 1. \tag{C.2b}$$

The inequalities (C.2) only involve the amplitudes of the doublets contained in the **10**, $\overline{\mathbf{126}}$, and **120**.

Taking into account that the **126** and the **210** have no Yukawa couplings, the Dirac mass matrices are given by

$$M_a = v \cos \beta \left[c_1^a \bar{\alpha}_1 Y_{10} + c_2^a \bar{\alpha}_2 Y_{\overline{126}} + (c_5^a \bar{\alpha}_5 + c_6^a \bar{\alpha}_6) Y_{120} \right] \quad (a = d, \ell), \quad (\text{C.3a})$$

$$M_b = v \sin \beta \left[c_1^b \alpha_1 Y_{10} + c_2^b \alpha_2 Y_{\overline{126}} + (c_5^b \alpha_5 + c_6^b \alpha_6) Y_{120} \right] \quad (b = u, D), \quad (\text{C.3b})$$

with Yukawa-coupling matrices Y_{10} , $Y_{\overline{126}}$, and Y_{120} and Clebsch–Gordan coefficients $c_j^{a,b}$; the latter derive from the $SO(10)$ -invariant Yukawa couplings [5,31]. The absolute values of the Clebsch–Gordan coefficients have no physical meaning and some of their phases are convention dependent. With our conventions,⁹ the required information reads

$$c_1^d = c_1^u = c_1^\ell = c_1^D, \quad (\text{C.4a})$$

$$3c_2^d = -3c_2^u = -c_2^\ell = c_2^D, \quad (\text{C.4b})$$

$$\sqrt{3}c_5^d = -\sqrt{3}c_5^u = \sqrt{3}c_5^\ell = -\sqrt{3}c_5^D = 3c_6^d = 3c_6^u = -c_6^\ell = -c_6^D. \quad (\text{C.4c})$$

In order to make contact with the mass formulas, we define

$$H \equiv c_1^d Y_{10}, \quad (\text{C.5a})$$

$$F \equiv c_2^d Y_{\overline{126}}, \quad (\text{C.5b})$$

$$G \equiv \sqrt{(c_5^d)^2 + (c_6^d)^2} Y_{120}. \quad (\text{C.5c})$$

Then, by using equations (C.4) and (C.5c) we derive

$$\begin{aligned} (c_5^d \bar{\alpha}_5 + c_6^d \bar{\alpha}_6) Y_{120} &= \frac{c_5^d \bar{\alpha}_5 + c_6^d \bar{\alpha}_6}{\sqrt{(c_5^d)^2 + (c_6^d)^2}} G \\ &= \left(\frac{\sqrt{3}}{2} \bar{\alpha}_5 + \frac{1}{2} \bar{\alpha}_6 \right) G, \end{aligned} \quad (\text{C.6a})$$

$$\begin{aligned} (c_5^\ell \bar{\alpha}_5 + c_6^\ell \bar{\alpha}_6) Y_{120} &= \frac{c_5^\ell \bar{\alpha}_5 + c_6^\ell \bar{\alpha}_6}{\sqrt{(c_5^\ell)^2 + (c_6^\ell)^2}} \sqrt{\frac{(c_5^\ell)^2 + (c_6^\ell)^2}{(c_5^d)^2 + (c_6^d)^2}} G \\ &= \left(\frac{1}{2} \bar{\alpha}_5 - \frac{\sqrt{3}}{2} \bar{\alpha}_6 \right) \sqrt{3} G, \end{aligned} \quad (\text{C.6b})$$

$$\begin{aligned} (c_5^u \alpha_5 + c_6^u \alpha_6) Y_{120} &= \frac{c_5^u \alpha_5 + c_6^u \alpha_6}{\sqrt{(c_5^u)^2 + (c_6^u)^2}} \sqrt{\frac{(c_5^u)^2 + (c_6^u)^2}{(c_5^d)^2 + (c_6^d)^2}} G \\ &= \left(-\frac{\sqrt{3}}{2} \alpha_5 + \frac{1}{2} \alpha_6 \right) G, \end{aligned} \quad (\text{C.6c})$$

⁹ See the appendix of Ref. [16].

$$\begin{aligned} (c_5^D \alpha_5 + c_6^D \alpha_6) Y_{120} &= \frac{c_5^D \alpha_5 + c_6^D \alpha_6}{\sqrt{(c_5^D)^2 + (c_6^D)^2}} \sqrt{\frac{(c_5^D)^2 + (c_6^D)^2}{(c_5^d)^2 + (c_6^d)^2}} G \\ &= \left(-\frac{1}{2} \alpha_5 - \frac{\sqrt{3}}{2} \alpha_6 \right) \sqrt{3} G. \end{aligned} \tag{C.6d}$$

Equations (C.5a), (C.5b), and (C.6) may now be plugged into the mass formulas (C.3). The result is

$$M_d = v \cos \beta \left[\bar{\alpha}_1 H + \bar{\alpha}_2 F + \left(\frac{\sqrt{3}}{2} \bar{\alpha}_5 + \frac{1}{2} \bar{\alpha}_6 \right) G \right], \tag{C.7a}$$

$$M_\ell = v \cos \beta \left[\bar{\alpha}_1 H - 3\bar{\alpha}_2 F + \left(\frac{1}{2} \bar{\alpha}_5 - \frac{\sqrt{3}}{2} \bar{\alpha}_6 \right) \sqrt{3} G \right], \tag{C.7b}$$

$$M_u = v \sin \beta \left[\alpha_1 H - \alpha_2 F + \left(-\frac{\sqrt{3}}{2} \alpha_5 + \frac{1}{2} \alpha_6 \right) G \right], \tag{C.7c}$$

$$M_D = v \sin \beta \left[\alpha_1 H + 3\alpha_2 F + \left(-\frac{1}{2} \alpha_5 - \frac{\sqrt{3}}{2} \alpha_6 \right) \sqrt{3} G \right]. \tag{C.7d}$$

By comparing equations (C.7) and (2) we obtain the identifications

$$k_d = v \cos \beta \bar{\alpha}_1, \tag{C.8a}$$

$$k_u = v \sin \beta \alpha_1, \tag{C.8b}$$

$$v_d = v \cos \beta \bar{\alpha}_2, \tag{C.8c}$$

$$v_u = -v \sin \beta \alpha_2, \tag{C.8d}$$

$$\kappa_d = v \cos \beta \left(\frac{\sqrt{3}}{2} \bar{\alpha}_5 + \frac{1}{2} \bar{\alpha}_6 \right), \tag{C.8e}$$

$$\kappa_\ell = v \cos \beta \left(\frac{1}{2} \bar{\alpha}_5 - \frac{\sqrt{3}}{2} \bar{\alpha}_6 \right) \sqrt{3}, \tag{C.8f}$$

$$\kappa_u = v \sin \beta \left(-\frac{\sqrt{3}}{2} \alpha_5 + \frac{1}{2} \alpha_6 \right), \tag{C.8g}$$

$$\kappa_D = v \sin \beta \left(-\frac{1}{2} \alpha_5 - \frac{\sqrt{3}}{2} \alpha_6 \right) \sqrt{3}. \tag{C.8h}$$

Computing $\bar{\alpha}_5$ and $\bar{\alpha}_6$ from equations (C.8e) and (C.8f) gives

$$|\bar{\alpha}_5|^2 + |\bar{\alpha}_6|^2 = \frac{1}{v^2 \cos^2 \beta} \left(|\kappa_d|^2 + \frac{1}{3} |\kappa_\ell|^2 \right), \tag{C.9}$$

while computing α_5 and α_6 from equations (C.8g) and (C.8h) leads to

$$|\alpha_5|^2 + |\alpha_6|^2 = \frac{1}{v^2 \sin^2 \beta} \left(|\kappa_u|^2 + \frac{1}{3} |\kappa_D|^2 \right). \tag{C.10}$$

Finally, the consistency conditions (C.2) may be translated into the following conditions for the VEVs:

$$|k_d|^2 + |v_d|^2 + |\kappa_d|^2 + \frac{1}{3} |\kappa_\ell|^2 \leq v^2 \cos^2 \beta, \quad (\text{C.11a})$$

$$|k_u|^2 + |v_u|^2 + |\kappa_u|^2 + \frac{1}{3} |\kappa_D|^2 \leq v^2 \sin^2 \beta. \quad (\text{C.11b})$$

References

- [1] R. Slansky, Group theory for unified model building, *Phys. Rep.* 79 (1981) 1.
- [2] G.G. Ross, *Grand Unified Theories*, Addison–Wesley Publishing Company, Redwood City, California, 1985.
- [3] C.S. Aulakh, R.N. Mohapatra, Implications of supersymmetric $SO(10)$ grand unification, *Phys. Rev. D* 28 (1983) 217;
T.E. Clark, T.K. Kuo, N. Nakagawa, A $SO(10)$ supersymmetric grand unified theory, *Phys. Lett. B* 115 (1982) 26;
K.S. Babu, R.N. Mohapatra, Predictive neutrino spectrum in minimal $SO(10)$ grand unification, *Phys. Rev. Lett.* 70 (1993) 2845, arXiv:hep-ph/9209215;
L. Lavoura, Predicting the neutrino spectrum in minimal $SO(10)$ grand unification, *Phys. Rev. D* 48 (1993) 5440, arXiv:hep-ph/9306297;
B. Bajc, G. Senjanović, F. Vissani, b – τ unification and large atmospheric mixing: a case for a noncanonical seesaw mechanism, *Phys. Rev. Lett.* 90 (2003) 051802, arXiv:hep-ph/0210207;
H.S. Goh, R.N. Mohapatra, S.P. Ng, Minimal SUSY $SO(10)$ model and predictions for neutrino mixings and leptonic CP violation, *Phys. Rev. D* 68 (2003) 115008, arXiv:hep-ph/0308197;
B. Dutta, Y. Mimura, R.N. Mohapatra, CKM CP violation in a minimal $SO(10)$ model for neutrinos and its implications, *Phys. Rev. D* 69 (2004) 115014, arXiv:hep-ph/0402113;
T. Fukuyama, A. Ilakovac, T. Kikuchi, S. Meljanac, N. Okada, Detailed analysis of proton decay rate in the minimal supersymmetric $SO(10)$ model, *JHEP* 0409 (2004) 052, arXiv:hep-ph/0406068;
S. Bertolini, M. Malinský, On CP violation in minimal renormalizable SUSY $SO(10)$ and beyond, *Phys. Rev. D* 72 (2005) 055021, arXiv:hep-ph/0504241;
K.S. Babu, C. Macesanu, Neutrino masses and mixings in a minimal $SO(10)$ model, *Phys. Rev. D* 72 (2005) 115003, arXiv:hep-ph/0505200.
- [4] C.S. Aulakh, B. Bajc, A. Melfo, G. Senjanović, F. Vissani, The minimal supersymmetric grand unified theory, *Phys. Lett. B* 588 (2004) 196, arXiv:hep-ph/0306242.
- [5] C.S. Aulakh, A. Girdar, *Nucl. Phys. B* 711 (2005) 275, arXiv:hep-ph/0405074.
- [6] C.S. Aulakh, S.K. Garg, MSGUT: from bloom to doom, *Nucl. Phys. B* 757 (2006) 47, arXiv:hep-ph/0512224;
S. Bertolini, T. Schwetz, M. Malinský, Fermion masses and mixings in $SO(10)$ models and the neutrino challenge to SUSY GUTs, *Phys. Rev. D* 73 (2006) 115012, arXiv:hep-ph/0605006.
- [7] C.S. Aulakh, S.K. Garg, The new minimal supersymmetric GUT: spectra, RG analysis and fermion fits, *Nucl. Phys. B* 857 (2012) 101, arXiv:0807.0917 [hep-ph].
- [8] N. Oshimo, Antisymmetric Higgs representation in $SO(10)$ for neutrinos, *Phys. Rev. D* 66 (2002) 095010, arXiv:hep-ph/0206239;
N. Oshimo, Model for neutrino mixing based on $SO(10)$, *Nucl. Phys. B* 668 (2003) 258, arXiv:hep-ph/0305166;
B. Dutta, Y. Mimura, R.N. Mohapatra, Suppressing proton decay in the minimal $SO(10)$ model, *Phys. Rev. Lett.* 94 (2005) 091804, arXiv:hep-ph/0412105;
S. Bertolini, M. Frigerio, M. Malinský, Fermion masses in SUSY $SO(10)$ with type II seesaw: a nonminimal predictive scenario, *Phys. Rev. D* 70 (2004) 095002, arXiv:hep-ph/0406117;
W.M. Yang, Z.G. Wang, Fermion masses and flavor mixing in supersymmetric $SO(10)$ model, *Nucl. Phys. B* 707 (2005) 87, arXiv:hep-ph/0406221;
B. Dutta, Y. Mimura, R.N. Mohapatra, Neutrino mixing predictions of a minimal $SO(10)$ model with suppressed proton decay, *Phys. Rev. D* 72 (2005) 075009, arXiv:hep-ph/0507319;
A.S. Joshipura, B.P. Kodrani, K.M. Patel, Fermion masses and mixings in a μ – τ symmetric $SO(10)$, *Phys. Rev. D* 79 (2009) 115017, arXiv:0903.2161 [hep-ph];
A.S. Joshipura, K.M. Patel, Quasi-degenerate neutrinos in $SO(10)$, *Phys. Rev. D* 82 (2010) 031701, arXiv:1005.0045 [hep-ph];
A.S. Joshipura, K.M. Patel, Viability of the exact tri–bimaximal mixing at M_{GUT} in $SO(10)$, *JHEP* 1109 (2011) 137, arXiv:1105.5943 [hep-ph].

- [9] F. Feruglio, K.M. Patel, D. Vicino, A realistic pattern of fermion masses from a five-dimensional $SO(10)$ model, JHEP 1509 (2015) 040, arXiv:1507.00669 [hep-ph].
- [10] P.O. Ludl, A.Yu. Smirnov, Lepton mixing from the hidden sector, Phys. Rev. D 92 (2015) 073010, arXiv:1507.03494 [hep-ph].
- [11] D.V. Forero, M. Tórtola, J.W.F. Valle, Neutrino oscillations refitted, Phys. Rev. D 90 (2014) 093006, arXiv:1405.7540 [hep-ph].
- [12] F. Capozzi, G.L. Fogli, E. Lisi, A. Marrone, D. Montanino, A. Palazzo, Status of three-neutrino oscillation parameters, circa 2013, Phys. Rev. D 89 (2014) 093018, arXiv:1312.2878 [hep-ph];
M.C. Gonzalez-Garcia, M. Maltoni, T. Schwetz, Updated fit to three neutrino mixing: status of leptonic CP violation, JHEP 1411 (2014) 052, arXiv:1409.5439 [hep-ph];
J. Bergström, M.C. Gonzalez-Garcia, M. Maltoni, T. Schwetz, Bayesian global analysis of neutrino oscillation data, JHEP 1509 (2015) 200, arXiv:1507.04366 [hep-ph].
- [13] G. Altarelli, G. Blankenburg, Different $SO(10)$ paths to fermion masses and mixings, JHEP 1103 (2011) 133, arXiv:1012.2697 [hep-ph];
A.S. Joshipura, K.M. Patel, Fermion masses in $SO(10)$ models, Phys. Rev. D 83 (2011) 095002, arXiv:1102.5148 [hep-ph];
A. Dueck, W. Rodejohann, Fits to $SO(10)$ grand unified models, JHEP 1309 (2013) 024, arXiv:1306.4468 [hep-ph].
- [14] L. Lavoura, H. Kühböck, W. Grimus, Charged-fermion masses in $SO(10)$: analysis with scalars in $\mathbf{10} \oplus \mathbf{120}$, Nucl. Phys. B 754 (2006) 1, arXiv:hep-ph/0603259.
- [15] K. Matsuda, T. Fukuyama, H. Nishiura, $SO(10)$ GUT and quark–lepton mass matrices, Phys. Rev. D 61 (2000) 053001, arXiv:hep-ph/9906433;
T. Fukuyama, K. Matsuda, H. Nishiura, Zero texture model and $SO(10)$ GUT, Int. J. Mod. Phys. A 22 (2007) 5325, arXiv:hep-ph/0702284;
S. Dev, S. Kumar, S. Verma, S. Gupta, R.R. Gautam, Four zero texture fermion mass matrices in $SO(10)$ GUT, Eur. Phys. J. C 72 (2012) 1940, arXiv:1203.1403 [hep-ph].
- [16] W. Grimus, H. Kühböck, Fermion masses and mixings in a renormalizable $SO(10) \times \mathbb{Z}_2$ GUT, Phys. Lett. B 643 (2006) 182, arXiv:hep-ph/0607197.
- [17] B. Dutta, Y. Mimura, R.N. Mohapatra, Neutrino masses and mixings in a predictive $SO(10)$ model with CKM CP violation, Phys. Lett. B 603 (2004) 35, arXiv:hep-ph/0406262;
W. Grimus, H. Kühböck, A renormalizable $SO(10)$ GUT scenario with spontaneous CP violation, Eur. Phys. J. C 51 (2007) 721, arXiv:hep-ph/0612132.
- [18] G. Senjanović, $SO(10)$: a theory of fermion masses and mixings, arXiv:hep-ph/0612312.
- [19] J.C. Pati, A. Salam, Lepton number as the fourth color, Phys. Rev. D 10 (1974) 275, Phys. Rev. D 11 (1975) 703.
- [20] M. Magg, C. Wetterich, Neutrino mass problem and gauge hierarchy, Phys. Lett. B 94 (1980) 61;
G. Lazarides, Q. Shafi, C. Wetterich, Proton lifetime and fermion masses in an $SO(10)$ model, Nucl. Phys. B 181 (1981) 287;
R.N. Mohapatra, G. Senjanović, Neutrino masses and mixings in gauge models with spontaneous parity violation, Phys. Rev. D 23 (1981) 165;
R.N. Mohapatra, P. Pal, Massive Neutrinos in Physics and Astrophysics, World Scientific, Singapore, 1991, p. 127;
E. Ma, U. Sarkar, Neutrino masses and leptogenesis with heavy Higgs triplets, Phys. Rev. Lett. 80 (1998) 5716, arXiv:hep-ph/9802445.
- [21] P. Minkowski, $\mu \rightarrow e\gamma$ at a rate of one out of 10^9 muon decays?, Phys. Lett. B 67 (1977) 421;
T. Yanagida, Horizontal gauge symmetry and masses of neutrinos, in: O. Sawata, A. Sugamoto (Eds.), Proceedings of the Workshop on Unified Theory and Baryon Number in the Universe, Tsukuba, Japan, 1979, 1979, KEK report 79-18;
S.L. Glashow, The future of elementary particle physics, in: M. Lévy, et al. (Eds.), Quarks and Leptons, Proceedings of the Advanced Study Institute, Cargèse, Corsica, 1979, Plenum Press, New York, U.S.A., 1980;
M. Gell-Mann, P. Ramond, R. Slansky, Complex spinors and unified theories, in: D.Z. Freedman, F. van Nieuwenhuizen (Eds.), Supergravity, North Holland, Amsterdam, The Netherlands, 1979;
R.N. Mohapatra, G. Senjanović, Neutrino mass and spontaneous parity violation, Phys. Rev. Lett. 44 (1980) 912.
- [22] H. Fritzsch, Weak-interaction mixing in the six-quark theory, Phys. Lett. B 73 (1978) 317;
H. Fritzsch, Quark masses and flavor mixing, Nucl. Phys. B 155 (1979) 189;
L.F. Li, Comments on the derivation of the mixing angles, Phys. Lett. B 84 (1979) 461.
- [23] Z.z. Xing, H. Zhang, S. Zhou, Updated values of running quark and lepton masses, Phys. Rev. D 77 (2008) 113016, arXiv:0712.1419 [hep-ph].

- [24] C.R. Das, M.K. Parida, New formulas and predictions for running fermion masses at higher scales in SM, 2HDM, and MSSM, *Eur. Phys. J. C* 20 (2001) 121, arXiv:hep-ph/0010004;
S. Antusch, M. Spinrath, Quark and lepton masses at the GUT scale including SUSY threshold corrections, *Phys. Rev. D* 78 (2008) 075020, arXiv:0804.0717 [hep-ph].
- [25] K.A. Olive, et al., Particle Data Group Collaboration, Review of particle physics, *Chin. Phys. C* 38 (2014) 090001.
- [26] R. Storn, K. Price, Differential evolution – a simple and efficient heuristic for global optimization over continuous spaces, *J. Glob. Optim.* 11 (1997) 341.
- [27] C.S. Aulakh, A. Girdhar, $SO(10)$ à la Pati–Salam, *Int. J. Mod. Phys. A* 20 (2005) 865, arXiv:hep-ph/0204097.
- [28] I. Garg, New minimal supersymmetric $SO(10)$ GUT phenomenology and its cosmological implications, Ph.D. thesis, University of Panjab, 2014, arXiv:1506.05204 [hep-ph];
I. Garg, S. Mohanty, No scale SUGRA $SO(10)$ derived Starobinsky model of inflation, *Phys. Lett. B* 751 (2015) 7, arXiv:1504.07725 [hep-ph].
- [29] N. Yamatsu, Finite-dimensional Lie algebras and their representations for unified model building, arXiv:1511.08771 [hep-ph].
- [30] B. Bajc, A. Melfo, G. Senjanović, F. Vissani, Minimal supersymmetric grand unified theory: symmetry breaking and the particle spectrum, *Phys. Rev. D* 70 (2004) 035007, arXiv:hep-ph/0402122.
- [31] C.S. Aulakh, MSGUTs from germ to bloom: towards falsifiability and beyond, arXiv:hep-ph/0506291.