

Mouse model of human infertility: Transient and local inhibition of endometrial STAT-3 activation results in implantation failure

Hitomi Nakamura^{a,1}, Tadashi Kimura^{a,*}, Shinsuke Koyama^a, Kazuhide Ogita^a, Tateki Tsutsui^a, Koichiro Shimoya^a, Takeshi Taniguchi^b, Masayasu Koyama^a, Yasufumi Kaneda^c, Yuji Murata^a

^a Division of Obstetrics and Gynecology, Osaka University Graduate School of Medicine, 2-2, Yamadaoka, Suita, Osaka 5650871, Japan

^b Taniguchi Hospital, 1-5-20, Ohnishi, Izumisano, Osaka 5980043, Japan

^c Division of Gene Therapy Science, Osaka University Graduate School of Medicine, 2-2, Yamadaoka, Suita, Osaka 5650871, Japan

Received 22 February 2006; revised 6 April 2006; accepted 7 April 2006

Available online 21 April 2006

Edited by Robert Barouki

Abstract Embryo implantation involves a series of biochemical reactions and its failure is an important therapeutic target of infertility treatment. We established an infertile mouse model using transient and local suppression of signal transducer and activator of transcription-3 (STAT-3) activity by STAT-3 decoy transfer into the uterine cavity during implantation, resulting in <30% implantation. This infertility is caused by suppression of decidualization, which is indispensable for implantation, and independent of progesterone. These conditions may mimic clinically unexplained infertility. Our results suggest that STAT-3 could be a useful target for diagnosis and therapy of human implantation failure.

© 2006 Federation of European Biochemical Societies. Published by Elsevier B.V. All rights reserved.

Keywords: STAT-3; Implantation failure; Infertility; In vivo gene transfer; Molecular therapy; HVJ-E vector

1. Introduction

Approximately 10–15% of all couples experience infertility during their reproductive lives [1]. Implantation failure is considered to be a major reason for their infertility. One third of implantation failures might be attributed to the embryo itself. The remaining 2/3 of failures appear to be the result of inadequate uterine receptivity [2]. It is thus important to improve uterine receptivity in infertility treatment. However, the pathophysiology of implantation failure is unexplained as is the molecular mechanism of implantation. Uterine receptivity has been clinically evaluated via serum progesterone levels, ultrasonographic endometrial thickness, and histological dating by endometrial biopsy [3]. A discrepancy between the hormonal milieu and endometrial morphology and/or function could be an important cause of implantation failure [4],

although there is no consensus description for the diagnosis to date. We hypothesize that a few multi-potential transcriptional factors might initiate the regulatory cascade and have pivotal roles in generating the implantation window. Therefore, we devised a local and transient in vivo gene transfer system to modulate the function of endometrial signaling molecules during the implantation window, without disturbing the course of pregnancy [5]. STATs are a family of latent cytoplasmic proteins involved in transmitting extra-cellular signals to the nucleus [6]. Signal transducer and activator of transcription-3 (STAT-3) is activated by gp130, a subunit of the receptor for the interleukin-6 (IL-6) family of cytokines, including leukemia inhibitory factor (LIF) and IL-11. As gene targeting analysis indicated LIF and IL-11 play crucial roles in implantation [7,8], the physiological function of STAT-3 activation in implantation is of great interest. However, STAT-3 deficiency in embryos is lethal, and obtaining adult STAT-3 null female mice impossible [9]. Here, we perturbed STAT-3 activation in the mouse uterus during the implantation period using a STAT3-decoy/transient in vivo DNA transfer system to establish an implantation failure model.

2. Materials and methods

2.1. Animals

Female ICR mice (SLC, Shizuoka, Japan) aged 8–10 weeks in estrus were bred with male ICR mice, and the morning when vaginal plugging was observed was designated as day 0.5 post coitus (p.c.). Animal experiments were performed according to the guidelines for animal use approved by the Institutional Animal Care and Use Committee of Osaka University Graduate School of Medicine.

2.2. STAT-3 decoy transfer

A STAT-3 binding consensus sequence which acts as a decoy *cis*-element (5'-CCTTCCGGGAATTCCTTCCGGGAATTC-3'; underlines indicate consensus element) [10] and a scramble sequence (5'-AGTCCATTCGGCAGGCCTCTGCTCTAT-3'), were synthesized as phosphorothioate-modified oligodeoxynucleotides (ODNs), and purified by HPLC (Hokkaido System Science, Sapporo, Japan). Decoy ODNs were prepared by annealing of sense and antisense ODNs. Double stranded decoy ODNs were transferred to the murine uterus by HVJ-E vector (GenomONE-NEO®, Ishihara Sangyo Co. Ltd. Osaka, Japan) as described previously [5,11].

2.3. Tissue sampling and analysis of implantation/course of pregnancy

The mice on day 5.0 p.c., after in vivo gene transfer, were sacrificed by administration of excess anesthetic. The uteri were removed and frozen in liquid nitrogen. Day 5.0 p.c. pregnant animals received an

*Corresponding author. Fax: +81 6 6879 3359.

E-mail address: tadashi@gyne.med.osaka-u.ac.jp (T. Kimura).

¹ Present address: Research Centre for Reproductive Health, Department of Obstetrics and Gynaecology, University of Adelaide, SA 5005, Australia.

Abbreviations: STAT-3, signal transducer and activator of transcription-3; LIF, leukemia inhibitory factor; IL, interleukin; HVJ-E, (hemagglutinating virus of Japan, also known as Sendai virus)-envelope; HTF, human tubal fluid medium; ODN, oligodeoxynucleotide

i.v. injection of 0.5% Evans Blue 15 min prior to sacrifice to visualize implantation sites [12]. The serum progesterone level on day 5.0 p.c. was measured using a DPC progesterone RIA kit (Diagnostic Production Co. Ltd., LA, USA).

2.4. Activity assay of active STAT-3

Nuclear proteins from uterine tissue were prepared as described [13]. Nuclear extract was aliquoted, frozen in liquid nitrogen and stored at -80°C . Protein concentration was determined using a protein assay kit (Bio-Rad Laboratories, Inc. CA) and the amount of activated STAT-3 was assayed using intracellular DuoSet[®] IC (R&D systems, Inc. MN), according to the manufacturer's instructions.

2.5. Detection of transfected DNA in embryos

The cDNA of murine STAT-3 Y705 F, a dominant negative mutant of STAT-3, was distributed from RIKEN BRC DNA Bank (RDB & 2354). The open reading frame of STAT-3 Y705F was excised and introduced into pcDNA3 (Invitrogen, San Diego, CA) (pcDNA3-STAT-3 Y705F). Plasmid DNA was purified using a Qiagen column (Tokyo, Japan). Transfection of pcDNA3-STAT-3 Y705F with HVJ-E vector was performed as previously described [11] and blastocysts were flushed out from the uterus on day 4.0 p.c. The DNA was extracted from each single blastocyst and PCR was performed as described previously [11].

2.6. Induction of decidualization

To induce experimental decidualization, pseudopregnant female mice mated with vasectomized male mice received an intraluminal infusion of 10 μl sesame oil in one uterine horn on day 3.5 p.c. and were sacrificed after 72 h as described by Rider et al. [14].

2.7. RT-PCR

Total RNA was extracted by using TRIZOL[®] (Invitrogen). Single-stranded cDNA was synthesized from 3 μg of total RNA using SuperScript[™] II (Invitrogen). The PCR for progesterone receptor-A was performed using rTaq DNA polymerase (Toyobo, Osaka, Japan) as described by Clemens et al. [15].

2.8. Statistical evaluation of results

Statistical analysis was performed using the Mann–Whitney *U* test.

3. Results and discussion

3.1. Transfection of STAT-3 decoy suppressed STAT-3 activity and pregnancy

STAT-3 activity in the uterine nuclear extract on day 5.0 p.c. after STAT-3 decoy transfection was significantly suppressed compared to control uteri (Fig. 1A). On day 15.5 p.c., we examined viable fetuses in utero. In the group treated with vehicle (human tubal fluid medium, HTF) and scramble decoy, all mice became pregnant. However, in mice treated with the STAT-3 decoy, approximately 70% of mice lacked viable fetuses (Fig. 1B). An alternative STAT-3 decoy with the single cis-binding sequence, (5'-GATCCTTCTGGGAATTCCTA-GATC-3', 3'-CTAGGAAGACCCTTAAGGATCTAG-5', consensus sequences) failed to suppress pregnancy (100% of mice became pregnant after transfection; data not shown). Using this gene transfer method, we previously showed that the transferred gene did not affect embryos and the course of pregnancy, i.e., in luciferase cDNA transferred mice, embryos developed and were delivered normally [11] and I κ B α M cDNA, a dominant negative mutant of I κ B to suppress the activation of nuclear factor κ B, transferred mice delivered their pups 1 day after the normal due date [5]. We flushed out normal blastocysts on 4.0 p.c. from STAT-3 decoy transfer uteri (Fig. 2A), although no embryo was recovered from scramble




Fig. 1. Effect of STAT-3 decoy transfer in mouse uterus. The mice had vehicle (HTF), scramble or STAT-3 decoy transferred on day 1.5 p.c. (A) STAT-3 activity in nuclear extract from whole uterus at 5.0 p.c. Each group consisted of at least 5 mice. All bars represent the mean \pm S.E.M. * $P < 0.0005$ by the Mann–Whitney *U* test. (B) Pregnancy rate on day 15.5 p.c. * $P < 0.05$ by the Mann–Whitney *U* test.

decoy or pcDNA3 transferred uterus on 4.0 p.c. The transferred gene after pcDNA3-STAT-3 Y705F was undetectable in flushed out embryos by PCR (Fig. 2B). These previous and present observations showed that STAT-3 decoy transfer had neither killed the blastocysts nor washed them out through the uterus.

In the STAT-3 decoy transfer group, 75% of mice did not exhibit implantation sites on day 5.0 p.c. detected by Evans Blue. In the scramble decoy transfer group, implantation sites were evident in all mice (Fig. 3A and B). On day 8.0 p.c., in control uteri, embryos had already formed primitive somites. However, decidualization was apparently disturbed in the STAT-3 decoy treated uterus (Fig. 3C), indicating that decreased implantation sites on day 5.0 p.c. were not due to delayed implantation of dormant embryos. Therefore, STAT-3 suppression led infertility was caused by inadequate uterine receptivity, i.e., implantation failure.

3.2. Decidualization response

As decidualization was totally inhibited in the STAT-3 decoy transfected uterus on day 8.0 p.c., we examined the effect of STAT-3 suppression on mechanical decidualization. The decidual response was obviously suppressed in the STAT-3 decoy transfected uterus (Fig. 4A and B).




Fig. 4. Experimentally induced decidualization. (A) Histological examination 72 h after oil injection revealed decidualization (a, c) many decidualized cells (c) in scramble decoy transfected uterus. STAT-3 decoy transfected uterus was smaller than control and (b) there were many glands and less characterized decidualization (d). Scale bars: (a, b) 500 μ m; (c, d) 100 μ m and (e, f) 50 μ m. (B) Fold-increase of uterine wet weights, measured 72 h after oil injection. All bars represent the mean \pm S.E.M. from at least five animals. * $P = 0.002$.

tions similar to those observed in LIF knockout mice [28,29]. Taken together with our present data, full activation of STAT-3, rather than signaling via STAT-1 and SHP2/ERK

are necessary for uterine receptivity. In a preliminary study we tested the activation of STAT-3 in human endometrium from unexplained infertile women. As activation of STAT-3




Fig. 5. Ovarian function. (A) On day 5.0 p.c. in both ovaries, corpus luteums were obvious and no significant difference was observed after scramble decoy (a) and STAT-3 decoy (b) transfection. (B) Progesterone levels in peripheral blood on day 5.0 p.c. were similar in STAT-3 and scramble decoy transfer groups. Bars represent the mean \pm S.E.M. from at least five animals. (C) The level of progesterone receptor (PR)-A transcripts in uterus on day 5.0 p.c. were similar in vehicle (HTF), scramble decoy and STAT-3 decoy transferred groups.

was disturbed in the mid-luteal phase endometrial tissue of some infertile patients (unpublished data), STAT-3 could be one of the diagnostic and therapeutic targets for unexplained infertility along with IL-11 [7,8,24] and Wnt/ β -catenin signaling [30]. Transcription factors comprise useful molecular therapies to improve fertility or alternatively to achieve contraception. The uterus is a tubular organ and its inner endometrial surface is accessible via the cervix, therefore the endometrium could be a good candidate for local, transient molecular target therapy. Our in vivo gene transfer system into the mouse uterus has proven to be a good experimental tool for mining the opportunities for molecular therapy of implantation failure.

Acknowledgements: We thank Ms. Satomi Okamoto for technical assistance, Drs. Virginia Rider (Pittsburg Univ.), Yasuo Kiso (Yamaguchi Univ.), Kazuhiko Imakawa (Tokyo Univ.), Richard Ivell and Sarah A Robertson (Adelaide Univ.) for critical advice about our experiments. HVJ-E vector (GenomONE-NEO®) was provided by Ishihara Sangyo Co. Ltd. This work was supported in part by

Grants-in-Aid for Scientific Research (Nos. 15209054, 16390476, 17591733, 17591734, and 17591735) from the Ministry of Education, Science and Culture of Japan (Tokyo, Japan).

References

- [1] Speroff, L., Glass, R.H. and Kase, N.G. (1999) Clinical Gynecologic Endocrinology and Infertility, 6th edn, Lippincott Williams & Wilkins, Baltimore, MD.
- [2] Simon, C., Moreno, C., Remohi, J. and Pellicer, A. (1998) Cytokines and embryo implantation. *J. Reprod. Immunol.* 39, 117–131.
- [3] Bukulmez, O. and Arici, A. (2004) Luteal phase defect: myth or reality. *Obstet. Gynecol. Clin. North Am.* 31, 727–744.
- [4] Li, T.C. et al. (1993) A study of endometrial morphology in women who failed to conceive in a donor insemination programme. *Br. J. Obstet. Gynaecol.* 100, 935–938.
- [5] Nakamura, H. et al. (2004) Alteration of the timing of implantation by in vivo gene transfer: delay of implantation by suppression of nuclear factor kappaB activity and partial rescue by leukemia inhibitory factor. *Biochem. Biophys. Res. Commun.* 321, 886–892.
- [6] Levy, D.E. and Darnell Jr., J.E. (2002) Stats: transcriptional control and biological impact. *Nat. Rev. Mol. Cell Biol.* 3, 651–662.
- [7] Stewart, C.L. et al. (1992) Blastocyst implantation depends on maternal expression of leukemia inhibitory factor. *Nature* 359, 76–79.
- [8] Robb, L. et al. (1998) Infertility in female mice lacking the receptor for interleukin 11 is due to a defective uterine response to implantation. *Nat. Med.* 4, 303–308.
- [9] Takeda, K. et al. (1997) Targeted disruption of the mouse Stat3 gene leads to early embryonic lethality. *Proc. Natl. Acad. Sci. USA* 94, 3801–3804.
- [10] Akira, S. et al. (1994) Molecular cloning of APRF, a novel IFN-stimulated gene factor 3 p91-related transcription factor involved in the gp130-mediated signaling pathway. *Cell* 77, 63–71.
- [11] Nakamura, H. et al. (2003) Highly efficient and minimally invasive in-vivo gene transfer to the mouse uterus using haemagglutinating virus of Japan (HVJ) envelope vector. *Mol. Hum. Reprod.* 9, 603–609.
- [12] Huet, Y.M. and Dey, S.K. (1987) Role of early and late oestrogenic effects on implantation in the mouse. *J. Reprod. Fert.* 81, 453–458.
- [13] Nakamura, H. et al. (2004) NF kappaB activation at implantation window of the mouse uterus. *Am. J. Reprod. Immunol.* 51, 16–21.
- [14] Rider, V., McRae, A., Heap, R.B. and Feinstein, A. (1985) Passive immunization against progesterone inhibits endometrial sensitization in pseudopregnant mice and has antifertility effects in pregnant mice which are reversible by steroid treatment. *J. Endocrinol.* 104, 153–158.
- [15] Clemens, J.W. et al. (1998) Hormone induction of progesterone receptor (PR) messenger ribonucleic acid and activation of PR promoter regions in ovarian granulosa cells: evidence for a role of cyclic adenosine 3',5'-monophosphate but not estradiol. *Mol. Endocrinol.* 12, 1201–1214.
- [16] Coutifaris, C. et al. NICHD National Cooperative Reproductive Medicine Network (2004) Histological dating of timed endometrial biopsy tissue is not related to fertility status. *Fertil. Steril.* 82, 1264–1272.
- [17] Myers, E.R. et al. NICHD National Cooperative Reproductive Medicine Network. (2004) Interobserver and intraobserver variability in the histological dating of the endometrium in fertile and infertile women. *Fertil. Steril.* 82, 1278–1282.
- [18] Hambartsumian, E. (1998) Endometrial leukemia inhibitory factor (LIF) as a possible cause of unexplained infertility and multiple failures of implantation. *Am. J. Reprod. Immunol.* 39, 137–143.
- [19] Mikołajczyk, M., Skrzypczak, J., Szymanowski, K. and Wirstlein, P. (2003) The assessment of LIF in uterine flushing—a possible new diagnostic tool in states of impaired fertility. *Reprod. Biol.* 3, 259–270.

- [20] Ledee-Bataille, N. et al. (2002) Concentration of leukemia inhibitory factor (LIF) in uterine flushing fluid is highly predictive of embryo implantation. *Hum. Reprod.* 17, 213–218.
- [21] Olivennes, F. et al. (2003) Assessment of leukemia inhibitory factor levels by uterine flushing at the time of egg retrieval does not adversely affect pregnancy rates with in vitro fertilization. *Fertil. Steril.* 79, 900–904.
- [22] Giess, R., Tanasescu, I., Steck, T. and Sendtner, M. (1999) Leukaemia inhibitory factor gene mutations in infertile women. *Mol. Hum. Reprod.* 5, 581–586.
- [23] Sherwin, J.R.A., Smith, S.K., Wilson, A. and Sharkey, A.M. (2002) Soluble gp130 is up-regulated in the implantation window and shows altered secretion in patients with primary unexplained infertility. *J. Clin. Endocrinol. Metab.* 87, 3953–3960.
- [24] Bilinski, P., Roopenian, D. and Gossler, A. (1998) Maternal IL-11R α function is required for maternal deciduas and fetoplacental development in mice. *Genes Dev.* 12, 2234–2243.
- [25] Ernst, M. et al. (2001) Defective gp130-mediated signal transducer and activator of transcription (STAT) signaling results in degenerative joint disease, gastrointestinal ulceration, and failure of uterine implantation. *J. Exp. Med.* 194, 189–203.
- [26] Cheng, J.G., Chen, J.R., Hernandez, L., Alvord, W.G. and Stewart, C.L. (2001) Dual control of LIF expression and LIF receptor function regulate Stat3 activation at the onset of uterine receptivity and embryo implantation. *Proc. Natl. Acad. Sci. USA* 98, 8680–8685.
- [27] Teng, C-B. et al. (2004) Signal transducer and activator of transcription 3 (Stat3) expression and activation in rat uterus during early pregnancy. *Reproduction* 128, 197–205.
- [28] Catalano, R.D. et al. (2005) Inhibition of Stat3 activation in the endometrium prevents implantation: a nonsteroidal approach to contraception. *Proc. Natl. Acad. Sci. USA* 102, 8585–8590.
- [29] Rodriguez, C.I., Cheng, J.G., Liu, L. and Stewart, C.L. (2004) Cochlin, a secreted von Willebrand factor type a domain-containing factor, is regulated by leukemia inhibitory factor in the uterus at the time of embryo implantation. *Endocrinology* 145, 1410–1418.
- [30] Mohamed, O.A. et al. (2005) Uterine Wnt/beta-catenin signaling is required for implantation. *Proc. Natl. Acad. Sci. USA* 102, 8579–8584.