



ELSEVIER

Contents lists available at SciVerse ScienceDirect

Results in Immunology

journal homepage: www.elsevier.com/locate/rinim

Rondonin an antifungal peptide from spider (*Acanthoscurria rondoniae*) haemolymph

K.C.T. Riciluca^{a,b}, R.S.R. Sayegh^a, R.L. Melo^a, P.I. Silva Jr.^{a,*}^a Laboratório Especial de Toxinologia Aplicada, Instituto Butantan, 05503-900 São Paulo – SP, Brazil^b Coordenadoria de Controle de Doenças – CCD, São Paulo, Brazil

ARTICLE INFO

Article history:

Received 30 January 2012

Received in revised form

20 March 2012

Accepted 22 March 2012

Available online 2 April 2012

Keywords:

Antimicrobial peptide

Haemocyanin fragment

Rondonin

Acanthoscurria rondoniae

Spider

ABSTRACT

Antimicrobial activities were detected in the haemolymph of the spider *Acanthoscurria rondoniae*. A novel antifungal peptide, rondonin, was purified by reverse phase high performance liquid chromatography (RP-HPLC). Rondonin has an amino acid sequence of IIIQYEGHKH and a molecular mass of 1236.776 Da. This peptide has identity to a C-terminal fragment of the “d” subunit of haemocyanin from the spiders *Eurypelma californicum* and *Acanthoscurria gomesiana*. A synthetic peptide mimicking rondonin had identical characteristics to those of the isolated material, confirming its sequence. The synthetic peptide was active only against fungus. These data led us to conclude that the antifungal activity detected in the plasma of these spiders is the result of enzymatic processing of a protein that delivers oxygen in the haemolymph of many chelicerate. Several studies have suggested that haemocyanins are involved in the arthropod immune system, and the activity of this haemocyanin fragment reinforces this idea.

© 2012 Elsevier B.V. Open access under [CC BY-NC-ND license](http://creativecommons.org/licenses/by-nc-nd/4.0/).

1. Introduction

Invertebrate animals can be found in almost every habitat in the world. Because many invertebrates live in environments in which microorganisms thrive, their widespread distribution and survival are primarily due to successful defences that efficiently recognise and combat potentially harmful microorganisms [37]. Invertebrates only possess innate immunity, which is considered to be an ancient defence mechanism [16].

One characteristic of innate immunity is the production of antimicrobial substances, which are often peptides or polypeptides [12]. Several of these antimicrobial peptides (AMPs) have been recognised as important components of the nonspecific host defence or innate immune system in a variety of organisms and have been isolated and characterised from plants and animals, including insects, molluscs, crustaceans, amphibians, birds, fish, mammals, and humans [2,17,3,11]. Several antimicrobial peptides were isolated from the venom and haemolymph of venomous arthropods such as scorpions and spiders [22]. The haemolymph of invertebrates are the main source of antimicrobial peptides [14]. The first biochemical study of an antimicrobial peptide in arachnids demonstrated the presence of an antibacterial peptide

in the haemolymph of the scorpion species *Leiurus quinquestriatus* [5] and *Androctonus australis* [8]. Gomesin was the first antimicrobial peptide isolated from spider blood cells [34]. Other antimicrobial peptides were found in the plasma of shrimp [6], freshwater crayfish [24], the plasma of the tarantula spider *Acanthoscurria gomesiana*, named theraphosinin [34], and crude haemolymph of *Agelena labyrinthica* [46]. These peptides are typically relatively short, positively charged (cationic), and amphiphilic [19,40] and generally interact with the outer membranes of microorganisms due to their negative charge [20]. Compared to cationic AMPs, much less is known about how anionic AMPs work [39,36,13].

While most AMPs are derived from a biologically inactive proprotein that is processed into an active state, some AMPs are derived from larger, functional proteins, such as haemoglobin [10,23,25,41] and haemocyanin [6,24]. Arthropod haemocyanins are composed of heterogeneous subunits in the 75-kDa range that combine to form either a regular cubic single hexamer (1 × 6) or multiple hexamers (2–8 × 6) depending upon the species and physiological conditions [29]. The haemocyanin of the North American tarantula *Eurypelma californicum* is a native 24-mer protein complex consisting of two identical dodecamers with an estimated total molecular mass of approximately 1800 kDa [33,26,27,29]. Formation of the 24-mer complex requires the aggregation of seven different subunits in a constant stoichiometric amount with four copies of each of the subunits *a*, *d*, *e*, *f*, and *g* and two copies of subunits *b* and *c* [28,29]. Since

* Correspondence to: Laboratório Especial de Toxinologia Aplicada, CAT/CEPID, Instituto Butantan, Av: Vital Brasil, 1500 CEP, 05503-900 São Paulo – SP, Brazil. Tel./fax: +55 11 3726 1024.

E-mail addresses: pisjr@usp.br, pisjr@butantan.gov.br (P.I. Silva Jr.).

antimicrobial peptides have been characterised in haemocytes of *A. gomesiana* and *Acanthoscurria rondoniae* which belongs to the same genera, we choose this species to look for the presence of these peptides. So, in the present study, we report the first isolation and characterisation of an antifungal fragment of haemocyanin from arachnids.

2. Experimental procedures

2.1. Microorganisms

Fungal and bacterial strains were obtained from various sources. *Escherichia coli* SBS363 and *Micrococcus luteus* A270 were from the Pasteur Institut, Paris; *Candida albicans* (MDM8) was from the Department of Microbiology from the University of São Paulo, Brazil; and *E. coli* ATCC 25922, *Pseudomonas aeruginosa* ATCC 27853 (Strain Boston 41501), *Staphylococcus aureus* ATCC 29213 and *S. epidermidis* ATCC 12228 were from the American Type Culture Collection (ATCC). The following human clinical yeast isolates, which can be agents of candidiasis disease, obtained from the Oswaldo Cruz Institute, Brazil, were also used: *Trichosporon* sp. IOC 4569, *Candida krusei* IOC 4559, *C. glabrata* IOC 4565, *C. albicans* IOC 4558, *C. parapsilosis* IOC 4564, *C. tropicalis* IOC 4560 and *C. guilliermondii* IOC 4557. The filamentous fungi *Aspergillus niger*, *Cladosporium* sp. and *Penicillium expansum* and *Beauveria bassiana*, an entomopathogenic fungus, were isolated from a mummified spider.

2.2. Spider blood

The spiders (*Acanthoscurria rondoniae*, a tarantula of the Theraphosidae family) were kept alive in the biotherium of the Special Laboratory of Toxinology Applied of the Institute Butantan (São Paulo, Brazil) (Fig. 1). These animals were collected under Licence Permanent Zoological Material no. 11024-3-IBAMA and Special Authorisation for Access to Genetic Patrimony no. 001/2008. The haemolymph (approximately 10 mL) from animals of either sex at different stages of development was collected by cardiac puncture with an apyrogenic syringe. To avoid haemocyte degranulation and coagulation, the haemolymph was collected in the presence of sodium citrate buffer (0.14 M NaCl, 0.1 M glucose, 30 mM trisodium citrate, 26 mM citric acid, 10 mM EDTA, pH 4.6 (2:1, v/v)) [38]. The plasma was separated, and the blood cells were removed by centrifugation at 800g for 10 min at 4 °C.

2.3. Antimicrobial activity assays

During the purification procedure, the antimicrobial activities of the samples were monitored by a liquid growth inhibition



Fig. 1. Specimen of the spider *Acanthoscurria rondoniae* (Theraphosidae, Mygalomorphae).

assay using the Gram-negative bacteria *Escherichia coli* SBS363, Gram-positive bacteria *Micrococcus luteus* A270 that were cultured in poor broth nutrient medium (PB: 1.0 g peptone in 100 ml of water containing 86 mM NaCl at pH 7.4; 217 mOsM), whereas yeast strain *Candida albicans* MDM8 was cultured in poor dextrose broth (1/2 PDB: 1.2 g potato dextrose in 100 ml of H₂O at pH 5.0; 79 mOsM). Determination of antimicrobial peptide was performed using 5-fold microtiter broth dilution assay in 96-well sterile plates at a final volume of 100 µL. Mid-log phase culture were diluted to a final concentration of 1×10^5 colony forming units/mL. Dried fractions were dissolved in 200 µL of water ultrapure and 20 µL applied into each well and added to 80 µL of the bacterium/yeast dilution. The fractions were tested in triplicate. The microtiter plates were incubated for 18 h at 30 °C; growth inhibition was determined by measuring absorbance at 595 nm.

2.4. Purification of rondonin from plasma

For purification of antimicrobial peptides, the plasma was homogenised and then trifluoroacetic acid was added to a final concentration of 0.05%. The sample was agitated on ice for 30 min and centrifuged at 16,000g at 4 °C. The acidic supernatant was loaded onto classic Sep-Pak C18 cartridges equilibrated in acidified water (TFA 0.05%). After washing with acidified water, three stepwise elutions were successively performed with 5%, 40% and 80% acetonitrile (ACN/TFA 0.05%). The 40% Sep-Pak fraction was concentrated in a vacuum centrifuge and reconstituted in Milli-Q water (Millipore™) and directly subjected to C18 reverse-phase on a semi-preparative Jupiter C18 column equilibrated at room temperature with 0.05% trifluoroacetic acid in water. The sample was purified using acetonitrile/water/0.05% trifluoroacetic acid gradients of 2–60% acetonitrile in 60 min at a flow rate 1.5 mL/min. Ultraviolet absorbance was monitored at 225 nm. The eluted peaks fractions were collected by hand and were vacuum dried (Speed-Vac Savant) and used for assay of antimicrobial activity and determination of amino acid sequence.

2.5. Rondonin mass analysis by MALDI-TOF and LC/MS

Briefly, 0.35 µL of sample in Milli-Q water was mixed with 0.35 µL of saturated matrix α -cyano-4-hydroxycinnamic acid solution deposited onto the sample slide and dried on the bench. The analysis was performed with the spectrometer operating in positive mode, which detects positively charged ions. To determine the amino acid sequence of rondonin, the doubly charged ions were subjected to “de novo” sequencing in a Q-TOF Ultima API (Micromass) spectrometer operating in positive ionisation mode. The spectrum was analysed, and the “y” and “b” fragments were used to elucidate the primary structure of the molecule.

2.6. Peptide synthesis of rondonin

Synthetic rondonin was synthesised by solid phase peptide synthesis using the Fmoc procedure [1]. The peptide was purified by reverse phase (Shim-pack Prep-ODS, 5 µ, 20 mm × 250 mm Shimadzu Co.) semi-preparative HPLC, and the purity and identity of the peptide were confirmed by MALDI-TOF mass spectrometry and analytical HPLC using the conditions described above.

2.7. Minimal inhibitory concentration (MIC) of rondonin

The minimal inhibitory concentration was determined using the synthetic peptide against the Gram-negative bacterial strains, the Gram-positive bacterial strains, the fungal strains and the yeast strains, as described above (experimental procedures 2.1 and 2.3).

The peptide was dissolved in sterile Milli-Q water at a final concentration of 670 μM . Determination of minimal inhibitory concentrations (MICs) for rondonin was performed using a 5-fold microtiter broth dilution assay of stock solution (670 μM) and serial dilution in 96-well sterile plates at a final volume of 100 μL where 20 μL of stock solution was applied into each well at serial dilution 2-fold microtiter broth dilution and added to 80 μL of the bacterium/yeast dilution. Microbial growth was measured by monitoring the increase in OD at 595 nm after incubation at 30 °C for 18 h (modified [8]). The rondonin was tested in duplicate. The MIC is defined as the minimal concentration of peptide that caused 100% growth inhibitions [47].

2.8. Antifungal assay of rondonin

The antifungal assay was performed using a 5-fold microtiter broth dilution assay and serial dilution in 96-well sterile plates at a final volume of 100 μL where 20 μL of stock solution (670 μM) was applied into each well at serial dilution 2-fold microtiter broth dilution and added to 80 μL of the yeast dilution. The inhibition growth curve of rondonin was determined by incubating twice the concentration of the MIC (67 μM) of rondonin with *C. albicans* MDM8 at 30 °C for various amounts of time (0, 10 min, 1 h, 3 h, 5 h, 8 h, 10 h, 12 h, 18 h, and 24 h) and counting the number of conidia present; the viability of the yeast was verified by incubating the colonies on a nutrient agar plate (1.5%). The rondonin was tested in triplicate.

2.9. Haemolytic assay to determine rondonin toxicity

Human erythrocytes from a healthy donor were collected in 0.15 M citrate buffer, pH 7.4, and washed three times by centrifugation (700g, 10 mins, 4 °C) with 0.15 M phosphate-buffered saline (PBS), pH 7.4. After the final centrifugation, the erythrocytes were suspended in PBS, pH 7.4. Aliquots of 50 μL containing rondonin at concentrations ranging from 0.2 to 134 μM were added to 50 μL of a 3% suspension of erythrocytes in the wells of

U-shaped bottom plates and incubated for 3 h at 37 °C. The supernatant was first collected and haemolysis was determined by measuring the absorbance at 414 nm of each well in a Victor³ (1420 Multilabel Counter/Victor³, Perkin Elmer). The haemolysis percentage was expressed in relation to a 100% lysis control (erythrocytes incubated with 0.1% triton X-100); PBS was used as a negative control. The rondonin was tested in triplicate.

3. Results

3.1. Antimicrobial activity of rondonin measured by liquid growth inhibition

Antimicrobial activity of synthetic rondonin was screened using 3 species of Gram-positive bacteria, 3 species of Gram-negative bacteria, 7 species of yeast and 2 species of fungi. Rondonin was tested against these organisms at concentrations ranging from 0.1 to 67 μM and demonstrated antimicrobial activity against all 7 species of yeast and one species of filamentous fungus (Table 1).

3.2. Purification of rondonin from plasma

The plasma of the spider *A. rondoniae* from the Theraphosidae family was collected (10 mL) and dissolved in acidified Milli-Q water as previously described. The supernatant obtained by centrifugation was applied to a Sep-Pak C18 column and subjected to three successive extractions of increasing concentrations of acetonitrile (5%, 40% and 80% ACN) to pre-purify antimicrobial peptides. The material eluted at 40% ACN was subjected to fractionation by RP-HPLC, which resulted in fractions with antimicrobial activity (Fig. 2). All fractions were analysed in the liquid growth inhibition assay using *M. luteus*, *E. coli*, and *C. albicans*. We found six fractions that showed antimicrobial activity only against *C. albicans*: 2, 3, 7, 11, 12, and 13. Only the fraction 2, named rondonin, was purified to homogeneity.

Table 1
Antimicrobial activity spectrum of synthetic rondonin compared to gomesin. The minimal inhibitory concentrations (MICs) of the synthetic peptides were determined by liquid growth inhibition. MICs are expressed as the (a) and (b) interval of concentrations where (a) is the highest concentration tested at which the microorganisms are growing and (b) is the lowest concentration that causes 100% growth inhibition.

Microorganisms	Minimal inhibitory concentration (μM)					
	Rondonin		Gomesin			
Gram-positive bacteria						
<i>Micrococcus luteus</i> A270	ND		0.4	–		0.8 ^a
<i>Staphylococcus aureus</i> ATCC 29213	ND		1.2	–		2.5 ^b
<i>Staphylococcus epidermidis</i> ATCC 12228	ND			NA		
Gram-negative bacteria						
<i>Escherichia coli</i> SBS363	ND		0.4	–		0.8 ^a
<i>Escherichia coli</i> ATCC 25922	ND		0.2	–		0.9 ^b
<i>Pseudomonas aeruginosa</i> ATCC 27853 (Strain Boston 41501)	ND			NA		
Fungi						
<i>Beauveria bassiana</i>	ND			NA		
<i>Trichosporon</i> sp IOC 4569	1.1	–	2.1	0.6	–	1.2 ^b
Yeasts						
<i>Candida albicans</i> MDM8	16.75	–	33.5	0.15	–	0.3 ^a
<i>Candida krusei</i> IOC 4559	16.75	–	33.5	5.0	–	10.0 ^b
<i>Candida glabrata</i> IOC 4565	8.37	–	16.5	12.5	–	25
<i>Candida albicans</i> IOC 4558	8.37	–	16.5	5.0	–	10.0 ^b
<i>Candida parapsilosis</i> IOC 4564	16.75	–	33.5	2.5	–	5.0 ^b
<i>Candida tropicalis</i> IOC 4560	8.75	–	16.5	0.3	–	0.6 ^b
<i>Candida guilliermondii</i> IOC 4557	16.75	–	33.5	2.5	–	5.0 ^b

The highest concentration tested was 67 μM . ND, activity not detected in the range of concentration tested; NA, activity not available.

^a Silva et al. [35].

^b Yamane [44].

3.3. Minimal inhibitory concentration

The MIC of synthetic rondonin was tested against Gram-negative bacterial strains, Gram-positive bacterial strains, fungal strains and yeast strains. Synthetic rondonin showed activity against all yeasts tested and one fungus. However, no activity could be detected against Gram-positive and Gram-negative bacteria and the fungus *Aspergillus niger*, *Cladosporium* sp.,

Penicillium expansum and *B. bassiana* strains tested in the range of concentration investigated (above 67 μ M). MICs are expressed as the (a) and (b) interval of concentrations where (a) is the highest concentration tested at which the microorganisms are growing and (b) is the lowest concentration that causes 100% growth inhibition. We compare our results with the synthetic gomesin performed by Silva et al. [35] and Yamane [44]. As shown in Table 1, rondonin could be a specific antifungal activity against yeasts.




Fig. 2. Purification of rondonin from spider plasma by reverse phase HPLC. An acidic extract obtained from *A. rondoniae* plasma was submitted to solid phase extraction on Sep-Pak C18 cartridges. The fraction that eluted at 40% acetonitrile was analysed on a semi-preparative Jupiter C₁₈ column with a linear gradient from 2% to 60% acetonitrile in acidified water over 60 min (dotted line) at a flow rate of 1.5 mL/min.




Fig. 3. MALDI-TOF spectrum of fraction 2 (rondonin). Analysis of fraction 2 by mass spectrometry showed a single molecule with an m/z of 1237.776.

3.4. MALDI-TOF and LC/MS analysis

Analysis by mass spectrometry MALDI-TOF revealed a single molecule with a mass of 1236.776 Da (Fig. 3). Following the methodology of Budnik et al. [4], “de novo” sequencing (Fig. 4) of this molecule revealed a sequence of 10 amino acids, III-QYEGHKH (Fig. 5), that showed identity to the C-terminus fragment of the subunit “d” of haemocyanin from the tarantula *Eurypelma californicum* (Theraphosidae) [42]. Furthermore, when compared to a database of partial genomes, rondonin showed identity with the C-terminus fragment of subunit “d” and 90% similarity to a fragment (IIIQYEGHKH) of subunit “f” of haemocyanin from the spider *A. gomesiana* (<http://www.compsysbio.org/partigene/>). Therefore, in the present study, we report the first isolation and characterisation of a fragment of haemocyanin with antifungal activity from arachnids.

3.5. Microbicidal activity of rondonin

The microbicidal properties of rondonin were determined by the Neubauer chamber and plate count method. When synthetic rondonin was incubated with *C. albicans* at a concentration two times higher than the respective MIC, no living yeast were detected

a	MGYPFDRKITADTHEEFLTGNMNIHVTVRFQD	631
b	MGFPFDRDIKADSIPEWLHPNMHFSEVTITHHQ	627
c	MGFPFDRPIPEGHASNLHQPNVSFSQIKIQHH	629
d	MGYPFDRPIQVVRTPSQFKTPNMAFQELIIQYEGHKH	627
e	MGFPFDRVIEGLTLEEFLLTPMSCTDVRIKYTDIK	624
f	MGFPFDRVIKARTVADFRTTNMSFTDVKIQFKDQV	629
g	MGFPFDRPTKIHTAEILTPNMSLTDVVIOYVGHE	628

Fig. 5. *Eurypelma californicum* (Theraphosidae) haemocyanin subunits (a–g). The peptide showed identity with the C-terminal fragment from subunit “d” [42].




Fig. 4. Representative “de novo” sequencing of fraction 2 from *A. rondoniae* plasma in a Q-TOF Micro™ (Micromass). The fragments shown correspond to the b and y series. The peptide sequence following the b and y series orientation is shown on the top of the graph. Mass spectrometric “de novo” peptide sequencing was performed in positive ionisation mode on a Q-TOF Micro™ fitted with an electrospray ion source.




Fig. 6. Growth inhibition of *Candida albicans* MDM8 by rondonin. Synthetic rondonin (67 μM , solid line) or water (control, dotted line) was added to an exponential phase culture of *C. albicans*. Aliquots were removed at various times, and the number of blastoconidia were counted in a Neubauer chamber. To test the viability of blastoconidia, 20 μL aliquots for each time point were incubated on Luria Bertani agar plates for 48 h at 30 $^{\circ}\text{C}$.




Fig. 7. Toxicity of rondonin with HRBCs. Synthetic rondonin was incubated with human erythrocytes ranging from 0.2 to 134 μM for 3 h at 37 $^{\circ}\text{C}$. The haemolysis percentage was expressed in relation to 0% haemolysis with PBS and 100% lysis control (0.1% triton X-100).

after 10 min (data not shown). Fig. 6 shows that rondonin inhibits the growth of *Candida albicans* MDM8 compared to the control.

3.6. Toxicity to human red blood cells

Haemolytic assays were used to assess the toxicity of peptides towards HRBCs *in vitro*. Incubating HRBCs (w/v) with various concentrations of rondonin for 3 h at 37 $^{\circ}\text{C}$ did not affect the OD at 414 nm. Triton X-100 was used as a positive control and taken as the 100% haemolysis value. PBS was used as a negative control and taken as the 0% haemolysis value. These results demonstrate that rondonin is not haemolytic (Fig. 7).

4. Discussion

Considering the studies of the immune system of invertebrates, the knowledge of the mechanisms involved in the immune response of arachnids is less studied than *Limulus polyphemus* [18]. The life expectancy of many invertebrates is as long as the lifespan of vertebrates, despite the continuous challenge of pathogens. All multicellular animals are subject to frequent microbial challenges and the attack of endo- and ectoparasites. In addition to defences against predators, survival depends on the presence of an efficient immune system that can quickly remove or inactivate pathogenic organisms. The immune system of the tarantula spider is particularly interesting because, in addition to

having a life expectancy of more than 20 years [9], tarantula spiders are also phylogenetically very old with fossil records dating from the Devonian period (400 million years ago) [32,31].

In this work, we purified six molecules with antimicrobial activity that have never been before described from the plasma of the tarantula spider *A. rondoniae*. Only one molecule, rondonin, which was named in honour of the studied species, was isolated and characterised. Rondonin, a peptide characterised with antifungal activity that inhibits the growth of *C. albicans* MDM8 in ten minutes and exhibited fungicidal activity, like a study previous with some antifungal agents like amphotericin B, fluconazole and LY303366 showed that fluconazole and LY303366 are fungistatic and amphotericin B exhibited fungicidal activity with a reduction in $\geq 3 \log_{10}$ compared to the starting inoculum [21]. Furthermore, rondonin showed no haemolytic activity, had identity with the C-terminal fragment of subunit “d” of haemocyanin from the tarantula *Eurypelma californicum* and *A. gomesiana* and showed 90% similarity with a fragment of subunit “f” of *A. gomesiana*, differing in only the second amino acid: ILIQYEGHKH. Lee et al. [24] also found a fragment of haemocyanin, named astacidin 1, in the plasma of the crayfish *Pacifastacus leniusculus*, which consists of 16 amino acids with the sequence FKVQNQHGVVVKIFHH-COOH, has a molecular mass of 1945.2 Da, and possesses antimicrobial activity.

Destoumieux-Garzon et al. [6] showed that the plasma of shrimp contains an original class of strictly antifungal (poly)peptides with molecular masses ranging from 2753.2 Da for a peptide from *Penaeus vannamei*, which consists of 23 amino acids with the sequence FEDLPNFGHIQVKVFNHGEIH, 7982.4 Da for a peptide of 31 amino acids from *P. stylirostris* (VTDGDADSAPNLHENTYNYHSGHGVYPDK) and 8362.8 Da for a 32 amino acid peptide, also from *P. stylirostris* (LVVAVTDGDADSAPNLHENTYNYHSGHGVY). The two peptides from *P. stylirostris* revealed perfect homology with the C-terminus of the haemocyanin of *Penaeus*.

In this case, the authors speculated that the penaeid shrimp can use haemocyanin, which is abundant and readily available in the plasma, to produce C-terminal fragments that possess broad antifungal activities within the first hours of an infection. Therefore, haemocyanin has a potential function in crustacean immunity by serving as a substrate for the generation of antifungal (poly)peptides that could contribute to microorganism elimination in plasma. It remains to be established whether the mechanism leading to the partial cleavage of haemocyanin is part of the shrimp immune reaction and how involved this process can be in an immediate and systemic antimicrobial response in shrimp.

This result suggests that, as observed in crustaceans, the cleavage of haemocyanin and the production of peptide fragments with antimicrobial activity also occur in spiders as a first line of defence against infection. Several studies suggest that haemocyanins are involved in the arthropod immune system. The activity of the haemocyanin fragment discovered in this study reinforces that idea.

The identification and characterisation of new substances can lead to the development of new drugs that kill resistant pathogenic microorganisms. This peptide has activity against clinical isolates that cause candidiasis, one of the opportunistic pathogens responsible for nosocomial infections that colonise human mucosal surfaces [30]. Yeasts of the genus *Candida* are significant due to the high frequency at which they colonise and infect a human host. *Candida* species are found in the gastrointestinal tract in 20–80% of the healthy adult population. In addition, approximately 20–30% of women have *Candida* colonies in their vaginas [7]. The increase in the prevalence of yeast infections is likely due to the AIDS epidemic, cancer chemotherapy, organ and bone-marrow transplants and invasive hospital procedures [43,45] because of the overuse of antifungal agents, such as fluconazole

[45]. Due to its small size, rondonin can be synthesised quickly and can kill yeast in ten minutes. Furthermore, no toxicity towards human erythrocytes was observed in this study. Therefore, rondonin may represent a new strategy for developing drugs that neutralise or inhibit pathogens.

Acknowledgements

We are grateful to Dr. Mirian A.F. Hayashi (Dept. Farmacology/ UNIFESP) for providing the clinical strains. We also appreciate the financial support of Fapesp (CAT/Cepid Project 98-14307-9), Capes and Cnpq.

References

- [1] Atherton, Sheppard. Solid phase peptide synthesis—a practical approach. Oxford: IRL Press; 1989.
- [2] Boman HG. Antibacterial peptides: key components needed in immunity. *Cell* 1991;65:205–7.
- [3] Boman HG. Peptide antibiotics and their role in innate immunity. *Annual Review of Immunology* 1995;13:61–92.
- [4] Budnik BA, Olsen JV, Egorov TA, Anisimova VE, Galkina TG, Musolyamov AK, Grishin EV, Zubarev RA. De novo sequencing of antimicrobial peptides isolated from the venom glands of the wolf spider *Lycosa singoriensis*. *Journal of Mass Spectrometry* 2004;39:193–201.
- [5] Cociancich S, Goyffon M, Bontems F, Bulet P, Bouet F, Menez A, Hoffmann JA. Purification and characterization of a scorpion defensin, a 4 kDa antibacterial peptide presenting structural similarities with insect defensins and scorpion toxins. *Biochemical and Biophysical Research Communications* 1993;194:17–22.
- [6] Destoumieux-Garzon D, Saulnier D, Garnier J, Jouffrey C, Bulet P, Bache're E. Crustacean immunity: antifungal peptides are generated from the C-terminus of shrimp hemocyanin in response to microbial challenge. *The Journal of Biological Chemistry* 2001;276:47070–7.
- [7] Dignani MC, Solomkin JS, Anaissie E. Candida. In: Anaissie E, McGinnis MR, Pfaller MA, editors. *Medical mycology*. 1ª Edição. Filadélfia: Churchill Livingstone; 2003. p. 195–239.
- [8] Ehret-Sabatier L, Loew D, Goyffon M, Fehlbaum P, Hoffmann JA, Van Dorsseleer A, Bulet P. Characterization of novel cysteine-rich antimicrobial peptides from scorpion blood. *The Journal of Biological Chemistry* 1996;271:29537–29544.
- [9] Foelix RF. *Biology of spiders*. Cambridge, Massachusetts/London: Harvard University Press; 1996.
- [10] Fogaça AC, Silva Jr PI, Miranda MT, Bianchi AG, Miranda A, Ribolla PE, Daffre S. Antimicrobial activity of a bovine hemoglobin fragment in the tick *Boophilus mirabilis*. *The Journal of Biological Chemistry* 1999;274:25330–4.
- [11] Haerberli S, Kuhn-Nentwig L, Schaller J, Nentwig W. Characterization of antibacterial activity of peptides isolated from the venom of the spider *Cupiennius salei* (Aranae: Ctenidae). *Toxicon* 2000;38:373–80.
- [12] Hancock REW, Diamond G. The role of cationic antimicrobial peptides in innate host defense. *Trends in Microbiology* 2000;8:402–10.
- [13] Harris F, Dennison SR, Phenix DA. Anionic antimicrobial peptide from eukaryotic organisms. *Current Protein and Peptide Science* 2009;10:585–606.
- [14] Hetru C, Bulet P. Strategies for the isolation and characterization of antimicrobial peptides of invertebrates. *Methods in Molecular Biology* 1997;78(1):35–48.
- [15] Hoffman IA, Kafatos FC, Janeway CA, Ezekowitz RAB. Phylogenetic perspectives in innate immunity. *Science* 1999;284:1313–8.
- [16] Hoffman JA, Hertu C. Insect defensins: inducible antibacterial peptides. *Immunology Today* 1992;13:411–5.
- [17] Iwanaga S, Kawabata S, Muta T. New types of clotting factors and defense molecules found in horseshoe crab hemolymph: their structures and functions. *The Journal of Biochemistry (Tokyo)* 1998;123:1–15.
- [18] Jenssen H, Hamill P, Hancock REW. Peptide antimicrobial agents. *Clinical Microbiology Reviews* 2006;19(3):491–511.
- [19] Kamysz W, Okrój M, Lukasiak J. Novel properties of antimicrobial peptides. *Acta Biochimica Polonica* 2003;50:461–9.
- [20] Klepser ME, Ernest EJ, Lewis RS, Ernest ME, Pfaller MA. Influence of test conditions on antifungal time-kill curve results: proposal for standardized methods. *Antimicrobial Agents and Chemotherapy* 1998;42(5):1207–12.
- [21] Kuhn-Nentwig L. Antimicrobial and cytotoxic peptides of venomous arthropods. *Cellular and Molecular Life Sciences* 2003;60:2651–68.
- [22] Lai R, Lomas LO, Jonczy J, Turner PC, Rees HH. Two novel noncationic defensin-like antimicrobial peptides from haemolymph of the female tick, *Amblyomma hebraeum*. *The Journal of Biochemistry* 2004;379:681–5.
- [23] Lee SY, Lee BL, Söderhäll K. Processing of an antibacterial peptide from hemocyanin of the freshwater crayfish *Pacifastacus leniusculus*. *The Journal of Biological Chemistry* 2002;278(10):7927–33.
- [24] Mak P, Wojcik K, Wicherek Ł, Suder P, Dubin A. Antibacterial hemoglobin peptides in human menstrual blood. *Peptides* 2004;25:1839–47.
- [25] Markl J, Markl A, Schartau W, Linzen B. Subunit heterogeneity in arthropod hemocyanins: I (Chelicerata). *Journal of Comparative Physiology B* 1979;130:283–292.
- [26] Markl J, Savel A, Decker H, Linzen B. Hemocyanins in spiders, IX. Homogeneity, subunit composition and the basic oligomeric structure of *Eurypelma californicum* hemocyanin. *Hoppe-Seyler's Zeitschrift für Physiologische Chemie* 1980;36:649–60.
- [27] Markl J, Decker H, Stöcker W, Savel A, Linzen B, Schutter WG, van Bruggen EFJ. On the role of dimeric subunits in the quaternary structure of arthropod hemocyanins. *Hoppe-Seyler's Zeitschrift für Physiologische Chemie* 1981;362:185–188.
- [28] Markl J. Evolution and function of structurally diverse subunits in the respiratory protein hemocyanin from arthropods. *The Biological Bulletin* 1986;171:90–115.
- [29] Odds FC, Gow NA, Brown AJ. Fungal virulence studies come of age. *Genome Biology* 2001;2 [p. 1009.1–4].
- [30] Selden P, Shea WA, Bonamo PM. A spider and other arachnids from Devonian of New York, and reinterpretations of Devonian Aeneae. *Paleontology* 1991;34:241.
- [31] Shear WA, Palmer JM, Coddington JA, Bonamo PM. A Devonian spinneret: early evidences of spiders and silk use. *Science* 1989;246:479.
- [32] Schneider H-J, Markl J, Schartau W, Linzen B. Subunit heterogeneity of *Eurypelma* (Dugesella) hemocyanin, and separation of polypeptide chains. *Hoppe-Seyler's Zeitschrift für Physiologische Chemie* 1977;358:1133–41.
- [33] Silva Jr PI. Sistema Imune de Aracnídeos: Estrutura química e atividade de peptídeos antimicrobianos da Hemolinfa de *Acanthoscurria gomesiana*. Tese apresentada ao Instituto de Ciências Biomédicas da USP para obtenção do título de doutor em Ciências; 2000.
- [34] Silva Jr PI, Daffre S, Bulet P. Isolation and characterization of gomesin, an 18-residue cysteine-rich defense peptide from the spider *Acanthoscurria gomesiana*. Hemocytes with sequence similarities to horseshoe crab antimicrobial peptides of the Tachyplesin family. *The Journal of Biological Chemistry* 2000;275(45):33464–70.
- [35] Silva FD, Rezende CA, Rossi DCP, Esteves E, Dyszy FH, Schreier S, Gueiros-Filho F, Campos CB, Pires JR, Daffre S. Structure and mode of action of microplusin, a copper II-chelating antimicrobial peptide from the cattle tick *Rhipicephalus (Boophilus) microplus*. *The Journal of Biological Chemistry* 2009;284:34735–46.
- [36] Söderhäll K, Cerenius L. Role of the prophenoloxidase-activating system in invertebrate immunity. *Current Opinion in Immunology* 1998;10:23–8.
- [37] Söderhäll K, Smith VJ. Separation of the haemocyte of *Carcinus maenas* and other decapod crustaceans and phenoloxidase distribution. *Developmental and Comparative Immunology* 1983;7:229–39.
- [38] Steffen H, Rieg S, Wiedemann I, Kalbacher H, Deeg M, Sahl HG, Peschel A, Götz F, Garbe C, Schitteck B. Naturally processed dermcidin-derived peptides do not permeabilize bacterial membranes and kill microorganisms irrespective of their charge. *Antimicrobial Agents and Chemotherapy* 2006;50:2608–20.
- [39] Strominger. Animal antimicrobial peptides: ancient players in innate immunity. *The Journal of Immunology* 2009;182:6633–4.
- [40] Ullal AJ, Litaker RW, Noga EJ. Antimicrobial peptides derived from haemoglobin are expressed in epithelium of channel catfish (*Ictalurus punctatus*, Rafinesque). *Developmental and Comparative Immunology* 2008;32:1301–12.
- [41] Voit R, Feldmaier-Fuchs G, Schweikardt T, Decker H, Burmester T. Complete sequence of the 24-mer hemocyanin of the tarantula *Eurypelma californicum*. *The Journal of Biological Chemistry* 2000;275(50):39339–44.
- [42] White TC, Man KA, Bowden RA. Clinical, cellular, and molecular factors that contribute to antifungal drug resistance. *Clinical Microbiology Reviews* 1998;11:382–402.
- [43] Yamane ES. Estudo da atividade antimicrobiana da crotamina nativa e sintética, do seu análogo recombinante e fragmentos peptídicos. Dissertação apresentada a Universidade Federal de São Paulo—Escola Paulista de Medicina para obtenção do título de mestre em Ciências; 2010.
- [44] Yang YL, Lo H-J. Mechanisms of antifungal agent resistance. *Journal of Microbiology, Immunology and Infection* 2001;34:79–86.
- [45] Yigit N, Benli M. The antibacterial activity of hemolymph of spider, *Agelena labyrinthica* (Aranae: Agelenidae). *Journal of Forestry Faculty* 2008;8:120–4.
- [46] Zhu WL, Song YM, Park Y, Park KH, Yang ST, Kim JI, et al. Substitution of the leucine zipper sequence in melittin with peptoid residues affects self-association, cell selectivity, and mode of action. *Biochimica et Biophysica Acta* 2007;1768(6):1506–17.