

Report

Salicylic Acid Inhibits Pathogen Growth in Plants through Repression of the Auxin Signaling Pathway

Dong Wang,^{1,3,4} Karolina Pajeroska-Mukhtar,^{1,4} Angela Hendrickson Culler,² and Xinnian Dong^{1,*}

¹Department of Biology
Duke University

Durham, North Carolina 27708

²Department of Horticultural Science and Microbial
and Plant Genomics Institute

University of Minnesota

Saint Paul, Minnesota 55108

Summary

The phytohormone auxin regulates almost every aspect of plant development. At the molecular level, auxin induces gene expression through direct physical interaction with the TIR1-like F box proteins, which in turn remove the Aux/IAA family of transcriptional repressors [1–4]. A growing body of evidence indicates that many plant pathogens can either produce auxin themselves or manipulate host auxin biosynthesis to interfere with the host's normal developmental processes [5–11]. In response, plants probably evolved mechanisms to repress auxin signaling during infection as a defense strategy. Plants overaccumulating the defense signal molecule salicylic acid (SA) frequently display morphological phenotypes that are reminiscent of auxin-deficient or auxin-insensitive mutants, indicating that SA might interfere with auxin responses. By using the Affymetrix ATH1 Gene-Chip for *Arabidopsis thaliana*, we performed a comprehensive study of the effects of SA on auxin signaling [12]. We found that SA causes global repression of auxin-related genes, including the *TIR1* receptor gene, resulting in stabilization of the Aux/IAA repressor proteins and inhibition of auxin responses. We demonstrate that this inhibitory effect on auxin signaling is a part of the SA-mediated disease-resistance mechanism.

Results and Discussion

To uncover the molecular basis of SA-mediated plant immunity, we began by analyzing transcriptome changes in the model plant *Arabidopsis thaliana* in response to an SA analog, benzothiadiazole S-methylester (BTH). A detailed description of this analysis has been previously published [12]. Interestingly, among the 1133 BTH-repressed genes, a significant number (21 genes) was related to auxin signal transduction (p value $< 2.12 \times 10^{-3}$, Table 1). These repressed genes include *AUX1* and *PIN7*, encoding an auxin importer and exporter, respectively

[13, 14]; genes for the auxin receptors TIR1 and AFB1; and the well-characterized auxin-inducible *SAUR* genes and *Aux/IAA*-family genes [15]. Additionally, among genes upregulated by BTH, two of them encode auxin-conjugating enzymes [16], implying that BTH might also affect auxin homeostasis by lowering the levels of free auxin. Together, these data suggest that BTH treatment results in an overall reduction in auxin responses, which might attenuate the effectiveness of pathogen-derived auxin and contribute to disease resistance. In a different microarray experiment, we found that 18 of these 21 auxin-related genes were also repressed in systemic tissue after induction of systemic acquired resistance (SAR) [17]; thus, downregulation of auxin-related genes might be a part of the SAR response. In the microarray for NPR1 direct transcriptional targets [18], many of these genes had low signal levels: Eight were called “absent.” Of the rest, only one gene was repressed more than 2-fold in both replicates, suggesting that these auxin-pathway genes are likely to be indirectly regulated by NPR1.

To investigate whether SA treatment affects auxin levels, we measured free auxin (indole-3-acetic acid; IAA) in SA-treated plants under conditions similar to those of the microarray experiment [12]. We found no significant changes in free-auxin levels 24 and 48 hr after SA treatment (Figure S1A in the Supplemental Data available online). This suggests that induction of the auxin-conjugating enzymes has no immediate effect on free-auxin levels and that SA and BTH repress auxin-mediated genes predominantly at the signaling level.

To validate this prediction, we tested the effect of SA on the inducibility of the auxin-signaling reporter *DR5:: β -glucuronidase (GUS)* [19] by using increasing concentrations of SA. As shown in Figure 1A, SA significantly inhibited auxin-mediated expression of this reporter at 0.5 mM, a concentration that is normally used for inducing *Pathogenesis-related (PR)* gene expression and disease resistance [20]. This result was also observed with *in situ* staining for reporter activity in roots, in which the *DR5* promoter is most active. As shown in Figure 1B, exogenous treatment of seedlings with 1 μ M synthetic auxin, 1-naphthalacetic acid (NAA), stimulated strong GUS activity, indicated by a bright-blue color in the root. Addition of 0.5 mM SA abolished this effect. Similarly, SA also blocked *DR5::GUS* induction by both the natural auxin IAA and another synthetic auxin, 2,4-dichlorophenoxyacetic acid (2,4-D) (data not shown). This effect was not observed with an inactive SA analog 4-hydroxybenzoic acid (4-HBA) or another plant hormone jasmonic acid (Figures S2A and S2B), confirming its SA specificity.

Several SA overaccumulating mutants, such as *cpr5*, *cpr6*, and *snc1*, exhibit morphologies, namely reduced apical dominance and stunted growth, that are reminiscent of auxin deficiency [21–23]. We first measured free-auxin levels in *cpr6* and *snc1* and found that they

*Correspondence: xdong@duke.edu

³Present address: Department of Biological Sciences, Stanford University, Stanford, California 94305.

⁴These authors contributed equally to this work.

Table 1. Expression of Annotated Auxin-Related Genes Modulated by BTH

Array Element	Locus Identifier	Annotation	8 Hr F.C.	24 Hr F.C.	p Value
264014_at	At2g21210	Auxin-responsive protein	-4.78	-11.47	1.81E-03
254746_at	At4g12980	Auxin-responsive protein, putative	-4.29	-4	2.18E-03
252965_at	At4g38860	Auxin responsive SAUR protein	-5.81	-7.52	2.23E-03
259773_at	At1g29500	Auxin responsive SAUR protein	-4.61	-5.78	2.27E-03
259787_at	At1g29460	Auxin responsive SAUR protein	-2.25	-2	2.39E-03
265319_at	At2g22670	Indoleacetic acid-induced protein 8 (IAA8)	-2.62	-3.53	2.55E-03
259783_at	At1g29510	Auxin responsive SAUR protein	-4.46	-5.85	2.80E-03
264900_at	At1g23080	Auxin efflux carrier PIN7	-2.67	-2.96	2.89E-03
267092_at	At2g38120	Amino acid permease (AUX1)	-2.6	-2.82	2.89E-03
252972_at	At4g38840	Auxin-inducible SAUR gene	-3.97	-4.12	3.02E-03
251199_at	At3g62980	Auxin receptor TIR1	-1.7	-2.04	3.23E-03
265454_at	At2g46530	Transcriptional factor B3 family protein / auxin-responsive factor AUX/IAA-related	-2.75	-2.92	3.47E-03
259790_s_at	At1g29430, At5g27780	Auxin responsive SAUR protein	-2.27	-1.66	3.83E-03
267310_at	At2g34680	Leucine-rich repeat protein	-2.16	-2.39	6.19E-03
266215_at	At2g06850	Endo-xyloglucan transferase (EXT) (EXGT-A1)	-10.66	-13.59	8.26E-03
245076_at	At2g23170	IAA-amido synthase that conjugates amino acids to auxin	1.46	2.11	8.65E-03
255417_at	At4g03190	F box family protein (FBL18), AFB1	-2.02	-2.03	1.14E-02
248801_at	At5g47370	Homeobox-leucine zipper protein 2 (HAT2)	-1.5	-2.07	1.32E-02
252970_at	At4g38850	Auxin-responsive protein/small auxin up RNA (SAUR-AC1)	-1.82	-2.16	2.14E-02
250012_x_at	At5g18060	Auxin-responsive protein, putative	-1.87	-3.32	2.16E-02
253908_at	At4g27260	IAA-amido synthase that conjugates amino acids to auxin	2.41	1.87	4.00E-02

Fold changes (F.C.) and BTH-dependency p values were calculated on the basis of gene-expression levels at 0, 8, and 24 hr after induction.

were significantly lower than those in the wild-type ($p \leq 0.05$; Figure 2A). This indicates that although exogenous treatment of plants with SA had little immediate effect on free-auxin levels, the presence of high endogenous SA did have a long-term effect on auxin homeostasis. To test whether these SA-accumulating mutants display altered responses to auxin, we introduced the *DR5::GUS* reporter into these mutants through genetic crosses. We then quantified the auxin sensitivity of these plants by measuring the activity of the *DR5::GUS* reporter. As shown in Figure 2B, induction of the reporter by NAA was clearly diminished in these mutants, indicating that SA accumulation causes insensitivity to exogenously applied auxin.

To rule out the possibility that SA inhibits the uptake of auxin, we crossed SA-accumulating mutants with an auxin-overproducing mutant, *yucca*. As a result of high endogenous auxin levels, *yucca* plants display long hypocotyls and epinastic cotyledons at the seedling stage, narrow rosette leaves with long petioles during the adult stage, and increased apical dominance after bolting [24]. Interestingly, SA-accumulating mutants suppressed most of the phenotypes associated with *yucca*. For example, the *snc1 yucca* double mutant had small rosette with curly leaves, typical of the *snc1* single mutant (Figure 2C and Figure S3), and the *cpr6 yucca* double mutant exhibited reduced apical dominance as in *cpr6* (Figure 2D). Only at the seedling stage were auxin phenotypes still evident, presumably

because SA had not accumulated to high levels early in development. Because the *yucca* mutant was generated through activation tagging, the *YUCCA* gene is no longer regulated by its endogenous promoter or sensitive to SA repression. Not surprisingly, free-IAA levels in the *yucca snc1* and *yucca cpr6* double mutants were not significantly different from that found in the *yucca* single mutant (Figure S1B), indicating that suppression of *yucca* phenotypes by *snc1* and *cpr6* is mainly due to repressed auxin response, not reduced auxin synthesis.

As suggested by our expression-profiling experiment, one mechanism by which SA might inhibit auxin signaling is through transcriptional repression of the auxin receptor genes (Table 1). Reduction in TIR1 and AFB1 levels would lead to reduced degradation of Aux/IAA repressors and thus inhibit auxin responses. To test this hypothesis, we used the reporter *HS::AXR3NT-GUS*, which encodes a fusion between the coding sequences of amino terminus (NT) of one auxin-response repressor, *AXR3*, and GUS-encoding *uidA* gene driven by a heat-shock-inducible promoter (*HS*). After heat shock, the level of the AXR3NT-GUS fusion protein can be measured using MUG assay [4]. To distinguish transcriptional induction from increased protein stability, we also measured the transcript levels of GUS by using semiquantitative RT-PCR. As shown in Figure 3A, transcription of the reporter was induced after heat shock, and this induction was further enhanced in the presence




Figure 1. SA Suppresses Auxin Response

(A) Seven-day-old *DR5::GUS* seedlings were incubated overnight with 1 μ M NAA and increasing concentrations of SA, as indicated. GUS activity was measured as previously described [20]. Error bars represent SDs.

(B) GUS staining in representative root segments after treatment with 1 μ M NAA with or without 0.5 mM SA. This experiment was repeated multiple times, with more than ten seedlings per treatment, with similar results.

of SA. GUS activities correlated with this increase in transcription. However, 40 min after the heat shock, transcript abundance dropped to background levels, and GUS activity could then be used as a measure of protein stability. For the sample treated with 1 μ M NAA alone, GUS activity started to decline. In contrast, in the sample treated with both NAA and SA (0.5 mM), GUS activity continued to increase, suggesting that SA has a stabilizing effect on AXR3NT-GUS. To confirm this result, we directly examined endogenous Aux/IAA repressor protein levels after SA treatment. Aux/IAA proteins are normally short lived and difficult to detect on a western blot [4]. However, SA treatment resulted in a drastic increase in the amount of Aux/IAA proteins detected with an antibody against AXR2/IAA7 (Figure 3B). Examination of AXR2 transcript levels showed that this gene was not affected by SAR induction [12, 18]. As expected, 4-HBA did not delay the degradation of AXR3NT-GUS (Figure S2C). From these results, we conclude that SA significantly increases Aux/IAA repressor protein levels, and such an increase leads to inhibition of auxin responses.

Stabilization of the auxin repressor proteins by SA might be due to transcriptional repression of the auxin receptor genes (Table 1) or posttranscriptional mechanisms. Navarro et al. reported a microRNA-mediated pathway that causes a rapid reduction in auxin receptor levels in response to the defense elicitor flg22 [25]. We found that the auxin induction of *DR5::GUS* was unaffected after a short SA treatment of 2 hr (data not

shown). The level of miR393, the microRNA involved in flg22-mediated response, was not altered 24 hr after SA treatment (Figure S4). Apparently, the mechanism by which SA downregulates auxin signaling is independent of miR393.

Besides repressing the transcription of auxin receptor genes, SA might interfere with auxin signaling in a more direct manner, namely by disrupting the interaction between Aux/IAA proteins and auxin receptors. To test this possibility, we examined the effect of SA on the ability of GST-IAA1 to pull down Myc-tagged TIR1 in vitro. *TIR1-myc* expression is driven by a dexamethasone (Dex)-inducible promoter, and this induction is not altered by SA (data not shown). As previously reported, GST-IAA1 could pull down TIR1 in an auxin-dependent manner [1, 4]. However, we found that this interaction was unaffected by SA, whether it was added at the point of TIR1 induction or only in the pulldown reaction (Figure 3C). These in vitro data strongly argue that SA stabilizes Aux/IAA proteins mainly by limiting auxin receptors that are required for Aux/IAA protein degradation. However, we cannot completely exclude the possibility that SA can modify Aux/IAA proteins in vivo to make them more resistant to degradation.

A growing body of evidence indicates that pathogen infection perturbs auxin homeostasis [5–11, 25]. During an infection experiment with the virulent oomycete pathogen *Hyaloperonospora parasitica* Noco2, we noticed that the auxin-responsive reporter *DR5::GUS* was induced. Intriguingly, the induction pattern of *DR5::GUS* coincided with the growth of conidiophores, whereas leaves free of pathogen growth only displayed background levels of staining (Figure 4A). Increases in free auxin and auxin sensitivity were also observed in plants expressing the *Pseudomonas syringae* type III effector AvrRpt2 and in response to *P. syringae* pv. *tomato* DC3000 (*Pst*DC3000) infection (Z. Chen and B. Kunkel, personal communication). Exogenous auxin application has also been shown to promote the development of *Pst*DC3000 disease symptoms ([25], Z. Chen and B. Kunkel, personal communication). We found that coinoculation of *P. syringae* pv. *maculicola* ES4326 (*Psm*4326) with 50 μ M of NAA not only enhanced disease symptoms but also moderately promoted pathogen growth (Figure 4B). Although the exact role of auxin in pathogenesis is unknown, we observed that auxin application could inhibit the full induction of SA-mediated *PR-1* expression (Figure 4C). Our results lend support to the hypothesis that auxin might downregulate host defense responses [26].

On the basis of the strong inhibitory effect of SA on auxin responses, as well as the likely role of auxin in promoting pathogenesis, it is logical to hypothesize that repressing the auxin pathway is an important aspect of the defense response. Therefore, we tested the disease-resistance phenotype of *axr2-1* plants, which are insensitive to auxin because of production of a nondegradable form of the AXR2/IAA7 repressor protein [27]. When infected with the bacterial pathogen *Psm*4326 at a high inoculum ($OD_{600} = 0.001$), wild-type plants were unable to prevent the proliferation of the pathogen. In contrast, the *axr2-1* mutant displayed a 10-fold reduction in bacterial growth (Figure 4D). This result clearly




Figure 2. SA-Accumulating Mutants Exhibit Lower Levels of Free IAA and Reduced Sensitivity to Auxin
(A) Free-IAA levels were measured in 4-week-old wild-type, *cpr6*, and *snc1* plants. Error bars represent SDs for three independent samples.
(B) The *DR5::GUS* reporter was introduced into SA-accumulating mutants, and the responsiveness of the reporter to exogenous auxin was determined for seven-day-old seedlings in a MUG assay. Error bars represent the SDs for three independently collected samples.
(C) Among the progeny from a *snc1-yucca* cross, seedlings with *yucca*-associated long cotyledons were transplanted. Plants homozygous for *snc1* started to suppress *yucca* phenotypes as the plants aged. Shown here are rosettes of six-week-old plants. Inflorescences were removed for a clearer view of the rosette.
(D) Flowering *cpr6 yucca* plants showed reduced apical dominance typical of the *cpr6* single mutant. Note that the short-silique phenotype of *yucca* was also suppressed.

shows that blocking auxin responses can lead to heightened disease resistance.

The importance of repressing auxin signaling to disease resistance might be more evident when other forms of SA-mediated defenses are absent. In *NahG*

transgenic plants, a bacterial salicylate hydroxylase, encoded by the *NahG* gene, degrades SA. As a result, these plants cannot defend against *Psm4326* even when challenged with a low bacterial inoculum ($OD_{600} = 0.0001$, Figure 4E). If inhibition of auxin responses is




Figure 3. SA Stabilizes Auxin-Response Repressors

(A) Seven-day-old seedlings containing *HS::AXR3NT-GUS* were incubated in either water or 0.5 mM SA for 24 hr. GUS activity was then examined before heat shock, immediately after a 2 hr heat shock, or after seedlings had been incubated for 40 min after heat shock in the presence of 1 μ M NAA. Error bars represent the SDs for three independent samples. *AXR3NT-GUS* transcript levels were measured for each sample with RT-PCR. β -Tubulin subunit 4 (*β 4-TUB*) transcript levels were used as loading controls.
(B) The level of the AXR2 protein after 24 hr water or SA treatment was examined with an antibody raised against the endogenous AXR2 protein. Because of constitutive degradation, the protein in water-treated samples usually cannot be detected. α -tubulin was probed as a loading control.
(C) Inducible *TIR1-myc* transgenic seedlings were treated with Dex overnight to turn on TIR1-Myc synthesis. Total protein was extracted and TIR1-Myc was pulled down with purified GST-IAA1 in the absence or the presence of 50 μ M IAA. SA was added either during Dex induction (lanes 3 and 4) or during the pulldown assay (lanes 7 and 8).

(B) The level of the AXR2 protein after 24 hr water or SA treatment was examined with an antibody raised against the endogenous AXR2 protein. Because of constitutive degradation, the protein in water-treated samples usually cannot be detected. α -tubulin was probed as a loading control.
(C) Inducible *TIR1-myc* transgenic seedlings were treated with Dex overnight to turn on TIR1-Myc synthesis. Total protein was extracted and TIR1-Myc was pulled down with purified GST-IAA1 in the absence or the presence of 50 μ M IAA. SA was added either during Dex induction (lanes 3 and 4) or during the pulldown assay (lanes 7 and 8).

(C) Inducible *TIR1-myc* transgenic seedlings were treated with Dex overnight to turn on TIR1-Myc synthesis. Total protein was extracted and TIR1-Myc was pulled down with purified GST-IAA1 in the absence or the presence of 50 μ M IAA. SA was added either during Dex induction (lanes 3 and 4) or during the pulldown assay (lanes 7 and 8).




Figure 4. Crosstalk between Auxin and SA Affects Plant Defense Responses

(A) Plants carrying the *DR5::GUS* reporter were either mock treated or infected with *Hyaloperonospora parasitica* Noco2 (1×10^4 spores/ml) and stained for GUS activity 7 days postinoculation (dpi). We photographed the same infected plant with different background lighting to reveal conidiophores. Plants shown here are representative of multiple samples.

(B) Four-week-old plants were inoculated with *Psm4326* (OD600 = 0.0005), 50 μ M NAA, or *Psm4326* together with 50 μ M NAA. Disease symptoms were photographed (left panel), and bacterial growth was measured (right panel) 3 dpi. Arrows indicate infiltrated leaves. Eight samples were taken for each treatment. The experiment has been repeated with similar results. Bacterial growth in NAA-treated plants is significantly different from control plants ($p = 0.0048$, Student's *t* test). Error bars represent 95% confidence intervals. "cfu" stands for colony-forming unit.

(C) Four-week-old plants were treated with water, SA (0.5 mM), or SA plus NAA (50 μ M) for 24 hr. SA response was examined by probing *PR-1* transcripts on a Northern blot. Ribosomal RNA (*rRNA*) is shown as a loading control.

(D) Four-week-old plants were infiltrated with a high dose of *Psm4326* (OD600 = 0.001). Bacterial growth was quantified 3 dpi. The absolute values of bacterial growth in *axr2-1* are significantly different from those of the wild-type ($p = 0.0027$, Student's *t* test). Error bars represent 95% confidence intervals.

(E) The assay was performed in the same manner as for (D), except with a lower dose of pathogen (OD600 = 0.0001). Error bars represent 95% confidence intervals.

a component of SA-mediated resistance, then blocking auxin signaling should restore some resistance in *NahG* plants. Indeed, as shown in Figure 4E, the auxin-insensitive *axr2-1* mutation drastically reduced *Psm4326* growth in *NahG* plants. SA is well known for activating multiple types of defense responses (See model in Figure 4F). It is therefore not surprising that *axr2-1* did not fully restore resistance in *NahG* plants. Our results are in agreement with those of a complementary study, in which overaccumulation of an auxin receptor protein compromised resistance [25]. Together, our results establish that inhibiting auxin sensitivity is an integral component of the SA-mediated defense response.

In summary, by using expression profiling, we discovered that SA inhibits the auxin signaling pathway as part of the plant defense mechanism. This inhibition is accomplished through SA-mediated stabilization of the auxin-response repressors. In addition, auxin might not be the only plant growth hormone targeted by SA. In the BTH-treatment microarray experiment [12], several genes known to be involved in the gibberellin pathway were also found to be downregulated, indicating the presence of an elaborate interaction network between developmental and defense pathways in plants.

Supplemental Data

Experimental Procedures and four figures are available at <http://www.current-biology.com/cgi/content/full/17/20/1784/DC1/>.

Acknowledgments

We thank Dr. Mark Estelle for the *HS::AXR3NT-GUS* transgenic line and the AXR2 antibody, Dr. Thomas Guilfoyle for the *DR5::GUS* transgenic line, Dr. Jerry Cohen and Dr. Jennifer Normanly for help in auxin measurement, and Dr. Yunde Zhao for the *yucca* mutant. We also thank Dr. Mark Estelle, Dr. Barbara Kunkel, and Dr. Ludmila Tyler for helpful discussions on the manuscript. This work was supported by a National Science Foundation 2010 Project grant to X.D. and the Hargitt postdoctoral fellowship to K.P.-M. The auxin measurements were performed through a National Science Foundation Plant Genome Program grant to Dr. Jerry Cohen at UMN and a National Science Foundation-MCB Small Grant for Exploratory Research supporting A.H.C.

Received: May 4, 2007

Revised: September 4, 2007

Accepted: September 5, 2007

Published online: October 4, 2007

References

- Dharmasiri, N., Dharmasiri, S., and Estelle, M. (2005). The F-box protein TIR1 is an auxin receptor. *Nature* 435, 441–445.
- Kepinski, S., and Leyser, O. (2005). The Arabidopsis F-box protein TIR1 is an auxin receptor. *Nature* 435, 446–451.
- Dharmasiri, N., Dharmasiri, S., Weijers, D., Lechner, E., Yamada, M., Hobbie, L., Ehrismann, J.S., Jurgens, G., and Estelle, M. (2005). Plant development is regulated by a family of auxin receptor F box proteins. *Dev. Cell* 9, 109–119.
- Gray, W.M., Kepinski, S., Rouse, D., Leyser, O., and Estelle, M. (2001). Auxin regulates SCF(TIR1)-dependent degradation of AUX/IAA proteins. *Nature* 414, 271–276.
- Manulis, S., Haviv-Chesner, A., Brandl, M.T., Lindow, S.E., and Barash, I. (1998). Differential involvement of indole-3-acetic acid biosynthetic pathways in pathogenicity and epiphytic fitness of *Erwinia herbicola* pv. *gypsophylae*. *Mol. Plant Microbe Interact.* 11, 634–642.
- Manulis, S., Shafrir, H., Epstein, E., Lichter, A., and Barash, I. (1994). Biosynthesis of indole-3-acetic acid via the indole-3-acetamide pathway in *Streptomyces* spp. *Microbiology* 140, 1045–1050.
- Glickmann, E., Gardan, L., Jacquet, S., Hussain, S., Elasri, M., Petit, A., and Dessaux, Y. (1998). Auxin production is a common feature of most pathovars of *Pseudomonas syringae*. *Mol. Plant Microbe Interact.* 11, 156–162.
- Ansari, M.M., and Sridhar, R. (2000). Some tryptophan pathways in the phytopathogen *Xanthomonas oryzae* pv. *oryzae*. *Folia Microbiol. (Praha)* 45, 531–537.
- Chung, K.R., Shilts, T., Erturk, U., Timmer, L.W., and Ueng, P.P. (2003). Indole derivatives produced by the fungus *Colletotrichum acutatum* causing lime anthracnose and postbloom fruit drop of citrus. *FEMS Microbiol. Lett.* 226, 23–30.
- Maor, R., Haskin, S., Levi-Kedmi, H., and Sharon, A. (2004). In planta production of indole-3-acetic acid by *Colletotrichum gloeosporioides* f. sp. *aeschynomene*. *Appl. Environ. Microbiol.* 70, 1852–1854.
- Vandeputte, O., Oden, S., Mol, A., Vereecke, D., Goethals, K., El Jaziri, M., and Prinsen, E. (2005). Biosynthesis of auxin by the gram-positive phytopathogen *Rhodococcus fascians* is controlled by compounds specific to infected plant tissues. *Appl. Environ. Microbiol.* 71, 1169–1177.
- Wang, D., Amornsiripanitch, N., and Dong, X. (2006). A genomic approach to identify regulatory nodes in the transcriptional network of systemic acquired resistance in plants. *PLoS Pathog.* 2, e123.
- Yang, Y., Hammes, U.Z., Taylor, C.G., Schachtman, D.P., and Nielsen, E. (2006). High-affinity auxin transport by the AUX1 influx carrier protein. *Curr. Biol.* 16, 1123–1127.
- Leyser, O. (2005). Auxin distribution and plant pattern formation: How many angels can dance on the point of PIN? *Cell* 121, 819–822.
- Hagen, G., and Guilfoyle, T. (2002). Auxin-responsive gene expression: Genes, promoters and regulatory factors. *Plant Mol. Biol.* 49, 373–385.
- Woodward, A.W., and Bartel, B. (2005). Auxin: Regulation, action, and interaction. *Ann. Bot. (Lond.)* 95, 707–735.
- Spivey, N.W., Wang, D., and Dong, X. (2007). Arabidopsis Information Resource (www.arabidopsis.org).
- Wang, D., Weaver, N.D., Kesarwani, M., and Dong, X. (2005). Induction of protein secretory pathway is required for systemic acquired resistance. *Science* 308, 1036–1040.
- Ulmasov, T., Murfett, J., Hagen, G., and Guilfoyle, T.J. (1997). Aux/IAA proteins repress expression of reporter genes containing natural and highly active synthetic auxin response elements. *Plant Cell* 9, 1963–1971.
- Cao, H., Bowling, S.A., Gordon, S., and Dong, X. (1994). Characterization of an Arabidopsis mutant that is nonresponsive to inducers of systemic acquired resistance. *Plant Cell* 6, 1583–1592.
- Bowling, S.A., Clarke, J.D., Liu, Y., Klessig, D.F., and Dong, X. (1997). The *cpr5* mutant of Arabidopsis expresses both NPR1-dependent and NPR1-independent resistance. *Plant Cell* 9, 1573–1584.
- Clarke, J.D., Liu, Y., Klessig, D.F., and Dong, X. (1998). Uncoupling PR gene expression from NPR1 and bacterial resistance: Characterization of the dominant Arabidopsis *cpr6-1* mutant. *Plant Cell* 10, 557–569.
- Li, X., Clarke, J.D., Zhang, Y., and Dong, X. (2001). Activation of an EDS1-mediated R-gene pathway in the *src1* mutant leads to constitutive, NPR1-independent pathogen resistance. *Mol. Plant Microbe Interact.* 14, 1131–1139.
- Zhao, Y., Christensen, S.K., Fankhauser, C., Cashman, J.R., Cohen, J.D., Weigel, D., and Chory, J. (2001). A role for flavin

(F) A model for SA-mediated inhibition of the auxin pathway as part of the defense response. Pathogens manipulate auxin responses to both promote susceptibility (solid arrow) and attenuate defense gene expression in the host (solid blocked arrow). To counter pathogen attack, SA stabilizes auxin-response repressors by downregulating auxin receptors (solid blocked arrow). Long-term SA accumulation might also reduce auxin levels (dashed blocked arrow). This antagonism of auxin signaling, together with other SA-mediated responses, leads to disease resistance.

monooxygenase-like enzymes in auxin biosynthesis. *Science* 297, 306–309.

25. Navarro, L., Dunoyer, P., Jay, F., Arnold, B., Dharmasiri, N., Estelle, M., Voinnet, O., and Jones, J.D. (2006). A plant miRNA contributes to antibacterial resistance by repressing auxin signaling. *Science* 312, 436–439.
26. Yamada, T. (1993). The role of auxin in plant-disease development. *Annu. Rev. Phytopathol.* 31, 253–273.
27. Nagpal, P., Walker, L.M., Young, J.C., Sonawala, A., Timppe, C., Estelle, M., and Reed, J.W. (2000). AXR2 encodes a member of the Aux/IAA protein family. *Plant Physiol.* 123, 563–574.