

THE EFFECTS OF THYROXINE TREATMENT, IN VIVO AND IN VITRO, ON Ca^{2+} EFFLUX FROM RAT LIVER MITOCHONDRIA

S. B. SHEARS* and J. R. BRONK

Department of Biology, University of York, York YO1 5DD, England

Received 28 January 1981

1. Introduction

We have proposed [1,2] that thyroid hormones act in vivo to stimulate mitochondrial state 4 [3] respiratory rates by inhibiting an ohmic component of the conductance of the inner mitochondrial membrane and thus elevating $\Delta\mu\text{H}^+$ and activating non-ohmic conductance pathways. Ca^{2+} recycling across the mitochondrial membrane necessitates respiratory-driven H^+ efflux to maintain the steady state and involves independent influx and efflux mechanisms [4] which exhibit ohmic [5] and non-ohmic properties [6]. Several hormones including insulin [7], glucagon [8–10] and the catecholamines [10] have been shown to influence these Ca^{2+} fluxes. Therefore, reports that Ca^{2+} efflux is enhanced by thyroid hormones added in vitro [11,12] suggested that Ca^{2+} efflux might be the non-ohmic pathway which was stimulated by our in vivo thyroxine treatment. We now report that this treatment does stimulate Ca^{2+} efflux although there are important differences between the in vitro and in vivo actions of thyroxine. However, we also show that the extra Ca^{2+} cycling does not contribute significantly to the increased respiration resulting from thyroxine treatment in vivo.

2. Methods

Male Wistar rats (250 g) were injected with either 8 mg thyroxine/kg body wt or isotonic saline, and after

Abbreviations: $\Delta\mu\text{H}^+$, mitochondrial proton electrochemical potential gradient; $\Delta\Psi$, transmembrane electrical potential; CCFP, carbonylcyanide *p*-trifluoromethoxyphenylhydrazone; EGTA, ethylenedis (oxonitrilo) tetraacetic acid

* Present address: Department of Biochemistry, University of Edinburgh, Edinburgh, Scotland

24 h liver mitochondria were isolated as in [1,2]. Protein was assayed by a biuret method [13]. Mitochondrial Ca^{2+} fluxes were determined in one of two media: medium (A) contained 80 mM sucrose, 20 mM KCl, 10 mM PO_4 (as Tris- PO_4 , pH 7.0), 5 mM MgCl_2 , and 5 mM Tris-succinate; (B) was identical with (A), except for the absence of KCl and MgCl_2 , since Mg^{2+} [14] and K^+ [15] modulate Ca^{2+} transport. Mitochondria (2 mg protein/ml) were incubated in the presence of $^{45}\text{Ca}^{2+}$ (0.6 $\mu\text{Ci/ml}$) as CaCl_2 at either 19 or 45 nats Ca^{2+}/mg protein. The final volume was 1.6 ml. At 0.75–4 min after the addition of the mitochondria, 0.2 ml samples were removed and spun at $10\,000 \times g$ for 0.5 min in a Beckman microfuge. The supernatant was immediately poured off and 0.05 ml was taken to 1.5 ml with scintillation fluid (Fisoflor 'mpc', Fisons, Loughborough) and counted in a Beckman LS-230. For each experiment, endogenous non-radioactive Ca^{2+} was determined [2] after adding 0.5 ml stock mitochondrial suspension to 0.5 ml 2 M perchloric acid. Full exchange ability of radioactive with non-radioactive pools of Ca^{2+} was assumed.

EGTA-induced Ca^{2+} efflux was estimated following the addition of 4 μM ruthenium red and 2 mM EGTA [16] to the medium 1 min after the mitochondria were added. Control experiments (not shown) indicated that Ca^{2+} uptake was inhibited by >98% during the time course of our incubations. The contribution of Ca^{2+} cycling to respiratory rates was estimated by assuming Ca^{2+} efflux via $\text{Ca}^{2+}/2 \text{H}^+$ antiport and Ca^{2+} influx by Ca^{2+} uniport [6], and a succinate supported $\text{H}^+:\text{O}$ stoichiometry of 6 [17].

Respiratory rates and ADP:O ratios were determined as in [1,2]. All incubations were stirred, and maintained at 30°C.

CCFP, ADP and ruthenium red were supplied by Sigma, Kingston Upon Thames. Ruthenium red was

purified [18]. $^{45}\text{CaCl}_2$ was purchased from Radiochemical Centre, Amersham.

3. Results and discussion

With mitochondria from euthyroid and thyroxine-treated rats, respiratory rates were unaffected when ruthenium red and EGTA were used to inhibit recycling of endogenous Ca^{2+} across the mitochondrial membrane (table 1). Thus our thyroxine treatment increased respiratory rates by up to 88% when recycling of endogenous Ca^{2+} was not making a significant contribution.

We also noted (table 1) that CCFP-stimulated respiratory rates were enhanced by thyroxine treatment, due presumably to a direct effect of the hormone upon the respiratory enzymes. It is not clear to what extent such a mechanism might be responsible for the stimulation of state 3 respiratory rate, which may be further improved by a hormone-mediated increase in the uptake of ADP and phosphate by the mitochondria [1,2,19]. Our measurements of Ca^{2+} efflux were made using <50 natoms Ca^{2+}/mg protein since above this level the rate of Ca^{2+} efflux increases dramatically [20]. It has been suggested [20] that at <50 natoms $\text{Ca}^{2+}/\text{protein}$, Ca^{2+} efflux occurred at a relatively slow rate because the matrix Ca^{2+} activity was significantly diminished as a consequence of extensive precipita-

tion and binding of this cation. We disagree with this explanation since, as pointed out in [21], it is difficult to understand how the matrix Ca^{2+} activity could be low under these conditions since retention of the ion is almost totally dependent upon $\Delta\Psi$ and virtually all the Ca^{2+} is lost from the mitochondria within seconds of the addition of CCFP (e.g., table 2). Perhaps the increased rate of Ca^{2+} efflux following addition of >50 natoms Ca^{2+}/mg protein [20], reflects some destabilization of the mitochondrial membrane with a consequent reduction in the value of $\Delta\Psi$. Under such conditions, Ca^{2+} leaves the mitochondria via a reversal of the Ca^{2+} influx carrier [21], a process apparently insensitive to ruthenium red [22], (line 4 in, table 2). We have also observed that thyroxine treatment *in vivo*, appears to protect against the destabilizing effects of the higher Ca^{2+} levels [2] and to avoid these complicating effects, we have carried out our experiments with <50 natoms Ca^{2+}/mg protein.

Our data (lines 1–3 in table 2) indicate that thyroxine treatment *in vivo* does increase the rate of Ca^{2+} efflux from isolated mitochondria by 27–51%. The effect cannot be a consequence of the hormone uncoupling the mitochondria since the thyroxine treatment does not modify ADP:O ratios (table 1). This conclusion is further substantiated by the increases we have observed in $\Delta\mu\text{H}^+$ and ion uptake [1,2]. In the steady state, the effect of Ca^{2+} efflux (table 2) on respiratory rate (section 2) would be ≤ 1 natom

Table 1
The effect of *in vivo* thyroxine treatment upon respiratory rates and ADP:O ratios of isolated mitochondria, and the influence of ruthenium red plus EGTA

	ADP:O	Respiratory rate (natoms O . mg protein ⁻¹ . min ⁻¹)		
		State 4	State 3	Uncoupled
Control	1.88 ± 0.2 (3)	13.5 ± 0.9 (7)	73.2 ± 3.2 (7)	59.4 ± 4.7 (4)
Control plus ruthenium red and EGTA	1.89 ± 0.17 (3)	12.6 ± 1.3 (4)	80.6 ± 11.2 (3)	n.d.
Thyroxine-treated	1.85 ± 0.16 (3)	25.4 ± 0.8 (7)	106.9 ± 9.3 (7)	98.4 ± 3.9 (4)
Thyroxine-treated plus ruthenium red and EGTA	1.85 ± 0.17 (3)	23.8 ± 1.7 (4)	123.0 ± 20.7 (3)	n.d.

Incubation conditions: medium (A) containing 2 mg mitochondrial protein/ml. ADP (0.2 $\mu\text{mol}/\text{mg}$ protein) added at 2 min. Uncoupled mitochondria were those treated with 0.3 μM CCFP. Where present, 4 μM ruthenium red and 2 mM EGTA were added immediately after the mitochondria. Data include standard errors of the mean, and the number of mitochondrial preparations in parentheses; n.d., not determined

Table 2
The effect of *in vivo* and *in vitro* thyroxine treatment upon mitochondrial Ca^{2+} fluxes and respiratory rates under various conditions

Line	Medium	Ca^{2+} added (natoms/ mg protein)	Control		Thyroxine-treated	
			-RR	+RR	-RR	+RR
ΔCa^{2+} (natoms . mg protein ⁻¹ . min ⁻¹)						
1	A	19	-0.08	-1.55	-0.07	-1.97
2	A	45	-0.18	-2.2	+0.09	-3.04
3	B	19	-0.12	-1.95	+0.08	-2.95
4	B + CCFP	19	-55	-58.2	-69.7	-79.4
5	B + Thyroxine	19	-57.2	- 4.76	n.d.	n.d.
Respiratory rate (natoms O . mg protein ⁻¹ . min ⁻¹)						
6	A	19	16.4	16.4	26.1	31
7	A	45	17.6	19.5	35.3	33.4
8	B	19	18.8	19.9	28.4	30.8
9	B + CCFP	19	50.1	48.9	90.5	70.5
10	B + Thyroxine	19	53.0	25.6	n.d.	n.d.

Incubation conditions: medium (A) or (B) as indicated. Where present, CCFP at 0.3 μM , thyroxine at 100 μM . RR represents presence of 4 μM ruthenium red plus 2 mM EGTA. Negative ΔCa^{2+} indicates loss from mitochondria. - is inserted where rapidity of flux is such that it exceeds the ability of our technique to record it accurately. All respiratory rates are means from two mitochondrial preparations; 2 or 3 preparations were used to record Ca^{2+} fluxes; n.d., not determined

O . mg protein⁻¹ . min⁻¹. Moreover, the different rates of Ca^{2+} efflux of the two populations of mitochondria account for respiratory differences of ≤ 0.3 natoms O . mg protein⁻¹ . min⁻¹, i.e., 3% of the respiratory enhancement due to thyroxine treatment (lines 3,8 in table 2). It is unlikely that we underestimated the extent of Ca^{2+} efflux, because of any inhibition by ruthenium red, since the latter did not inhibit respiratory rate (lines 6-8 in table 2).

Our data may be compared with those in [12]; following the accumulation by heart mitochondria of 20 natoms Ca/mg protein in the absence of Mg^{2+} , ruthenium red and ~50 nmol thyroxine/mg protein were added. The latter induced a >10-fold stimulation in the rate of Ca^{2+} efflux and a similar phenomenon was observed in liver mitochondria [11]. We have confirmed this *in vitro* action of thyroxine (compare lines 3 and 5 in table 2) but the interpretation of these results is likely to be complicated. For instance, in the absence of ruthenium red, the thyroxine-induced increase in respiratory rate (line 10) was not further enhanced by CCFP (not shown). Moreover, the rate of Ca^{2+} efflux from mitochondria treated with thyroxine *in vitro* (line 5) was similar to that

from uncoupled mitochondria (line 4). Thus thyroxine *in vitro* may uncouple the mitochondria [23] inducing a reversal of the Ca^{2+} influx mechanism [21]. Alternatively, if the hormone activates the independent Ca^{2+} efflux carrier, subsequent Ca^{2+} recycling may uncouple the organelles. In any case, these events differ from the effects of thyroxine treatment *in vivo* (see above). Note also that ruthenium red largely reverses or inhibits the increases in rates of Ca^{2+} efflux and respiration caused by thyroxine *in vitro* (lines 5 and 10). In conclusion, thyroxine treatment *in vivo* stimulates Ca^{2+} efflux in contrast to the inhibitory effect of glucagon and catecholamines [10] on this process. However, the thyroid hormone effects on Ca^{2+} efflux do not appear to contribute significantly to the enhancement of respiratory rate by the hormone.

References

- [1] Shears, S. B. and Bronk, J. R. (1979) *Biochem. J.* 178, 505-507.
- [2] Shears, S. B. and Bronk, J. R. (1981) *J. Bioenerg. Biomembr.* in press.

- [3] Chance, B. and Williams, G. R. (1955) *J. Biol. Chem.* 217, 409–427.
- [4] Puskin, J. S., Gunter, T. E., Gunter, K. K. and Russell, P. R. (1976) *Biochemistry* 15, 3834–3842.
- [5] Akerman, K. E. O. (1980) *Biochem. Soc. Trans.* 8, 262–264.
- [6] Heaton, G. M. and Nicholls, D. G. (1976) *Biochem. J.* 156, 635–646.
- [7] Dorman, D. M., Barritt, G. J. and Bygrave, F. L. (1975) *Biochem. J.* 150, 389–395.
- [8] Prpic, V., Spencer, T. L. and Bygrave, F. L. (1978) *Biochem. J.* 176, 705–714.
- [9] Hughes, B. P. and Barritt, G. J. (1978) *Biochem. J.* 176, 295–304.
- [10] Taylor, W. M., Prpic, V., Exton, J. H. and Bygrave, F. L. (1980) *Biochem. J.* 188, 443–450.
- [11] Al-Shaikhaly, M. H. and Baum, H. (1979) *Biochem. Soc. Trans.* 7, 215–216.
- [12] Harris, E. J., Al-Shaikhaly, M. and Baum, H. (1979) *Biochem. J.* 182, 455–464.
- [13] Layne, E. (1957) *Methods Enzymol.* 3, 447–454.
- [14] Sordahl, L. A. (1975) *Arch. Biochem. Biophys.* 167, 104–115.
- [15] Crompton, M., Heid, I. and Carafoli, E. (1980) *FEBS Lett.* 115, 257–259.
- [16] Reed, K. C. and Bygrave, F. L. (1975) *Anal. Biochem.* 67, 44–54.
- [17] Brand, M. D., Reynafarje, B. and Lehninger, A. L. (1976) *J. Biol. Chem.* 251, 5670–5679.
- [18] Fletcher, J. M., Greenfield, B. F., Hardy, C. J., Scargill, D. and Woodhead, J. L. (1961) *J. Chem. Soc. London* 2000–2006.
- [19] Babior, B. M., Greagan, S., Inbar, S. H. and Kipnes, R. S. (1973) *Proc. Natl. Acad. Sci. USA* 70, 98–102.
- [20] Dawson, A. P. and Fulton, D. V. (1980) *Biochem. J.* 188, 749–755.
- [21] Nicholls, D. G. (1978) *Biochem. J.* 176, 463–474.
- [22] Pozzan, T., Brogandin, M. and Azzone, G. F. (1977) *Biochemistry* 16, 5618–5625.
- [23] Bronk, J. R. (1965) *Biochim. Biophys. Acta* 97, 9–15.