

Silica Crystals and Aluminum Salts Regulate the Production of Prostaglandin in Macrophages via NALP3 Inflammasome-Independent Mechanisms

Etsushi Kuroda,^{1,*} Ken J. Ishii,^{3,4} Satoshi Uematsu,⁵ Keiichi Ohata,⁶ Cevayir Coban,⁶ Shizuo Akira,⁵ Kosuke Aritake,⁷ Yoshihiro Urade,⁷ and Yasuo Morimoto²

¹Department of Immunology and Parasitology

²Department of Occupational Pneumology

University of Occupational and Environmental Health, Japan, Kitakyushu, Fukuoka, 807-8555, Japan

³Laboratory of Adjuvant Innovation, National Institute of Biomedical Innovation, Ibaraki, Osaka, 567-0085, Japan

⁴Laboratory of Vaccine Science

⁵Laboratory of Host Defense

⁶Laboratory of Malaria Immunology WPI Immunology Frontier Research Center

Osaka University, Suita, Osaka, 565-0871, Japan

⁷Department of Molecular Behavioral Biology, Osaka Bioscience Institute, Suita, Osaka, 565-0874, Japan

*Correspondence: kuroetu@med.uoeh-u.ac.jp

DOI 10.1016/j.immuni.2011.03.019

SUMMARY

Particulates such as silica crystal (silica) and aluminum salts (alum) activate the inflammasome and induce the secretion of proinflammatory cytokines in macrophages. These particulates also induce the production of immunoglobulin E via a T helper 2 (Th2) cell-associated mechanism. However, the mechanism involved in the induction of type 2 immunity has not been elucidated. Here, we showed that silica and alum induced lipopolysaccharide-primed macrophages to produce the lipid mediator prostaglandin E₂ (PGE₂) and interleukin-1 β (IL-1 β). Macrophages deficient in the inflammasome components caspase 1, NALP3, and ASC revealed that PGE₂ production was independent of the NALP3 inflammasome. PGE₂ expression was markedly reduced in PGE synthase-deficient (*Ptges*^{-/-}) macrophages, and *Ptges*^{-/-} mice displayed reduced antigen-specific serum IgE concentrations after immunization with alum or silica. Our results indicate that silica and alum regulate the production of PGE₂ and that the induction of PGE₂ by particulates controls the immune response in vivo.

INTRODUCTION

Some particulates and crystals can stimulate the innate immune system to induce inflammatory responses. In particular, aluminum salts (referred to as alum) and silica crystals can induce type 2 inflammatory responses, which are characterized by the accumulation of eosinophils at the site of injection and the elevation of antigen-specific serum IgE and IgG1 amounts in vivo (Aimanianda et al., 2009; Marrack et al., 2009; Kumar et al., 2009). However, the basis for the adjuvanticity of these particu-

lates and the mechanisms by which they elicit type 2 immunity remain poorly understood.

In the innate immune system, macrophages and dendritic cells (DCs) function as the first line of defense against foreign antigens. These cells can recognize pathogen-associated molecular patterns (PAMPs) through pattern-recognition receptors (PPRs) and can induce inflammatory responses (Akira et al., 2006). NOD-like receptors (NLRs) are intracellular PRRs (Ye and Ting, 2008). Among the known NLRs, NALP3 (also known as NLRP3, Cryopyrin, CIAS1, or PYPAF1) is one of the best characterized. Upon activation, NALP3 forms a multiprotein complex with apoptosis-associated speck-like protein containing a caspase recruitment domain (ASC) and caspase-1. This complex, referred to as the NALP3 inflammasome, promotes the secretion of the proinflammatory cytokines interleukin-1 β (IL-1 β) and IL-18 by the action of caspase-1 (Franchi et al., 2009; Martinon et al., 2009; Schroder and Tschopp, 2010; Schroder et al., 2010). The activated NALP3 inflammasome contributes to antifungal host defense, antitumor immunity, and inflammation (Düewell et al., 2010; Ghiringhelli et al., 2009; Gross et al., 2009; Halle et al., 2008; Watanabe et al., 2008). In addition to being activated by PAMPs, the NALP3 inflammasome is also activated by ATP, various crystals (i.e., silica, asbestos and monosodium urate), and alum (Cassel et al., 2008; Dostert et al., 2008; Eisenbarth et al., 2008; Hornung et al., 2008). The NALP3 inflammasome has been shown to induce type 2 immune responses, and *Nalp3*^{-/-} and *Asc*^{-/-} mice display reduced type 2 immune responses to alum (Eisenbarth et al., 2008; Kool et al., 2008; Li et al., 2008). However, other reports have shown that the NALP3 inflammasome is dispensable for alum adjuvanticity (Franchi and Núñez, 2008; McKee et al., 2009). In addition, it has been reported that the NALP3 inflammasome is not required for antibody production in response to vaccination by the particulate adjuvant (Sharp et al., 2009). Thus, the role of the NALP3 inflammasome in the induction of type 2 immunity is still open to further investigation.

In addition to proinflammatory cytokines, lipid mediators such as prostaglandins (PGs) are also involved in the induction of

inflammatory responses (Narumiya, 2009). PGE₂, a well-characterized proinflammatory lipid mediator, is an arachidonic acid metabolite that is produced by various types of cells including antigen-presenting cells. Previous reports have shown that PGE₂ suppresses T helper 1 (Th1) cell-type responses by elevating intracellular cAMP concentrations in DCs, macrophages, and Th1 cells, thus inhibiting their ability to produce type 1 cytokines such as IL-12 and IFN- γ (Fabricius et al., 2010; Koga et al., 2009; Kuroda and Yamashita, 2003). In addition, PGE₂ can enhance IL-23 production by DCs and favors Th17 cell polarization and IL-17 production (Boniface et al., 2009; Yao et al., 2009). More recently, PGE₂ was shown to facilitate Th1 cell differentiation in the presence of IL-12 and high doses of the costimulatory CD28 antibody, via activation of the PI3-kinase pathway (Yao et al., 2009). Thus, PGE₂ has various functions in the regulation of immune responses. However, the involvement of PGE₂ in particulate-mediated adjuvanticity has not been investigated.

In this study, we showed that alum, silica, and ATP, which activate the NALP3 inflammasome, induced macrophages to produce IL-1 β , IL-18 and PGE₂. Interestingly, PGE₂ production in macrophages was regulated by the spleen tyrosine kinase (Syk) and p38 MAP kinase pathway but did not depend on inflammasome activation. In addition, particulate-induced PGE₂ regulated antigen-specific serum IgE production in vivo. Our results suggest that the activation of the PGE₂ pathway by particulates may be an important signal for the induction of type 2 immune responses.

RESULTS

Silica, Alum, and ATP Induce PGE₂ Production in LPS-Primed Macrophages

Many reports have shown that alum, silica, and ATP stimulate macrophages to produce the caspase-1-dependent cytokines IL-1 β and IL-18 by activating the NALP3 inflammasome (Cassel et al., 2008; Dostert et al., 2008; Eisenbarth et al., 2008; Hornung et al., 2008). These cytokines are thought to be important for regulation of immune responses. However, some studies have shown that caspase-1-dependent cytokines are dispensable for immune regulation (Franchi and Núñez, 2008; McKee et al., 2009; Sharp et al., 2009). Thus, we first ascertained whether inflammasome activators could induce the production of caspase-1-independent factor(s) in macrophages. To this end, we examined the production of several cytokines, chemokines and lipid mediators by macrophages in response to silica, alum and ATP. Silica, alum and ATP induced LPS-primed macrophages to produce IL-1 β and IL-18, which is in agreement with previous reports (Cassel et al., 2008; Dostert et al., 2008; Eisenbarth et al., 2008; Hornung et al., 2008). These inflammasome activators also induced LPS-primed macrophages to produce PGE₂ (Figure 1A). A time-course analysis revealed that silica stimulation of LPS-primed macrophages for 2 hr was sufficient for detecting similar amounts of PGE₂ and IL-1 β , in terms of pg/ml secreted (Figure 1B). In our experiments, we tested three different alum compounds: Imject alum (a mixture of aluminum hydroxide and magnesium hydroxide; Thermo Scientific), alhydrogel (aluminum hydroxide gel; Sigma-Aldrich) and LSL alum (aluminum hydroxide hydrate gel suspension; LSL). The experiments

described in Figures 1A and 1B were performed with LSL alum as a stimulator. To test which alum compound was the most potent PGE₂ inducer in macrophages, we stimulated LPS-primed macrophages with the three different compounds. Each alum compound stimulated the production of PGE₂, but LSL alum (referred to hereafter as alum) induced the highest PGE₂ production in LPS-primed macrophages. Alhydrogel induced modest production of PGE₂ in LPS-primed macrophages, whereas Imject alum induced low amounts of PGE₂ (Figure S1A). It is important to note that LPS-primed macrophages did not produce any other well-characterized inflammatory cytokines or chemokines in response to silica (Figure S1B) or alum (data not shown).

Activated macrophages are known to produce PGD₂ (Mohri et al., 2003). Indeed, we found that LPS-primed macrophages produced PGD₂ in response to silica (Figure 1C). We also examined the effect of titanium dioxide (TiO₂), which does not cause severe inflammation on inhalational exposure and does not activate the NALP3 inflammasome (Cassel et al., 2008), on PGE₂ production by macrophages. As shown in Figure 1D, TiO₂ did not induce IL-1 β or PGE₂ production in LPS-primed macrophages. Activation of the NALP3 inflammasome in DCs contributes to particulate-mediated adjuvanticity (Kool et al., 2008). Thus, we compared PGE₂ production by bone marrow (BM)-derived DCs and macrophages. BM-derived DCs produced IL-1 β and PGE₂ in response to silica and alum, but the amounts of PGE₂ were markedly higher in macrophages than in DCs (Figure 1E). We have previously reported that macrophages from BALB/c mice produce higher amounts of PGE₂ than those from C57BL/6 mice (Kuroda et al., 2007; Kuroda and Yamashita, 2003). Macrophages from BALB/c mice produced ~2-fold greater amounts of IL-1 β and PGE₂ in response to silica than macrophages from C57BL/6 mice (Figure 1F). Our observations are not restricted to mouse studies, given that human peripheral blood mononuclear cells (PBMCs) also produced PGE₂ and IL-1 β in response to silica (Figure 1G). These results indicate that in addition to IL-1 β and IL-18, macrophages also produce PG in response to silica, alum, and ATP.

Silica- and Alum-Induced PGE₂ Production Is Independent of the Activity of Caspase-1

IL-1 β and IL-18 are known to induce PG production (Dinarello, 2002; Lee et al., 2004). However, simple exposure of LPS-primed macrophages to IL-1 β and/or IL-18 did not induce the production of PGE₂ (Figure 2A). BM cells from WT and IL-1 receptor-deficient mice were cultured for 7 days with macrophage colony stimulating factor (M-CSF), and adherent macrophages were analyzed for silica- and alum-induced PGE₂ production. As shown in Figure 2B, WT and *Il1r1*^{-/-} macrophages produced comparable amounts of PGE₂. In addition, treatment of silica- or alum-activated macrophages with a caspase-1 inhibitor significantly reduced their ability to produce IL-1 β but did not affect their ability to produce PGE₂ (Figure 2C). Because silica and alum can induce cell death, we wanted to rule out the possibility that the enhanced production of PGE₂ in silica- and alum-activated macrophages was due to cell death. For this purpose, we measured cell death in silica- and alum-activated macrophages by quantifying the release of lactose dehydrogenase (LDH) into the culture media. Treatment of macrophages with a caspase-1 inhibitor reduced silica- and alum-induced




Figure 1. Inflammation Activators Induce Macrophages to Produce PGE₂

(A) Peritoneal macrophages from BALB/c mice were primed for 3 hr with or without low-dose LPS (1 ng/ml) and then stimulated with silica, alum or ATP for 2 hr. (B) Macrophages with or without LPS priming were stimulated with silica for 2, 6, or 18 hr. Macrophages stimulated with high-dose LPS (1 μg/ml) for 6 hr were used as controls. (C) The culture supernatants of macrophages stimulated for 2 hr with silica were analyzed for PGD₂ production as indicated in (B). (D) LPS-primed macrophages were stimulated with silica or TiO₂ for 2 hr. (E) BM-derived macrophages and DCs were primed with LPS and then stimulated with 100 μg/ml silica or 400 μg/ml alum for 6 hr. (F) Peritoneal macrophages from C57BL/6 (B6) and BALB/c (BALB) mice were primed with LPS and then stimulated with 100 μg/ml silica or 400 μg/ml alum for 2 hr. (G) Human PBMCs were primed for 3 hr with LPS and then stimulated with silica for 3 hr. For all experiments, the amounts of IL-1β and PGE₂ in the culture supernatants were determined by ELISA. Data represent mean ± SE of three to five independent experiments (*p < 0.01).




Figure 2. Silica- and Alum-Induced Lysosomal Damage Triggers PGE₂ Production in Macrophages via Caspase-1-Independent Mechanisms
 (A) LPS-primed peritoneal macrophages were stimulated with IL-1 β or IL-1 β plus IL-18 for 2 or 6 hr. Macrophages stimulated with high-dose LPS were used as controls.
 (B) M-CSF-derived BM macrophages from WT (C57BL/6) and *Il1r1*^{-/-} mice were primed with low-dose LPS and then stimulated with 50 μ g/ml silica or 200 μ g/ml alum for 6 hr.
 (C) LPS-primed macrophages were stimulated with 100 μ g/ml silica or 400 μ g/ml alum in the presence or absence of caspase-1 inhibitors for 2 hr.
 (D and E) LPS-primed macrophages were stimulated with 100 μ g/ml silica or 400 μ g/ml alum in the presence or absence of cytochalasin D (D) or cathepsin B inhibitor (E) for 2 hr.
 (F) LPS-primed macrophages were incubated with Leu-Leu-OMe for 2 hr. Data represent mean \pm SE of three (A and C–E) or two (B and F) independent experiments (* p < 0.01).

cell death (Figure S2). This result was expected because caspase-1 is involved in promoting cell death. However, we have already shown that PGE₂ production in macrophages is independent of caspase-1 activity (Figure 2C and Figure S2). Therefore, these results indicate that caspase-1-dependent cytokine release and cell death are not responsible for silica- and alum-induced PGE₂ production in macrophages.

Silica and Alum Induce PGE₂ Production in Macrophages through Phagosomal Destabilization

Phagocytes engulf particulates such as silica or alum, which leads to lysosomal damage and rupture, followed by the release of lysosomal enzymes, such as cathepsin B, into the cytoplasm. Studies suggest that lysosomal damage can activate the NALP3 inflammasome and induce the production of IL-1 β (Düewell et al., 2010; Hornung et al., 2008). To address whether the engulfment of particulates, lysosomal rupture, and release of lysosomal enzymes could induce PGE₂ production in macrophages, we cultured macrophages in the presence of cytochalasin D, which inhibits actin filament assembly and phagocytosis. Treatment of macrophages with cytochalasin D significantly reduced their ability to produce IL-1 β and PGE₂ in response to silica and alum (Figure 2D).

The release of lysosomal enzymes such as cathepsin B is thought to trigger the activation of the NALP3 inflammasome. In addition, the cathepsin B inhibitor CA-074 has been shown to inhibit the NALP3 inflammasome (Duncan et al., 2009; Hornung et al., 2008; Sharp et al., 2009). We showed the production of IL-1 β and PGE₂ by macrophages was partially inhibited by treating the cells with CA-074 (Figure 2E). Lysosomal damage has also been shown to trigger the activation of the NALP3 inflammasome. To investigate this process, we treated macrophages with leucyl-leucine methyl ester (Leu-Leu-OMe) to induce lysosomal damage (Hornung et al., 2008). Leu-Leu-OMe-treated macrophages produced large amounts of IL-1 β (Figure 2F) as previously reported. Leu-Leu-OMe-treated macrophages also produced PGE₂. Altogether, these results suggest that, similar to the stimulatory effect of NALP3 inflammasome activation and IL-1 β release, lysosomal damage and rupture triggers PGE₂ production in macrophages.

Silica- and Alum-Induced PGE₂ Production in Macrophages Is Independent of the NALP3 Inflammasome

We have shown that the NALP3 inflammasome activators silica, alum, and ATP induce LPS-primed macrophages to produce PGE₂. We have also demonstrated that PGE₂ production is independent of caspase-1 activity. To investigate whether the NALP3 inflammasome is involved in silica- and alum-induced PGE₂ production, we performed experiments similar to those described above using *Nalp3*^{-/-}, *Asc*^{-/-}, and *Casp1*^{-/-} macrophages. M-CSF-derived BM macrophages from WT and inflammasome-deficient mice were used in this experiment. As shown in Figure 3A, macrophages deficient in NALP3, ASC, or caspase-1 failed to secrete IL-1 β in response to silica and alum, which is consistent with previously published reports. However, macrophages from inflammasome-deficient mice produced slightly higher amounts of PGE₂ in response to silica and alum than cells from WT mice. IL-6 production was previously shown to be inde-

pendent of the NALP3 inflammasome (Kumar et al., 2009; Yamamoto et al., 2004). Indeed, WT and inflammasome-deficient macrophages produced comparable amounts of IL-6. Similar results were obtained when we performed the same experiments with granulocyte-macrophage colony stimulating factor (GM-CSF)-derived macrophages (Figure 3B). These results indicate that silica and alum induce macrophages to produce PGE₂ through NALP3 inflammasome-independent mechanisms.

PGE₂ synthesis is regulated by cyclooxygenase (COX) and PGE synthase. In particular, COX-2 and PTGES (also known as mPGES-1) have been reported to regulate stimulation-dependent PGE₂ production in macrophages (Kuroda and Yamashita, 2003; Uematsu et al., 2002). As shown in Figure 3C, treatment with NS-398, which is a COX-2-specific inhibitor, significantly suppressed PGE₂ production in silica-activated macrophages. Similar to COX-2 inhibition, *Ptges*^{-/-} macrophages did not produce detectable amounts of PGE₂ upon stimulation with silica (Figure 3D). However, neither COX-2 inhibition nor PTGES deficiency had an effect on silica-induced IL-1 β production in macrophages, suggesting that IL-1 β production and activation of the inflammasome are independent of PGE₂ production (Figure S3). Similar results were obtained when we performed the same experiments with alum-activated macrophages (data not shown). We then assessed the expression of COX-2 and PTGES in macrophages upon stimulation with silica and alum. As shown in Figure 3E, priming of macrophages with LPS induced the expression of the COX-2 and PTGES proteins. However, silica, alum, and ATP stimulation had no effect on COX-2 and PTGES expression in macrophages. This result indicates that silica- and alum-induced PGE₂ production in macrophages does not involve increased expression of COX-2 or PTGES. Collectively, these results indicate that silica- and alum-induced PGE₂ production in macrophages is mediated by the COX-2 and PTGES pathways.

Silica- and Alum-Induced Production of PGE₂ by Macrophages Regulates Immune Responses In Vivo

Given that *Ptges*^{-/-} macrophages cannot produce PGE₂ while retaining inflammasome function and the ability to produce IL-1 β in response to silica and alum (Figure 3 and Figure S3), we determined whether alum-induced PGE₂ production plays a role in regulating immune responses in vivo. We immunized *Ptges*^{+/+} and *Ptges*^{-/-} mice with alum plus OVA twice (day 0 and 7). Ten days after the last immunization, sera were collected and analyzed for amounts of OVA-specific IgE, IgG1, and IgG2c antibodies. In *Ptges*^{+/+} mice, OVA-alum immunization stimulated the generation of OVA-specific IgE, IgG1, and IgG2c. In contrast, *Ptges*^{-/-} mice displayed reduced amounts of OVA-specific IgE (Figure 4A). In contrast, the amounts of IgG1 and IgG2c in the sera taken from *Ptges*^{+/+} and *Ptges*^{-/-} mice were comparable (Figure 4A).

Because previous reports have shown that OVA-silica immunization induces OVA-specific antibody responses (Kumar et al., 2009), we assessed whether silica could trigger IgE responses. As shown in Figure 4B, OVA-silica immunization induced the generation of OVA-specific IgE, IgG1, and IgG2c antibodies. As observed with OVA-alum immunization, OVA-silica-immunized *Ptges*^{-/-} mice displayed reduced amounts of OVA-specific IgE. In contrast, amounts of OVA-IgG1 and IgG2c antibodies were comparable between *Ptges*^{+/+} and




Figure 3. The Mechanisms of Particulate-Induced PGE₂ Production

(A and B) Silica- and alum-induced PGE₂ production is independent of the NALP3 inflammasome. M-CSF- (A) or GM-CSF- (B) derived BM macrophages from WT (C57BL/6), *Asc*^{-/-}, *Nalp3*^{-/-}, and *Casp1*^{-/-} mice were primed with low-dose LPS and then stimulated with 100 μg/ml silica or 400 μg/ml alum for 6 hr. (C–E) Silica-induced PGE₂ production is mediated by COX-2 and PTGES. (C) LPS-primed BALB/c peritoneal macrophages were stimulated with 100 μg/ml silica in the presence or absence of the COX-2 inhibitor NS-398 for 2 hr. (D) Peritoneal macrophages from *Ptges*^{+/+} and *Ptges*^{-/-} mice were primed with low-dose LPS and then stimulated with 100 μg/ml silica for 2 hr. (E) *Ptges*^{+/+} and *Ptges*^{-/-} macrophages were primed with or without LPS and stimulated with 100 μg/ml silica, 400 μg/ml alum, or 1 mM ATP for 2 hr. Cell lysates were analyzed for COX-2, PTGES, and GAPDH (loading control) expression by western blotting. Data represent mean ± SE of two (A and B) or three (C and D) independent experiments (*p < 0.01).

Ptges^{-/-} mice. We carried out similar in vivo experiments using *Nalp3*^{-/-} and *Casp1*^{-/-} mice, and we found that the amounts of OVA-IgE from WT and mutant mice were comparable (Figure S4A). We also assessed the effect of a Th1 cell adjuvant, *Propionibacterium* (*P.*) *acnes*, on antigen-specific IgE, IgG1, and IgG2c production in vivo. We found that *P. acnes* induced comparable amounts of OVA-IgG1 and IgG2c in *Ptges*^{+/+} and *Ptges*^{-/-} mice and did not induce OVA-IgE in either genotype (Figure S4B). Although *Ptges*^{-/-} mice displayed reduced amounts of OVA-IgE after immunization, the total IgE concentrations were similar in unimmunized *Ptges*^{+/+} and *Ptges*^{-/-} mice (Figure S4C). This result indicates that *Ptges*^{-/-} mice, unlike *Ii4*^{-/-} or *Stat6*^{-/-} mice, are not Th2 cell type prone.

We also immunized hematopoietic PGD synthase-deficient (*Ptgsd*^{-/-}) mice, in which macrophages cannot produce PGD₂, with OVA-alum and analyzed sera for amounts of OVA-specific IgE. As shown in Figure 4C, the amounts of OVA-specific IgE in

the sera were similar in WT and *Ptgsd*^{-/-} mice. These results indicate that silica- and alum-induced PGE₂ production contributes to the generation of IgE antibodies in vivo. In fact, we found that PGE₂ promoted the production of IgE in spleen cells stimulated with LPS plus IL-4 or anti-CD40 plus IL-4 in vitro (Figure S4D).

Mice that received OVA alone did not exhibit an increase in OVA-specific serum IgE levels and, in fact, exhibited 50-fold lower amounts of OVA-specific IgG1 and IgG2c (Figure 5C and data not shown).

Particulate Nickel Oxide Induces Macrophages to Produce PGE₂, but Not IL-1β, and Enhances IgE production In Vivo

Nickel oxide (NiO) could induce macrophages to produce PGE₂, but not IL-1β or IL-18. NiO is a nanoparticle known to cause lung inflammation when inhaled (Nishi et al., 2009; Ogami et al., 2009). To determine whether NiO could activate macrophages, we




Figure 4. *Ptges*^{-/-} Mice Display Reduced Antigen-Specific IgE Levels after Immunization with Silica or Alum

(A and B) *Ptges*^{+/+} and *Ptges*^{-/-} mice (n = 6) were immunized twice (day 0 and 7) with OVA plus alum (A) or OVA plus silica (B). Ten days after the last immunization, sera were collected and analyzed for OVA-specific IgE, IgG1 and IgG2c antibodies by ELISA.

(C) *Ptgds*^{+/+} and *Ptgds*^{-/-} mice (n = 4) were immunized twice (day 0 and 7) with OVA plus alum. Ten days after the last immunization, sera were collected and analyzed for OVA-specific IgE antibodies by ELISA.

carried out experiments similar to those in which we used silica and alum. We found that, as observed with TiO₂ (Figure 1D), NiO failed to induce IL-1β production in LPS-primed macrophages (Figure 5A). However, NiO-activated macrophages produced significant amounts of PGE₂ at amounts comparable to silica-activated macrophages. As expected, NiO-induced PGE₂ production in macrophages is independent of the NALP3 inflammasome (Figure 5B). In addition, mice immunized with OVA plus NiO exhibited a significant increase in OVA-specific serum IgE concentrations (Figure 5C). In contrast, TiO₂, which did not activate the NALP3 inflammasome or induce PGE₂ production in macrophages in vitro (Figures 1D and 5A), stimulated much weaker IgE responses than immunization with alum and NiO (Figure 5C). These results indicate that particulates that induce PGE₂ production, but not inflammasome activation in macrophages, positively regulate the generation of IgE antibodies in vivo.

Silica-Dependent PGE₂ Production Is Regulated by the Syk and the p38 MAP Kinase Pathway

The mechanisms through which silica, alum, and ATP induce the production of PGE₂ in macrophages are unclear. Therefore, we

sought to determine which signaling pathway was involved in the production of PGE₂. To this end, we stimulated LPS-treated macrophages with silica in the presence or absence of various signaling inhibitors, and we then determined which inhibitor(s) suppressed PGE₂ but had no effect on the production of IL-1β. As shown in Figure 6A, only wortmannin suppressed silica-induced production of IL-1β. In contrast, SB203580, U0126, and SP600125 suppressed silica-induced production of PGE₂. We also found that cyclosporin A, rapamycin, and wedelolactone partially suppressed silica-induced PGE₂ production. We conducted similar experiments by using ATP-activated macrophages. SB203580 and wortmannin enhanced ATP-induced IL-1β production, whereas the other inhibitors reduced IL-1β production. With the exception of rapamycin, all the inhibitors reduced PGE₂ production in ATP-activated macrophages (Figure 6B). It is worth noting that SB203580 had no effect on the amounts of COX-2 and PTGES expressed by macrophages (Figure S5A). Taken together, these data suggest that the p38 MAP kinase inhibitor SB203580 preferentially suppresses the production of PGE₂ in both silica- and ATP-activated macrophages. We also sought to ascertain whether silica and ATP




Figure 5. NiO Functions as a Th2 Adjuvant

(A) LPS-primed BALB/c peritoneal macrophages were stimulated with 100 μ g/ml silica, 50 or 100 μ g/ml NiO, or 50 or 100 μ g/ml TiO₂ for 2 hr. (B) BM-derived macrophages from WT (C57BL/6), *Asc*^{-/-}, *Nalp3*^{-/-} or *Casp1*^{-/-} mice were primed with low-dose LPS and then stimulated with 100 μ g/ml NiO for 6 hr. Data represent mean \pm SE of two (B) or three (A) independent experiments (**p* < 0.01). (C) C57BL/6 mice (*n* = 4) were immunized twice (day 0 and 7) with OVA alone, OVA plus TiO₂, OVA plus alum, or OVA plus NiO. Ten days after the last immunization, sera were collected and analyzed for OVA-specific IgE antibodies by ELISA.

could activate p38 MAP kinase in macrophages and found that stimulation of LPS-primed macrophages with silica and ATP induced p38 MAP kinase phosphorylation and activation (Figure 6C). Several reports have shown that ATP stimulates p38 MAP kinase, which then activates cytosolic phospholipase A₂ (cPLA₂) (Gijón et al., 2000; Ulmann et al., 2010). As shown in Figure 6D, treatment with a cPLA₂ inhibitor preferentially suppressed PGE₂ production, but not IL-1 β production (data not shown), in both ATP- and silica-activated macrophages.

Next, we examined the mechanisms involved in silica-mediated activation of p38 MAP kinase. We have shown that lysosomal rupture by Leu-Leu-OME activated macrophages to produce PGE₂ (Figure 2E). However, SB203580 significantly suppressed the production of PGE₂ but not IL-1 β in Leu-Leu-OME-treated macrophages (Figure 6E). In addition, treatment

of macrophages with poly-2-vinylpyridine-N-oxide (PVNO), which is a lysosomal stabilizing agent (Allison et al., 1966; Von Behren et al., 1983), also suppressed silica-induced PGE₂ production, which was similar to SB203580 treatment (Figure 6F). Treatment with PVNO had no effect on ATP-induced PGE₂ production, given that ATP is not involved in lysosome damage (Hornung et al., 2008). These results indicate that lysosomal damage is involved in PGE₂ production via the activation of p38 MAP kinase in silica-activated macrophages.

Several reports indicate that Syk plays an important role in antifungal responses by activating the NALP3 inflammasome (Gross et al., 2009). In addition, Syk has been shown to be involved in malarial hemozoin-mediated and monosodium urate crystal-mediated inflammasome activation (Ng et al., 2008; Shio et al., 2009). A Syk inhibitor markedly suppressed the production of PGE₂ in both silica- and Leu-Leu-OME-treated macrophages (Figure 6G). In contrast, Syk inhibition partially suppressed ATP-induced PGE₂ production, suggesting that the signaling pathway involved in PGE₂ production is different between particulate- and ATP-activated macrophages. Syk was partially associated with IL-1 β production in silica-stimulated macrophages (Figure 6G). In addition, knockdown of Syk by siRNA in macrophages significantly reduced PGE₂ and IL-1 β production compared to cells transfected with control siRNA (Figures S5B and S5C). Syk might act upstream of p38 MAP kinase because Syk inhibition suppressed phosphorylation of p38 MAP kinase in silica-stimulated macrophages (Figure 6H). Taken together, these results suggest that lysosomal damage triggers Syk activation, and then activated Syk upregulates cPLA₂ activity via the phosphorylation of p38 MAP kinase.

DISCUSSION

The NALP3 inflammasome has been reported to be activated by alum and involved in alum adjuvanticity and IgE production (Eisenbarth et al., 2008; Kool et al., 2008; Li et al., 2008). However, whether the NALP3 inflammasome is required for alum adjuvanticity is controversial. Here, we found that silica, alum, and ATP, which normally activate the NALP3 inflammasome, stimulate macrophages to produce PGE₂ through mechanisms that do not involve the NALP3 inflammasome. We also found that PGE₂ production by macrophages regulates the generation of antigen-specific IgE antibody in vivo.

The cells of the innate immune system can sense cellular danger and stress via the NALP3 inflammasome. Particulates, such as silica and alum, function as danger signals to activate the NALP3 inflammasome. We found that lysosomal damage and rupture and the subsequent leakage of lysosomal enzymes into the cytoplasm promoted the production of PGE₂ by macrophages. These results demonstrate that the danger signal caused by particulate-induced phagosomal destabilization activates the NALP3 inflammasome and induces macrophages to produce PGE₂. However, *Nalp3*^{-/-}, *Asc*^{-/-}, and *Casp1*^{-/-} macrophages produced PGE₂ in amounts similar to WT macrophages, suggesting that the lysosomal damage caused by silica and alum activates at least two different pathways, the NALP3 inflammasome pathway and the PGE₂-inducing pathway. We also examined the pathway involved in PGE₂ induction and found that lysosomal damage triggered the production of PGE₂ via the




Figure 6. Silica-Induced PGE $_2$ Production in Macrophages Is Regulated by the Syk/p38 MAP Kinase Pathway

(A and B) LPS-primed BALB/c peritoneal macrophages were stimulated with 100 μ g/ml silica (A) or 1 mM ATP (B) for 2 hr in the presence or absence of signaling inhibitors as described in the [Experimental Procedures](#). The results are expressed as the percentage (%) of IL-1 β and PGE $_2$ produced, and the amounts of IL-1 β and PGE $_2$ produced by macrophages stimulated with silica or ATP in the absence of inhibitors were used as the 100% controls.

(C) Macrophages were stimulated with 100 μ g/ml silica or 1 mM ATP for the indicated time. Cell lysates were subjected to western blot analysis with anti-phospho-p38 MAP kinase and total p38 MAP kinase antibodies.

(D) LPS-primed macrophages were stimulated with silica or ATP for 2 hr in the presence or absence of 10 μ M cPLA $_2$ inhibitor (cPLA $_2$ i).

activation of Syk and the p38 MAP kinase. Syk is known to play an important role in adaptive immune receptor signaling (Mócsai et al., 2010), and it is involved in malarial hemozoin-mediated and monosodium urate crystal-mediated inflammasome activation (Ng et al., 2008; Shio et al., 2009). Monosodium urate crystals have been reported to induce prostaglandin synthesis in phagocytic cells (Gordon et al., 1985), and we also found that monosodium urate crystals stimulates macrophages to produce PGE $_2$ and IL-1 β in our experimental system (data not shown). In addition, cPLA $_2$ activation and prostaglandin production are regulated by the activation of Syk (Suram et al., 2006). We also found that curdlan, which activates Dectin-1 and Syk, stimulated macrophages to produce higher amounts of PGE $_2$ in a Syk- and p38 MAP kinase-dependent manner (data not shown). Thus, Syk activation triggered by lysosomal damage promotes the production of PGE $_2$ in macrophages.

We demonstrated that the p38 MAP kinase pathway is critical for PGE $_2$ (but not IL-1 β) production in silica- and ATP-stimulated macrophages. Several reports have shown that p38 MAP kinase activates cPLA $_2$ and induces arachidonic acid release and PGE $_2$ secretion in a similar manner as Syk (Gijón et al., 2000; Umann et al., 2010). In addition, Syk has been reported to be required for p38 MAP kinase activation under the stress conditions (He et al., 2002). We showed that Syk inhibition suppressed phosphorylation of the p38 MAP kinase. These reports and our findings suggest that Syk activates cPLA $_2$ and PGE $_2$ production via the p38 MAP kinase pathway in particulate-activated macrophages.

We also found that ATP stimulated macrophages to induce PGE $_2$ production. However, the required signaling pathway might be different between particulate- and ATP-activated macrophages. Inhibition of the p38 MAP kinase and cPLA $_2$ significantly suppressed PGE $_2$ production in both silica- and ATP-activated macrophages. In contrast, Syk inhibition partially suppressed ATP-induced PGE $_2$ production. We have shown that lysosomal damage triggers Syk activation, and ATP has been reported to not be involved in lysosome damage (Hornung et al., 2008). However, ATP stimulation regulates Syk in osteoclasts (Hazama et al., 2009). Hazama et al. and our findings suggest that ATP-activated Syk is partially involved in PGE $_2$ production in macrophages independent of lysosomal damage. The detailed mechanism involved in lysosomal damage-triggered activation of Syk and p38 MAP kinase remains unclear. In addition, the involvement of cathepsin B in PGE $_2$ production should be clarified. As such, we are currently investigating which

(E) LPS-primed macrophages were incubated with 1 mM Leu-Leu-OMe for 2 hr in the presence or absence of p38 MAP kinase inhibitor.

(F) Macrophages were incubated with or without 20 μ g/ml PVNO for 5 hr. Then cells were primed with LPS. Primed macrophages were stimulated with silica or ATP for 2 hr.

(G) LPS-primed macrophages were stimulated with 100 μ g/ml silica, 1 mM ATP, or 1 mM Leu-Leu-OMe for 2 hr in the presence or absence of 1 μ M Syk inhibitor (Syki).

(H) Macrophages were stimulated with 100 μ g/ml silica for 30 min in the presence or absence of 1 μ M Syk inhibitor. Cell lysates from stimulated macrophages were subjected to western blot analysis. Data represent mean \pm SE of two (C and H) or three (A, B, and D–G) independent experiments (* p < 0.05, ** p < 0.01).

intracellular factor(s) or event(s) are associated with activation of Syk and p38 MAP kinase in particulate-activated macrophages.

We have herein shown that *Nalp3*^{-/-} and *Casp1*^{-/-} mice produce similar amounts of OVA-IgE to WT mice. These results suggest that NALP3 inflammasome-dependent cytokines, such as IL-1 β and IL-18, are not required for particulate-induced IgE responses. Our data in this study demonstrate that inflammasome activation and PGE₂ production are not required for the induction of antigen-specific IgG1 and IgG2c production. We found that treatment of macrophages with PVNO suppressed the production of PGE₂; however, a previous study has reported that treatment with PVNO prevent the adjuvant effect of silica on both IgE and IgG1 antibody production in vivo (Mancino et al., 1983). Taken together, these reports and our findings suggest that factor(s) induced by phagosomal destabilization other than IL-1 β , IL-18, and PGE₂ are also associated with alum and silica adjuvant activity. Furthermore, these results also suggest that the properties of individual particulates are very important for the control of acquired immune responses when particulates are used as adjuvants.

The effect of PGE₂ on Th1 and Th2 cell responses is a complex issue. Previous studies have reported that PGE₂ induces type 2 immunity by suppressing the production of cytokines from Th1 cells, macrophages, and DCs (Koga et al., 2009; Kuroda and Yamashita, 2003; Fabricius et al., 2010). However, PGE₂ facilitates the differentiation of Th1 cells in the presence of IL-12 and high doses of CD28 antibody through the activation of the PI3-kinase pathway (Yao et al., 2009). Why does PGE₂ function as an activator of Th1 and Th2 cells? We speculate that the cytokine milieu, containing IL-4 and IL-12, among many other cytokines, influences the effects of PGE₂ on the immune systems. PGE₂ facilitates Th1 cell responses in the presence of IL-12 (Yao et al., 2009). In contrast, herein we show that silica and alum induce macrophages to produce only caspase-1-dependent cytokines and PGE₂, but not other inflammatory cytokines, including IL-12. Similar to the findings of others (Fedyk and Phipps, 1996; Roper et al., 1995), our findings also showed that PGE₂ cooperates with IL-4 to promote IgE production in spleen cells in vitro. Our results suggest that PGE₂ facilitates the generation of either Th1 or Th2 cell responses, depending on the balance of IL-12 and IL-4 amounts.

Our in vivo and in vitro experiments suggest that PGE₂ functions as an activator of IgE production in B cells. Consistent with these results, previous reports have shown that PGE₂ facilitates IgE production in LPS plus IL-4-stimulated B cells (Fedyk and Phipps, 1996; Roper et al., 1995). However, contrary to our findings, Garrone et al. showed that PGE₂ suppresses IgE production in anti-CD40 plus IL-4-stimulated B cells (Garrone et al., 1994). This discrepancy between their results and ours might be due to the different types of cells used. We carried out our experiments with unsorted spleen cells, but Garrone et al. used purified B cells (Garrone et al., 1994). Our findings suggest that non-B cells affected by PGE₂ might stimulate B cells to induce IgE.

In conclusion, we have found that silica and alum stimulate macrophages to produce PGE₂ through a pathway that is dependent on Syk and p38 MAP kinase. PGE₂ generated by this mechanism regulates type 2 immune responses in vivo. Our results suggest that manipulating particulate-induced cyto-

kines and PGE₂ production could open new possibilities for the treatment of allergic inflammation, infectious diseases, and cancer.

EXPERIMENTAL PROCEDURES

Animals

Female C57BL/6 and BALB/c mice were purchased from Charles River Laboratories Japan (Yokohama, Japan). *Ptges*^{-/-} mice were generated by S. Uematsu and S. Akira at Osaka University (Uematsu et al., 2002) and were backcrossed to the C57BL/6 background for five generations; their WT littermates were used as controls. *Ptgs*^{-/-} mice (BALB/c background) were established by Y. Urade at the Osaka Bioscience Institute (Mohri et al., 2006). *Asc*^{-/-}, *Nalp3*^{-/-} and *Casp1*^{-/-} mice on C57BL/6 background were described previously (Mariathasan et al., 2004, 2006). *IL1r1*^{-/-} mice were obtained from Jackson Laboratories. All the animal experiments were carried out in accordance to the guidelines for the care and use of animals approved by the University of Occupational and Environmental Health.

Reagents

We used three different types of alum compounds in this study. LSL alum was purchased from Cosmo Bio Co. Ltd. (Tokyo, Japan); alhydrogel was purchased from Sigma Aldrich (St. Louis, MO); and Imject alum was purchased from Pierce (Rockford, IL). All the cytokines except IL-18 were purchased from PeproTech (Rocky Hill, NJ). Recombinant mouse IL-18 and the mouse IL-18 ELISA kit were purchased from MBL (Nagoya, Japan). The cytokine and chemokine ELISA kits were purchased from PeproTech. The PGE₂ and PGD₂-MOX EIA kits were purchased from Cayman Chemical (Ann Arbor, MI). The silica crystals (Min-u-sil 5 silica) were purchased from U.S. Silica (Berkeley Springs, WV). NiO was purchased from Vacuum Metallurgical (Chiba, Japan). TiO₂ and ATP were purchased from Wako Chemical (Osaka, Japan). The PI3 kinase inhibitor wortmannin was purchased from Sigma Aldrich. The caspase-1 inhibitor I (YVAD-CHO), cathepsin B inhibitor (CA-074 Me), p38 MAP kinase inhibitor (SB203580), cyclosporine A, MEK1/2 (Erk) inhibitor (U0126), rapamycin, JNK inhibitor (SP600125), NF- κ B inhibitor (wedelolactone), and Syk inhibitor were purchased from Calbiochem (Merck; Darmstadt, Germany). Cytochalasin D was purchased from Enzo Life Science (Plymouth Meeting, PA). Cytosolic PLA₂ inhibitor (Arachidonyl Trifluoromethyl Ketone) was purchased from Cayman Chemical. Leu-Leu-OMe was purchased from Chem-Impex International (Wood Dale, IL). Poly-2-vinylpyridine N-oxide (PVNO) was purchased from Polysciences (Warrington, PA). The following antibodies were used for western blot analysis: anti-GAPDH (Fitzgerald Industries International, Concord, MA); anti-COX-2 and anti-PTGES (mPGES-1) (Cayman Chemical); and anti-p38 MAP kinase and anti-phospho-p38 MAP kinase (Thr180/Tyr182) (Cell Signaling, Danvers, MA). The cells were cultured in RPMI 1640 medium (Nissui Pharmaceutical, Tokyo, Japan) supplemented with 10% FBS (BioWhittaker, Walkersville, MD, USA), 2 mM glutamine, 50 U/ml penicillin, and 50 μ g/ml streptomycin (all from Life Technologies, Rockville, MD, USA).

Cell Preparation

Peritoneal macrophages were generated by injecting mice i.p. with 2 ml 4% thioglycolate broth (Eiken, Tokyo, Japan) and harvested as described previously (Kuroda and Yamashita, 2003). BM-derived macrophages were prepared by culturing BM cells in the presence of 10 ng/ml of M-CSF or GM-CSF as described previously (Kuroda et al., 2007; Kuroda et al., 2009). The purity of the BM-derived macrophages was more than 95% F4/80 and Mac-1 positive as determined by flow cytometric analysis. BM-derived DCs were prepared by culturing BM cells in the presence of 10 ng/ml of GM-CSF for 5 days. CD11c⁺ cells were then enriched from the culture with the MACS (Miltenyi Biotec, Bergisch Gladbach, Germany). The enriched CD11c⁺ BM-derived DCs were 80% CD11c pure, as determined by flow cytometry. Human PBMC were purified by density gradient using Lymphoprep (Axis-Shield Poc AS, Oslo, Norway).

In Vitro Stimulation

In all the experiments, the cells were cultured at a density of 5 \times 10⁵/ml/well in 24-well plates (Falcom 3047; BD Biosciences, Franklin Lake, NJ), unless

otherwise specified. Macrophages and BM-derived DCs were primed with or without 1 ng/ml LPS (low dose) for 3 hr and then stimulated with 50 or 100 $\mu\text{g/ml}$ silica, 200 or 400 $\mu\text{g/ml}$ alum, 1 mM ATP, 50 or 100 $\mu\text{g/ml}$ TiO_2 , or 50 or 100 $\mu\text{g/ml}$ NiO for 2, 6, or 18 hr. In other experiments, 10 ng/ml of IL-1 or IL-18 or 1 or 2 mM of Leu-Leu-OMe was used instead of silica or alum. In some experiments, inhibitors were added to LPS-primed macrophages together with silica, alum, or ATP. The working concentrations of the inhibitors used were as follows: 10 μM caspase-1 inhibitor I, 10 μM CA-074 Me, 2 μM cytochalasin D, 1 μM NS-398, 10 μM SB203580, 100 nM wortmannin, 100 nM cyclosporin A, 10 μM U0126, 100 nM rapamycin, 10 μM SP600125, 20 μM wedelolactone, 10 μM cPLA₂ inhibitor, and 1 μM Syk inhibitor. After stimulation, cell-free supernatants were collected and used for ELISA. In some experiments, macrophages stimulated with 1 $\mu\text{g/ml}$ LPS (high dose) for 6 hr were used as controls. For the PVNO treatment, macrophages were incubated with or without PVNO for 5 hr. Cells were washed and then primed with low-dose LPS. LPS-primed macrophages were stimulated with silica, alum, or ATP for 30 and 60 min. The cells were then lysed with RIPA lysis buffer and used for western blot analysis as described previously (Kuroda et al., 2009). Human PBMCs were cultured at a density of $1 \times 10^6/\text{ml}$ in 24-well plates and primed with 0.1 ng/ml LPS for 3 hr. Primed PBMC were then stimulated with 50 $\mu\text{g/ml}$ silica for 3 hr.

Immunization

The mice were immunized twice (day 0 and 7) i.p. with 100 μg OVA plus 2 mg alum, 100 μg OVA plus 0.5 mg silica or 100 μg OVA plus 0.5 mg NiO in 200 μl of PBS. Ten days after the last immunization, sera were collected and analyzed for the OVA-specific IgE, IgG1, and IgG2c antibodies by ELISA.

ELISA

The amounts of cytokines, chemokines, and PGs were measured with either cytokine/chemokine ELISA kits or PGE₂/PGD₂ EIA kit in accordance with the manufacturer's instructions. The amounts of OVA-IgE in the sera were determined with the DS mouse IgE ELISA (OVA) kit (DS Pharma Biochemical, Osaka, Japan). For the analysis of OVA-IgG1 and IgG2c, serial dilutions of sera were prepared in 96-well plates coated with 10 $\mu\text{g/ml}$ OVA. HRP-conjugated goat anti-mouse IgG1 or IgG2c (Bethyl Laboratories, Montgomery, TX) were used as secondary antibodies. The reciprocal value of serum dilution being absorbance (OD 405 nm) at 0.5 was defined as the titer of antigen-specific serum IgG1 and IgG2c.

Statistical analysis

All the experiments were repeated between two to five times and representative results are shown. The statistical analyses were performed with the Student's *t* test. *p* values of < 0.05 were considered statistically significant and marked with an asterisk.

SUPPLEMENTAL INFORMATION

Supplemental Information includes five figures and Supplemental Experimental Procedures and can be found with this article online at doi:10.1016/j.immuni.2011.03.019.

ACKNOWLEDGMENTS

We thank G. Krystal (British Columbia Cancer Research Agency) for helpful discussion. This work was supported in part by a UOEH Research Grant for Promotion of Occupational Health (to E.K.) and a Grant-in-Aid for Scientific Research from the Ministry of Education, Science, Sports and Culture of Japan (21890297 and 22791094 to E.K.).

Received: June 10, 2010

Revised: March 14, 2011

Accepted: March 24, 2011

Published online: April 14, 2011

REFERENCES

- Aimanianda, V., Haensler, J., Lacroix-Desmazes, S., Kaveri, S.V., and Bayry, J. (2009). Novel cellular and molecular mechanisms of induction of immune responses by aluminum adjuvants. *Trends Pharmacol. Sci.* 30, 287–295.
- Akira, S., Uematsu, S., and Takeuchi, O. (2006). Pathogen recognition and innate immunity. *Cell* 124, 783–801.
- Allison, A.C., Harington, J.S., and Birbeck, M. (1966). An examination of the cytotoxic effects of silica on macrophages. *J. Exp. Med.* 124, 141–154.
- Boniface, K., Bak-Jensen, K.S., Li, Y., Blumenschein, W.M., McGeachy, M.J., McClanahan, T.K., McKenzie, B.S., Kastelein, R.A., Cua, D.J., and de Waal Malefyt, R. (2009). Prostaglandin E2 regulates Th17 cell differentiation and function through cyclic AMP and EP2/EP4 receptor signaling. *J. Exp. Med.* 206, 535–548.
- Cassel, S.L., Eisenbarth, S.C., Iyer, S.S., Sadler, J.J., Colegio, O.R., Tephly, L.A., Carter, A.B., Rothman, P.B., Flavell, R.A., and Sutterwala, F.S. (2008). The Nalp3 inflammasome is essential for the development of silicosis. *Proc. Natl. Acad. Sci. USA* 105, 9035–9040.
- Dinarello, C.A. (2002). The IL-1 family and inflammatory diseases. *Clin. Exp. Rheumatol.* 20 (5, Suppl 27), S1–S13.
- Dostert, C., Pétrilli, V., Van Bruggen, R., Steele, C., Mossman, B.T., and Tschopp, J. (2008). Innate immune activation through Nalp3 inflammasome sensing of asbestos and silica. *Science* 320, 674–677.
- Duewell, P., Kono, H., Rayner, K.J., Sirois, C.M., Vladimer, G., Bauernfeind, F.G., Abela, G.S., Franchi, L., Nuñez, G., Schnurr, M., et al. (2010). NLRP3 inflammasomes are required for atherogenesis and activated by cholesterol crystals. *Nature* 464, 1357–1361.
- Duncan, J.A., Gao, X., Huang, M.T., O'Connor, B.P., Thomas, C.E., Willingham, S.B., Bergstralh, D.T., Jarvis, G.A., Sparling, P.F., and Ting, J.P. (2009). *Neisseria gonorrhoeae* activates the proteinase cathepsin B to mediate the signaling activities of the NLRP3 and ASC-containing inflammasome. *J. Immunol.* 182, 6460–6469.
- Eisenbarth, S.C., Colegio, O.R., O'Connor, W., Sutterwala, F.S., and Flavell, R.A. (2008). Crucial role for the Nalp3 inflammasome in the immunostimulatory properties of aluminium adjuvants. *Nature* 453, 1122–1126.
- Fabricius, D., Neubauer, M., Mandel, B., Schütz, C., Viardot, A., Vollmer, A., Jahrsdörfer, B., and Debatin, K.M. (2010). Prostaglandin E₂ inhibits IFN- α secretion and Th1 costimulation by human plasmacytoid dendritic cells via E-prostanoid 2 and E-prostanoid 4 receptor engagement. *J. Immunol.* 184, 677–684.
- Fedyk, E.R., and Phipps, R.P. (1996). Prostaglandin E₂ receptors of the EP2 and EP4 subtypes regulate activation and differentiation of mouse B lymphocytes to IgE-secreting cells. *Proc. Natl. Acad. Sci. USA* 93, 10978–10983.
- Franchi, L., and Nuñez, G. (2008). The Nlrp3 inflammasome is critical for aluminium hydroxide-mediated IL-1 β secretion but dispensable for adjuvant activity. *Eur. J. Immunol.* 38, 2085–2089.
- Franchi, L., Eigenbrod, T., Muñoz-Planillo, R., and Nuñez, G. (2009). The inflammasome: A caspase-1-activation platform that regulates immune responses and disease pathogenesis. *Nat. Immunol.* 10, 241–247.
- Garrone, P., Galibert, L., Rousset, F., Fu, S.M., and Banchereau, J. (1994). Regulatory effects of prostaglandin E2 on the growth and differentiation of human B lymphocytes activated through their CD40 antigen. *J. Immunol.* 152, 4282–4290.
- Ghiringhelli, F., Apetoh, L., Tesniere, A., Aymeric, L., Ma, Y., Ortiz, C., Vermaelen, K., Panaretakis, T., Mignot, G., Ullrich, E., et al. (2009). Activation of the NLRP3 inflammasome in dendritic cells induces IL-1 β -dependent adaptive immunity against tumors. *Nat. Med.* 15, 1170–1178.
- Gijón, M.A., Spencer, D.M., Siddiqi, A.R., Bonventre, J.V., and Leslie, C.C. (2000). Cytosolic phospholipase A₂ is required for macrophage arachidonic acid release by agonists that do and do not mobilize calcium. *J. Biol. Chem.* 275, 20146–20156.
- Gordon, T.P., Kowanko, I.C., James, M., and Roberts-Thomson, P.J. (1985). Monosodium urate crystal-induced prostaglandin synthesis in the rat subcutaneous air pouch. *Clin. Exp. Rheumatol.* 3, 291–296.

- Gross, O., Poeck, H., Bscheider, M., Dostert, C., Hanneschläger, N., Endres, S., Hartmann, G., Tardivel, A., Schweighoffer, E., Tybulewicz, V., et al. (2009). Syk kinase signalling couples to the Nlrp3 inflammasome for anti-fungal host defence. *Nature* 459, 433–436.
- Halle, A., Hornung, V., Petzold, G.C., Stewart, C.R., Monks, B.G., Reinheckel, T., Fitzgerald, K.A., Latz, E., Moore, K.J., and Golenbock, D.T. (2008). The NALP3 inflammasome is involved in the innate immune response to amyloid- β . *Nat. Immunol.* 9, 857–865.
- Hazama, R., Qu, X., Yokoyama, K., Tanaka, C., Kinoshita, E., He, J., Takahashi, S., Tohyama, K., Yamamura, H., and Tohyama, Y. (2009). ATP-induced osteoclast function: The formation of sealing-zone like structure and the secretion of lytic granules via microtubule-deacetylation under the control of Syk. *Genes Cells* 14, 871–884.
- He, J., Takano, T., Ding, J., Gao, S., Noda, C., Sada, K., Yanagi, S., and Yamamura, H. (2002). Syk is required for p38 activation and G2/M arrest in B cells exposed to oxidative stress. *Antioxid. Redox Signal.* 4, 509–515.
- Hornung, V., Bauernfeind, F., Halle, A., Samstad, E.O., Kono, H., Rock, K.L., Fitzgerald, K.A., and Latz, E. (2008). Silica crystals and aluminum salts activate the NALP3 inflammasome through phagosomal destabilization. *Nat. Immunol.* 9, 847–856.
- Koga, K., Takaesu, G., Yoshida, R., Nakaya, M., Kobayashi, T., Kinjo, I., and Yoshimura, A. (2009). Cyclic adenosine monophosphate suppresses the transcription of proinflammatory cytokines via the phosphorylated c-Fos protein. *Immunity* 30, 372–383.
- Kool, M., Pétrilli, V., De Smedt, T., Rolaz, A., Hammad, H., van Nimwegen, M., Bergen, I.M., Castillo, R., Lambrecht, B.N., and Tschopp, J. (2008). Cutting edge: Alum adjuvant stimulates inflammatory dendritic cells through activation of the NALP3 inflammasome. *J. Immunol.* 181, 3755–3759.
- Kumar, H., Kumagai, Y., Tsuchida, T., Koenig, P.A., Satoh, T., Guo, Z., Jang, M.H., Saitoh, T., Akira, S., and Kawai, T. (2009). Involvement of the NLRP3 inflammasome in innate and humoral adaptive immune responses to fungal β -glucan. *J. Immunol.* 183, 8061–8067.
- Kuroda, E., and Yamashita, U. (2003). Mechanisms of enhanced macrophage-mediated prostaglandin E_2 production and its suppressive role in Th1 activation in Th2-dominant BALB/c mice. *J. Immunol.* 170, 757–764.
- Kuroda, E., Noguchi, J., Doi, T., Uematsu, S., Akira, S., and Yamashita, U. (2007). IL-3 is an important differentiation factor for the development of prostaglandin E_2 -producing macrophages between C57BL/6 and BALB/c mice. *Eur. J. Immunol.* 37, 2185–2195.
- Kuroda, E., Ho, V., Ruschmann, J., Antignano, F., Hamilton, M., Rauh, M.J., Antov, A., Flavell, R.A., Sly, L.M., and Krystal, G. (2009). SHIP represses the generation of IL-3-induced M2 macrophages by inhibiting IL-4 production from basophils. *J. Immunol.* 183, 3652–3660.
- Lee, J.K., Kim, S.H., Lewis, E.C., Azam, T., Reznikov, L.L., and Dinarello, C.A. (2004). Differences in signaling pathways by IL-1 β and IL-18. *Proc. Natl. Acad. Sci. USA* 101, 8815–8820.
- Li, H., Willingham, S.B., Ting, J.P., and Re, F. (2008). Cutting edge: Inflammasome activation by alum and alum's adjuvant effect are mediated by NLRP3. *J. Immunol.* 181, 17–21.
- Mancino, D., Buono, G., Cusano, M., and Minucci, M. (1983). Adjuvant effects of a crystalline silica on IgE and IgG1 antibody production in mice and their prevention by the macrophage stabilizer poly-2-vinylpyridine N-oxide. *Int. Arch. Allergy Appl. Immunol.* 71, 279–281.
- Mariathasan, S., Newton, K., Monack, D.M., Vucic, D., French, D.M., Lee, W.P., Roose-Girma, M., Erickson, S., and Dixit, V.M. (2004). Differential activation of the inflammasome by caspase-1 adaptors ASC and Ipaf. *Nature* 430, 213–218.
- Mariathasan, S., Weiss, D.S., Newton, K., McBride, J., O'Rourke, K., Roose-Girma, M., Lee, W.P., Weinrauch, Y., Monack, D.M., and Dixit, V.M. (2006). Cryopyrin activates the inflammasome in response to toxins and ATP. *Nature* 440, 228–232.
- Marrack, P., McKee, A.S., and Munks, M.W. (2009). Towards an understanding of the adjuvant action of aluminium. *Nat. Rev. Immunol.* 9, 287–293.
- Martinon, F., Mayor, A., and Tschopp, J. (2009). The inflammasomes: Guardians of the body. *Annu. Rev. Immunol.* 27, 229–265.
- McKee, A.S., Munks, M.W., MacLeod, M.K., Fleenor, C.J., Van Rooijen, N., Kappler, J.W., and Marrack, P. (2009). Alum induces innate immune responses through macrophage and mast cell sensors, but these sensors are not required for alum to act as an adjuvant for specific immunity. *J. Immunol.* 183, 4403–4414.
- Mócsai, A., Ruland, J., and Tybulewicz, V.L. (2010). The SYK tyrosine kinase: A crucial player in diverse biological functions. *Nat. Rev. Immunol.* 10, 387–402.
- Mohri, I., Eguchi, N., Suzuki, K., Urade, Y., and Taniike, M. (2003). Hematopoietic prostaglandin D synthase is expressed in microglia in the developing postnatal mouse brain. *Glia* 42, 263–274.
- Mohri, I., Taniike, M., Taniguchi, H., Kanekiyo, T., Aritake, K., Inui, T., Fukumoto, N., Eguchi, N., Kushi, A., Sasai, H., et al. (2006). Prostaglandin D_2 -mediated microglia/astrocyte interaction enhances astrogliosis and demyelination in twitcher. *J. Neurosci.* 26, 4383–4393.
- Narumiya, S. (2009). Prostanoids and inflammation: A new concept arising from receptor knockout mice. *J. Mol. Med.* 87, 1015–1022.
- Ng, G., Sharma, K., Ward, S.M., Desrosiers, M.D., Stephens, L.A., Schoel, W.M., Li, T., Lowell, C.A., Ling, C.C., Amrein, M.W., and Shi, Y. (2008). Receptor-independent, direct membrane binding leads to cell-surface lipid sorting and Syk kinase activation in dendritic cells. *Immunity* 29, 807–818.
- Nishi, K., Morimoto, Y., Ogami, A., Murakami, M., Myojo, T., Oyabu, T., Kadoya, C., Yamamoto, M., Todoroki, M., Hirohashi, M., et al. (2009). Expression of cytokine-induced neutrophil chemoattractant in rat lungs by intratracheal instillation of nickel oxide nanoparticles. *Inhal. Toxicol.* 21, 1030–1039.
- Ogami, A., Morimoto, Y., Myojo, T., Oyabu, T., Murakami, M., Todoroki, M., Nishi, K., Kadoya, C., Yamamoto, M., and Tanaka, I. (2009). Pathological features of different sizes of nickel oxide following intratracheal instillation in rats. *Inhal. Toxicol.* 21, 812–818.
- Roper, R.L., Brown, D.M., and Phipps, R.P. (1995). Prostaglandin E_2 promotes B lymphocyte Ig isotype switching to IgE. *J. Immunol.* 154, 162–170.
- Schroder, K., and Tschopp, J. (2010). The inflammasomes. *Cell* 140, 821–832.
- Schroder, K., Zhou, R., and Tschopp, J. (2010). The NLRP3 inflammasome: A sensor for metabolic danger? *Science* 327, 296–300.
- Sharp, F.A., Ruane, D., Claass, B., Creagh, E., Harris, J., Malyala, P., Singh, M., O'Hagan, D.T., Pétrilli, V., Tschopp, J., et al. (2009). Uptake of particulate vaccine adjuvants by dendritic cells activates the NALP3 inflammasome. *Proc. Natl. Acad. Sci. USA* 106, 870–875.
- Shio, M.T., Eisenbarth, S.C., Savaria, M., Vinet, A.F., Bellemare, M.J., Harder, K.W., Sutterwala, F.S., Bohle, D.S., Descoteaux, A., Flavell, R.A., and Olivier, M. (2009). Malarial hemozoin activates the NLRP3 inflammasome through Lyn and Syk kinases. *PLoS Pathog.* 5, e1000559.
- Suram, S., Brown, G.D., Ghosh, M., Gordon, S., Loper, R., Taylor, P.R., Akira, S., Uematsu, S., Williams, D.L., and Leslie, C.C. (2006). Regulation of cytosolic phospholipase A2 activation and cyclooxygenase 2 expression in macrophages by the β -glucan receptor. *J. Biol. Chem.* 281, 5506–5514.
- Uematsu, S., Matsumoto, M., Takeda, K., and Akira, S. (2002). Lipopolysaccharide-dependent prostaglandin E_2 production is regulated by the glutathione-dependent prostaglandin E_2 synthase gene induced by the Toll-like receptor 4/MyD88/NF- κ B pathway. *J. Immunol.* 168, 5811–5816.
- Ulmann, L., Hirbec, H., and Rassendren, F. (2010). P2X4 receptors mediate PGE2 release by tissue-resident macrophages and initiate inflammatory pain. *EMBO J.* 29, 2290–2300.
- Von Behren, L.A., Chaudhary, S., Rabinovich, S., Shu, M.D., and Tewari, R.P. (1983). Protective effect of poly-2-vinylpyridine-N-oxide on susceptibility of silica-treated mice to experimental histoplasmosis. *Infect. Immun.* 42, 818–823.
- Watanabe, H., Gehrke, S., Contassot, E., Roques, S., Tschopp, J., Friedmann, P.S., French, L.E., and Gaide, O. (2008). Danger signaling through the

inflammasome acts as a master switch between tolerance and sensitization. *J. Immunol.* *180*, 5826–5832.

Yamamoto, M., Yaginuma, K., Tsutsui, H., Sagara, J., Guan, X., Seki, E., Yasuda, K., Yamamoto, M., Akira, S., Nakanishi, K., et al. (2004). ASC is essential for LPS-induced activation of procaspase-1 independently of TLR-associated signal adaptor molecules. *Genes Cells* *9*, 1055–1067.

Yao, C., Sakata, D., Esaki, Y., Li, Y., Matsuoka, T., Kuroiwa, K., Sugimoto, Y., and Narumiya, S. (2009). Prostaglandin E₂-EP4 signaling promotes immune inflammation through Th1 cell differentiation and Th17 cell expansion. *Nat. Med.* *15*, 633–640.

Ye, Z., and Ting, J.P. (2008). NLR, the nucleotide-binding domain leucine-rich repeat containing gene family. *Curr. Opin. Immunol.* *20*, 3–9.