

Contents lists available at ScienceDirect

Physics Letters B

www.elsevier.com/locate/physletb

New mechanisms for double charmed meson production at the LHCb

Rafał Maciuła^a, Vladimir A. Saleev^b, Alexandra V. Shipilova^b, Antoni Szczurek^{a,c,*}^a Institute of Nuclear Physics, Polish Academy of Sciences, Radzikowskiego 152, PL-31-342 Kraków, Poland^b Samara State Aerospace University, Moscow Highway, 34, 443086, Samara, Russia^c University of Rzeszów, PL-35-959 Rzeszów, Poland

ARTICLE INFO

Article history:

Received 8 February 2016

Received in revised form 22 April 2016

Accepted 17 May 2016

Available online 19 May 2016

Editor: A. Ringwald

ABSTRACT

We discuss production of $D^0 D^0$ (and $\bar{D}^0 \bar{D}^0$) pairs related to the LHCb Collaboration results for $\sqrt{s} = 7$ TeV in proton–proton scattering. We consider double-parton scattering (DPS) mechanisms of double $c\bar{c}$ production and subsequent $cc \rightarrow D^0 D^0$ hadronization as well as double g and mixed $g\bar{c}$ production with $gg \rightarrow D^0 D^0$ and $gc \rightarrow D^0 D^0$ hadronization calculated with the help of the scale-dependent hadronization functions of Kniehl et al. Single-parton scattering (SPS) mechanism of digluon production is also taken into account. We compare our results with several correlation observables in azimuthal angle $\varphi_{D^0 D^0}$ between D^0 mesons or in dimeson invariant mass $M_{D^0 D^0}$. The inclusion of new mechanisms with $g \rightarrow D^0$ fragmentation leads to larger cross sections, than when including only DPS mechanism $cc \rightarrow D^0 D^0$ with standard scale-independent fragmentation functions. Some consequences of the presence of the new mechanisms are discussed. In particular a larger σ_{eff} is needed to describe the LHCb data. There is a signature that σ_{eff} may depend on transverse momentum of c quarks and/or \bar{c} antiquarks.

© 2016 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP³.

1. Introduction

Some time ago two of us predicted that at large energies relevant for the LHC the production of double charm should be dominated by the double-parton scattering (DPS) mechanism [1]. In the first calculation the cross section for each step was calculated in the leading-order (LO) collinear approach. However, the LO collinear approach is not sufficient for a detailed description of actual cross section for the $c\bar{c}$ production. The double $c\bar{c}$ production was extended next to the k_t -factorization approach which includes effectively higher-order QCD effects [2,3]. A relatively good description of the LHCb experimental data [4] was achieved for both the total yield and the dimeson correlation observables. In these calculations the standard scale-independent Peterson fragmentation function (FF) [5] was used. The single-parton scattering (SPS) $gg \rightarrow c\bar{c}c\bar{c}$ contribution was discussed carefully in both collinear [3] and k_t -factorization [6] approaches. Their contribution to the $c\bar{c}c\bar{c}$ cross section was found to be rather small and was not able to describe details of the LHCb data [4].

Studies of inclusive D meson production at the LHC based on scale-independent FFs have been done in next-to-leading order (NLO) collinear approach within the FONLL scheme [7] as well as in the k_t -factorization [8]. In turn, in Ref. [9] the calculation was done according to the GM-VFNS NLO collinear scheme together with the several scale-dependent FFs of a parton (gluon, $u, d, s, \bar{u}, \bar{d}, \bar{s}, c, \bar{c}$) to D mesons proposed by Kniehl et al. [10,11], that undergo DGLAP evolution equations. It has been found that important contribution to inclusive production of D mesons comes from gluon fragmentation (see also Ref. [12]). Similar calculation were done recently also in the k_t -factorization approach with parton Reggeization hypothesis by two of us [13]. They have also shown there that the new mechanism constitutes a big fraction of the cross section for D meson production and a good description of the inclusive D meson production at the LHC was achieved.

In the present paper we wish to investigate how important is the gluon fragmentation mechanism for the double D -meson production, i.e. double fragmentation of each of the gluons in the gluon dijets in SPS production and double fragmentation of each of the gluons in the gluon jet in DPS production mechanism. Here the gluon and digluon production is considered in the k_t -factorization approach with Reggeized gluons in the t -channel [14] via subprocesses $RR \rightarrow g$ and $RR \rightarrow gg$, where R is the Reggeized gluon. In our analysis we shall use scale-dependent fragmentation functions

* Corresponding author.

E-mail addresses: rafal.maciula@ifj.edu.pl (R. Maciuła), saleev@samsu.ru (V.A. Saleev), alexshipilova@samsu.ru (A.V. Shipilova), antoni.szczurek@ifj.edu.pl (A. Szczurek).




Fig. 1. A diagrammatic illustration of the considered mechanisms.

of Kneesch–Kniehl–Kramer–Schienbein (KKKS08) [15] as implemented in the code available on the Web [16].

2. A sketch of the theoretical formalism

We will compare results with first (old) and second (new) approach. In the second (new) scenario with $g \rightarrow D$ fragmentation the number of contributing processes grows compared to the first (old) scenario with $c \rightarrow D$ fragmentation only. Naturally a new single-parton scattering mechanism (called here SPS $gg \rightarrow DD$) appears (top-right panel in Fig. 1). Since here the two produced gluons are correlated in azimuth, the mechanism will naturally lead to an azimuthal correlation between the DD (or $\bar{D}\bar{D}$) mesons. Such a correlation was actually observed in the LHCb experimental data [4] and could not be explained by the SPS $2 \rightarrow 4$ perturbative $gg \rightarrow c\bar{c}c\bar{c}$ contribution (see e.g. Ref. [6]) which turned out to be rather small. In the new scenario we have more processes for single D meson production (two components) and as a consequence many more processes for the pair production in double-parton scattering. Now (in the new scenario) there are three classes of DPS contributions. In addition to the conventional DPS $cc \rightarrow DD$ (top-left panel in Fig. 1) considered in Refs. [2,3,6] there is a double $g \rightarrow D$ (or double $g \rightarrow \bar{D}$) fragmentation mechanism, called here DPS $gg \rightarrow DD$ (bottom-left panel in Fig. 1) as well as the mixed DPS $gc \rightarrow DD$ contribution (bottom-right panel in Fig. 1).

DPS cross section for production of cc , gg or gc system, assuming factorization of the DPS model, can be written as:

$$\frac{d\sigma^{DPS}(pp \rightarrow ccX)}{dy_1 dy_2 d^2 p_{1,t} d^2 p_{2,t}} = \frac{1}{2\sigma_{eff}} \cdot \frac{d\sigma^{SPS}(pp \rightarrow c\bar{c}X_1)}{dy_1 d^2 p_{1,t}} \cdot \frac{d\sigma^{SPS}(pp \rightarrow c\bar{c}X_2)}{dy_2 d^2 p_{2,t}}, \quad (2.1)$$

$$\frac{d\sigma^{DPS}(pp \rightarrow ggX)}{dy_1 dy_2 d^2 p_{1,t} d^2 p_{2,t}} = \frac{1}{2\sigma_{eff}} \cdot \frac{d\sigma^{SPS}(pp \rightarrow gX_1)}{dy_1 d^2 p_{1,t}} \cdot \frac{d\sigma^{SPS}(pp \rightarrow gX_2)}{dy_2 d^2 p_{2,t}}, \quad (2.2)$$

$$\frac{d\sigma^{DPS}(pp \rightarrow gcX)}{dy_1 dy_2 d^2 p_{1,t} d^2 p_{2,t}} = \frac{1}{\sigma_{eff}} \cdot \frac{d\sigma^{SPS}(pp \rightarrow gX_1)}{dy_1 d^2 p_{1,t}} \cdot \frac{d\sigma^{SPS}(pp \rightarrow c\bar{c}X_2)}{dy_2 d^2 p_{2,t}}. \quad (2.3)$$

When integrating over kinematical variables one recovers the commonly used pocket formula:

$$\sigma^{DPS}(pp \rightarrow ccX) = \frac{1}{2\sigma_{eff}} \sigma^{SPS}(pp \rightarrow c\bar{c}X_1) \cdot \sigma^{SPS}(pp \rightarrow c\bar{c}X_2), \quad (2.4)$$

$$\sigma^{DPS}(pp \rightarrow ggX) = \frac{1}{2\sigma_{eff}} \sigma^{SPS}(pp \rightarrow gX_1) \cdot \sigma^{SPS}(pp \rightarrow gX_2), \quad (2.5)$$

$$\sigma^{DPS}(pp \rightarrow gcX) = \frac{1}{\sigma_{eff}} \sigma^{SPS}(pp \rightarrow gX_1) \cdot \sigma^{SPS}(pp \rightarrow c\bar{c}X_2). \quad (2.6)$$

The often called pocket-formula is a priori a severe approximation. The flavor, spin and color correlations lead, in principle, to interference effects that result in its violation as discussed e.g. in Refs. [17, 18]. Even for unpolarized proton beams, the spin polarization of the two partons from one hadron can be mutually correlated, especially when the partons are relatively close in phase space (having comparable x 's). Moreover, in contrast to the standard single PDFs, the two-parton distributions have a nontrivial color structure which also may lead to a non-negligible correlations effects. Such effects are usually not included in phenomenological analyses. They were exceptionally discussed in the context of double charm production [19]. However, the effect on e.g. azimuthal correlations between charmed quarks was found there to be very small, much smaller than effects of the SPS contribution associated with double gluon fragmentation discussed in the present paper. In addition, including perturbative parton splitting mechanism also leads to a breaking of the pocket-formula [20–22]. This formalism was so far formulated for the collinear leading-order approach which for charm (double charm) may be a bit academic as this leads to underestimation of the cross section. Imposing sum rules also leads to a breaking of the factorized Ansatz but the effect almost vanishes for small longitudinal momentum fractions [23]. Taken the above we will use the pocket-formula in the following.

In the k_t -factorization approach, the cross sections for SPS mechanisms can be presented as follows:

$$\frac{d\sigma^{SPS}(pp \rightarrow c\bar{c}X)}{dy_1 dy_2 d^2 p_{1,t} d^2 p_{2,t}} = \frac{1}{16\pi^2 (x_1 x_2 S)^2} \int \frac{d^2 k_{1t}}{\pi} \frac{d^2 k_{2t}}{\pi} \overline{|\mathcal{M}_{RR \rightarrow c\bar{c}}|^2} \times \delta^2(\vec{k}_{1t} + \vec{k}_{2t} - \vec{p}_{1t} - \vec{p}_{2t}) \mathcal{F}(x_1, k_{1t}^2, \mu^2) \times \mathcal{F}(x_2, k_{2t}^2, \mu^2), \quad (2.7)$$

$$\frac{d\sigma^{SPS}(pp \rightarrow ggX)}{dy_1 dy_2 d^2 p_{1,t} d^2 p_{2,t}} = \frac{1}{16\pi^2 (x_1 x_2 S)^2} \int \frac{d^2 k_{1t}}{\pi} \frac{d^2 k_{2t}}{\pi} \overline{|\mathcal{M}_{RR \rightarrow gg}|^2} \times \delta^2(\vec{k}_{1t} + \vec{k}_{2t} - \vec{p}_{1t} - \vec{p}_{2t}) \mathcal{F}(x_1, k_{1t}^2, \mu^2) \times \mathcal{F}(x_2, k_{2t}^2, \mu^2), \quad (2.8)$$

$$\frac{d\sigma^{SPS}(pp \rightarrow gX)}{dy d^2 p_t} = \frac{\pi}{(x_1 x_2 S)^2} \int \frac{d^2 k_{1t}}{\pi} \frac{d^2 k_{2t}}{\pi} \overline{|\mathcal{M}_{RR \rightarrow g}|^2} \times \delta^2(\vec{k}_{1t} + \vec{k}_{2t} - \vec{p}_t) \mathcal{F}(x_1, k_{1t}^2, \mu^2) \times \mathcal{F}(x_2, k_{2t}^2, \mu^2). \quad (2.9)$$

Here the four-momenta of the initial-state gluons are parameterized as a sum of longitudinal and transverse parts $k_{1,2} =$

$x_{1,2}P_{1,2} + k_{t1,2}$, $k_{t1,2} = (0, \vec{k}_{t1,2}, 0)$, $k_{t1,2}^2 = -\vec{k}_{t1,2}^2$, $P_{1,2}$ are the four-momenta of the protons, $2P_1 P_2 = S$, $|\overline{\mathcal{M}}_{RR \rightarrow g, gg, c\bar{c}}|^2$ are the partonic cross sections with Reggeized gluons in the initial state.

Fully gauge invariant treatment of the initial-state off-shell gluons can be achieved in k_t -factorization approach only when they are considered as Reggeized gluons or Reggeons. The relevant Reggeized amplitudes can be presented using Fadin–Kuraev–Lipatov effective vertices: $C_{RR}^{g,\mu}$, $C_{RR}^{gg,\mu\nu}$ and $C_{RR}^{q\bar{q}}$ [24]. The squared amplitude of the partonic subprocess $RR \rightarrow g$ is very simple and can be presented as

$$|\overline{\mathcal{M}}_{RR \rightarrow g}|^2 = \frac{3}{2} \pi \alpha_S \vec{p}_t^2. \quad (2.10)$$

The useful analytical formulae for $|\overline{\mathcal{M}}_{RR \rightarrow gg}|^2$ and $|\overline{\mathcal{M}}_{RR \rightarrow c\bar{c}}|^2$ squared amplitudes are more complicated and we use the ones as they have been written in Ref. [14].

In the approach used here, the gluon unintegrated parton distribution function (unPDF) $\mathcal{F}(x, k_t^2, \mu^2)$ is normalized with respect to the collinear parton distribution function (PDF) by the following condition

$$\int dk_t^2 \mathcal{F}(x, k_t^2, \mu^2) = x \mathcal{G}(x, \mu^2).$$

A few phenomenological schemes to compute unPDFs of a proton where proposed. In the present paper we use the LO Kimber–Martin–Ryskin (KMR) unPDFs [25], generated from the LO set of a up-to-date Martin–Motylinski–Harland–Lang–Thorne (MMHT2014) collinear PDFs [26] fitted also to the LHC data.

In the perturbative part of the calculations here we use a running LO α_S provided with the MMHT2014 PDFs. The charm quark mass used in the numerical calculations is $m_c = 1.5$ GeV. We set both the renormalization and factorization scales equal to $\mu^2 = p_t^2$ for $RR \rightarrow g$ subprocess, to the averaged transverse momentum $\mu^2 = (p_{1t}^2 + p_{2t}^2)/2$ for $RR \rightarrow gg$, and to the averaged transverse mass $\mu^2 = (m_{1t}^2 + m_{2t}^2)/2$ for $RR \rightarrow c\bar{c}$ case, where $m_t = \sqrt{p_t^2 + m_c^2}$.

In order to calculate correlation observables for two mesons we follow here, similar as in the single meson case, the fragmentation function technique for hadronization process:

$$\begin{aligned} \frac{d\sigma^{DPS}(pp \rightarrow DDX)}{dy_1 dy_2 d^2 p_{1t}^D d^2 p_{2t}^D} &= \int \frac{D_{c \rightarrow D}(z_1, \mu)}{z_1} \cdot \frac{D_{c \rightarrow D}(z_2, \mu)}{z_2} \\ &\cdot \frac{d\sigma^{DPS}(pp \rightarrow ccX)}{dy_1 dy_2 d^2 p_{1t}^c d^2 p_{2t}^c} dz_1 dz_2 \\ &+ \int \frac{D_{g \rightarrow D}(z_1, \mu)}{z_1} \cdot \frac{D_{g \rightarrow D}(z_2, \mu)}{z_2} \\ &\cdot \frac{d\sigma^{DPS}(pp \rightarrow ggX)}{dy_1 dy_2 d^2 p_{1t}^g d^2 p_{2t}^g} dz_1 dz_2 \\ &+ \int \frac{D_{g \rightarrow D}(z_1, \mu)}{z_1} \cdot \frac{D_{c \rightarrow D}(z_2, \mu)}{z_2} \\ &\cdot \frac{d\sigma^{DPS}(pp \rightarrow gcX)}{dy_1 dy_2 d^2 p_{1t}^g d^2 p_{2t}^c} dz_1 dz_2, \end{aligned} \quad (2.11)$$

where: $p_{1t}^{g,c} = \frac{p_{1t}^D}{z_1}$, $p_{2t}^{g,c} = \frac{p_{2t}^D}{z_2}$ and meson momentum fractions $z_1, z_2 \in (0, 1)$.

The same formula for SPS DD -production via digluon fragmentation reads

$$\begin{aligned} \frac{d\sigma_{gg}^{SPS}(pp \rightarrow DDX)}{dy_1 dy_2 d^2 p_{1t}^D d^2 p_{2t}^D} &\approx \int \frac{D_{g \rightarrow D}(z_1, \mu)}{z_1} \cdot \frac{D_{g \rightarrow D}(z_2, \mu)}{z_2} \\ &\cdot \frac{d\sigma^{SPS}(pp \rightarrow ggX)}{dy_1 dy_2 d^2 p_{1t}^g d^2 p_{2t}^g} dz_1 dz_2, \end{aligned} \quad (2.12)$$

where: $p_{1t}^g = \frac{p_{1t}^D}{z_1}$, $p_{2t}^g = \frac{p_{2t}^D}{z_2}$ and meson momentum fractions $z_1, z_2 \in (0, 1)$.

In the case of calculations with the scale-independent Peterson FF the parameter $\varepsilon_c = 0.5$ is taken, which is the averaged value extracted from different e^+e^- experiments and is commonly used in the literature. In turn, for predictions with the KKKS08 FF the evolution scale is set to the charm quark transverse mass $\mu^2 = m_t^2$ and gluon transverse momentum $\mu^2 = p_t^2$ for the $c \rightarrow D$ and $g \rightarrow D$ components, respectively.

In e^+e^- collisions it is assumed naturally that gluons do not fragment to D mesons for scales smaller than $\mu^2 = \hat{s} = 4m_c^2$. In our calculation we have also tried to take $\mu^2 = \hat{s}$ as the hadronization scale, which is an alternative to the typical choice of $\mu^2 = m_t^2$. In the first case naturally such initial scale is $4m_c^2$ while for the second case the initial scale is set to m_c^2 so the effect of the evolution of fragmentation functions is present all over the phase space, however is very small for small transverse momenta of gluons. In summary, both the choices of the hadronization scale lead to fairly similar final results for DD correlations, which we have checked numerically.

3. Comparison to the LHCb data

We start the presentation of our new results with a revision of inclusive single D^0 meson production measured some time ago by the LHCb collaboration [27]. We already performed corresponding theoretical studies of the inclusive LHCb charm data based on both, the first (only $c \rightarrow D$) [8] and the second ($c + g \rightarrow D$) scenario [13] in two separate papers. However, a direct comparison of the theoretical predictions based on these two scenarios for single D meson production, calculated with the same set of α_S , scales, unPDFs and other details, can be helpful for drawing definite conclusions in the following discussion of double D meson production. As shown in Fig. 2, both prescriptions give a very good description of the LHCb experimental data. Some small differences between them can be observed for both very small and large meson transverse momenta. The latter effect can be recognized as a result of the DGLAP evolution which makes the slope of the transverse momentum distribution in the second scenario a bit steeper than in the case of the first scenario, which is more favorable by the experimental data points. In the region of very small p_t 's the second scenario gives larger cross sections and slightly overestimates the experimental data points. This may come from the $g \rightarrow D$ fragmentation component which approaches a problematic region where $p_t \sim 2m_c$. Then the treatment of charm quarks as massless in the DGLAP evolution of fragmentation function for very small evolution scale can be a bit questionable and may lead to a small overestimation of the integrated cross sections (especially in the case of $RR \rightarrow g \rightarrow D$ mechanism). We will come back to possible consequences of this effect when discussing DD correlation observables.

Now we wish to compare results of our theoretical approach for double charm production described briefly in the previous section with the LHCb experimental data for $D^0 D^0$ pair production. In Fig. 3 we compare results of our calculation with experimental distribution in transverse momentum of one of the meson from the $D^0 D^0$ (or $\bar{D}^0 \bar{D}^0$) pair. We show results for the first scenario when standard Peterson FF is used for the $c \rightarrow D^0$ (or $\bar{c} \rightarrow \bar{D}^0$)




Fig. 2. Charm meson transverse momentum distribution within the LHCb acceptance for inclusive single D^0 mesons (plus their conjugates) production. Left and right panels correspond to two different rapidity intervals. Theoretical predictions for the Peterson $c \rightarrow D$ fragmentation function (solid lines) are compared to the second scenario calculations with the KKKS08 fragmentation functions (long-dashed lines) with $c \rightarrow D$ (dotted) and $g \rightarrow D$ (short-dashed) components that undergo DGLAP evolution equation.




Fig. 3. D^0 meson transverse momentum distribution within the LHCb acceptance region. The left panel is for the first scenario and for Peterson $c \rightarrow D$ fragmentation function while the right panel is for the second scenario and for the fragmentation function that undergoes DGLAP evolution equation.

fragmentation (left panel) as well as the result for the second scenario when the KKKS08 FFs with DGLAP evolution for $c \rightarrow D^0$ (or $\bar{c} \rightarrow \bar{D}^0$) and $g \rightarrow D^0$ (or $g \rightarrow \bar{D}^0$) are used. The results are almost independent of the scale of the fragmentation function. The dependence on factorization scale of parton distributions and on renormalization scale was discussed e.g. in Ref. [2]. One can observe that the DPS $cc \rightarrow D^0D^0$ contribution in the new scenario is much smaller than in the old scenario. In addition, the slope of the distribution in transverse momentum changes. Both the effects are due to evolution of corresponding fragmentation function in the second scenario, compared to lack of such an effect in the first scenario. The different new mechanisms shown in Fig. 1 give contributions of similar size. We can obtain agreement in the second case provided σ_{eff} parameter is increased from conventional 15 mb to 30 mb. Even then we overestimate the LHCb data for $3 < p_T < 5$ GeV.

Can the increased value of $\sigma_{eff} = 30$ mb be understood? First of all the LHCb experiment measures charmed mesons (charm quarks/antiquarks) in forward directions. As shown in Ref. [22] at larger charm quark/antiquark rapidities the relative contribution of perturbative partonic splitting increases. The σ_{eff} parameter includes both conventional $2\nu 2$ uncorrelated and correlated single

parton $2\nu 1$ splitting contribution. As shown in Ref. [22] the smaller perturbative single parton splitting contribution the larger σ_{eff} . Also the conventional uncorrelated parton picture may be too simplistic. The nonperturbative correlations may lead to the effective dependence of σ_{eff} on c and/or \bar{c} transverse momentum (see a recent model analysis for jet production in Ref. [28]).

In Fig. 4 we show dimeson invariant mass distribution $M_{D^0D^0}$ again for the two cases considered. In the first scenario we get a good agreement only for small invariant masses while in the second scenario we get a good agreement only for large invariant masses. The large invariant masses are strongly correlated with large transverse momenta, so the situation here (for the invariant mass distribution) is quite similar as in Fig. 3 for the transverse momentum distribution.

In Fig. 5 we show azimuthal angle correlation $\varphi_{D^0D^0}$ between D^0 and D^0 (or \bar{D}^0 and \bar{D}^0 mesons). While the correlation function in the first scenario is completely flat, the correlation function in the second scenario shows some tendency similar as in the experimental data. The increase at small $\Delta\varphi$ for the SPS $gg \rightarrow D^0D^0$ contribution is due to s-channel pole in the amplitude for $RR \rightarrow gg$ which we regularize by $\hat{s} > 4m_c^2$ condition. In the k_t -factorization, initial partons have transverse momenta, but final




Fig. 4. $M_{D^0D^0}$ dimeson invariant mass distribution within the LHCb acceptance region. The left panel is for the first scenario and for the Peterson $c \rightarrow D$ fragmentation function while the right panel is for the second scenario and for the fragmentation function that undergo DGLAP evolution equation.




Fig. 5. Distribution in azimuthal angle $\varphi_{D^0D^0}$ between the two D^0 mesons within the LHCb acceptance region. The left panel is for the first scenario and for Peterson $c \rightarrow D$ fragmentation function while the right panel is for the second scenario and for the fragmentation function that undergo DGLAP evolution equation.

gluons may have equal rapidities even when $\Delta\varphi$ is far from π . The observed overestimation comes from the region of small transverse momenta. The situation may be improved when a proper transverse momentum dependence of σ_{eff} will be included, but this needs further studies.

Finally we wish to summarize the present situation for the second scenario. In Fig. 6 we show the different distributions discussed above for different values of σ_{eff} . Good description can be obtained only for extremely large values of σ_{eff} which goes far beyond the geometrical picture [22] and that are much larger than for other reactions and in this sense is inconsistent with the factorized Ansatz. We think that the solution of the inconsistency is not only in the DPS sector as already discussed in this paper.

4. Conclusions

In the present paper we have discussed production of D^0D^0 or $\bar{D}^0\bar{D}^0$ meson–meson pairs in proton–proton collisions at the LHC. We have considered the double-parton scattering mechanism of double $c\bar{c}$ production and subsequent double hadronization of two c quarks or two \bar{c} antiquarks using $c \rightarrow D^0$ or $c \rightarrow \bar{D}^0$ fragmentation functions that undergo DGLAP evolution equation with one of the traditional, scale-independent fragmentation function used as an input at the initial scale that is set to $\mu^2 = 4 m_c^2$.

In addition we have included also production of gluonic dijets and their subsequent hadronization to the neutral pseudoscalar D mesons. The $g \rightarrow D$ fragmentation function is assumed to be zero at the initial scale that is set to $\mu^2 = 4 m_c^2$. Also mixed $g \rightarrow D$ and $c \rightarrow D$ mechanisms occur naturally.

We find that at $\sqrt{s} = 7$ TeV the two mechanisms give similar contribution for the LHCb experimental acceptance. While the DPS mechanism dominates at small D meson transverse momenta, the SPS double gluon fragmentation takes over for larger transverse momenta.

When added together the new mechanisms give similar result as the first scenario with one subprocess ($cc \rightarrow DD$) and fixed (scale-independent) fragmentation function. However, some correlation observables, such as dimeson invariant mass or azimuthal correlations between D mesons, are slightly better described.

In our calculation within the second scenario a larger value of σ_{eff} is needed to describe the LHCb data than found from the review of several experimental studies of different processes. This can be partially understood by a lower contribution of perturbative parton splitting as found in Ref. [22] and/or due to nonperturbative correlations in the nucleon which may lead to transverse momentum dependent σ_{eff} . Clearly more involved studies are needed to understand the situation in details. Some problem may be also related to the fact that the fragmentation function used in the second scenario were obtained in the DGLAP formalism with massless




Fig. 6. The dependence of the results of the second scenario on the parameter σ_{eff} used in the calculation of the DPS contributions. Here the three lines correspond to σ_{eff} equal to 15, 30, and 60 mb, from top to bottom, respectively.

c quarks and \bar{c} antiquarks which may be a too severe approximation, especially for low factorization scales (i.e. low transverse momenta) for fragmentation functions. We expect that including mass effects in the evolution would lower the $g \rightarrow c$ (or $g \rightarrow \bar{c}$) fragmentation. Such a study would be useful but clearly goes beyond the scope of the present paper. At present one may only expect that the final (fully consistent) result should be in between the old and new (not completely consistent at present) approach.

In this context we remind a trial to describe the correlation observables in more involved non-factorized approach to DPS [19]. The authors there neglected hadronization and worked in leading-order collinear approach. However, they were not able to describe the details of the LHCb distributions.

The presence of the new SPS mechanism may mean that the extraction of σ_{eff} directly from the LHCb experimental data [4] may be not correct.

We expect that at higher energies (for example for Future Circular Collider) the proportions will change and at asymptotically high energies, much above the LHC energies, the DPS mechanism will win.

Acknowledgements

We are particularly indebted to Anton Karpishkov for discussion of several technical issues and Ingo Schienbein for explaining some details of KKKS08 fragmentation functions. This study was partially supported by the Polish National Science Centre

grant DEC-2014/15/B/ST2/02528 and by the Center for Innovation and Transfer of Natural Sciences and Engineering Knowledge in Rzeszów. The work was supported by Russian Foundation for Basic Research through the Grant No 14-02-00021 and by the Ministry of Education and Science of the Russian Federation under Competitiveness Enhancement Program of SSAU for 2013-2020.

References

- [1] M. Łuszczak, R. Maciula, A. Szczurek, Phys. Rev. D 85 (2012) 094034, arXiv:1111.3255 [hep-ph].
- [2] R. Maciula, A. Szczurek, Phys. Rev. D 87 (2013) 074039, arXiv:1301.4469 [hep-ph].
- [3] A. van Hameren, R. Maciula, A. Szczurek, Phys. Rev. D 89 (2014) 094019, arXiv:1402.6972 [hep-ph].
- [4] R. Aaij, et al., LHCb Collaboration, J. High Energy Phys. 06 (2012) 141; J. High. Energy Phys. 03 (2014) 108.
- [5] C. Peterson, D. Schlatter, I. Schmitt, P.M. Zerwas, Phys. Rev. D 27 (1983) 105.
- [6] A. van Hameren, R. Maciula, A. Szczurek, Phys. Lett. B 748 (2015) 167, arXiv:1504.06490 [hep-ph].
- [7] M. Cacciari, et al., J. High Energy Phys. 10 (2012) 137.
- [8] R. Maciula, A. Szczurek, Phys. Rev. D 87 (9) (2013) 094022, <http://dx.doi.org/10.1103/PhysRevD.87.094022>, arXiv:1301.3033 [hep-ph].
- [9] B.A. Kniehl, et al., Eur. Phys. J. C 72 (2012) 2082.
- [10] B.A. Kniehl, G. Kramer, Phys. Rev. D 71 (2005) 094013, arXiv:hep-ph/0504058.
- [11] B.A. Kniehl, G. Kramer, Phys. Rev. D 74 (2006) 037502, arXiv:hep-ph/0607306.
- [12] B.A. Kniehl, G. Kramer, I. Schienbein, H. Spiesberger, Phys. Rev. Lett. 96 (2006) 012001, arXiv:hep-ph/0508129.
- [13] A.V. Karpishkov, M.A. Nefedov, V.A. Saleev, A.V. Shipilova, Phys. Rev. D 91 (2015) 054009, arXiv:1410.7139 [hep-ph].

- [14] M.A. Nefedov, V.A. Saleev, A.V. Shipilova, *Phys. Rev. D* 87 (2013) 094030, arXiv:1304.3549 [hep-ph].
- [15] T. Kneesch, B.A. Kniehl, G. Kramer, I. Schienbein, *Nucl. Phys. B* 799 (2008) 34, arXiv:0712.0481 [hep-ph].
- [16] KKKS08-package: <http://lapth.cnrs.fr/ffgenerator/>.
- [17] M. Diehl, A. Schafer, *Phys. Lett. B* 698 (2011) 389, arXiv:1102.3081 [hep-ph].
- [18] M. Diehl, D. Ostermeier, A. Schafer, *J. High Energy Phys.* 03 (2012) 089, arXiv:1111.0910 [hep-ph].
- [19] M.G. Echevarria, T. Kasemets, P.J. Mulders, C. Pisano, *J. High Energy Phys.* 04 (2015) 034, arXiv:1501.07291 [hep-ph].
- [20] M.G. Ryskin, A.M. Snigirev, *Phys. Rev. D* 83 (2011) 114047, arXiv:1103.3495 [hep-ph].
- [21] J.R. Gaunt, *J. High Energy Phys.* 1301 (2013) 042, [http://dx.doi.org/10.1007/JHEP01\(2013\)042](http://dx.doi.org/10.1007/JHEP01(2013)042), arXiv:1207.0480 [hep-ph].
- [22] J.R. Gaunt, R. Maciula, A. Szczurek, *Phys. Rev. D* 90 (5) (2014) 054017, <http://dx.doi.org/10.1103/PhysRevD.90.054017>, arXiv:1407.5821 [hep-ph].
- [23] K. Golec-Biernat, E. Lewandowska, M. Serino, Z. Snyder, A.M. Stasto, *Phys. Lett. B* 750 (2015) 559, arXiv:1507.08583 [hep-ph].
- [24] V.S. Fadin, E.A. Kuraev, L.N. Lipatov, *Sov. Phys. JETP* 44 (1976) 443–450, *Zh. Eksp. Teor. Fiz.* 71 (1976) 840–855.
- [25] M.A. Kimber, A.D. Martin, M.G. Ryskin, *Eur. Phys. J. C* 12 (2000) 655, arXiv:hep-ph/9911379; *Phys. Rev. D* 63 (2001) 114027, arXiv:hep-ph/0101348; G. Watt, A.D. Martin, M.G. Ryskin, *Eur. Phys. J. C* 31 (2003) 73, arXiv:hep-ph/0306169; *Phys. Rev. D* 70 (2004) 014012; 70, 079902(E), arXiv:hep-ph/0309096, 2004.
- [26] L.A. Harland-Lang, A.D. Martin, P. Motylinski, R.S. Thorne, *Eur. Phys. J. C* 75 (5) (2015) 204, arXiv:1412.3989 [hep-ph].
- [27] R. Aaij, et al., LHCb Collaboration, *Nucl. Phys. B* 871 (2013) 1, arXiv:1302.2864 [hep-ex].
- [28] S. Ostapchenko, M. Bleicher, arXiv:1511.06784 [hep-ph].