

HOSTED BY



Contents lists available at ScienceDirect

## HAYATI Journal of Biosciences

journal homepage: <http://www.journals.elsevier.com/hayati-journal-of-biosciences>

Original research article

Two Newly Recorded Species of the Lobster Family Scyllaridae (*Thenus indicus* and *Scyllarides haanii*) From South of Java, Indonesia

Yusli Wardiatno,\* Agus Alim Hakim, Ali Mashar, Nurlisa Alias Butet, Luky Adrianto

Department of Aquatic Resources Management, Faculty of Fisheries and Marine Science, Bogor Agricultural University, Bogor, West Java, Indonesia.

## ARTICLE INFO

## Article history:

Received 5 February 2016

Received in revised form

30 April 2016

Accepted 9 May 2016

Available online xxx

## KEYWORDS:

Crustacean,  
Decapoda,  
first record,  
Indian Ocean,  
slipper lobster

## ABSTRACT

Two species of slipper lobster, *Thenus indicus* Leach, 1815, and *Scyllarides haanii* De Haan, 1841, are reported for the 1<sup>st</sup> time from the coastal waters of South of Java, part of the Indian Ocean. A total of two specimens, one specimen of *T. indicus* from Palabuhanratu Bay and one specimen of *S. haanii* from Yogyakarta coastal waters, were collected in April and September 2015, respectively. Descriptions and illustrations of the morphological characteristics of the two species and their habitat are presented.

Copyright © 2016 Institut Pertanian Bogor. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## 1. Introduction

From a fishery point of view, slipper lobsters of the family Scyllaridae are less popular than palinurid lobsters in Indonesia. Much information on palinurid lobsters [i.e. *Panulirus penicillatus* (Chow *et al.* 2011; Kalih 2012; Abdullah *et al.* 2014), *Panulirus versicolor* (Ongkers *et al.* 2014), *Panulirus homarus*, *Panulirus longipes*, and *Panulirus ornatus* (Kalih 2012)] has been reported, but little information is available on slipper lobsters [e.g. *Parribacus antarcticus* (Kalih 2012)]. The family Scyllaridae consists of four subfamilies, 19 genera, 88 species, and two subspecies worldwide (Holthuis 1991; Chan 1998; Chan 2010; Yang *et al.* 2011; Yang & Chan 2012). The four subfamilies are Arctidinae, Ibacinae, Scyllarinae, and Theninae. The subfamily Arctidinae comprises two genera (i.e. *Arctides* and *Scyllarides*) and the subfamily Theninae only one genus (i.e. *Thenus*). All slipper lobster species are bottom-dwelling and found in very shallow water to a depth of more than 484 m (Chan 1998).

Before the recognition of three new species by Burton & Davie (2007), Jones (1993) and Davie & Burton (2000) resurrected one species of the genus *Thenus* Leach, 1815, a monotypic genus with

only one species, *Thenus orientalis*. Taxonomic and biodiversity studies on Indonesian lobsters resulted in the collection of two different species of family Scyllaridae from South of Java. The present observation is the 1<sup>st</sup> record of these two species from South of Java, Indonesia. The morphological characteristics of the two species are illustrated and described.

## 2. Materials and Methods

A total of two specimens were collected: one specimen of *Thenus indicus* was collected in April 2015 from Palabuhanratu Bay and one specimen of *Scyllarides haanii* in September 2015 from Yogyakarta coastal waters (Figure 1). They were bought from fishermen and preserved in 96% alcohol and transported to the laboratory for analysis. The fishermen caught them with 2- and 3-in nets. The two specimens are lodged in Laboratory of Aquatic Biomolecular, Department of Aquatic Resources Management, Faculty of Fisheries and Marine Science, Bogor Agricultural University. The voucher number is #TIP01 for *T. indicus* and #SHY01 for *S. haanii*.


Identification of the genus *Scyllarides* was based on the morphological characteristics using the taxonomic key book from Food and Agriculture Organization (Holthuis 1991; Chan 1998), i.e. carapace, 4<sup>th</sup> abdominal segment, posterior margin of the 2<sup>nd</sup> abdominal, and spots on the 1<sup>st</sup> abdominal segment. Whereas, the genus *Thenus* was identified based on the

\* Corresponding author.

E-mail address: [yusli@ipb.ac.id](mailto:yusli@ipb.ac.id) (Y. Wardiatno).

Peer review under responsibility of Institut Pertanian Bogor.

<http://dx.doi.org/10.1016/j.hjb.2016.05.001>1978-3019/Copyright © 2016 Institut Pertanian Bogor. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).



**Figure 1.** Locations from which *Thenus indicus* and *Scyllarides haanii* were collected. The red circle indicates Palabuhanratu Bay and the blue circle indicates Yogyakarta coastal waters, South of Java, Indonesia.

morphological characteristics using the taxonomic key from FAO (Burton & Davie 2007), i.e. spotting on pereopods, outer face of the 2<sup>nd</sup> propodus, merus of the 3<sup>rd</sup> maxilliped, and several morphometric ratios. Information on habitat of the two species was recorded by interview with the fishermen.

### 3. Results

The systematics of these two species from the order are as follow:

Order DECAPODA Latreille, 1803

Infraorder ACHELATA Scholts and Richter, 1995

Family SCYLLARIDAE Latreille, 1825

Genus *Thenus* Leach, 1816

*Thenus indicus* Leach, 1815

(Figures 2A and 3)

Genus *Scyllarides* Gill, 1898

*Scyllarides haanii* De Haan, 1841

(Figures 2B and 4)


Synonyms: *Scyllarus haanii* De Haan, 1841

#### 3.1. Materials examined

*T. indicus*. #TIP01. Male: carapace length (CL), 50.29 mm; total length, 84.06 cm; weight, 280 g. April 2015, Palabuhanratu Bay, South of Java, Indonesia.



**Figure 2.** (A) Species of *Thenus indicus* (male) from Palabuhanratu Bay, South of Java, Indonesia, April 2015. (B) Species of *Scyllarides haanii* (male) from Yogyakarta coastal waters, South of Java, Indonesia, September 2015.



**Figure 3.** *Thenus indicus* (Lund 1793), Palabuhanratu Bay, South of Java (Indian Ocean), male (carapace length 50.29 mm). (A) First pereopod; (B) 2<sup>nd</sup> pereopod; (C) 3<sup>rd</sup> maxilliped; (D) abdomen, lateral view. Scale bars, 10 mm.


*S. haanii*. #SHY01. Male: CL, 68.28 mm; total length, 158.64 mm; weight, 164 g. September 2015, Yogyakarta coastal waters, South of Java, Indonesia.

### 3.2. Distinctive characteristics

*T. indicus*. No spotting on the pereopods (Figure 3A,B); the outer face of the propodus of the 2<sup>nd</sup> pereopod had an uppermost longitudinal groove bearing obvious setae over at least the proximal half (Figure 3B). Merus of the 3<sup>rd</sup> maxilliped with a small spine proximally on the inner ventral margin; the inner margin of the ischium was prominently dentate along the entire length (Figure 3C). Abdominal segments, each with lateral margins expanded downward concealing the pleopods (Figure 3D). No single morphometric ratio that would exclusively identify this

species has been identified, but only *T. indicus* can have ratios that fall outside the following maximum and minimum values: 1<sup>st</sup> pereopod merus width less than 0.07 CL and 3<sup>rd</sup> pereopod merus length more than 0.45 CL.

*S. haanii*. Body vaulted, pubescent, and covered with rounded tubercles. Eyes small and subspherical. Carapace somewhat rectangular, but the posterior half distinctly wider than the anterior half; cervical groove strong; pregastric, gastric, and cardiac teeth all strongly protruding. Antennae broad, flattened, and plate-like, with distal margin finely crenate (Figure 4A). All legs without pincers and similar in size. Abdomen uniformly granulated and not particularly sculptured; dorsal midline of the 2<sup>nd</sup> and 3<sup>rd</sup> segments strongly ridged and that of the 4<sup>th</sup> segment producing a remarkable hump; pleura directed downward, with the posterior margin of the



**Figure 4.** *Scyllarides haanii* (De Haan 1841), Yogyakarta coastal waters, South of Java (Indian Ocean), male (carapace length 68.28 mm). (A) Carapace and eye, lateral view; (B) abdomen, lateral view. Scale bars, 10 mm.

2<sup>nd</sup> pleuron somewhat concave (Figure 4B). The posterior half of the tail fan is soft and flexible. The 1<sup>st</sup> abdominal segment was yellowish with three diffuse purplish red spots. The soft part of the tail fan is light brown with numerous purple dots.

### 3.3. Habitat and distribution

As per the information from the fisherman, the two lobster species are actually present in both Palabuhanratu Bay and Yogyakarta coastal waters. In Palabuhanratu Bay, *T. indicus* was caught from shallow sandy habitat with small portion of silt fraction near rocky coast line with clear waters at 5–15 m depth, whereas habitat of *S. haanii* in Yogyakarta coastal waters is sandy substrate.

According to Burton and Davie (2007), *T. indicus* occurred in Pakistan, India, the Gulf of Thailand, Singapore, Indonesia (west coast of Sumatra Island), and Taiwan. Nevertheless, this study showed that *T. indicus* could be found in Palabuhanratu Bay, South of Java, as a new distribution record. Meanwhile, Holthuis (1991) and Chan (1998) demonstrated the presence of *S. haanii* in Indo-West Pacific from Mauritius to the Red Sea, Japan, Indonesia, Australia, and Hawaii. In the works by Holthuis (1991) and Chan (1998), the distribution map of *S. haanii* in Indonesia showed its presence in west coast of Sumatra Island, Banten coastal waters to the Sunda Strait, and Aru archipelago. The occurrence of this species in Yogyakarta coastal waters, South of Java is a new distribution record.

## 4. Discussion

As targets in marine fishery, the economic value of slipper lobsters has increased since the 1990s, and its consumption as seafood has become more popular in some parts of the world (Spanier and Lavalli 2006). *Thenus* is the most economically important genus of

the family Scyllaridae (Jones 1993), despite the fact that in some countries, the slipper lobster is a bycatch product, such as in southern and southeastern Brazil (Oliveira et al. 2008, Duarte et al., 2010) and Australia (Courtney et al. 2001; Brewer et al. 2006). In Indonesia, the slipper lobster species *P. antarcticus* was one of six economically valuable lobsters in the coastal waters of Lombok Island (Kalih 2012).

Slipper lobster fisheries on the east coast of Queensland, Australia, face a problem in that *T. orientalis* and *T. indicus* could not be differentiated in logbook records and the market place as their taxonomy and distribution differ only slightly (Courtney et al. 2001). Before Jones (1993), *T. indicus* was named *T. orientalis*. *Thenus* is a complex of five species, and the species called *T. indicus* by Jones (1993) is in fact the new species *Thenus parindicus* (Burton and Davie 2007). *S. haanii* is closely related to *Scyllarides squammosus* and has similarly shaped, blunt dactyls (Hwang & Yu 1983; Lau 1988). However, it can be distinguished by its larger body and distinct protuberances on the 2<sup>nd</sup> to 4<sup>th</sup> terga (Hwang and Yu 1983).

According to the distribution maps (Holthuis 1991; Chan 1998; Burton and Davie 2007), *T. indicus* and *S. haanii* occurred in the west coast of Sumatra Island and Banten coastal waters to the Sunda Strait. It is interesting that the two species were found in coastal waters of South Java. The water mass circulation of south east Asian waters was clearly demonstrated by Wyrтки (1961). With the circulation pattern, the coastal waters of west Sumatra and the Sunda Strait have a connectivity with the coastal waters of South Java. Because phyllosoma larvae period of the scyllarid lobsters is long (3 until 6 months; Sekiguchi et al. 2007), the larvae may be taken away with long distances by currents. As a result, the settlement may be taking place far away from their original spawning grounds (Lewis 1951). So, it is not impossible for the two species to exist in south of Java waters with the assumption that west coast of Sumatra, Banten coastal waters, and Sunda Strait was the breeding grounds.

In terms of habitat, the habitat characteristics of *T. indicus* and *S. haanii* as informed by the fisherman are similar to the explanation of Jones (2007) and Chan (1998). Jones (2007) found *T. indicus* on fine mud or “silty inshore substrates” between 10 and 30 m. Whereas, Chan (1998) informed the habitat of *S. haanii* in coral or rocky reefs at depths from 10 to 135 m, usually less than 50 m.

The number of Indonesian crustaceans reported as 1<sup>st</sup> records has increased [e.g. *Albunea symmista* (Mashar et al. 2015), *Hippa marmorata* (Wardiatno et al. 2015), *Hippa adactyla* (Ardika et al. 2015), *Puerulus mesodontus* (Wardiatno et al. 2016)]. Studies on lobsters in Indonesia mostly focused on highly economically valuable species, such as *P. penicillatus* (Chow et al. 2011; Kalih 2012; Abdullah et al. 2014), *Linuparus somniosus* (Wowor 1999), *P. versicolor* (Ongkers et al. 2014), *P. homarus*, *P. longipes*, *P. ornatus* (Kalih 2012), and *P. mesodontus* (Wardiatno et al. 2016). A study on the occurrence of the slipper lobster, *P. antarcticus*, in the coastal waters of Lombok Island, Indonesia, was reported by Kalih (2012). The presence of these two species in two coastal areas of the southern part of Java Island complements the extant crustacean biodiversity in Indonesian waters. Moreover, this strengthens the notion that Indonesia is a hotspot of marine biodiversity. Information on the biological aspects of these two species is needed to facilitate their sustainable management; therefore, studies of the biology of these slipper lobsters are warranted.

## Acknowledgements

Mr Agus is thanked for his help during specimen collection in Palabuhanratu Bay. The authors are indebted to three anonymous

reviewers for their constructive comments to improve the submitted manuscript.

### Conflict of Interest

None.

### Funding

The research was funded by Directorate General of Higher Education, Ministry of Research, Technology, and Higher Education, Republic of Indonesia from Fiscal 2015 under the research umbrella entitled "The analysis of lobster population (*Panulirus* spp.) in southern waters of Java to support the implementation of ecosystem based lobster fisheries management".

### References

- Abdullah MF, Alimuddin Muththalib M, Salama AJ, Imai H. 2014. Genetic isolation among the Northwestern, Southwestern and Central-Eastern Indian Ocean populations of the pronghorn spiny lobster *Panulirus penicillatus*. *Int J Mol Sci* 15:9242–54.
- Ardika PU, Farajallah A, Wardiatno Y. 2015. First record of *Hippa adactyla* (Fabricius, 1787; Crustacea, Anomura, Hippidae) from Indonesian Waters. *Trop Life Sci Res* 26:105–10.
- Brewer D, Heales D, Milton D, Dell Q, Fry G, Venables B, Jones P. 2006. The impact of turtle excluder devices and bycatch reduction devices on diverse tropical marine communities in Australia's northern prawn trawl fishery. *Fish Res* 81: 176–88.
- Burton TE, Davie PJF. 2007. A revision of the shovel-nosed lobsters of the genus *Thenus* (Crustacea: Decapoda: Scyllaridae), with descriptions of three new species. *Zootaxa* 1429:1–38.
- Chan TY. 1998. Lobster. In: Carpenter KE, Niem VH (Eds.). *FAO Species Identification Guide for Fishery Purposes. The Living Marine Resources of the Western Central Pacific. Volume 2. Cephalopods, Crustaceans, Holothurians and Sharks*. Rome: FAO. pp. 687–1396.
- Chan TY. 2010. Annotated checklist of the world's marine lobsters (Crustacea: Decapoda: Astacidea, Glypheidea, Achelata, Polychelida). *Raffles Bull Zool (Suppl)* 23:153–81.
- Chow S, Jeffs A, Miyake Y, Konishi K, Okazaki M, Suzuki N, Abdullah MF, Imai H, Wakabayashi T, Sakai M. 2011. Genetic isolation between the Western and Eastern Pacific populations of pronghorn spiny lobster *Panulirus penicillatus*. *PLoS One* 6:e29280.
- Courtney AJ, Cosgrove MG, Die DJ. 2001. Population dynamics of Scyllarid lobsters of the genus *Thenus* spp. on the Queensland (Australia) east coast I. Assessing the effects of tagging. *Fish Res* 53:251–61.
- Davie PJF, Burton TE. 2000. Case 3135. *Scyllarus orientalis* Lund, 1793 (currently *Thenus orientalis*; Crustacea, Decapoda): proposed designation of a neotype. *Bull Zool Nomencl* 57:84–6.
- Duarte LFA, Severino-Rodrigues E, Gasalla MA. 2010. Slipper lobster (Crustacea, Decapoda, Scyllaridae) fisheries off the southeastern coast of Brazil: I. Exploitation patterns between 23°00'U and 29°65'S. *Fish Res* 102:141–51.
- Holthuis LB. 1991. Marine lobsters of the world. An annotated and illustrated catalogue of species of interest to fisheries known to date. FAO species catalogue. Volume 13. *FAO Fish Synop* 125. viii + 1–292.
- Hwang JJ, Yu HP. 1983. Report of the Scyllarid lobsters (Crustacea: Decapoda: Scyllaridae) from Taiwan. *Bull Inst Zool Acad Sin* 22:261–7.
- Jones CM. 1993. Population structure of *Thenus orientalis* and *T. indicus* (Decapoda: Scyllaridae) in northeastern Australia. *Mar Ecol Prog Ser* 97:143–55.
- Jones CM. 2007. Biology and fishery of the bay lobster, *Thenus* spp. In: Lavalli KL, Spanier E (Eds.). *The Biology and Fisheries of the Slipper Lobster*, vol. 17. Boca Raton, F. L.: CRC Press. pp. 325–58. Crustacean Issues.
- Kalih LATTWS. 2012. Diversity and Distribution of Palinurid and Scyllarid Lobsters in the Lombok Island Coastal Waters (Master Thesis). Yogyakarta, Indonesia: Graduate School, Gadjah Mada University.
- Lau CJ. 1988. Dietary comparison of two slow-moving crustacean (Decapoda: Scyllaridae) predators by a modified index of relative importance. In: Choat JH, Barnes D, Bo MA (Eds.). *Proceedings of the 6th International Coral Reef Symposium, Townsville, Australia, 8–12 August 1988*, vol. 2. pp. 95–100.
- Lewis JB. 1951. The phyllosoma larvae of the spiny lobster *Panulirus argus*. *Bull Mar Sci Gulf Caribb* 1:89–95.
- Mashar A, Wardiatno Y, Boer M, Butet NA, Farajallah A, Ardika PU. 2015. First record of *Albunea symmysta* (Crustacea: Decapoda: Albuneidae) from Sumatra and Java, Indonesia. *ACL* 8:611–5.
- Oliveira G, Freire AS, Bertuol PRK. 2008. Reproductive biology of the slipper lobster *Scyllarides deceptor* (Decapoda: Scyllaridae) along the southern Brazilian coast. *J Mar Biol Assoc UK* 8:1433–40.
- Ongkers OTS, Pattiasina BJ, Tetelepta JMS, Natan Y, Pattikawa JA. 2014. Some biological aspects of painted spiny lobster (*Panulirus versicolor*) in Latuhalat waters, Ambon Island, Indonesia. *ACL* 7:469–74.
- Sekiguchi H, Booth JD, Webber RW. 2007. Early life histories of slipper lobsters, Chapter 4. In: Lavalli KL, Spanier E (Eds.). *The Biology and Fisheries of the Slipper Lobster*. United State of America: CRC Press. pp. 69–90.
- Spanier E, Lavalli KL. 2006. Scyllarides species, Chapter 14. In: Phillips BF (Ed.). *Lobsters: Biology, Management, Aquaculture and Fisheries. Part 2: Lobsters of Commercial Importance*. United Kingdom: Oxford. pp. 462–96.
- Wardiatno Y, Ardika PU, Farajallah A, Mashar A, Ismail. 2015. The mole crab *Hippa marmorata* (Hombron et Jacquinot, 1846) (Crustacea Anomura Hippidae): a first record from Indonesian waters. *Biodivers J* 6:517–20.
- Wardiatno Y, Hakim AA, Mashar A, Butet NA, Adrianto L, Farajallah A. 2016. First record of *Puerulus mesodontus* Chan, Ma & Chu, 2013 (Crustacea, Decapoda, Achelata, Palinuridae) from south of Java, Indonesia. *BDJ* 4:e8069. <http://dx.doi.org/10.3897/BDJ.4.e8069>.
- Wyrarki K. 1961. Physical Oceanography of the Southeast Asian Waters. NAGA Report Volume 2. Scientific Results of Marine Investigations of the South China Sea and the Gulf of Thailand, 1959–1961. La Jolla, Calif: S.I.O..
- Yang CH, Chan TY. 2012. On the taxonomy of the slipper lobster *Chelarctus cultrifer* (Ortmann, 1897) (Crustacea: Decapoda: Scyllaridae), with description of a new species. *Raffles Bull Zool* 60:449–60.
- Yang CH, Chen IS, Chan TY. 2011. A new slipper lobster of the genus *Galearctus* Holthuis, 2002 (Crustacea, Decapoda, Scyllaridae) from New Caledonia. *Zoosystema* 33:207–17.