

- Griendling KK, Sorescu D, Ushio-Fukai M. NAD(P)H oxidase. Role in cardiovascular biology and disease. *Circ Res* 2000;86:494-501.
- Bendall JK, Cave AC, Heymes C, Gall N, Shah AM. Pivotal role of gp91^{phox}-containing NADPH oxidase in angiotensin II induced cardiac hypertrophy. *Circulation* 2002;105:293-6.

Depression as Another Possible Explanation for Worse Outcomes in Myocardial Infarction During Off-Hours

We read with great interest the findings of Henriques et al. (1) regarding worse clinical outcomes for acute myocardial infarction (MI) patients treated with primary angioplasty during off-hours. In their otherwise excellent study, the investigators failed to measure depression or to mention it as another possible reason for the increased rates of failed angioplasty and 30-day mortality observed in those patients presenting between the hours of 1800 and 0800. Depression is common among patients with coronary disease (2), and depressed patients are more likely to develop initial MI symptoms off-hours (3). Depressed patients admitted for MI are at significantly increased risk for mortality (2) and for repeat coronary revascularization (4). Moreover, depression is associated with perturbations of circadian rhythms in patients with and without coronary disease (5). Thus, in addition to the reasons offered by the investigators (1), depression may be another explanation for the worse outcomes observed in patients presenting off-hours.

Donald C. Haas, MD, MPH

Karina W. Davidson, PhD

Thomas G. Pickering, MD, DPhil

The Zena and Michael A. Wiener Cardiovascular Institute

Mount Sinai School of Medicine

One Gustave L. Levy Place

Box 1030

New York, NY 10029-6574

E-mail: donald.haas@msnyuhealth.org

doi:10.1016/j.jacc.2003.10.005

REFERENCES

- Henriques JPS, Haasdijk AP, Zijlstra F, on behalf of the Zwolle Myocardial Infarction Study Group. Outcome of primary angioplasty for acute myocardial infarction during routine duty hours versus during off-hours. *J Am Coll Cardiol* 2003;41:2138-42.
- Rozanski A, Blumenthal JA, Kaplan J. Impact of psychological factors on the pathogenesis of cardiovascular disease and the implications for therapy. *Circulation* 1999;99:2192-217.
- Carney RM, Freedland KE, Jaffe AS. Altered circadian pattern of acute myocardial infarction in patients with depression. *Coron Artery Dis* 1991;2:61-5.
- Sullivan MD, LaCroix AZ, Spertus JA, Hecht J, Russo J. Depression predicts revascularization procedures for 5 years after coronary angiography. *Psychosom Med* 2003;65:229-36.
- Carney RM, Freedland KE, Jaffe AS. Insomnia and depression prior to myocardial infarction. *Psychosom Med* 1990;52:603-9.

Circadian Variations in Outcome of Primary Percutaneous Coronary Intervention

We read with interest the report by Henriques et al. (1) regarding circadian variations in outcomes of primary angioplasty for acute

myocardial infarction. The investigators cite a higher angioplasty failure and a higher mortality in patients treated during the "off-hours" despite similar baseline characteristics and procedural delays compared with patients treated during daytime. The researchers also point out that only 23% of the total cohort was treated during the night. This suggests a potential selection bias of the patients treated at night, particularly among the 11 community hospitals referring patients to the single percutaneous transluminal coronary angioplasty (PTCA) center involved. In a previous smaller study (2), addressing a cohort of consecutive ST-elevation myocardial infarction patients admitted to a single center performing primary percutaneous coronary intervention (PCI), we found that 61% of the patients were treated during off-hours (also counting weekend hours), a percentage perfectly in line with the proportion of off-hours during the week (64%). In that study, despite worse baseline characteristics of patients admitted during off-hours, patients experienced almost identical procedural success rates (98% vs. 96% TIMI [Thrombolysis In Myocardial Infarction] 3 flow rates for "off" vs. business hours), predischARGE radionuclide left ventricular ejection fraction (51 ± 14 vs. $53 \pm 12\%$) and in-hospital death rates (7% vs. 5%) compared with patients treated during business hours ($p = \text{NS}$ for all comparisons).

As pointed out in the accompanying editorial in the *Journal* (3), the variations in patterns of care and outcomes reported by Henriques et al. (1) may be related to biological circadian variations but also to variations in health care delivery, particularly prehospital referral patterns and delays. We submit that the latter are more likely than the former. To explore this issue further and to minimize biases related to prehospital referral, we suggest that data be analyzed from several centers with a "captive" patient population (i.e., only one hospital serving a given population) and using primary PCI as the sole reperfusion therapy for ST-elevation myocardial infarction.

Ph. Gabriel Steg, MD, FACC

Department of Cardiology

Hôpital Bichat

46 rue Henri Huchard

75877 Paris Cedex 18

France

E-mail: gabriel.steg@bch.ap-hop-paris.fr

Philippe Garot, MD

doi:10.1016/j.jacc.2003.10.006

REFERENCES

- Henriques JPS, Haasdijk AP, Zijlstra F, on behalf of the Zwolle myocardial infarction study group. Outcome of primary angioplasty for acute myocardial infarction during routine duty hours versus during off-hours. *J Am Coll Cardiol* 2003;41:2138-42.
- Garot PH, Juliard JM, Benamer H, Steg PG. Are the results of primary PTCA for acute myocardial infarction different during the "off" hours? *Am J Cardiol* 1997;79:1527-9.
- Spencer FA, Becker RC. Circadian variations in acute myocardial infarction: patients or health care delivery? *J Am Coll Cardiol* 2003;41:2143-6.

REPLY

We appreciate Dr. Haas's as well as Dr. Steg's and their colleagues' interest and comments on our report concerning the different