

Linear choosability of sparse graphs

Daniel W. Cranston^{a,*}, Gexin Yu^{b,1}

^a Department of Mathematics and Applied Mathematics, Virginia Commonwealth University, Richmond, VA 23284, United States

^b Department of Mathematics, College of William & Mary, Williamsburg, VA 23185, United States

ARTICLE INFO

Article history:

Received 8 October 2010

Received in revised form 7 May 2011

Accepted 15 May 2011

Available online 12 June 2011

Keywords:

Linear coloring

List coloring

Planar graph

Discharging method

Maximum average degree

ABSTRACT

A linear coloring is a proper coloring such that each pair of color classes induces a union of disjoint paths. We study the linear list chromatic number, denoted $lc_\ell(G)$, of sparse graphs. The maximum average degree of a graph G , denoted $mad(G)$, is the maximum of the average degrees of all subgraphs of G . It is clear that any graph G with maximum degree $\Delta(G)$ satisfies $lc_\ell(G) \geq \lceil \Delta(G)/2 \rceil + 1$. In this paper, we prove the following results: (1) if $mad(G) < 12/5$ and $\Delta(G) \geq 3$, then $lc_\ell(G) = \lceil \Delta(G)/2 \rceil + 1$, and we give an infinite family of examples to show that this result is best possible; (2) if $mad(G) < 3$ and $\Delta(G) \geq 9$, then $lc_\ell(G) \leq \lceil \Delta(G)/2 \rceil + 2$, and we give an infinite family of examples to show that the bound on $mad(G)$ cannot be increased in general; (3) if G is planar and has girth at least 5, then $lc_\ell(G) \leq \lceil \Delta(G)/2 \rceil + 4$.

© 2011 Elsevier B.V. All rights reserved.

1. Introduction

In 1973, Grünbaum introduced *acyclic colorings* [3], which are proper colorings with the additional property that each pair of color classes induces a forest. In 1997, Hind, Molloy, and Reed introduced *frugal colorings* [4]. A proper coloring is *k-frugal* if the subgraph induced by each pair of color classes has maximum degree less than k . Yuster [8] combined the ideas of acyclic coloring and 3-frugal coloring in the notion of a *linear coloring*, which is a proper coloring such that each pair of color classes induces a union of disjoint paths—also called a *linear forest*. We write $lc(G)$ to denote the *linear chromatic number* of G , which is the smallest integer k such that G has a proper k -coloring in which every pair of color classes induces a linear forest.

We begin by noting easy upper and lower bounds on $lc(G)$. If G is a graph with maximum degree $\Delta(G)$, then we have the naive lower bound $lc(G) \geq \lceil \Delta(G)/2 \rceil + 1$, since each color can appear on at most two neighbors of a vertex of maximum degree. Observe that $lc(G) \leq \chi(G^2) \leq \Delta(G^2) + 1 \leq \Delta(G)^2 + 1$, where $\chi(G)$ denotes the chromatic number of G and G^2 is the square graph of G . Yuster [8] constructed an infinite family of graphs such that $lc(G) \geq C_1 \Delta(G)^{3/2}$, for some constant C_1 . He also proved an upper bound of $lc(G) \leq C_2 \Delta(G)^{3/2}$, for some constant C_2 and for sufficiently large $\Delta(G)$.

Note that trees with maximum degree $\Delta(G)$ have linear chromatic number $\lceil \Delta(G)/2 \rceil + 1$, i.e., the naive lower bound holds with equality (for example, we can color greedily in the order of a breadth-first search from an arbitrary vertex). This equality for trees suggests that sparse graphs might have linear chromatic number close to the naive lower bound. To be more precise: Does there exist a constant C such that every sparse graph G satisfies $lc(G) \leq \lceil \Delta(G)/2 \rceil + C$? To state the previous results related to this question, we first introduce some more notation.

We start with linear list colorings, which are linear colorings from assigned lists. Formally, let $lc_\ell(G)$ be the *linear list chromatic number* of G , that is, the smallest integer k such that if each vertex $v \in V(G)$ is given a list $L(v)$ with $|L(v)| \geq k$,

* Corresponding author.

E-mail addresses: dcranston@vcu.edu, dcransto@gmail.com (D.W. Cranston), gyu@wm.edu (G. Yu).

¹ Research supported in part by the NSF grant DMS-0852452.

then G has a linear coloring such that each vertex v gets a color $c(v)$ from its list $L(v)$. When all the lists are the same, linear list coloring is the same as linear coloring. General list coloring was first introduced by Erdős, Rubin, and Taylor [1] and independently by Vizing [7] in the 1970s, and it has been well explored since then [5].

Linear list colorings were first studied by Esperet, Montassier, and Raspaud [2]. The *maximum average degree* of a graph G , denoted $\text{mad}(G)$, is the maximum of the average degrees of all of its subgraphs, i.e., $\text{mad}(G) = \max_{H \subseteq G} \frac{2|E(H)|}{|V(H)|}$. Observe that the family of all trees is precisely the set of connected graphs with $\text{mad}(G) < 2$ (so indeed we are generalizing our motivating example, trees). The following results were shown in [2]:

Theorem A ([2]). *Let G be a graph:*

- (1) If $\text{mad}(G) < 8/3$, then $\text{lc}_\ell(G) \leq \lceil \Delta(G)/2 \rceil + 3$.
- (2) If $\text{mad}(G) < 5/2$, then $\text{lc}_\ell(G) \leq \lceil \Delta(G)/2 \rceil + 2$.
- (3) If $\text{mad}(G) < 16/7$ and $\Delta(G) \geq 3$, then $\text{lc}_\ell(G) = \lceil \Delta(G)/2 \rceil + 1$.

The *girth* of a graph G , denoted $g(G)$, or simply g , is the length of its shortest cycle. By an easy application of Euler’s formula, we see that every planar graph G with girth g satisfies $\text{mad}(G) < 2g/(g - 2)$. So we can obtain some results on planar graphs from the above results. Raspaud and Wang [6] proved somewhat stronger results for planar graphs.

Theorem B ([6]). *Let G be a planar graph:*

- (1) If $g(G) \geq 5$, then $\text{lc}(G) \leq \lceil \Delta(G)/2 \rceil + 14$.
- (2) If $g(G) \geq 6$, then $\text{lc}(G) \leq \lceil \Delta(G)/2 \rceil + 4$.
- (3) If $g(G) \geq 13$ and $\Delta(G) \geq 3$, then $\text{lc}(G) = \lceil \Delta(G)/2 \rceil + 1$.

Our goal in the paper is to improve the results in the above two theorems. We prove the following.

Theorem 1. *Let G be a graph:*

- (1) If G is planar and has $g(G) \geq 5$, then $\text{lc}_\ell(G) \leq \lceil \Delta(G)/2 \rceil + 4$.
- (2) If $\text{mad}(G) < 3$ and $\Delta(G) \geq 9$, then $\text{lc}_\ell(G) \leq \lceil \Delta(G)/2 \rceil + 2$.
- (3) If $\text{mad}(G) < 12/5$ and $\Delta(G) \geq 3$, then $\text{lc}_\ell(G) = \lceil \Delta(G)/2 \rceil + 1$.

Raspaud and Wang [6] conjectured that the bound in Theorem 1(2) holds for all planar graphs with girth at least 6. Since every such graph G has $\text{mad}(G) < 3$, our result proves their conjecture for graphs with $\Delta(G) \geq 9$. Since $\text{mad}(K_{3,3}) = 3$ and $\text{lc}(K_{3,3}) = 5$, we can construct an infinite family of sparse graphs G containing $K_{3,3}$ such that $\text{mad}(G) = 3$, $\Delta(G) = 4$, and $\text{lc}(G) > \lceil \Delta(G)/2 \rceil + 2$. Thus, the maximum degree condition in Theorem 1(2) cannot be lower than 5.

We also note that $\text{lc}(K_{2,3}) = 4 > \lceil \Delta(K_{2,3})/2 \rceil + 1$ and $\text{mad}(K_{2,3}) = 12/5$. Thus, we can construct an infinite family of sparse graphs containing $K_{2,3}$ with maximum degree at most 4. All such graphs have $\text{lc}(G) = \lceil \Delta(G)/2 \rceil + 2$ and can be made sparse enough so that $\text{mad}(G) = \text{mad}(K_{2,3}) = 12/5$. So the bound on $\text{mad}(G)$ in Theorem 1(3) is sharp.

The proofs of our three results all follow the same outline. First, we prove a structural lemma; this says that each graph under consideration must contain at least one from a list of “configurations”. Second, we prove that any minimal counterexample to our theorem cannot contain any of these configurations. In this second step, we begin with a linear list coloring of part of the graph, and show how to extend it to the whole graph. As we extend the coloring, we often say that we “choose $c(v) \in L(v)$ ”; by this we mean that we pick some color $c(v)$ from $L(v)$ and use $c(v)$ to color vertex v . In the following three sections, we will prove our three main results, respectively.

For convenience, we introduce the following notation. A k -vertex is a vertex of degree k . A k^+ -vertex (k^- -vertex) is a vertex of degree at least (at most) k . A k -thread is a path of $k + 2$ vertices, where each of the k internal vertices has degree 2, and each of the end vertices has degree at least 3.

2. Planar with girth at least 5 implies $\text{lc}_\ell(G) \leq \lceil \frac{\Delta(G)}{2} \rceil + 4$

Lemma 1. *If G is a planar graph with $\delta(G) \geq 2$ and with girth at least 5, then G contains one of the following two configurations:*

- (RC1) a 2-vertex adjacent to a 5^- -vertex,
- (RC2) a 5-face with four incident 3-vertices and the fifth incident vertex of degree at most 5.

Proof. We use the discharging method, with initial charge $\mu(f) = d(f) - 5$ for each face f and initial charge $\mu(v) = \frac{3}{2}d(v) - 5$ for each vertex v . By Euler’s formula, we have $\sum_{v \in V(G)} \mu(v) + \sum_{f \in F(G)} \mu(f) = (3|E| - 5|V|) + (2|E| - 5|F|) = -5(|F| - |E| + |V|) = -10$. We redistribute charge via the following two discharging rules:

- (R1) Each 4^+ -vertex v sends charge $\frac{\frac{3}{2}d(v) - 5}{d(v)}$ to each incident face (for each time they touch).
- (R2) Each face sends charge 1 to each incident 2-vertex and charge $\frac{1}{6}$ to each incident 3-vertex (for each time they touch).

When we write “for each time they touch”, we mean that if a vertex v is a cut-vertex and appears k times on a facial walk around face f , then the rule for sending charge between v and f should be applied k times. Otherwise, (for example) a 2-vertex that is a cut-vertex would not receive sufficient charge from its one incident face.

Now we will show that if G contains neither configuration (RC1) nor (RC2), then each vertex and each face finishes with nonnegative charge. This is a contradiction, since the discharging rules preserve the sum of the charges (which begins negative). We write $\mu^*(v)$ and $\mu^*(f)$ to denote the charge at vertex v or face f after we apply all discharging rules. If $d(v) = 2$, then $\mu^*(v) = (\frac{3}{2}(2) - 5) + 2(1) = 0$. If $d(v) = 3$, then $\mu^*(v) = (\frac{3}{2}(3) - 5) + 3(\frac{1}{6}) = 0$. By design, each 4^+ -vertex finishes with charge 0. So, we now consider the final charge on each face.

Let f be a face of G . For each pair, u_1 and u_2 , of adjacent vertices on f , we compute the net charge given from f to u_1 and u_2 . If neither of u_1 and u_2 is a 2-vertex, then each vertex receives charge at most $\frac{1}{6}$ from f , so the net charge given from f to u_1 and u_2 is at most $2(\frac{1}{6}) = \frac{1}{3}$. If one of u_1 and u_2 , say u_1 , is a 2-vertex, then, since G does not contain (RC1), we have $d(u_2) \geq 6$. Hence, the net charge given from f to u_1 and u_2 is at most $1 - \frac{2}{3} = \frac{1}{3}$. (This is true because as the degree of a vertex increases beyond 6, the charge it gives to each incident face increases beyond $\frac{2}{3}$.) By a simple counting argument, we see that the net total charge given from f to all incident vertices is at most $\frac{1}{2}(\frac{1}{3}d(f)) = \frac{1}{6}d(f)$. Since $\mu(f) = d(f) - 5$, we see that $\mu^*(f) \geq 0$ when $d(f) \geq 6$. Now we consider 5-faces.

Suppose f is a 5-face. Let n_2, n_3 , and n_{6^+} denote the number of 2-vertices, 3-vertices, and 6^+ -vertices incident to f . Note that $\mu^*(f) \geq -n_2 - \frac{1}{6}n_3 + \frac{2}{3}n_{6^+}$. From (RC1), we have $n_2 \leq \lfloor d(f)/2 \rfloor = 2$. If $n_2 = 2$, then $n_3 = 0$ and $n_{6^+} = 3$, so $\mu^*(f) \geq -2 - \frac{1}{6}(0) + \frac{2}{3}(3) = 0$. If $n_2 = 1$, then $n_{6^+} \geq 2$, so $n_3 \leq 2$. Hence, $\mu^*(f) \geq -1 - \frac{1}{6}(2) + \frac{2}{3}(2) = 0$.

Suppose now that f is a 5-face and $n_2 = 0$. Since we have no copy of (RC2), we have either $n_3 = 4$ and $n_{6^+} = 1$, or we have $n_3 \leq 3$. In the first case, we get $\mu^*(f) \geq -0 - \frac{1}{6}(4) + \frac{2}{3}(1) = 0$. In the second case, note that f has at least two 4^+ -vertices, each of which gives f charge at least $\frac{1}{4}$. Thus $\mu^*(f) \geq -0 - \frac{1}{6}(3) + \frac{1}{4}(2) = 0$. Hence, every face and every vertex has nonnegative charge. This contradiction completes the proof. \square

In Sections 3 and 4, we will only assume bounded maximum average degree (rather than planarity and a girth bound). However, in the proof of the preceding lemma, we needed the stronger hypothesis of planar with girth at least 5. Specifically, we used this hypothesis when considering 5-faces. Our proof relied heavily on the fact that for a 5-face f we have $n_2 \leq \lfloor d(f)/2 \rfloor < d(f)/2$.

Now we use Lemma 1 to prove the following linear list coloring result, which immediately implies Theorem 1(1). For technical reasons, we phrase all of our theorems in terms of an integer M such that $\Delta(G) \leq M$. (Without this technical strengthening, when we consider a subgraph H such that $\Delta(H) < \Delta(G)$, we get complications.) Of course, the interesting case is when $M = \Delta(G)$.

Theorem 2. *Let M be an integer. If G is a planar graph with $\Delta(G) \leq M$ and girth at least 5, then $lc_\ell(G) \leq \lceil \frac{M}{2} \rceil + 4$.*

Proof. Suppose the theorem is false. Let G be a minimal counterexample and let the list assignment L of size $\lceil \frac{M}{2} \rceil + 4$ be such that G has no linear list coloring from L . Note that G must be connected. Suppose G has a 1-vertex u with neighbor v . By minimality, $G - u$ has a linear list coloring from L . Let $L'(u)$ denote the list of colors in $L(u)$ that neither appear on v , nor appear twice in $N(v)$. Note that $|L'(u)| \geq (\lceil \frac{M}{2} \rceil + 4) - (\lfloor \frac{M-1}{2} \rfloor + 1) = 4$. Thus, if G has a 1-vertex u , we can extend a linear list coloring of $G - u$ to G . So we may assume that $\delta(G) \geq 2$. Since G is a planar graph with $\delta(G) \geq 2$ and girth at least 5, G contains one of the two configurations specified in Lemma 1.

Case (RC1): First, suppose that G contains a 2-vertex u adjacent to a 5^- -vertex v . Let w be the other neighbor of u . By minimality, $G - u$ has a linear list coloring from L . In order to avoid creating any 2-colored cycles and to also avoid creating any vertices that have three neighbors with the same color, it is sufficient to avoid coloring u with any color that appears two or more times in $N(v) \cup N(w)$. Furthermore, u must not receive a color used on v or on w . Let $L'(u)$ denote the list of colors in $L(u)$ that may still be used on u . We have $|L'(u)| \geq (\lceil \frac{M}{2} \rceil + 4) - (\lfloor \frac{(M-1)+(5-1)}{2} \rfloor + 2) = (\lceil \frac{M}{2} \rceil + 4) - (\lceil \frac{M}{2} \rceil + 3) = 1$. Thus, we can extend a linear list coloring of $G - u$ to a linear list coloring of G .

Case (RC2): Suppose instead that G contains a 5-face f with four incident 3-vertices and with the fifth incident vertex of degree at most 5. We label the vertices as follows: let u_1, u_2, u_3 , and u_4 denote successive 3-vertices, and let v_2 and v_3 denote the neighbors of u_2 and u_3 not on f .

By minimality, $G - \{u_2, u_3\}$ has a linear list coloring from L . Now we will extend the coloring to u_2 and u_3 . Let $L'(u_2)$ and $L'(u_3)$ denote the colors in $L(u_2)$ and $L(u_3)$ that are still available for use on u_2 and u_3 . When we color u_2 , we clearly must avoid the colors on u_1 and v_2 . We also want to avoid creating a 2-colored cycle or a vertex that has three neighbors with the same color. To do this, it suffices to avoid any color that appears on two or more vertices at distance two from u_2 . This gives us an upper bound on the number of forbidden colors: $2 + \lfloor \frac{(M-1)+2+2}{2} \rfloor = \lceil \frac{M}{2} \rceil + 3$. So $|L'(u_2)| = \lceil \frac{M}{2} \rceil + 4 - (\lceil \frac{M}{2} \rceil + 3) \geq 1$. An analogous count shows that $|L'(u_3)| \geq 1$. However, we might have $L'(u_2) = L'(u_3)$. Thus, we now refine this argument to show that $|L'(u_2)| \geq 2$ or $|L'(u_3)| \geq 2$.

First suppose that $c(u_1) = c(v_2)$. Since the colors on u_1 and v_2 are the same, these two vertices only forbid a single color from use on u_2 , rather than the two colors we accounted for above. Thus we get $|L'(u_2)| \geq 2$. As above, $|L'(u_3)| \geq 1$, so we first color u_3 , then color u_2 with a color not on u_3 . This gives the desired linear coloring of G . Hence, we conclude that $c(u_1) \neq c(v_2)$.

Since $c(u_1) \neq c(v_2)$, when we color u_3 , we need not fear creating three neighbors of u_2 with the same color. Further, we need not worry about giving u_3 the same color as either u_1 or v_2 , for the following reason. Any 2-colored cycle that contains u_3 and either u_1 or v_2 must also contain u_2 and either u_4 or v_3 . Thus, by requiring that u_2 does not get a color that appears on two or more vertices at distance two, we avoid such a 2-colored cycle. So in fact, u_3 only needs to avoid colors that appear on v_3 , on u_4 , or on at least two vertices of $N(u_4) \cup N(v_3)$. This observation gives us $|L'(u_3)| \geq (\lceil \frac{M}{2} \rceil + 4) - (\lfloor \frac{M-1}{2} \rfloor + 2) = (\lceil \frac{M}{2} \rceil + 4) - (\lceil \frac{M}{2} \rceil + 2) = 2$. So we can color u_2 , then color u_3 with a color not on u_2 . This gives the desired linear list coloring, and completes the proof. \square

A similar, but more detailed, argument proves that if G is a planar graph with girth at least 5 and $\Delta(G) \geq 15$, then $lc_\ell(G) \leq \lceil \frac{\Delta(G)}{2} \rceil + 3$. A brief sketch of this proof is as follows. First, we can refine Lemma 1 to show that if $\Delta(G) \geq 15$, then in (RC2) at most two neighbors of u_1, u_2, u_3 , and u_4 can have high degree. (The key insight is that our present argument only requires that each 6^+ -vertex give charge $\frac{2}{3}$ to each incident face; not charge $(\frac{3}{2}d(v) - 5)/d(v)$. Thus, these high degree vertices have lots of extra charge that they can send to adjacent 3-vertices.) With a more careful analysis, we can show that both the original configuration (RC1) and this strengthened version of (RC2) are reducible even with only $\lceil \frac{\Delta(G)}{2} \rceil + 3$ colors.

3. $\text{mad}(G) < 3$ and $\Delta(G) \geq 9$ imply $lc_\ell(G) \leq \lceil \frac{\Delta(G)}{2} \rceil + 2$

Lemma 2. *If G is a graph with $\text{mad}(G) < 3$, $\delta(G) \geq 2$, and $\Delta(G) \geq 9$, then G contains one of the following five configurations:*

- (RC1) a 2-vertex u adjacent to vertices v and w such that $\lceil \frac{d(v)}{2} \rceil + \lceil \frac{d(w)}{2} \rceil < \lceil \frac{\Delta(G)}{2} \rceil + 2$,
- (RC2) a 3-vertex u adjacent to a 2-vertex and to two other vertices v and w , such that $d(v) + d(w) \leq 8$,
- (RC3) a 3-vertex adjacent to two 2-vertices,
- (RC4) a 4-vertex adjacent to four 2-vertices,
- (RC5) a 5-vertex u that is adjacent to four 2-vertices, each of which is adjacent to another 8^- -vertex; and u is also adjacent to a fifth 3^- -vertex.

In fact, the hypothesis $\Delta(G) \geq 9$ cannot be omitted (though the lower bound can possibly be reduced), as we show after we prove the lemma.

Proof. We use discharging, with initial charge $\mu(v) = d(v) - 3$ for each vertex v . Since $\text{mad}(G) < 3$, the sum of the initial charges is negative. Note that only the 2-vertices have negative charge, so we design our discharging rules to pass charge to the 2-vertices. We redistribute the charge via the following three discharging rules:

- (R1) Every 4-vertex gives charge $\frac{1}{3}$ to each adjacent 2-vertex.
- (R2) Every 5-vertex gives charge $\frac{3}{7}$ to each adjacent 2-vertex that is also adjacent to another 8^- -vertex, and it gives charge $\frac{5}{14}$ to every adjacent 3-vertex and every other adjacent 2-vertex.
- (R3) Every 6^+ -vertex v gives charge $\frac{d(v)-3}{d(v)}$ to each adjacent 2-vertex and 3-vertex.
- (R4) Every 3-vertex gives its charge (that it received from rules (R2) and (R3)) to its adjacent 2-vertex (if it has one).

We will show that if G contains none of the five configurations (RC1)–(RC5), then each vertex finishes with nonnegative charge, which is a contradiction. The following observation is an immediate corollary of the fact that G contains no copy of (RC1). We will use this observation below, to show that every vertex finishes with nonnegative charge.

Observation 1. *Suppose that a 2-vertex u has neighbors v and w .*

- (i) *If $d(v) \in \{3, 4\}$, then $d(w) = \Delta(G)$ if $\Delta(G)$ is odd, and $d(w) \geq \Delta(G) - 1$ if $\Delta(G)$ is even.*
- (ii) *If $d(v) \in \{5, 6\}$, then $d(w) \geq \Delta(G) - 2$ if $\Delta(G)$ is odd, and $d(w) \geq \Delta(G) - 3$ if $\Delta(G)$ is even.*

We now use Observation 1 to show that every vertex finishes with nonnegative charge. It is clear from (R3) that every 6^+ -vertex finishes with nonnegative charge. The same is true for 3-vertices. So we consider 4-vertices, 5-vertices, and 2-vertices.

Suppose $d(u) = 4$. Since G contains no copy of (RC4), every 4-vertex u is adjacent to at most three 2-vertices. Thus, we have $\mu^*(u) \geq \mu(u) - 3(\frac{1}{3}) = 1 - 3(\frac{1}{3}) = 0$.

Suppose $d(u) = 5$. If u has two or more neighbors that each receive charge at most $\frac{5}{14}$ from u , then $\mu^*(u) \geq \mu(u) - 3(\frac{3}{7}) - 2(\frac{5}{14}) = 2 - \frac{14}{7} = 0$. Similarly, if u has one neighbor that receives no charge from u , then $\mu^*(u) \geq \mu(u) - 4(\frac{3}{7}) > 0$. Hence, we may assume that u sends charge to each neighbor, and that it sends charge $\frac{3}{7}$ to at least four of its neighbors. However, this assumption implies that G contains a copy of configuration (RC5), which is a contradiction.

Finally, suppose $d(u) = 2$. Let the neighbors of u be v and w . Since $\mu(u) = -1$, it suffices to show that u always receives charge at least 1. If $d(v) \geq 6$ and $d(w) \geq 6$, then v and w each give u charge at least $\frac{1}{2}$. So we may assume that $d(v) \leq 5$. Suppose $d(v) = 5$. Since $\Delta(G) \geq 9$, Observation 1 implies that $d(w) \geq 7$. If $d(w) \in \{7, 8\}$, then u receives charge at least $\frac{3}{7} + \frac{4}{7} = 1$. If $d(w) \geq 9$, then u receives charge at least $\frac{5}{14} + \frac{6}{9} > 1$.




Fig. 1. Configurations (RC3), (RC4), and (RC5) from Lemma 2 and Theorem 3.

If $d(v) = 4$, then Observation 1 implies that $d(w) \geq 9$, so u receives charge at least $\frac{1}{3} + \frac{6}{9} = 1$. If $d(v) = 3$, then the absence of (RC2) implies that at least one neighbor x of v has degree at least 5, so v receives charge at least $\frac{5}{14}$ from x . Since v can have at most one adjacent 2-vertex, u gets charge at least $\frac{5}{14}$ from v . Hence, the total charge that u receives is at least $\frac{6}{9} + \frac{5}{14} > 1$. \square

Now we give two examples to show that the hypothesis $\Delta(G) \geq 9$, in Lemma 2 above, cannot be omitted. (We do suspect, however, that this hypothesis can be replaced by $\Delta(G) \geq 7$, or perhaps even by $\Delta(G) \geq 5$.) We first give an example with maximum degree 3. Let G be the dodecahedron, and let E be a matching in G of size 6, such that every face of G contains one edge of E . Form \widehat{G} from G by subdividing each edge of the matching. The girth of \widehat{G} is 6, so (by an easy application of Euler’s formula), $\text{mad}(\widehat{G}) < 3$. Despite having $\text{mad}(\widehat{G}) < 3$, \widehat{G} does not contain any of the five configurations (RC1)–(RC5) in Lemma 2. Now we give an example with maximum degree 4. Let G be the octahedron, and let E be a perfect matching in G . Form \widehat{G} from G by subdividing every edge of G except the three edges of E . The average degree of \widehat{G} is $(4 \times 6 + 2 \times 9)/(6 + 9) = \frac{14}{5}$; it is an easy exercise to verify that $\text{mad}(\widehat{G}) = \frac{14}{5}$. Again \widehat{G} contains none of the configurations (RC1)–(RC5).

Now we use Lemma 2 to prove the following linear list coloring result, which immediately implies Theorem 1(2).

Theorem 3. *Let $M \geq 9$ be an integer. If G is a graph with $\text{mad}(G) < 3$ and $\Delta(G) \leq M$, then $\text{lc}_\ell(G) \leq \lceil \frac{M}{2} \rceil + 2$.*

Proof. Suppose the theorem is false. Let G be a minimal counterexample and let the list assignment L of size $\lceil \frac{M}{2} \rceil + 2$ be such that G has no linear list coloring from L . Since $M \geq 9$, we have $|L(v)| = \lceil \frac{M}{2} \rceil + 2 \geq 7$ for every $v \in V$. Note that G must be connected. Suppose G has a 1-vertex u with neighbor v . By minimality, $G - u$ has a linear list coloring from L . Let $L'(u)$ denote the list of colors in $L(u)$ that neither appear on v , nor appear twice in $N(v)$. Note that $|L'(u)| \geq (\lceil \frac{M}{2} \rceil + 2) - (\lfloor \frac{M-1}{2} \rfloor + 1) = 2$. Thus, if G has a 1-vertex u , we can extend a linear list coloring of $G - u$ to G . So we may assume that $\delta(G) \geq 2$.

Since G is a graph with $\delta(G) \geq 2$ and $\text{mad}(G) < 3$, G contains one of the five configurations (RC1)–(RC5) specified in Lemma 2. We consider each of these five configurations in turn, and in each case we construct a linear coloring of G from L . *Case (RC1):* Suppose that G contains configuration (RC1). Let u be a 2-vertex adjacent to vertices v and w such that $\lceil \frac{d(v)}{2} \rceil + \lceil \frac{d(w)}{2} \rceil < \lceil \frac{M}{2} \rceil + 2$. By the minimality of G , subgraph $G - u$ has a linear list coloring c .

If $c(v) \neq c(w)$, then u can receive any color except for $c(v)$, $c(w)$, and those colors that appear twice on $N(v)$ or twice on $N(w)$. So the number of colors forbidden is at most $2 + \lfloor \frac{d(v)-1}{2} \rfloor + \lfloor \frac{d(w)-1}{2} \rfloor = \lceil \frac{d(v)}{2} \rceil + \lceil \frac{d(w)}{2} \rceil$. Since $|L(u)| = \lceil \frac{M}{2} \rceil + 2$, and since $\lceil \frac{d(v)}{2} \rceil + \lceil \frac{d(w)}{2} \rceil < \lceil \frac{M}{2} \rceil + 2$, we can extend the coloring to u . So we assume instead that $c(v) = c(w) = 1$.

If $c(v) = c(w)$, then (similar to that above), u can receive any color except for $c(v)$ and those colors that appear twice on $N(v) \cup N(w)$. The number of forbidden colors is at most $1 + \lfloor \frac{(d(v)-1)+(d(w)-1)}{2} \rfloor \leq \lceil \frac{d(v)}{2} \rceil + \lceil \frac{d(w)}{2} \rceil$. So, once again, we can extend the coloring to u .

Case (RC2): Suppose that G contains configuration (RC2). Let u be a 3-vertex adjacent to a 2-vertex and to two other neighbors v and w with $d(v) + d(w) \leq 8$. By the minimality of G , subgraph $G - u$ has a linear list coloring from L . If all three neighbors of u have the same color, then we will not get a linear coloring of G no matter how we color u . In this case, we can recolor the 2-vertex and still have a linear coloring of $G - u$. Now we will extend the coloring to u .

Let $L'(u)$ denote the colors in $L(u)$ that are still available for use on u . When we color u , we clearly must avoid the colors on its three neighbors. We also want to avoid creating a 2-colored cycle or a vertex that has three neighbors with the same color. To do this, it suffices to avoid any color that appears on two or more vertices at distance two from u . This gives us an upper bound on the number of forbidden colors: $3 + \lfloor \frac{(d(v)-1)+(d(w)-1)+1}{2} \rfloor = 3 + \lfloor \frac{d(v)+d(w)-1}{2} \rfloor \leq 3 + \lfloor \frac{7}{2} \rfloor = 6$. Since $|L(u)| \geq 7$, we have $|L'(u)| \geq 1$. Thus, we can extend the coloring to u .

Case (RC3): Suppose that G contains configuration (RC3), shown in Fig. 1. Let u be a 3-vertex that has neighbors v_1, v_2 , and v_3 with $d(v_1) = d(v_2) = 2$ and $d(v) = 3$. Let $N(v_i) = \{w_i, u\}$ for $i \in \{1, 2\}$. By the minimality of G , subgraph $G - \{u, v_1, v_2\}$ has a linear list coloring c from L . For each uncolored vertex $z \in \{u, v_1, v_2\}$, let $L'(z)$ denote the colors in $L(z)$ that are still available for use on z . Note that $|L'(z)| \geq 2$ for each uncolored vertex z .

Suppose that $L'(u) = \{c(w_1), c(w_2)\}$; this means that $c(v_3) \notin \{c(w_1), c(w_2)\}$. Color u with $c(w_1)$. Now choose $c(v_1) \in L'(v_1) - c(v_3)$ and $c(v_2) \in L'(v_2) - c(w_1)$. This is a valid linear coloring of G .

Suppose instead that $L'(u) \setminus \{c(w_1), c(w_2)\} \neq \emptyset$. Choose $c(u) \in L'(u) - \{c(w_1), c(w_2)\}$, choose $c(v_1) \in L'(v_1) - c(u)$, and choose $c(v_2) \in L'(v_2) - c(u)$. This coloring is proper and contains no 2-alternating path through u . Hence, it is a linear coloring unless $c(v_1) = c(v_2) = c(v_3)$. If no other choice of $c(v_1)$ and $c(v_2)$ can avoid this problem, then we can conclude that $L'(v_1) = L'(v_2) = \{c(v_3), c_1\}$ (for some color c_1); further $L'(u) - \{c(w_1), c(w_2)\} = \{c_1\}$. Suppose we are in this case.

If $c(w_1) \neq c(w_2)$, then, without loss of generality, $L'(u) = \{c(w_1), c_1\}$. Now let $c(u) = c(w_1)$, $c(v_1) = c_1$, and $c(v_2) = c(v_3)$. This is a valid linear coloring. So, by relabeling, we may assume that $c(w_1) = c(w_2) = 1$, $c(v_3) = 2$, and $c_1 = 3$. Thus $L'(v_1) = L'(v_2) = \{2, 3\}$ and $L'(u) = \{1, 3\}$.

Note that $\{2, 3\} \subseteq L'(v_i)$ implies that 2 and 3 each appear at most once in $N(w_i)$ (for $i \in \{1, 2\}$). If 3 does not appear on both $N(w_1)$ and $N(w_2)$, then let $c(v_1) = 2$ and $c(v_2) = 3$ and $c(u) = 1$. If 2 does not appear on both $N(w_1)$ and $N(w_2)$, then let $c(u) = 1$, $c(v_1) = 2$, $c(v_2) = 3$ (or $c(u) = 1$, $c(v_1) = 3$, $c(v_2) = 2$). So, we can assume that 2 and 3 each appear once on both $N(w_1)$ and $N(w_2)$. However, now $|L'(v_i)| \geq (\lceil \frac{M}{2} \rceil + 2) - (\lfloor \frac{M-3}{2} \rfloor + 1) \geq 3$, which is a contradiction.

Case (RC4): Suppose that G contains configuration (RC4), shown in Fig. 1. Let u be a 4-vertex and let $N(u) = \{v_i : 1 \leq i \leq 4 \text{ such that } d(v_i) = 2\}$. Also let $N(v_i) = \{u, w_i\}$ for $1 \leq i \leq 4$. By the minimality of G , subgraph $G - \{u, v_1, v_2, v_3, v_4\}$ has a linear list coloring from L . For each uncolored vertex z , let $L'(z)$ denote the list of colors still available for z . Note that $|L'(v_i)| \geq 2$ and $|L'(u)| = |L(u)| = \lceil \frac{M}{2} \rceil + 2 \geq 7$, since $M \geq 9$.

We can color the v_i 's from their lists so that every color is used on at most two v_i 's, as follows. If some color c is available for use on two or more v_i 's, then use c on exactly two of them, and color each of the remaining v_i 's with another color (which could be the same for both of them). Otherwise, all the v_i 's have disjoint lists of available colors, so color them arbitrarily.

If the four colors on the v_i 's are all distinct, then color u with a fifth color. If $c(v_1) = c(v_2)$ but $c(v_1), c(v_3)$, and $c(v_4)$ are all distinct, then choose $c(u)$ so that $c(u) \notin \{c(v_1), c(v_3), c(v_4), c(w_1)\}$. Finally, if $c(v_1) = c(v_2)$ and $c(v_3) = c(v_4)$ (which together imply $c(v_1) \neq c(v_3)$), then choose $c(u)$ so that $c(u) \notin \{c(v_1), c(v_3), c(w_1), c(w_3)\}$.

Case (RC5): Suppose that G contains configuration (RC5), shown in Fig. 1. Let u be a 5-vertex and let $N(u) = \{v_i : 1 \leq i \leq 5\}$, such that $d(v_i) = 2$ for $1 \leq i \leq 4$ and $d(v_5) \leq 3$. Also let $N(v_i) = \{u, w_i\}$ for $1 \leq i \leq 4$, where $d(w_i) \leq 8$. By the minimality of G , subgraph $G - \{u, v_1, v_2, v_3, v_4\}$ has a linear coloring c from L . For each uncolored vertex $z \in \{u, v_1, v_2, v_3, v_4\}$, let $L'(z)$ denote the list of colors still available for z . Since $d(w_i) \leq 8$, we have $|L'(v_i)| \geq 3$. Conversely, $|L'(u)| \geq \lceil \frac{M}{2} \rceil + 2 - (\lfloor \frac{2}{2} \rfloor + 1) = \lceil \frac{M}{2} \rceil \geq 5$, since $M \geq 9$. Now let $L''(v_i) = L'(v_i) - c(v_5)$; note that $|L''(v_i)| \geq 2$. We now extend the coloring by using the lists $L'(u)$ and $L''(v_i)$. We can completely ignore v_5 (since we deleted $c(v_5)$ from the lists), so the analysis is exactly the same as in Case (RC4). □

As we explained in the Introduction, this theorem immediately yields the following corollary.

Corollary 1. *If graph G is planar, has girth at least 6, and $\Delta(G) \geq 9$, then $lc_\ell(G) \leq \lceil \frac{\Delta(G)}{2} \rceil + 2$.*

Although our proof of Theorem 3 relies heavily on the hypothesis $\Delta(G) \geq 9$, we suspect that the Theorem is true even when this hypothesis is removed. Namely, we conjecture that every graph G with $\text{mad}(G) < 3$ satisfies $lc_\ell(G) \leq \lceil \frac{\Delta(G)}{2} \rceil + 2$. If true, this result is best possible, as shown by the graph $K_{3,3}$, since $lc_\ell(K_{3,3}) = 5$. Furthermore, every graph G with $K_{3,3} \subseteq G$, $\text{mad}(G) = 3$, and $\Delta(G) \in \{3, 4\}$ shows that this result is best possible.

4. $\text{mad}(G) < \frac{12}{5}$ implies $lc_\ell(G) = \lceil \frac{\Delta(G)}{2} \rceil + 1$

In this section, we prove that if G is a graph with $\Delta(G) \geq 3$ and $\text{mad}(G) < \frac{12}{5}$, then $lc_\ell(G) = \lceil \frac{\Delta(G)}{2} \rceil + 1$. For such graphs, we prove an upper bound that matches the trivial lower bound $lc_\ell(G) \geq \lceil \frac{\Delta(G)}{2} \rceil + 1$. Recall (from the Introduction) that our bound on $\text{mad}(G)$ is best possible, as demonstrated by $K_{2,3}$, since $\text{mad}(K_{2,3}) = \frac{12}{5}$ and $lc_\ell(K_{2,3}) > \lceil \frac{\Delta(K_{2,3})}{2} \rceil + 1$.

Lemma 3. *If G is a graph with $\text{mad}(G) < \frac{12}{5}$ and $\delta(G) \geq 2$, then G contains one of the following four configurations:*

- (RC1) a 3^+ -thread,
- (RC2) a 3-vertex v incident to two 1^+ -threads and one 2-thread, such that the vertex at distance two from v along each 1^+ -thread is a 3^- -vertex,
- (RC3) adjacent 3-vertices with at least seven 2-vertices in their incident threads,
- (RC4) a path of three vertices uwv with $d(u) = d(w) = d(v) = 3$ such that w is incident to a 2-thread and u and v are each incident to two 2-threads.

Proof. We use discharging, with initial charge $\mu(v) = d(v) - \frac{12}{5}$ for each vertex v . Since $\text{mad}(G) < \frac{12}{5}$, the sum of the initial charges is negative. We use the following three discharging rules:

- (R1) Every 2-vertex gets charge $\frac{1}{5}$ from each of the endpoints of its thread.
- (R2) Every 3-vertex incident to two 2-threads gets charge $\frac{1}{5}$ from its 3^+ -neighbor.
- (R3) Every 3-vertex incident to a 1-thread gets charge $\frac{1}{5}$ from the other endpoint of the 1-thread if it is a 4^+ -vertex.

Now we will show that if G contains none of the configurations (RC1)–(RC4), then every vertex finishes with nonnegative charge, which is a contradiction. If $d(v) = 2$, then $\mu^*(v) = d(v) - \frac{12}{5} + 2(\frac{1}{5}) = 0$. If $d(v) \geq 4$, then, since G contains no




Fig. 2. Configurations (RC2), (RC3), and (RC4) from Lemma 3 and Theorem 4.

3^+ -threads (by (RC1)), v gives away charge $\frac{1}{5}$ to each of at most $2d(v)$ 2-vertices. Note further that if v gives away charge $\frac{1}{5}$ to t 3-vertices via (R2) and/or (R3), for some constant t , then v gives away charge $\frac{1}{5}$ to at most $2d(v) - t$ 2-vertices. Thus, we have $\mu^*(v) \geq d(v) - \frac{12}{5} - \frac{1}{5}(2d(v)) = \frac{3}{5}(d(v) - 4) \geq 0$. So we only need to consider 3-vertices.

Let $d(v) = 3$. Suppose v has at most three 2-vertices in its incident threads. If v does not give away charge by (R2), then v gives away charge at most $3(\frac{1}{5})$, so $\mu^*(v) \geq 3 - \frac{12}{5} - 3(\frac{1}{5}) = 0$. If v does give charge by (R2), then, since G contains no copy of (RC3), v has at most two 2-vertices in its incident threads. Thus v gives away charge at most $3(\frac{1}{5})$, unless both v is incident to a 2-thread and also v gives away charge by (R2) to two distinct vertices. However, this situation cannot occur, since it implies that G contains a copy of (RC4), which is a contradiction.

Suppose instead that v has at least four 2-vertices in its incident threads. Since G contains no copy of (RC2), either v is incident to two 2-threads and also adjacent to a 3^+ -vertex, or v is incident to two 1-threads and one 2-thread and the other end of at least one 1-thread is a 4^+ -vertex. In each case, v gives away charge $4(\frac{1}{5})$ and receives charge at least $\frac{1}{5}$, so $\mu^*(v) \geq 3 - \frac{12}{5} - 4(\frac{1}{5}) + \frac{1}{5} = 0$. \square

Now we use Lemma 3 to prove the following linear list coloring result.

Theorem 4. Let $M \geq 3$ be an integer. If G is a graph with $\text{mad}(G) < \frac{12}{5}$ and $\Delta(G) \leq M$, then $\text{lc}_\ell(G) = \lceil \frac{M}{2} \rceil + 1$.

Proof. Suppose the theorem is false. Let G be a minimal counterexample and let list assignment L , of size $\lceil \frac{M}{2} \rceil + 1$, be such that G has no linear list coloring from L . Since $M \geq 3$, we have $|L(v)| = \lceil \frac{M}{2} \rceil + 1 \geq 3$ for all $v \in V$. Note that G must be connected. Suppose that G contains a 1-vertex u with neighbor v . By the minimality of G , subgraph $G - \{u\}$ has a linear list coloring from L . Let $L'(u)$ denote the list of colors in $L(u)$ that neither appear on v nor appear twice in $N(v)$. Note that $|L'(u)| \geq (\lceil \frac{M}{2} \rceil + 1) - \lfloor \frac{M-1}{2} \rfloor - 1 \geq 1$. Thus, if G has a 1-vertex u , we can extend a linear list coloring of $G - u$ to G . So we may assume that $\delta(G) \geq 2$.

Since G has $\delta(G) \geq 2$ and $\text{mad}(G) < \frac{12}{5}$, G contains one of the four configurations specified in Lemma 3. We consider each of these four configurations in turn, and in each case we construct a linear coloring of G from L .

Case (RC1): Suppose that G contains (RC1): a 3^+ -thread. Let u, u_1, u_2, u_3, u_4 be part of the thread, that is, $d(u) \geq 3$, $d(u_1) = d(u_2) = d(u_3) = 2$, and $d(u_4) \geq 2$. By the minimality of G , subgraph $G - \{u_2\}$ has a linear coloring from L . If $c(u_1) = c(u_3)$, then $|L(u_2)| \geq 2$, so we choose $c(u_2) \in L(u_2) - \{c(u)\}$. If $c(u_1) \neq c(u_3)$, then $|L(u_2)| \geq 1$, so we choose $c(u_2) \in L(u_2)$. Note that either $c(u_2) \neq c(u)$ or $c(u_1) \neq c(u_3)$, so we have not created a 2-colored cycle.

Case (RC2): Suppose instead that G contains (RC2), shown in Fig. 2. Let u be a 3-vertex that is incident to one 2-thread u, u_1, u'_1, u''_1 with $d(u'_1) \geq 3$ and incident to two 1^+ -threads u, u_2, u'_2 and u, u_3, u'_3 with $2 \leq d(u'_2) \leq 3$ and $2 \leq d(u'_3) \leq 3$. By the minimality of G , subgraph $G - \{u, u_1, u_2, u_3\}$ has a linear coloring from L . Now we will extend the coloring to G .

For each uncolored vertex $z \in \{u, u_1, u_2, u_3\}$, let $L'(z)$ denote the colors in $L(z)$ that are still available for use on z . When we extend the coloring, we obviously must get a proper coloring. In addition, we must avoid creating 2-colored cycles and avoid creating vertices with 3 neighbors of the same color. Note that $|L'(u_1)| \geq 2$, $|L'(u_2)| \geq 1$, and $|L'(u_3)| \geq 1$.

Suppose $|L'(u_2) \cup L'(u_3)| \geq 2$. We choose $c(u_2) \in L'(u_2)$ and $c(u_3) \in L'(u_3)$ such that $c(u_2) \neq c(u_3)$. Next we choose $c(u) \in L'(u) - \{c(u_2), c(u_3)\}$. If $c(u) \neq c(u'_1)$, then we choose $c(u_1) \in L'(u_1) - \{c(u)\}$. If instead $c(u) = c(u'_1)$, then we choose $c(u_1) \in L'(u_1) - \{c(u'_1)\}$. This gives a valid linear coloring.

Suppose instead that $|L'(u_2) \cup L'(u_3)| = 1$. Thus $L'(u_2) = L'(u_3) = \{a\}$, for some color a . Clearly, we must choose $c(u_2) = c(u_3) = a$. Note that this happens only if both $d(u'_2) = d(u'_3) = 3$ and the two other neighbors of u'_2 (and u'_3) have the same color. Now we choose $c(u_1) \in L(u_1) - \{a, c(u'_1)\}$ and $c(u) \in L(u) - \{a\}$.

Since $c(u_1) \neq a$, we have not created any vertex with 3 neighbors of the same color, and we have not created any 2-colored cycle passing through u_1 . Since $c(u_2)$ does not appear on the other neighbors of u'_2 , we have not created any 2-colored cycle passing through u_2 .

Case (RC3): Now suppose instead that G contains (RC3): two adjacent 3-vertices with at least seven 2-vertices in their incident threads (shown in Fig. 2). We label the vertices as follows: let u and v be the adjacent 3-vertices, u is incident to two 2-threads u, u_1, u'_1, u''_1 and u, u_2, u'_2, u''_2 and v is incident to one 2-thread v, v_1, v'_1, v''_1 and one 1^+ -thread v, v_2, v'_2 .

By the minimality of G , subgraph $G - \{u, v, u_1, u_2, v_1\}$ has a linear coloring from L . Now we will extend the coloring to G . For each vertex $z \in \{u, v, u_1, u_2, v_1\}$, let $L'(z)$ denote the colors in $L(z)$ that are still available for use on z . When we

extend the coloring, we obviously must get a proper coloring. In addition, we must avoid creating 2-colored cycles and avoid creating vertices with 3 neighbors of the same color. Note that $|L'(u_1)| \geq 2$, $|L'(u_2)| \geq 2$, $|L'(v_1)| \geq 2$, $|L'(u)| \geq 3$, and $|L'(v)| \geq 3$; we may assume that equality holds in each case.

Since $|L'(u)| = 3 > 2 = |L'(u_1)|$, we can choose $c(u) \in L'(u) - L'(u_1)$. If $c(u) = c(v_2)$, then choose $c(v_1) \in L'(v_1) - \{c(u)\}$ and $c(v) \in L'(v) - \{c(v_1)\}$. If instead $c(u) \neq c(v_2)$, then choose $c(v) \in L'(v) - \{c(u)\}$.

Now if $c(v) \neq c(v'_1)$, then choose $c(v'_1) \in L'(v_1) - \{c(v)\}$; if $c(v) = c(v'_1)$, then choose $c(v'_1) \in L'(v_1) - \{c(v'_1)\}$. Next, choose $c(u_1) \in L'(u_1) - \{c(v)\}$. Finally, if $c(u) = c(u'_2)$, then choose $c(u_2) \in L'(u_2) - \{c(u'_2)\}$; otherwise, choose $c(u_2) \in L'(u_2) - \{c(u)\}$.

Recall that $c(u_1) \neq c(v)$ and either $c(u) \neq c(v_2)$ or $c(v_1) \neq c(v_2)$; thus, we do not create any vertices with three neighbors of the same color. By construction, we have no 2-colored cycles through u_2 or v_1 . Further, $c(u_1) \neq c(v)$, so we do not create any 2-colored cycles.

Case (RC4): Suppose that G contains (RC4). We label the vertices as follows: let u, w, v be the path; let u, u_1, u'_1, u''_1 and u, u_2, u'_2, u''_2 be the 2-threads incident to u ; let v, v_1, v'_1, v''_1 and v, v_2, v'_2, v''_2 be the 2-threads incident to v ; and let w, w_1, w'_1, w''_1 be the 2-thread incident to w .

By the minimality of G , subgraph $G - \{u, u_1, u_2, v, v_1, v_2, w, w_1\}$ has a linear coloring from L . Now we will extend the coloring to G . For each vertex $z \in \{u, u_1, u_2, v, v_1, v_2, w, w_1\}$, let $L'(z)$ denote the colors in $L(z)$ that are still available for use on z . When we extend the coloring, we obviously must get a proper coloring. In addition, we must avoid creating 2-colored cycles and avoid creating vertices with 3 neighbors of the same color. We will show explicitly how to color u, u_1, u_2, w , and w_1 (and we will color v, v_1 , and v_2 , analogously). We consider two subcases. In fact, we may have one “side” (u, u_1, u'_1, u_2 , and u'_2) that is in Subcase (i) and the other side that is in Subcase (ii); this is not a problem, since we color the sides independently.

Subcase (i): Suppose that $c(u'_1) = c(u'_2)$. If $c(u'_1) \notin L'(u)$, then we can choose $c(u_1) \in L'(u_1)$ and $c(u_2) \in L'(u_2)$ such that $c(u_1) \neq c(u_2)$, and afterward we choose $c(u) \in L'(u) - \{c(u'_1), c(u_1), c(u_2)\}$. If $c(u'_1) \in L'(u)$, then let $c(u) = c(u'_1)$. Choose $c(v)$ analogously. In this instance, we wait to choose $c(u_1)$ and $c(u_2)$ until after we choose $c(w)$.

If $c(u) = c(v)$, then choose $c(w_1) \in L'(w_1) - \{c(u)\}$ and $c(w) \in L'(w) - \{c(w_1), c(u)\}$. If $c(u) \neq c(v)$, then choose $c(w) \in L'(w) - \{c(u), c(v)\}$ and $c(w_1) \in L'(w_1) - \{c(w)\}$. Finally, choose $c(u_1) \in L'(u_1) - \{c(u), c(u'_1)\}$ and $c(u_2) \in L'(u_2) - \{c(u), c(w)\}$ (if we have not chosen these colors yet; recall that $c(u) = c(u'_1)$, so $c(u_1) \neq c(u)$; analogously, $c(u_2) \neq c(w)$).

Subcase (ii): $c(u'_1) \neq c(u'_2)$. Choose $c(u) \in L'(u) - \{c(u'_1), c(u'_2)\}$. Choose $c(v)$ analogously. Now color w and w_1 as above. Finally, we will color u_1, u_2, v_1 , and v_2 , as below.

If we can, we choose $c(u_1) \in L'(u_1) - \{c(u)\}$, and $c(u_2) \in L'(u_2) - \{c(u)\}$ such that either $c(u_1) \neq c(w)$ or $c(u_2) \neq c(w)$. If this is impossible, then $L'(u_1) = L'(u_2) = \{c(u), c(w)\}$; furthermore, $L(u) = \{c(u), c(u'_1), c(u'_2)\}$. Now let $c(u_1) = c(u_2) = c(u)$ and recolor u with a new color in $L'(u) - \{c(u_1), c(w_1), c(w)\}$ (note that $c(w) \notin L'(u)$). Finally, color v_1, v_2 , and v analogously.

It is clear that we have created a proper coloring. It is also straightforward to verify that we did not create any vertices with 3 neighbors of the same color, and we did not create any 2-colored cycles. \square

This theorem immediately yields the following corollary.

Corollary 2. *If graph G is planar with girth at least 12 and $\Delta(G) \geq 3$, then $lc_\ell(G) = \lceil \frac{\Delta(G)}{2} \rceil + 1$.*

References

- [1] P. Erdős, A.L. Rubin, H. Taylor, Choosability in graphs, *Congr. Numer.* 26 (1979) 125–157.
- [2] L. Esperet, M. Montassier, A. Raspaud, Linear choosability of graphs, *Discrete Math.* 308 (2008) 3938–3950.
- [3] B. Grünbaum, Acyclic colorings of planar graphs, *Israel J. Math.* 14 (1973) 390–408.
- [4] H. Hind, M. Molloy, B. Reed, Colouring a graph frugally, *Combinatorica* 17 (4) (1997) 469–482.
- [5] T.R. Jensen, B. Toft, *Graph Coloring Problems*, John Wiley & Sons, New York, 1995.
- [6] A. Raspaud, W. Wang, Linear coloring of planar graphs with large girth, *Discrete Math.* 309 (2009) 5678–5686.
- [7] V.G. Vizing, Coloring the vertices of a graph in prescribed colors, *Metody Diskret. Analiz.* 19 (1976) 3–10 (in Russian).
- [8] R. Yuster, Linear coloring of graphs, *Discrete Math.* 185 (1998) 293–297.