

Coagulation, diffusion and the continuous Smoluchowski equation

Mohammad Reza Yaghouti^a, Fraydoun Rezakhanlou^{b,*}, Alan Hammond^c

^a *Mathematics and Computer Science Faculty, Amirkabir University, Iran*

^b *Department of Mathematics, University of California Berkeley, Evans Hall, 94720-384 Berkeley, CA, United States*

^c *Courant Institute, New York University, United States*

Received 11 November 2008; received in revised form 5 April 2009; accepted 8 April 2009

Available online 21 April 2009

Abstract

The Smoluchowski equations are a system of partial differential equations modelling the diffusion and binary coagulation of a large collection of tiny particles. The mass parameter may be indexed either by positive integers or by positive reals, these corresponding to the discrete or the continuous form of the equations. For dimension $d \geq 3$, we derive the continuous Smoluchowski PDE as a kinetic limit of a microscopic model of Brownian particles liable to coalesce, using a method similar to that used to derive the discrete form of the equations in [A. Hammond, F. Rezakhanlou, The kinetic limit of a system of coagulating Brownian particles, *Arch. Ration. Mech. Anal.* 185 (2007) 1–67]. The principal innovation is a correlation-type bound on particle locations that permits the derivation in the continuous context while simplifying the arguments of the cited work. We also comment on the scaling satisfied by the continuous Smoluchowski PDE, and its potential implications for blow-up of solutions of the equations.

© 2009 Elsevier B.V. All rights reserved.

MSC: primary 74A25; secondary 60K35

Keywords: Continuous Smoluchowski equation; Coagulating Brownian particles

1. Introduction

It is a common practice in statistical mechanics to formulate a microscopic model with simple dynamical rules in order to study a phenomenon of interest. In a colloid, a population of

* Corresponding author. Tel.: +1 510 642 6550; fax: +1 510 642 8204.

E-mail address: rezakhan@Math.Berkeley.EDU (F. Rezakhanlou).

comparatively massive particles is agitated by the bombardment of much smaller particles in the ambient environment: the motion of the colloidal particles may then be modelled as Brownian motion. Smoluchowski’s equation provides a macroscopic description for the evolution of the cluster densities in a colloid whose particles are prone to binary coagulation. Smoluchowski’s equation comes in two flavours: discrete and continuous. In the discrete version, the cluster mass may take values in the set of positive integers, whereas in the continuous version, the cluster mass takes values in \mathbb{R}^+ . Writing $f_n(x, t) = f(x, n, t)$ for the density of clusters (or particles) of size n , this density evolves according to

$$\frac{\partial f_n}{\partial t} = d(n)\Delta f_n(x, t) + Q_+^n(f)(x, t) - Q_-^n(f)(x, t), \tag{1.1}$$

where

$$Q_+^n(f) = \frac{1}{2} \int_0^n \beta(m, n - m) f_m f_{n-m} dm, \tag{1.2}$$

and

$$Q_-^n(f) = \int_0^\infty \beta(m, n) f_m f_n dm, \tag{1.3}$$

in the case of the continuous Smoluchowski equation. (See [8] for some results on the existence of weak solutions of (1.1).) In the discrete case, the integrations in (1.2) and (1.3) are replaced with summations.

In [4,5], we derived the discrete Smoluchowski equation as a many-particle limit of a microscopic model of coagulating Brownian particles. (See also [7,9] and [2] for similar results.) The main purpose of the present article is the derivation of (1.1) for the continuous case. We introduce an approach that is simpler than that used in [4,5]. We will present a robust argument that allows us to circumvent some induction-based steps of [4,5] (which anyway could not be applied in the continuous case). Thus, an auxiliary purpose of this article is to present a shorter proof of the kinetic limit derivations of Smoluchowski’s equation given in [4,5]. The main technical tool is a correlation-type bound on the particle distribution that seems to be applicable to general systems of Brownian particles. To explain this further, we need to sketch the derivation of Smoluchowski’s equation and explain the essential role of the correlation bounds.

The microscopic model that we study in this article consists of a large number of particles which move according to independent Brownian motions whose diffusion rates $2d(m)$ depend on their mass $m \in (0, \infty)$. Any pair of particles that approach to within a certain range of interaction are liable to coagulate, at which time they disappear from the system, to be replaced by a particle whose mass is equal to the sum of the masses of the colliding particles, and whose location is a specific point in the vicinity of the location of the coagulation. This range of interaction is taken to be equal to a parameter ϵ , whose dependence on the mean initial total number N of particles is given by $N = k_\epsilon Z$ for a constant Z , where

$$k_\epsilon = \begin{cases} \epsilon^{2-d} & \text{if } d \geq 3, \\ |\log \epsilon| & \text{if } d = 2. \end{cases}$$

This choice will ensure that a particle experiences an expected number of coagulations in a given unit of time that remains bounded away from zero and infinity as N is taken to be high.

Our main result is conveniently expressed in terms of empirical measures on the locations $x_i(t)$ and the masses $m_i(t)$ of particles. We write $g^\epsilon(dx, dn, t)$ for the measure on $\mathbb{R}^d \times [0, \infty)$

given by

$$g^\varepsilon(dx, dn, t) = k_\varepsilon^{-1} \sum_i \delta_{(x_i(t), m_i(t))}(dx, dn).$$

Our goal is to show that, in the low ε limit, the measure g^ε converges to $f_n(x, t) dx dn$, where f_n solves the system (1.1). The main step in the proof requires the replacement of the microscopic coagulation propensity $\alpha(n, m)$ (that we will shortly describe precisely) of particles of masses n and m with its macroscopic analogue $\beta(n, m)$. The main technical tool for this is a correlation bound which reads as follows, in the case where the coefficient $d(m)$ is non-increasing in m :

$$\begin{aligned} &\mathbb{E} \int_0^\infty \sum_{i_1, \dots, i_k} K(x_{i_1}(t), \dots, x_{i_k}(t)) \prod_{r=1}^k d(m_{i_r}(t))^{\frac{d}{2}} m_{i_r}(t) dt \\ &\leq \text{const.} \mathbb{E} \sum_{i_1, \dots, i_k} \hat{K}(x_{i_1}(0), m_{i_1}(0), \dots, x_{i_k}(0), m_{i_k}(0)) \prod_{r=1}^k d(m_{i_r}(0))^{\frac{d}{2}} m_{i_r}(0). \end{aligned} \tag{1.4}$$

Here, \mathbb{E} denotes the expectation with respect to the underlying randomness, $K : (\mathbb{R}^d)^k \rightarrow \mathbb{R}$ is any non-negative bounded continuous function, and $\hat{K} = -(d(m_{i_1})\Delta_{x_{i_1}} + \dots + d(m_{i_k})\Delta_{x_{i_k}})^{-1} K$. We refer the reader to Section 3 for the corresponding correlation inequality when the function $d(\cdot)$ is not non-increasing.

In fact, we need (1.4) only for certain examples of K with $k = 2, 3$ and 4. It was these examples that were treated in [4,5] with rather ad hoc arguments based on an inductive procedure on the mass of the particles. Those arguments seem to be specific to the discrete case and cannot be generalized to the continuous setting. Moreover, the bound (1.4) implies that the macroscopic particle densities belong to L^p for given $p \geq 2$, provided that a similar bound is valid initially. This rather straightforward consequence of (1.4) is crucial for the derivation of the macroscopic equation. The corresponding step in [4,5] is also carried out with a method that is very specific to the discrete case and does not apply to the continuous setting. This important consequence of (1.4) simplifies the proof drastically and renders the whole of Section 4 of [4] redundant.

We state and prove our results for when the dimension is at least 3. However, our proof for the correlation bound (1.4) works in any dimension, and an interested reader may readily check that, as in this article, the approach of [5] may be modified to establish Theorem 1.1 in dimension two.

We continue with the description of the microscopic model and the statement of the main result.

As a matter of convenience, we introduce two different microscopic models, that differ only in whether the number of particles is initially deterministic or random. We will refer to the model as deterministic or random accordingly. In either case, we define a sequence of microscopic models, indexed by a positive integer N .

A countable set I of symbols is provided. A configuration \mathbf{q} is an $\mathbb{R}^d \times (0, \infty)$ -valued function on a finite subset $I_{\mathbf{q}}$ of I . For any $i \in I_{\mathbf{q}}$, the component $q(i)$ may be written as (x_i, m_i) . The particle labelled by i has mass m_i and location x_i .

In the deterministic case, the index N of the model specifies the total number of particles present at time zero. Their placement is given as follows. There is a given function $h : \mathbb{R}^d \times (0, \infty) \rightarrow [0, \infty)$, with $h_n(x) := h(x, n)$, where $\int_0^\infty \int_{\mathbb{R}^d} h(x, n) dx dn < \infty$. We set $Z = \int_0^\infty \int_{\mathbb{R}^d} h_n(x) dx dn \in (0, \infty)$ and choose N points in $(0, \infty) \times \mathbb{R}^d$ independently according to a law whose density at (x, n) is equal to $h_n(x)/Z$. Selecting arbitrarily a set of N symbols

$\{i_j : j \in \{1, \dots, N\}\}$ from I , we define the initial configuration $\mathbf{q}(0)$ by insisting that $q_{i_j}(0)$ is equal to the j th of the randomly chosen members of $(0, \infty) \times \mathbb{R}^d$.

In the random case, the index N gives the mean number of initial particles. We suppose given some measure γ_N on positive integers that satisfies $\mathbb{E}(\gamma_N) = N$ and $\text{Var}(\gamma_N) = o(N^2)$. The initial particle number, written \mathcal{N} , is a sample of γ_N . The particles present at time zero are scattered in the same way as they are in the deterministic case. The subsequent evolution, whose randomness is independent of the sampling of \mathcal{N} , is also the same as in the deterministic setting.

To describe this dynamics, set a parameter $\varepsilon > 0$ according to $N = k_\varepsilon Z$, as described earlier. Let $F : \{\mathbb{R}^d \times [0, \infty)\}^I \rightarrow [0, \infty)$ denote a smooth function. The action on F of the infinitesimal generator \mathbb{L} is given by

$$(\mathbb{L}F)(\mathbf{q}) = \mathbb{A}_0 F(\mathbf{q}) + \mathbb{A}_c F(\mathbf{q}),$$

where the diffusion and collision operators are given by

$$\mathbb{A}_0 F(\mathbf{q}) = \sum_{i \in I_q} d(m_i) \Delta_{x_i} F$$

and

$$\begin{aligned} \mathbb{A}_c F(\mathbf{q}) &= \frac{1}{2} \sum_{i,j \in I_q} \varepsilon^{-2} V\left(\frac{x_i - x_j}{\varepsilon}\right) \alpha(m_i, m_j) \\ &\times \left[\frac{m_i}{m_i + m_j} F(S_{i,j}^1 \mathbf{q}) + \frac{m_j}{m_i + m_j} F(S_{i,j}^2 \mathbf{q}) - F(\mathbf{q}) \right]. \end{aligned} \tag{1.5}$$

Note that:

- The function $V : \mathbb{R}^d \rightarrow [0, \infty)$ is assumed to be Hölder continuous, of compact support, and with $\int_{\mathbb{R}^d} V(x) dx = 1$.
- We denote by $S_{i,j}^1 \mathbf{q}$ that configuration formed from \mathbf{q} by removing the indices i and j from I_q , and adding a new index from I to which $S_{i,j}^1 \mathbf{q}$ assigns the value $(x_i, m_i + m_j)$. The configuration $S_{i,j}^2 \mathbf{q}$ is defined in the same way, except that it assigns the value $(x_j, m_i + m_j)$ to the new index. The specifics of the collision event then are that the new particle appears in one of the locations of the two particles being removed, with the choice being made randomly with weights proportional to the mass of the two colliding particles.

Convention. Unless stated otherwise, we will adopt a notation whereby all the index labels appearing in sums should be taken to be distinct.

We refer the reader to [4,10] for the reasons for choosing $N = \varepsilon^{d-2} Z$, the form of the collision term in (1.5), and the interpretations of the various terms.

Let us write $\mathcal{M}_Z(\mathbb{R}^d \times [0, \infty))$ for the space of non-negative measures π on $\mathbb{R}^d \times [0, \infty)$ such that

$$\pi(\mathbb{R}^d \times [0, \infty)) \leq Z.$$

This space is equipped with the topology of vague convergence which turns \mathcal{M}_Z into a compact metric space. We also write $\mathcal{M}_Z(\mathbb{R}^d \times [0, \infty)^2)$ for the space of non-negative measures μ such that for every positive T , $\mu(\mathbb{R}^d \times [0, \infty) \times [0, T]) \leq TZ$, which is also compact with respect to the topology of vague convergence. This space has a closed subspace \mathcal{X} which consists of measures μ such that $\mu(\mathbb{R}^d \times [0, \infty) \times [t_1, t_2]) \leq (t_2 - t_1)Z$, for every $t_1 \leq t_2$. As we will show

in Lemma 6.2 of Section 6, the space \mathcal{X} consists of measures $\mu(dx, dn, dt) = g(dx, dn, t)dt$ with $t \mapsto g(dx, dn, t)$ a Borel-measurable function from $[0, \infty)$ to $\mathcal{M}_Z(\mathbb{R}^d \times [0, \infty))$. We will denote by $\mathbb{P}_N = \mathbb{P}^\varepsilon$ the probability measure on functions from $t \in [0, \infty)$ to the configurations determined by the process at time t . Its expectation will be denoted \mathbb{E}_N . Setting

$$g^\varepsilon(dx, dn, t) = \varepsilon^{d-2} \sum_i \delta_{(x_i(t), m_i(t))}(dx, dn),$$

the law of

$$\mathbf{q} \mapsto \int g^\varepsilon(dx, dn, t) dt$$

with respect to \mathbb{P}^ε induces a probability measure \mathcal{P}^ε on the space \mathcal{X} . We note that, since the space \mathcal{X} is a compact metric space, the sequence \mathcal{P}^ε is precompact with respect to the topology of weak convergence.

For the main result of this article, we need the following assumptions on $\alpha(\cdot, \cdot)$ and $d(\cdot)$:

Hypothesis 1.1.

- The diffusion coefficient $d : (0, \infty) \rightarrow (0, \infty)$ is a bounded continuous function and there exists a uniformly positive continuous function $\phi : (0, \infty) \rightarrow (0, \infty)$ such that both $\phi(\cdot)$ and $\phi(\cdot)d(\cdot)$ are non-increasing.
- The function $\alpha : (0, \infty) \times (0, \infty) \rightarrow (0, \infty)$ is a bounded symmetric continuous function satisfying

$$\sup_{n \leq L} \sup_m \frac{\alpha(n, m)}{md(m)^{\frac{d}{2}} \phi(m)^{d-1}} < \infty,$$

for every $L > 0$.

Remark 1.1.

- The condition that the function $\phi : (0, \infty) \rightarrow (0, \infty)$ exists is rather mild and is satisfied if $d(\cdot)$ is non-increasing. This condition requires that heavier particles diffuse more slowly which is natural from the physical point of view. In fact when $d(\cdot)$ is non-increasing, then we can simply choose $\phi(m) \equiv 1$. Also, if $d(\cdot)$ is non-decreasing, then the function ϕ exists and can be chosen to be $\phi(m) = d(m)^{-1}$. From these two cases, we guess that the first condition is related to the variation of the function $d(\cdot)$. As we will show in Lemma 2.1 of Section 2, the existence of such a function ϕ is equivalent to assuming that the total negative variation of $\log d(\cdot)$ over each interval $[n, \infty)$, $n > 0$, is finite.
- We note that if the function $d(\cdot)$ is non-increasing, then the second condition for small m and n is equivalent to saying that $\alpha(m, n) \leq C \min(m, n)$. However, when m and n are large, the second condition is satisfied if for example $\alpha(m, n) \leq C m d(m)^{d/2} n d(n)^{d/2}$. Our stipulation that d be bounded is more restrictive in the case for values of its argument close to zero, since it is reasonable to assume that very light particles diffuse rapidly.

We also need the following assumptions on the initial data h :

Hypothesis 1.2.

- $\int_0^\infty \int h_n(x) dx dn < \infty$.
- $\bar{h}_k * \lambda_k \in L_{loc}^\infty(\mathbb{R}^d)$, for $k = 2, 3$ and 4 , where $\bar{h}_k = \int_0^\infty nd(n)^{\frac{d}{2}-\frac{1}{k}} \phi(n)^{\frac{dk}{2}-1} h_n dn$ and $\lambda_k(x) = |x|^{\frac{2}{k}-d}$.

•

$$\int \hat{h}(x)\hat{h}(y)|x - y|^{2-d} dx dy < \infty$$

where $\hat{h} = \int_0^\infty (n + 1)h_n dn$.

Remark 1.2. Recall that if $d(\cdot)$ is non-increasing, then we may choose $\phi = 1$. In this case, Hypothesis 1.2 is satisfied if $\hat{h} \in L^1 \cap L^\infty$.

To prepare for the statement of our main result, we now recall the weak formulation of the system (1.1). Firstly, recall that a non-negative measurable function $f : \mathbb{R}^d \times [0, \infty) \times [0, \infty) \rightarrow [0, \infty)$ is a **weak solution** of (1.1) subject to the initial condition $f(x, n, 0) = h_n(x)$ if for every smooth function $J : \mathbb{R}^d \times (0, \infty) \times [0, \infty) \rightarrow \mathbb{R}$ of compact support,

$$\begin{aligned} \int_0^\infty \int_{\mathbb{R}^d} f(x, n, t)J(x, n, t) dx dn &= \int_0^\infty \int_{\mathbb{R}^d} h_n(x)J(x, n, 0) dx dn \\ &+ \int_0^t \int_0^\infty \int_{\mathbb{R}^d} \frac{\partial J}{\partial t}(x, n, s) f(x, n, s) dx dn ds \\ &+ \int_0^t \int_0^\infty \int_{\mathbb{R}^d} d(n)\Delta J(x, n, s) f(x, n, s) dx dn ds \\ &+ \frac{1}{2} \int_0^t \int_0^\infty \int_0^\infty \int_{\mathbb{R}^d} \beta(m, n) f(x, n, s) f(x, m, s) \\ &\times \tilde{J}(x, m, n, s) dx dn dm ds, \end{aligned}$$

where

$$\tilde{J}(x, m, n, s) = J(x, m + n, s) - J(x, m, s) - J(x, n, s).$$

Following Norris [9], we define an analogous measure-valued notion of a weak solution.

Definition 1.1. Let us write $M[0, \infty)$ for the space of non-negative measures on the interval $[0, \infty)$. We equip this space with the topology of vague convergence. A measurable function $f : \mathbb{R}^d \times [0, \infty) \rightarrow M[0, \infty)$ is called a measure-valued weak solution of (1.1) if, firstly, for each $\ell > 0$, the functions $g_\ell, h_\ell \in L^1_{loc}$, where

$$g_\ell(x, t) = \int_0^\ell f(x, t, dn), \quad h_\ell(x, t) = \int_0^\infty \int_0^\ell \beta(m, n) f(x, t, dn) f(x, t, dm),$$

and, secondly,

$$\begin{aligned} \int_{\mathbb{R}^d} \int_0^\infty J(x, n, t) f(x, t, dn) dx &= \int_0^\infty \int_{\mathbb{R}^d} h_n(x)J(x, n, 0) dx dn \\ &+ \int_0^t \int_0^\infty \int_{\mathbb{R}^d} \frac{\partial J}{\partial t}(x, n, s) f(x, s, dn) dx ds \\ &+ \int_0^t \int_{\mathbb{R}^d} \int_0^\infty d(n)\Delta J(x, n, s) f(x, s, dn) dx ds \\ &+ \frac{1}{2} \int_0^t \int_{\mathbb{R}^d} \int_0^\infty \int_0^\infty \beta(m, n) \tilde{J}(x, m, n, s) f(x, s, dn) f(x, s, dm) dx ds. \end{aligned} \tag{1.6}$$

Remark 1.3. The requirement $g_\ell, h_\ell \in L^1_{\text{loc}}$ is imposed in order to guarantee the existence of the integrals in (1.6).

We are now ready to state the main result of this article.

Theorem 1.1. Consider the deterministic or random model for some dimension $d \geq 3$. Assume Hypotheses 1.1 and 1.2. If \mathcal{P} is any limit point of \mathcal{P}^ε , then \mathcal{P} is concentrated on the space of measures $g(dx, dn, t) dt = f(x, t, dn) dx dt$ which are absolutely continuous with respect to Lebesgue measure $dx \times dt$, with f solving the system of partial differential equations (1.1) in the sense of (1.6). The quantities $\beta : (0, \infty) \times (0, \infty) \rightarrow (0, \infty)$ are specified by the formula

$$\beta(n, m) = \alpha(n, m) \int_{\mathbb{R}^d} V(x) [1 + u(x; n, m)] dx,$$

where, for each pair $(n, m) \in (0, \infty) \times (0, \infty)$, $u(\cdot) = u(\cdot; n, m) : \mathbb{R}^d \rightarrow \mathbb{R}$ is the unique solution of

$$\Delta u(x) = \frac{\alpha(n, m)}{d(n) + d(m)} V(x)[1 + u(x)], \tag{1.7}$$

satisfying $u(x) \rightarrow 0$ as $|x| \rightarrow \infty$.

Remark 1.4.

- The continuity with respect to m and n and other important properties of $u(\cdot; n, m)$ will be stated in Lemma 4.2 of Section 4. In particular $u \in [-1, 0]$, which implies that $\beta > 0$ because u is not identically -1 . It follows from Lemma 4.2 that β is a continuous function. We also refer to the last section of [4] in which several properties of β are established. In particular, it is shown that $\beta \leq \alpha$ and $\beta(n, m) \leq \text{Cap}(K)(d(n) + d(m))$, where K denotes the support of the function V and $\text{Cap}(K)$ denotes the Newtonian capacity of the set K . (See [4] for the definition of Newtonian capacity.)
- To simplify our presentation, we assume that all particles have the same “radius”. However, in a more realistic model, we may replace $\varepsilon^{-2}V(\varepsilon^{-1}(x_i - x_j))$ with $\varepsilon^{-2}V(\varepsilon^{-1}(x_i - x_j); m_i, m_j)$, where $V(a; n, m) = (r(n) + r(m))^{-2}V(a/(r(n) + r(m)))$ and $r(n)$ is interpreted as the radius of a particle of mass n . Our method of proof applies even when we allow such a radial dependence and we can prove Theorem 1.1 provided that $r(n) = n^\chi$ with $\chi < (d - 2)^{-1}$ (when $d \geq 3$). In fact, we anticipate that if $\chi > (d - 2)^{-1}$, then, at least in the case of a sufficiently large initial condition, the particle densities no longer approximate a solution of (1.1) in which the mass $\int_0^\infty \int_{\mathbb{R}^d} mf(x, m, t) dx dm$ is conserved throughout time. We note that when $\chi = 0$, then β is bounded and in this case the mass is conserved. We refer the reader to [10] and the introduction of [4] for a more thorough discussion.
- We note that because of the factor $1/2$ in the definition of \mathbb{A}_c , we are practically summing over unordered pairs $\{i, j\}$ in Eq. (1.5). This is responsible for the factor $1/2$ which appears in the definition of Q_+^n . This corrects our minor mistake in the earlier works [4] and [5]. In these works, the factor $1/2$ is missing from the definition of \mathbb{A}_c even though in the proof of the main result an unordered summation was used.

Our second result shows that the macroscopic density is absolutely continuous with respect to Lebesgue measure dn . We will require:

Hypothesis 1.3. There exists a continuous function $\tau : (0, \infty) \rightarrow (0, \infty)$ for which $\int_0^\infty \tau(n) \, dn = 1$, with

$$\int_0^\infty \int_{\mathbb{R}^d} (|x|^2 + |\log \tau(n)| + |\log h_n|) h_n \, dx \, dn < \infty,$$

and

$$\int_0^\infty \int_{\mathbb{R}^d} \rho(n) h_n(x) \, dx \, dn < \infty, \tag{1.8}$$

where

$$\rho(n) = \int_0^n \alpha(m, n - m) \frac{\tau(m)\tau(n - m)}{\tau(n)} \, dm.$$

We also assume that $D = \sup_m d(m) < \infty$.

Remark 1.5. For a simple example for τ , consider $\tau(n) = (n + 1)^{-2}$. If for example $\alpha(m, n) \leq C(m + n)$, then $\rho(n) \leq Cn$ and (1.8) requires that the total mass to be finite initially.

Theorem 1.2. Assume that the model is random, and that the law γ_N of the initial total particle number \mathcal{N} has a Poisson distribution. Assume also Hypothesis 1.3. Then every limit point \mathcal{P} of the sequence \mathcal{P}^ε is concentrated on measures that take the form $g(dx, dn, t) \, dt = f_n(x, t) \, dn \, dx \, dt$, where f solves (1.1) with β as in Theorem 1.1. Moreover, there exists a constant C , that may be chosen independently of \mathcal{P} , such that

$$\int_{\mathcal{X}} \left[\int_0^\infty \int_{\mathbb{R}^d} \psi(f_n(x, t)) r(x, n) \, dx \, dn \right] \mathcal{P}(d\mu) \leq C, \tag{1.9}$$

for every t , where $\psi(f) = f \log f - f + 1$ and $r(x, n) = (2\pi)^{-d/2} \exp(-|x|^2/2)\tau(n)$.

Remark 1.6. At the expense of discussing some extra technicalities, the proof of Theorem 1.2 might include the random model with some other choice of γ_N . We only need to assume that for every positive λ , there exists a constant $a(\lambda)$ such that $\log \mathbb{E}_N \exp(\lambda \mathcal{N}) \leq Na(\lambda)$.

Theorem 1.2 is proved by firstly establishing an entropy bound for the distribution of $\mathbf{q}(t)$, and then using large deviation techniques to deduce that any limit point \mathcal{P} of the sequence \mathcal{P}^ε is concentrated on the space of measures $g(dx, dn, t) \, dt = f_n(x, t) \, dx \, dn \, dt$. For this, we simply follow the classical work of Guo, Papanicolaou and Varadhan [3]. Even though our result is valid for more general initial randomness, we prefer to state and prove our results for Poisson-type distributions, thereby focusing on the main idea of the method of proof.

The function $\tau : (0, \infty) \mapsto (0, \infty)$ appearing in Hypothesis 1.3 is used to define a reference measure with respect to which the corresponding entropy per particle is uniformly finite as $\varepsilon \rightarrow 0$. For simplicity, we take the reference measure ν_N which induces a Poisson law of intensity 1 for the initial number of particles \mathcal{N} and whose conditional measure $\nu_N(\cdot | \mathcal{N}(\mathbf{q}) = k)$ is given by

$$\prod_{i=1}^k r(x_i, m_i) \, dx_i \, dm_i. \tag{1.10}$$

The entropy per particle is uniformly finite, because the first part of Hypothesis 1.3 implies that

$$\sup_N \varepsilon^{d-2} \int F^0 \log F^0 \, d\nu_N < \infty,$$

where $F^0(\mathbf{q})\nu_N(d\mathbf{q})$ denotes the law of $\mathbf{q}(0)$. The second part of Hypothesis 1.3 will be used to control the time derivative of the entropy.

We now comment on the possible uniqueness of the solution that the microscopic model approximates. We expect to have a unique solution of the system (1.1) for the initial condition h as above. However, with the aid of the arguments of [6,11], we know how to establish this uniqueness only if we assume that the initial condition satisfies the bound

$$\int_0^\infty n^b \|h_n\|_{L^\infty} dn < \infty, \tag{1.11}$$

for sufficiently large $b = b(a)$ (see [6,11] for an expression for $b(a)$). Using this uniqueness, we can assert that in fact the limit \mathcal{P} of \mathcal{P}^ε exists and is concentrated on the single measure $\mu(dx, dn, dt) = f(x, n, t) dx dn dt$, where f is the unique solution to (1.1). As a corollary we have:

Corollary 1.1. *Assume that Hypotheses 1.1–1.3 hold and that (1.11) holds for sufficiently large b . Let $J : \mathbb{R}^d \times (0, \infty) \times [0, \infty) \rightarrow \mathbb{R}$ be a bounded continuous function of compact support. Then,*

$$\limsup_{N \rightarrow \infty} \mathbb{E}_N \left| \int_{\mathbb{R}^d} \int_0^\infty \int_0^\infty J(x, n, t)(g^\varepsilon(dx, dn, t) dt - f(x, n, t) dx dn dt) \right| = 0. \tag{1.12}$$

In (1.12), $f : \mathbb{R}^d \times [0, \infty) \times [0, \infty) \rightarrow [0, \infty)$ denotes the unique solution to the system (1.1) with the initial data $f_n(\cdot, 0) = h_n(\cdot)$.

The paper contains an appendix that discusses the scalings available in the Smoluchowski equations in their continuous form. Examining these scalings produces a heuristic argument for the regime of choices of the asymptotic behaviour of the input parameters $\beta : (0, \infty)^2 \rightarrow (0, \infty)$ and $d : (0, \infty) \rightarrow (0, \infty)$ for which a solution (1.1) will see most of the mass depart from any given compact subset of $(0, \infty)$ as time becomes high.

To outline the remainder of the paper: In Section 2, we explain the strategy of the proof, giving an overview alternative to that presented in [4]. In this section, we also show how the microscopic coagulation rate is comparable to the product of densities and may be replaced with an expression that is similar to the term Q in (1.1) (see Theorem 2.1). The main technical step for such a replacement is a regularity property of the coagulation and is stated as Proposition 2.1. In Section 2 the proof of Proposition 2.1 is reduced to a collection of bounds that are stated as Lemma 2.1. In Section 3, we establish the crucial correlation bound (1.4). In Section 4, the proof of Lemma 2.1 is carried out with the aid of the correlation bounds of Section 3. In Section 5, we show how the correlation bounds can be used to establish L^p -type bounds on the macroscopic densities. Sections 6 and 7 are devoted to the proofs of Theorems 1.1 and 1.2 respectively.

2. An outline of the proof of the main theorem

Our aim in this section is to outline the proof of the principal result, Theorem 1.1. The overall scheme of the proof is the same as that presented in [4], and the reader may wish to consult Section 2 of that paper for another overview.

Our goal is to show that the empirical measures $g^\varepsilon(dx, dn, t) dt$ converge to $f(x, t, dn)dx dt$, where f is some measure-valued weak solution of Smoluchowski’s equation (1.1). To this end,

we choose a smooth test function $J : \mathbb{R}^d \times (0, \infty) \times [0, \infty) \rightarrow \mathbb{R}$ of compact support and consider the expression

$$Y(\mathbf{q}, t) = \varepsilon^{d-2} \sum_{i \in I_{\mathbf{q}}} J(x_i, m_i, t).$$

Evidently,

$$Y(\mathbf{q}(t), t) = \int J(x, n, t) g^\varepsilon(dx, dn, t).$$

Note that

$$Y(\mathbf{q}(T), T) = Y(\mathbf{q}(0), 0) + \int_0^T \left(\frac{\partial Y}{\partial t} + \mathbb{A}_0(Y) + \mathbb{A}_c(Y) \right) (\mathbf{q}(t), t) dt + M_T, \tag{2.1}$$

where M_T is a martingale, where the free motion term $\mathbb{A}_0 Y$ equals

$$\mathbb{A}_0 Y(\mathbf{q}, t) = \varepsilon^{d-2} \sum_{i \in I_{\mathbf{q}}} d(m_i) \Delta_{x_i} J(x_i, m_i, t) = \int d(n) \Delta_x J(x, n, t) g^\varepsilon(dx, dn, t)$$

and where the collision term $\mathbb{A}_c Y$ is equal to

$$\mathbb{A}_c Y(\mathbf{q}, t) = \frac{1}{2} \varepsilon^{d-2} \sum_{i, j \in I_{\mathbf{q}}} \alpha(m_i, m_j) V_\varepsilon(x_i - x_j) \hat{J}(x_i, m_i, x_j, m_j, t), \tag{2.2}$$

with $V_\varepsilon(x) = \varepsilon^{-2} V(x/\varepsilon)$, and $\hat{J}(x_i, m_i, x_j, m_j, t)$ given by

$$\begin{aligned} & \frac{m_i}{m_i + m_j} J(x_i, m_i + m_j, t) + \frac{m_j}{m_i + m_j} J(x_j, m_i + m_j, t) \\ & - J(x_i, m_i, t) - J(x_j, m_j, t). \end{aligned} \tag{2.3}$$

Our approach is simply to understand which terms dominate in (2.1) when the initial particle number N is high, and, in this way, to see that Eq. (1.6) emerges from considering (2.1) in the high N limit. Clearly, we expect the last two terms in (1.6), corresponding to free motion and collision, to arise from the terms in (2.1) in which the operators \mathbb{A}_0 or \mathbb{A}_c act. The time derivative terms in (1.6) and (2.1) also naturally correspond. And indeed, the sum of the second and third terms on the right-hand side of (2.1) is already expressed in terms of the empirical measure and corresponds to the macroscopic expression

$$\int_0^T \int_0^\infty \int \left(\frac{\partial}{\partial t} + d(n) \Delta_x \right) J(x, n, t) f(x, t, dn) dx dt.$$

As we will see in Section 6, the term martingale M_T vanishes as $\varepsilon \rightarrow 0$. The main challenge comes from the fourth term on the right-hand side of (2.1), the collision term. How does its counterpart in (1.6) emerge in the limit of high initial particle number? To answer this, we need to understand how to express the time integral of changes to $Y(\mathbf{q}, t)$ resulting from all the collisions occurring in the microscopic model. To do this, it is natural to introduce the quantity

$$f^\delta(x, dn; \mathbf{q}) = \varepsilon^{d-2} \sum_{i \in I_{\mathbf{q}}} \delta^{-d} \xi \left(\frac{x_i - x}{\delta} \right) \delta_{m_i}(dn),$$

where $\xi : \mathbb{R}^d \rightarrow [0, \infty)$ is a smooth function of compact support with $\int_{\mathbb{R}^d} \xi dx = 1$. For $\delta > 0$ fixed and small, f^δ in essence counts the number of particles in a small macroscopic region about any given point, this region having diameter of order δ . To find the analytic collision term in (1.6) from its microscopic counterpart in (2.1), we must approximate the time integral of $\mathbb{A}_c Y(\mathbf{q}(t), t)$ by some functional of the macroscopically smeared particle count f^δ , in such a way that the approximation becomes good if we take the smearing parameter $\delta \rightarrow 0$ after taking the initial particle number N to be high. This is achieved by the following important result, in which we write $\Gamma(\mathbf{q}, t) = 2\mathbb{A}_c Y(\mathbf{q}, t)$.

Theorem 2.1. *Assume that the function $\hat{J}(x, m, y, n, t)$ vanishes when $t > T$, or $m + n < L^{-1}$, or $\max(m, n) > L$. Then*

$$\lim_{\delta \rightarrow 0} \limsup_{N \rightarrow \infty} \mathbb{E}_N \left| \int_0^T \left[\Gamma(\mathbf{q}(t), t) - \hat{\Gamma}^\delta(\mathbf{q}(t), t) \right] dt \right| = 0,$$

with

$$\begin{aligned} \hat{\Gamma}^\delta(\mathbf{q}, t) &= \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \int_0^\infty \int_0^\infty \alpha(m, n) U_{m,n}^\varepsilon(w_1 - w_2) \hat{J}(w_1, m, w_2, n, t) \\ &\quad \times f^\delta(w_1, dm; \mathbf{q}) f^\delta(w_2, dn; \mathbf{q}) dw_1 dw_2, \end{aligned} \tag{2.4}$$

where we set

$$U_{m,n}(x) = V(x)[1 + u(x; m, n)], \quad U_{m,n}^\varepsilon(x) = \varepsilon^{-d} U_{m,n}(x/\varepsilon),$$

with $u(\cdot; m, n)$ being given in Theorem 1.1.

Remark 2.1.

- Note that even though J is of compact support, the function \hat{J} given in (2.3) is not in general of compact support. In fact, if x_j which appears in (2.2) belongs to the bounded support of J , then x_i belongs to a bounded set because of the presence of the term V_ε . The same reasoning does not work for m_i or m_j . Of course if $J(x, n, t)$ vanishes if either $n > L$ or $n < L^{-1}$, then $\hat{J}(x, m, y, n, t)$ vanishes if $m + n \leq L^{-1}$, or $\max(m, n) > L$. However, for Theorem 2.1 we assume that in fact \hat{J} vanishes even if one of m or n is larger than L . Because of this, we need to show that the contribution of particles with large sizes is small. We leave this issue for Section 6. (See Lemma 6.1.)
- As we mentioned in Section 1, the continuity with respect to m and n and other properties of $u(\cdot; m, n)$ will be stated in Lemma 4.2.

We now explain heuristically why the relation between the cumulative microscopic coagulation rate $\Gamma(\mathbf{q}(t), t)$ and its macroscopically smeared counterpart $\hat{\Gamma}^\delta(\mathbf{q}(t), t)$ holds.

Here is a naive argument that proposes a form for $\hat{\Gamma}^\delta(\mathbf{q}(t), t)$. In the microscopic model, particles at (w_1, m) and (w_2, n) are liable to coagulate if their locations differ on the scale of ε , $|w_1 - w_2| = O(\varepsilon)$. If two particles are so located, they coagulate at a Poisson rate of $\alpha(m, n)V(w_1 - w_2)$. When a pair does so, it effects a change in $Y(\mathbf{q}, t)$ of $\hat{J}(x, m, y, n, t)$. The density for the presence of a particle of mass m at location w_1 should be well approximated by the particle count $f^\delta(w_1, dm)$ computed on a small macroscopic scale. Multiplying the factors, and integrating over space and time, we seem to show that the expression for $\int_0^T \hat{\Gamma}^\delta(\mathbf{q}(t), t) dt$

should be given by

$$\int_0^T \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \int_0^\infty \int_0^\infty \alpha(m, n) V^\varepsilon(w_1 - w_2) \hat{J}(w_1, m, w_2, n, t) \times f^\delta(w_1, dm; \mathbf{q}) f^\delta(w_2, dn; \mathbf{q}) dw_1 dw_2 dt,$$

where $V^\varepsilon(x) = \varepsilon^{-d} V(x/\varepsilon)$. The integrand differs from the correct expression in (2.4) by the lack of a factor of $1 + u(w_1 - w_2; m, n)$. Why is the preceding argument wrong? The reason is the following. The joint density for particle presence (of masses m and n) at w_1 and w_2 (with $|w_2 - w_1| = O(\varepsilon)$) is not well approximated by the product $f^\delta(w_1, dm) f^\delta(w_2, dn)$, because some positive fraction of particle pairs at displacement of order ε do not in fact contribute, since such pairs were liable to coagulate in the preceding instants of time, and, had they done so, they would no longer exist in the model. The correction factor $1 + u(w_1 - w_2; m, n) \in (0, 1)$ measures the fraction of pairs of particles, one with diffusion rate $d(m)$, the other, $d(n)$, that survives without coagulating to reach a relative displacement $w_1 - w_2$, and is bounded away from 1 in a neighbourhood of the origin of order ε .

We note that in Theorem 1.2 we have reached our main goals, namely we have produced a quadratic expression of the densities and a function αU which has the macroscopic coagulation propensity β for its average.

The following proposition is the key to proving Theorem 2.1.

Proposition 2.1. *Choose T large enough that $\hat{J}(\cdot, t) = 0$ when $t \geq T$. We have*

$$\lim_{|z| \rightarrow 0} \limsup_{\varepsilon \downarrow 0} \mathbb{E}_N \left| \int_0^T [\Gamma(\mathbf{q}(t), t) - \bar{\Gamma}_z(\mathbf{q}(t), t)] dt \right| = 0, \tag{2.5}$$

where

$$\bar{\Gamma}_z(\mathbf{q}, t) = \varepsilon^{2(d-2)} \sum_{i, j \in I_{\mathbf{q}}} \alpha(m_i, m_j) U_{m_i, m_j}^\varepsilon(x_i - x_j + z) \hat{J}(x_i, m_i, x_j, m_j, t). \tag{2.6}$$

In the statement, z plays the role of a small macroscopic displacement, taken to zero after the limit of high initial particle number is taken in the microscopic model. The proposition shows that the cumulative influence of coagulations in space and time on $Y(\mathbf{q}(t), t)$ is similar to that computed by instead considering pairs of particles at the fixed small macroscopic distance z , with a modification in the coagulation propensity in the expression (2.6) being made for the reason just described.

It is not hard to deduce Theorem 2.1 from Proposition 2.1. We refer to Section 3.5 of [4] for a proof of Theorem 2.1 assuming Proposition 2.1. See also [10] for a repetition of this proof and more heuristic discussions about the strategy of the proof.

We will prove Proposition 2.1 in the following way. Define

$$X_z(\mathbf{q}, t) = \varepsilon^{2(d-2)} \sum_{i, j \in I_{\mathbf{q}}} u^\varepsilon(x_i - x_j + z; m_i, m_j) \hat{J}(x_i, m_i, x_j, m_j, t),$$

where $u^\varepsilon(x; m, n) = \varepsilon^{2-d} u(x/\varepsilon; m, n)$. Note that $u^\varepsilon(x) = u^\varepsilon(x; m, n)$ solves

$$(d(m) + d(n)) \Delta_x u^\varepsilon = \alpha(m, n) (V_\varepsilon u^\varepsilon + V^\varepsilon), \tag{2.7}$$

with

$$V_\varepsilon(x) = \varepsilon^{-2} V(x/\varepsilon), \quad V^\varepsilon(x) = \varepsilon^{-d} V(x/\varepsilon).$$

The process $\{(X_z - X_0)(\mathbf{q}(t), t) : t \geq 0\}$ satisfies

$$\begin{aligned} (X_z - X_0)(\mathbf{q}(T), T) &= (X_z - X_0)(\mathbf{q}(0), 0) + \int_0^T \left(\frac{\partial}{\partial t} + \mathbb{A}_0 \right) (X_z - X_0)(\mathbf{q}(t), t) dt \\ &\quad + \int_0^T \mathbb{A}_c (X_z - X_0)(\mathbf{q}(t), t) dt + M(T), \end{aligned} \tag{2.8}$$

with $\{M(t) : t \geq 0\}$ being a martingale. We will see that the form (2.5) emerges from the dominant terms in (2.8), those that remain after the limit of high initial particle number $N \rightarrow \infty$ is taken. To see this, we label the various terms which appear on the right-hand side of (2.8). Firstly, those terms arising from the action of the diffusion operator:

$$\left(\frac{\partial}{\partial t} + \mathbb{A}_0 \right) (X_z - X_0) = H_{11} + H_{12} + H_{13} + H_{14} + H_2 + H_3 + H_4,$$

with

$$\begin{aligned} H_{11}(\mathbf{q}, t) &= \varepsilon^{2(d-2)} \sum_{i,j \in I_{\mathbf{q}}} \alpha(m_i, m_j) \left[V^\varepsilon(x_i - x_j + z) - V^\varepsilon(x_i - x_j) \right] \\ &\quad \times \hat{J}(x_i, m_i, x_j, m_j, t), \\ H_{12}(\mathbf{q}, t) &= -\varepsilon^{2(d-2)} \sum_{i,j \in I_{\mathbf{q}}} \alpha(m_i, m_j) V_\varepsilon(x_i - x_j) u^\varepsilon(x_i - x_j; m_i, m_j) \\ &\quad \times \hat{J}(x_i, m_i, x_j, m_j, t), \\ H_{13}(\mathbf{q}, t) &= \varepsilon^{2(d-2)} \sum_{i,j \in I_{\mathbf{q}}} \alpha(m_i, m_j) V_\varepsilon(x_i - x_j + z) u^\varepsilon(x_i - x_j + z; m_i, m_j) \\ &\quad \times \hat{J}(x_i, m_i, x_j, m_j, t), \\ H_{14}(\mathbf{q}, t) &= \varepsilon^{2(d-2)} \sum_{i,j \in I_{\mathbf{q}}} d(m_i) \left[u^\varepsilon(x_i - x_j + z; m_i, m_j) - u^\varepsilon(x_i - x_j; m_i, m_j) \right] \\ &\quad \times \hat{J}_t(x_i, m_i, x_j, m_j, t), \end{aligned}$$

along with

$$\begin{aligned} H_2(\mathbf{q}, t) &= 2\varepsilon^{2(d-2)} \sum_{i,j \in I_{\mathbf{q}}} d(m_i) \left[u_x^\varepsilon(x_i - x_j + z; m_i, m_j) - u_x^\varepsilon(x_i - x_j; m_i, m_j) \right] \\ &\quad \cdot \hat{J}_x(x_i, m_i, x_j, m_j, t), \\ H_3(\mathbf{q}, t) &= -2\varepsilon^{2(d-2)} \sum_{i,j \in I_{\mathbf{q}}} d(m_j) \left[u_x^\varepsilon(x_i - x_j + z; m_i, m_j) - u_x^\varepsilon(x_i - x_j; m_i, m_j) \right] \\ &\quad \cdot \hat{J}_y(x_i, m_i, x_j, m_j, t), \end{aligned}$$

and

$$\begin{aligned} H_4(\mathbf{q}, t) &= \varepsilon^{2(d-2)} \sum_{i,j \in I_{\mathbf{q}}} \left[u^\varepsilon(x_i - x_j + z; m_i, m_j) - u^\varepsilon(x_i - x_j; m_i, m_j) \right] \\ &\quad \times \left[d(m_i) \Delta_x \hat{J}(x_i, m_i, x_j, m_j, t) + d(m_j) \Delta_y \hat{J}(x_i, m_i, x_j, m_j, t) \right], \end{aligned}$$

where \hat{J}_x denotes the gradient of \hat{J} with respect to its first spatial argument, \hat{J}_y the gradient of \hat{J} with respect to its second spatial argument, and \cdot the scalar product. As for those terms arising

from the action of the collision operator,

$$\mathbb{A}_c(X_z - X_0)(\mathbf{q}, t) = G_z^1(\mathbf{q}, t) + G_z^2(\mathbf{q}, t) - G_0^1(\mathbf{q}, t) - G_0^2(\mathbf{q}, t),$$

where $G_z^1(\mathbf{q}, t)$ is set equal to

$$\begin{aligned} & \frac{1}{2} \sum_{k, \ell \in I_{\mathbf{q}}} \alpha(m_k, m_\ell) V_\varepsilon(x_k - x_\ell) \varepsilon^{2(d-2)} \sum_{i \in I_{\mathbf{q}}} \left\{ \frac{m_k}{m_k + m_\ell} \right. \\ & \times \left[u^\varepsilon(x_k - x_i + z; m_k + m_\ell, m_i) \hat{J}(x_k, m_k + m_\ell, x_i, m_i, t) \right. \\ & \quad \left. + u^\varepsilon(x_i - x_k + z; m_i, m_k + m_\ell) \hat{J}(x_i, m_i, x_k, m_k + m_\ell, t) \right] \\ & \quad + \frac{m_\ell}{m_k + m_\ell} \left[u^\varepsilon(x_\ell - x_i + z; m_k + m_\ell, m_i) \hat{J}(x_\ell, m_k + m_\ell, x_i, m_i, t) \right. \\ & \quad \left. + u^\varepsilon(x_i - x_\ell + z; m_i, m_k + m_\ell) \hat{J}(x_i, m_i, x_\ell, m_k + m_\ell, t) \right] \\ & \quad - \left[u^\varepsilon(x_k - x_i + z; m_k, m_i) \hat{J}(x_k, m_k, x_i, m_i, t) \right. \\ & \quad \left. + u^\varepsilon(x_i - x_k + z; m_i, m_k) \hat{J}(x_i, m_i, x_k, m_k, t) \right] \\ & \quad - \left[u^\varepsilon(x_\ell - x_i + z; m_\ell, m_i) \hat{J}(x_\ell, m_\ell, x_i, m_i, t) \right. \\ & \quad \left. + u^\varepsilon(x_i - x_\ell + z; m_i, m_\ell) \hat{J}(x_i, m_i, x_\ell, m_\ell, t) \right] \left. \right\}, \end{aligned}$$

and where

$$\begin{aligned} G_z^2(\mathbf{q}, t) &= -\varepsilon^{2(d-2)} \sum_{k, \ell \in I_{\mathbf{q}}} \alpha(m_k, m_\ell) V_\varepsilon(x_k - x_\ell) u^\varepsilon(x_k - x_\ell + z; m_k, m_\ell) \\ & \quad \times \hat{J}(x_k, m_k, x_\ell, m_\ell, t). \end{aligned}$$

The terms in G_z^1 arise from the changes in the functional X_z when a collision occurs due to the influence of the appearance and disappearance of particles on other particles that are not directly involved. Those in G_z^2 are due to the absence after collision of the summand in X_z indexed by the colliding particles.

As we take a high N limit in (2.8), note that the quantity

$$\int_0^T \Gamma(\mathbf{q}(t), t) dt = \varepsilon^{2(d-2)} \sum_{i, j \in I_{\mathbf{q}}} \alpha(m_i, m_j) V^\varepsilon(x_i - x_j) \hat{J}(x_i, m_i, x_j, m_j, t)$$

appears, with a negative sign, in the term H_{11} . The term H_{12} also remains of unit order in the high N limit, and would disrupt our aim of approximating $\int_0^T \Gamma(\mathbf{q}(t), t) dt$ by z -displayed expressions. However, our definition of u^ε (see (2.7)) ensures that

$$H_{12} - G_0^2 = 0,$$

so this unwanted term disappears. The definition of u^ε was made in order to achieve this. The other term of unit order remaining in the high N limit is the z -displaced H_{13} . Rearranging (2.8),

we obtain

$$\begin{aligned} & \left| \int_0^T H_{11}(\mathbf{q}(t), t) dt + \int_0^T H_{13}(\mathbf{q}(t), t) dt \right| \leq |X_z - X_0|(\mathbf{q}(T), T) + |X_z - X_0|(\mathbf{q}(0), 0) \\ & + \int_0^T (|H_{14}| + |H_2| + |H_3| + |H_4|)(\mathbf{q}(t), t) dt \\ & + \int_0^T |G_z^1 - G_0^1|(\mathbf{q}(t), t) dt + \int_0^T |G_z^2|(\mathbf{q}(t), t) dt + |M(T)|. \end{aligned} \tag{2.9}$$

We have succeeded in writing $\bar{T}_z - \Gamma$ in the form $H_{11} + H_{13}$, so, for Proposition 2.1, it remains to prove that the right-hand side of (2.9) is small enough. Firstly, recall that, by our assumption, the function \hat{J} is of compact support. We now choose T sufficiently large that $\hat{J}(x, m, y, n, T) = 0$. As a result, the first term on the right-hand side vanishes. The other bounds that we require are now stated.

Lemma 2.1. *There exists a constant $C_2 = C_2(\hat{J}, T)$ such that*

$$\int_0^T \mathbb{E}_N (|H_2| + |H_3|)(\mathbf{q}(t), t) dt \leq C_2 |z|^{\frac{1}{d+1}}, \tag{2.10}$$

$$\int_0^T \mathbb{E}_N (|H_4| + |H_{14}|)(\mathbf{q}(t), t) dt \leq C_2 |z|^{\frac{2}{d+1}}, \tag{2.11}$$

$$\int_0^T \mathbb{E}_N |G_z^1 - G_0^1|(\mathbf{q}(t), t) dt \leq C_2 |z|^{\frac{2}{d+1}}, \tag{2.12}$$

$$\int_0^T \mathbb{E} |G_z^2|(\mathbf{q}(t), t) dt \leq C_2 \left(\frac{\varepsilon}{|z|}\right)^{d-2}, \tag{2.13}$$

$$\mathbb{E}_N |X_z - X_0|(\mathbf{q}(0)) \leq C_2 |z|, \tag{2.14}$$

$$\mathbb{E}_N [M(T)^2] \leq C_2 \varepsilon^{d-2}. \tag{2.15}$$

These bounds are furnished by the correlation inequality Theorem 3.1 that is the main innovation of this paper, to whose proof we now turn.

3. Correlation bounds

This section is devoted to the proof of the correlation bound which appeared as (1.4) when $d(\cdot)$ is non-increasing and takes the form (3.1) in general. Recall the function ϕ which appeared in Hypothesis 1.1. The main result of this section is Theorem 3.1.

Theorem 3.1. *For every non-negative bounded continuous function $K : (\mathbb{R}^d)^k \rightarrow \mathbb{R}$,*

$$\begin{aligned} & \mathbb{E}_N \int_0^\infty \sum_{i_1, \dots, i_k \in I_{\mathbf{q}(t)}} K(x_{i_1}(t), \dots, x_{i_k}(t)) \prod_{r=1}^k \gamma_k(m_{i_r}(t)) dt \\ & \leq \mathbb{E}_N \sum_{i_1, \dots, i_k \in I_{\mathbf{q}(0)}} (\Lambda^{m_{i_1}(0), \dots, m_{i_k}(0)} K)(x_{i_1}(0), \dots, x_{i_k}(0)) \prod_{r=1}^k \gamma_k(m_{i_r}(0)), \end{aligned} \tag{3.1}$$

where all summations are over distinct indices i_1, \dots, i_k , the function $\gamma_k(m) = md(m)^{d/2} \phi(m)^{\frac{kd}{2}-1}$, and the operator Λ is defined by

$$\begin{aligned} \Lambda^{n_1, \dots, n_k} K(y_1, \dots, y_k) &= c_0(kd) \int \left(\frac{|y_1 - z_1|^2}{d(n_1)} + \dots + \frac{|y_k - z_k|^2}{d(n_k)} \right)^{1-\frac{kd}{2}} \\ &\quad \times K(z_1, \dots, z_k) \prod_{r=1}^k d(n_r)^{-d/2} dz_r, \end{aligned} \tag{3.2}$$

where $c_0(kd) = (kd - 2)^{-1} \omega_{kd}^{-1}$, with ω_{kd} denoting the surface area of the unit sphere in \mathbb{R}^{kd} .

Let us make a comment about the form of (3.1) before embarking on its proof. Observe that if there were no coagulation, then it would have been straightforward to bound the left-hand side of (3.1) with the aid of the diffusion semigroup even if we allowed a function K that depends on the masses of particles. Indeed, if $S_t^{m_{i_1}, \dots, m_{i_k}}$ denotes the diffusion semigroup associated with particles $(x_{i_1}, m_{i_1}), \dots, (x_{i_k}, m_{i_k})$, then $\int_0^\infty S_t^{m_{i_1}, \dots, m_{i_k}} dt$ is exactly the operator $\Lambda^{m_{i_1}, \dots, m_{i_k}}$. What (3.1) asserts is that a similar bound is valid in spite of coagulation provided that we allow only a very special dependence on the masses of particles.

Proof of Theorem 3.1. Let us define

$$G(\mathbf{q}) = \sum_{i_1, \dots, i_k \in I_{\mathbf{q}}} (\Lambda^{m_{i_1}, \dots, m_{i_k}} K)(x_{i_1}, \dots, x_{i_k}) \prod_{r=1}^k \gamma_k(m_{i_r}).$$

Recall that the process $\mathbf{q}(t)$ is a Markov process with generator $\mathbb{L} = \mathbb{A}_0 + \mathbb{A}_c$ where $\mathbb{A}_0 = \sum_{i \in I_{\mathbf{q}}} d(m_i) \Delta_{x_i}$. By semigroup theory,

$$\mathbb{E}_N G(\mathbf{q}(t)) = \mathbb{E}_N G(\mathbf{q}(0)) + \mathbb{E}_N \int_0^t \mathbb{L}G(\mathbf{q}(s)) ds. \tag{3.3}$$

We have

$$\mathbb{A}_0 G(\mathbf{q}) = - \sum_{i_1, \dots, i_k \in I_{\mathbf{q}}} K(x_{i_1}, \dots, x_{i_k}) \prod_{r=1}^k \gamma_k(m_{i_r}). \tag{3.4}$$

This and the assumption $K \geq 0$ would imply (3.1) provided that we can show

$$\mathbb{A}_c G \leq 0. \tag{3.5}$$

To prove (3.5), let us study the effect of a coagulation between the i th and j th particles on G . We need to study three cases separately:

- $i, j \notin \{i_1, \dots, i_k\}$,
- $i, j \in \{i_1, \dots, i_k\}$,
- only one of i, j belongs to $\{i_1, \dots, i_k\}$.

If the first case occurs, then (i, j) -coagulation does not affect the term indexed by $\{i_1, \dots, i_k\}$ in $G(\mathbf{q})$.

If the second case occurs, then we need to remove those terms in the summation for which $\{i, j\} \subseteq \{i_1, \dots, i_k\}$. This contributes negatively to $\mathbb{A}_c G(\mathbf{q})$, because $K \geq 0$. This total

contribution for this case is given by

$$-\frac{1}{2} \sum_{i,j \in I_q} V_\varepsilon(x_i - x_j) \alpha(m_i, m_j) \times \sum_{i_1, \dots, i_k} \mathbb{1}(i, j \in \{i_1, \dots, i_k\}) (\Lambda^{m_{i_1}, \dots, m_{i_k}} K)(x_{i_1}, \dots, x_{i_k}) \prod_{r=1}^k \gamma_k(m_{i_r}).$$

If the third case occurs, then only one of i, j belongs to $\{i_1, \dots, i_k\}$. For example, either $i = i_1$, and $j \notin \{i_1, \dots, i_k\}$, or $j = i_1$, and $i \notin \{i_1, \dots, i_k\}$. In this case, the contribution is still non-positive because after the coagulation the expression

$$Y_1 = \sum_{i_2, \dots, i_k} (\Lambda^{m_i, m_{i_2}, \dots, m_{i_k}} K)(x_i, x_{i_2}, \dots, x_{i_k}) \gamma_k(m_i) \prod_{r=2}^k \gamma_k(m_{i_r}) + \sum_{i_2, \dots, i_k} (\Lambda^{m_j, m_{i_2}, \dots, m_{i_k}} K)(x_j, x_{i_2}, \dots, x_{i_k}) \gamma_k(m_j) \prod_{r=2}^k \gamma_k(m_{i_r}),$$

is replaced with the expression Y_2 which is given by

$$\frac{m_i}{m_i + m_j} \sum_{i_2, \dots, i_k} (\Lambda^{m_i + m_j, m_{i_2}, \dots, m_{i_k}} K)(x_i, x_{i_2}, \dots, x_{i_k}) \gamma_k(m_i + m_j) \prod_{r=2}^k \gamma_k(m_{i_r}) + \frac{m_j}{m_i + m_j} \sum_{i_2, \dots, i_k} (\Lambda^{m_i + m_j, m_{i_2}, \dots, m_{i_k}} K)(x_j, x_{i_2}, \dots, x_{i_k}) \gamma_k(m_i + m_j) \prod_{r=2}^k \gamma_k(m_{i_r}).$$

For (3.5), it suffices to show that $Y_2 \leq Y_1$. For this, it suffices to show that for every positive m, n, A and B ,

$$\phi(m+n)^{\frac{kd}{2}-1} \left[A \frac{d(m)}{d(m+n)} + B \right]^{1-\frac{kd}{2}} \leq \phi(m)^{\frac{kd}{2}-1} [A+B]^{1-\frac{kd}{2}}. \tag{3.6}$$

We are done because the assertion (3.6) for fixed m, n and all positive A and B is equivalent to the inequalities

$$\phi(m)d(m) \geq \phi(m+n)d(m+n),$$

and

$$\phi(m) \geq \phi(m+n),$$

both being satisfied, and these are true for all choices of m and n by Hypothesis 1.1. \square

Corollary 3.1. For every non-negative bounded continuous function K ,

$$\varepsilon^{k(d-2)} \mathbb{E}_N \int_0^T \sum_{i_1, \dots, i_k \in I_{\mathbf{q}(t)}} K(x_{i_1}(t), \dots, x_{i_k}(t)) \prod_{r=1}^k \gamma_k(m_{i_r}(t)) dt \leq c_0(kd) \int K(x_1, \dots, x_k) \prod_{r=1}^k (\bar{h}_k * \lambda_k)(x_r) dx_r, \tag{3.7}$$

where $\bar{h}_k = \int_0^\infty n\phi(n)^{\frac{kd}{2}-1}d(n)^{\frac{d}{2}-\frac{1}{k}}h_n dn$ and $\lambda_k(w) = |w|^{\frac{2}{k}-d}$.

Proof. From the elementary inequality $a_1 \dots a_k \leq (a_1^2 + \dots + a_k^2)^{k/2}$, we deduce that the kernel $\lambda^{n_1, \dots, n_k}$ of the operator A^{n_1, \dots, n_k} is bounded above by

$$\lambda^{n_1, \dots, n_k}(z_1, \dots, z_k) \leq c_0(kd) \prod_{r=1}^k |z_r|^{\frac{2}{k}-d} d(n_r)^{-\frac{1}{k}}.$$

This and (3.1) imply (3.7). \square

Remark 3.1. Corollary 3.1 will be applied in four places in coming sections and except for the last application, the function K to which this corollary applied is only bounded and continuous off a neighborhood of a set of zero Lebesgue measure. For such unbounded K , we pick a large positive parameter ℓ and apply (3.1) to $K'_\ell = \min\{\ell, K\}$. We then replace K'_ℓ with K on the right-hand side and pass to the limit $\ell \rightarrow \infty$ on the left-hand side. From this and the monotone convergence theorem we deduce (3.1) for such a singular function K .

We end this section with two lemmas concerning the first condition in Hypothesis 1.1.

Lemma 3.1. *Suppose the function $d(\cdot)$ has a finite negative variation in an interval $[a, b] \subset (0, \infty)$. Then there exists a positive continuous function ϕ such that ϕ and ϕd are non-increasing in the interval $[a, b]$.*

Proof. *Step 1.* Firstly, we assume that there exist points $a_0 = b > a_1 > \dots > a_{\ell-1} > a_\ell = a$ such that $d(\cdot)$ is monotone on each interval $[a_i, a_{i-1}]$, $i = 1, \dots, \ell$. For the sake of definiteness, let us assume that $d(\cdot)$ is non-decreasing (non-increasing) in $[a_i, a_{i-1}]$, if i is odd (even). In this case, we can construct a continuous ϕ as follows: Define $A_0 = A$ and $A_k = A \prod_{i=1}^k \frac{d(a_{2i})}{d(a_{2i-1})}$ for $k \geq 1$. For $x \in [a_{2k+1}, a_{2k}]$ and $k \geq 0$, we set $\phi(x) = \frac{A_k}{d(x)}$. For $x \in [a_{2k}, a_{2k-1}]$ and $k \geq 1$, we set $\phi(x) = \frac{A_{k-1}}{d(a_{2k-1})}$.

Step 2. Let d be a continuous positive function. Approximate d in L^∞ by a sequence of continuous piecewise monotone functions $\{d_n\}$. To simplify the presentation, we assume that each d_n is as in Step 1. That is, d_n increases near the end point b . Let us write ϕ_n for the corresponding ϕ , and let c_n denote the number of intervals in the partition (so that $a_{c_n} = a$). It remains to show that the sequence $\{\phi_n\}$ has a convergent subsequence. Since each ϕ_n is non-increasing, we may appeal to the Helly Selection Theorem. For this we need to make sure that the sequence $\{\phi_n\}$ is bounded. Note that $\sup_{x \in [a, b]} \phi_n(x) = \phi_n(a) = \phi_n(a_{c_n})$. Set $D_n = A_{\frac{c_n-1}{2}}$ if c_n is odd and $D_n = A_{\frac{c_n}{2}-1}$ if c_n is even. We readily see that $\phi_n(c_n) \leq (\inf_{x \in [a, b]} d(x))^{-1} D_n$, whatever the parity of c_n . The infimum being positive, we require that $\sup_{n \in \mathbb{N}} D_n < \infty$. For any $k \in \mathbb{N}$ for which A_k is defined, we may take the logarithm of A_k to produce a sum and observe that $d(\cdot)$ is non-increasing on the intervals $[a_{2i}, a_{2i-1}]$. Hence, $\log A_k$ measures the negative variation of the function $\log d$ on the interval $[a_{2k}, b]$. Since d is uniformly positive, $\sup_n D_n < \infty$ is implied by the function d having a finite negative variation. \square

Lemma 3.2. *Suppose the function $\log d(\cdot)$ has a finite negative variation in an interval $[n_0, \infty)$ with $n_0 > 0$. Then there exists a positive continuous function ϕ such that ϕ and ϕd are non-increasing in the interval $[n_0, \infty)$.*

Proof. The proof is very similar to the proof of Lemma 3.1. First we assume that d is piecewise monotone. This time we set $\phi(n_0) = A$ and define ϕ continuously so that ϕ is constant when d decreases and ϕ is a constant multiple of d^{-1} when d increases. Since ϕ is non-increasing, we may end with a function which crosses 0 and becomes negative. This can be fixed by adjusting $A = \phi(n_0)$ only if ϕ is bounded below. As in the proof of Lemma 3.1, we can readily see that ϕ is bounded below if the total negative variation of d is finite. \square

Note that in the statement of Lemma 3.2 we cannot drop log because on the infinite interval $[n_i, \infty)$ the function $d(\cdot)$ could take arbitrarily small values.

4. Proof of Lemma 2.1

The strategy of the proof of Lemma 2.1 is the same as the one used to prove the analogous inequalities in [4]. The only difference is that we only need to use our correlation bound Corollary 3.1 to get the bounds (2.10–2.15). For (2.10) and (2.11) we need to apply Corollary 3.1 for $k = 2$. Corollary 3.1 in the case $k = 3$ will be used for (2.12). As for (2.15) all cases $k = 2, 3, 4$ will be employed. We omit the proof of the inequalities (2.13) and (2.14) because they can be established by a verbatim argument as in [4]. In fact the proof (2.14) is straightforward because we are dealing with a calculation involving the initial configuration. For this, however, a suitable bound on the function u^ε would be needed that will be stated as a part of Lemma 4.2 below. The same bound and Lemma 4.1 below will imply (2.13).

The main ingredients for the proof of inequalities (2.10) and (2.11) are Corollary 3.1 (with $k = 2$), certain bounds on u^ε and its spatial gradient u_x^ε (which will appear in Lemma 4.2), and Lemma 4.1 below. The straightforward proof of Lemma 4.1 is also omitted and can be proved in exactly the same way as we proved Lemma 3.1 of [4].

Lemma 4.1. For any $T \in [0, \infty)$,

$$\mathbb{E}_N \int_0^T \varepsilon^{d-2} \sum_{i,j \in I_q(t)} \alpha(m_i(t), m_j(t)) V_\varepsilon(x_i(t) - x_j(t)) dt \leq Z.$$

As for the remaining inequalities, we only establish (2.12) and (2.15) because these are the most technically involved cases and the same idea of proof applies to (2.10) and (2.11).

We now state our lemma as regards the functions u and u^ε . Recall that $u^\varepsilon(x; n, m) = \varepsilon^{2-d} u(x/\varepsilon; n, m)$ where u satisfies

$$\Delta u(x; n, m) = \alpha'(n, m) V(x) [1 + u(x; n, m)],$$

with $u(x; n, m) \rightarrow 0$ as $|x| \rightarrow \infty$, and

$$\alpha'(n, m) := \frac{\alpha(n, m)}{d(n) + d(m)}.$$

For our purposes, let us write w^a for the unique solution of

$$\Delta w^a(x) = a V(x) [1 + w^a(x)],$$

with $w^a(x) \rightarrow 0$ as $|x| \rightarrow \infty$. Of course, if we choose $a = \alpha'(n, m)$, then we obtain $u(x; n, m)$. We choose the constant C_0 so that $V(x) = 0$ whenever $|x| \geq C_0$.

Lemma 4.2. There exists a constant C_3 for which the following bounds hold.

- $-1 \leq w^a(x) \leq 0$ and for $x \in \mathbb{R}^d$,

$$|w^a(x)| \leq C_3 a \min\{|x|^{2-d}, 1\},$$

$$|w_x^a(x)| \leq C_3 a \min\{|x|^{1-d}, 1\},$$

where w_x^a denotes the spatial gradient of w^a .

- For $x \in \mathbb{R}^d$ satisfying $|x| \geq \max\{2|z| + C_0\varepsilon, 2C_0\varepsilon\}$,

$$|u^\varepsilon(x+z; n, m) - u^\varepsilon(x; n, m)| \leq C_3\alpha'(n, m)|z||x|^{1-d} \tag{4.1}$$

and

$$|u_x^\varepsilon(x+z; n, m) - u_x^\varepsilon(x; n, m)| \leq C_3\alpha'(n, m)|z||x|^{-d}. \tag{4.2}$$

- The function w^a is differentiable with respect to a and $a^{-1}w^a \leq \frac{\partial w^a}{\partial a} \leq 0$.

Proof. The proof of the first and second parts can be found in Section 3.2 of [4] and we do not repeat it here. As for the third part, recall that the function w^a is uniquely determined by the equation

$$w^a(x) = -c_0 a \int_{\mathbb{R}^d} |x-y|^{2-d} V(y)(1+w^a(y)) dy, \tag{4.3}$$

where $c_0 = c_0(d) = (d-2)^{-1}\omega_d^{-1}$, with ω_d denoting the surface area of the unit sphere S^{d-1} . We wish to show the regularity of the function w^a with respect to the variable a . In fact the existence of the unique solution to (4.3) was established in [4] using the Fredholm Alternative Theorem. To explain this, let us pick a bounded continuous function R such that $R > 0$, with

$$\int_{\mathbb{R}^d} R(x) dx = \infty, \quad \int_{|x| \geq 1} R(x)|x|^{4-2d} dx < \infty.$$

Define

$$\mathcal{H} = \left\{ u : \mathbb{R}^d \rightarrow \mathbb{R} : u \text{ is measurable and } \int_{\mathbb{R}^d} u^2(x)R(x) dx < \infty \right\}.$$

Observe that \mathcal{H} is a Hilbert space with respect to the inner product

$$\langle u, v \rangle = \int_{\mathbb{R}^d} u(x)v(x)R(x) dx.$$

Note that if w^a solves (4.3), then, defining $\mathcal{F} : \mathcal{H} \mapsto \mathcal{H}$ by

$$\mathcal{F}(\omega) = c_0 \int |x-y|^{2-d} V(y)\omega(y) dy,$$

we have that

$$(id + a\mathcal{F})(w^a) = -a\Gamma \tag{4.4}$$

where

$$\Gamma(x) = c_0 \int_{\mathbb{R}^d} |x-y|^{2-d} V(y) dy,$$

and id means the identity transformation. We wish to show the differentiability of w^a with respect to $a > 0$. This is clear heuristically because we have a candidate for $v^a := \frac{\partial w^a}{\partial a}$; if we differentiate both sides of (4.4), then v^a solves

$$(id + a\mathcal{F})(v^a) = -\Gamma - \mathcal{F}w^a = a^{-1}w^a. \tag{4.5}$$

This provides us with a candidate for $\frac{\partial w^a}{\partial a}$, because the operator $id + a\mathcal{F}$ has a bounded inverse (see Section 6 of [4]). The rigorous proof of the differentiability of w^a goes as follows. First define $v^{a,h} = (w^{a+h} - w^a)/h$ and observe that $v^{a,h}$ satisfies

$$(id + a\mathcal{F})(v^{a,h}) = -\Gamma - \mathcal{F}w^{a+h}. \tag{4.6}$$

We would like to show that $v^{a,h}$ has a limit in \mathcal{H} , as $h \rightarrow 0$. One can readily show that the right-hand side of (4.6) is bounded in \mathcal{H} because $|w^a(x)| \leq C_2 a \min\{|x|^{2-d}, 1\}$ by the first part of the lemma. Hence $v^{a,h}$ stays bounded as $h \rightarrow 0$. If v^a is any weak limit, then v^a must satisfy (4.5). Since (4.5) has a unique solution, the weak limit of $v^{a,h}$ exists. In [4], it is shown that \mathcal{F} is a compact operator. From this and (4.6), we can readily deduce that the strong limit of $v^{a,h}$ exists. As a consequence, w^a is weakly differentiable in a and its derivative satisfies (4.5). Using Sobolev’s inequalities and the fact that V is Hölder continuous, we can deduce by standard arguments that v^a is indeed C^2 and satisfies

$$\Delta v^a = av^a V + (1 + w^a)V. \tag{4.7}$$

This means that $w^a(x)$ is continuously differentiable with respect to (x, a) .

We now want to use (4.7) or equivalently (4.5) to conclude that $a^{-1}w^a \leq v^a \leq 0$. In fact, by (4.5), we have that $v^a = -a\mathcal{F}v^a - a^{-1}w^a$, which implies that

$$|v^a(x)| \leq c'_a c_0 \int |x - y|^{2-d} dy + a^{-1}|w^a(x)|,$$

where c'_a is an upper bound for $|v^a(x)|$ with x in the support of the function V . From this, it is not hard to deduce that there exists a constant c''_a such that

$$|v^a(x)| \leq c''_a \max\{|x|^{2-d}, 1\}. \tag{4.8}$$

In a similar fashion, we can show that there exists a constant c'''_a such that

$$|\nabla v^a(x)| \leq c'''_a \max\{|x|^{1-d}, 1\}. \tag{4.9}$$

We now demonstrate that $v^a \leq 0$. Take a smooth function $\varphi_\delta : \mathbb{R} \rightarrow [0, \infty)$ such that $\varphi'_\delta, \varphi_\delta \geq 0$ and

$$\varphi_\delta(r) = \begin{cases} 0 & r \leq 0, \\ r & r \geq \delta. \end{cases}$$

We then have

$$-\int_{\mathbb{R}^d} \varphi'_\delta(v^a) |\nabla v^a|^2 dx = \int_{\mathbb{R}^d} \varphi_\delta(v^a) \Delta v^a dx = \int_{\mathbb{R}^d} V(1 + w^a + av^a) \varphi_\delta(v^a) dx, \tag{4.10}$$

the second equality by (4.7). Integration by parts was performed in the first equality: we write the analogue of (4.10) which is integrated over a bounded set $\{x : |x| \leq R\}$. We may obtain (4.10) by sending $R \rightarrow \infty$ but for this we need to make sure that the boundary contribution coming from the set $\{x : |x| = R\}$ goes away as $R \rightarrow \infty$. This is readily achieved with the aid of (4.9). Since $1 + w^a \geq 0$ by the first part of the lemma, and $v^a \varphi_\delta(v^a) \geq 0$, we deduce that the right-hand side of (4.10) is non-negative. Since the left-hand side is non-positive, we deduce that

$$\int_{\mathbb{R}^d} \varphi'_\delta(v^a) |\nabla v^a|^2 dx = \int_{\mathbb{R}^d} V(1 + w^a + av^a) \varphi_\delta(v^a) dx = 0.$$

We now send $\delta \rightarrow 0$ to deduce

$$0 = \int_{\mathbb{R}^d} |\nabla v^a|^2 \mathbb{1}(v^a \geq 0) \, dx = \int_{\mathbb{R}^d} V(1 + w^a + av^a)v^a \mathbb{1}(v^a \geq 0) \, dx.$$

As a result, on the set $A = \{x : v^a > 0\}$ we have $\nabla v^a = 0$. Hence v^a is constant on each component B of A . But this constant can only be 0 because on the boundary of A we have $v^a = 0$. This is impossible unless A is empty. Hence, $v^a \leq 0$ everywhere.

It remains to prove that $v^a \geq a^{-1}w^a$. For this observe that if $\gamma^a = a^{-1}w^a - v^a$, then

$$\Delta \gamma^a = aV\gamma^a + V(-w^a).$$

We can now repeat the proof of $v^a \leq 0$ to deduce that $\gamma^a \leq 0$ because $-w^a \geq 0$. This completes the proof of the third part of the lemma. \square

Proof of (2.12). Note that

$$\int_0^T \mathbb{E}_N |G_z^1 - G_0^1|(\mathbf{q}(t), t) \, dt \leq \frac{1}{2} \sum_{i=1}^8 D_i,$$

where the first four of the D_i are given by

$$\begin{aligned} D_1 &= \mathbb{E}_N \int_0^T dt \sum_{k, \ell \in I_q} \alpha(m_k, m_\ell) V_\varepsilon(x_k - x_\ell) \frac{m_k}{m_k + m_\ell} \varepsilon^{2(d-2)} \\ &\quad \times \sum_{i \in I_q} |u^\varepsilon(x_k - x_i + z; m_k + m_\ell, m_i) - u^\varepsilon(x_k - x_i; m_k + m_\ell, m_i)| \\ &\quad \times |\hat{J}(x_k, m_k + m_\ell, x_i, m_i, t)|, \end{aligned}$$

$$\begin{aligned} D_2 &= \mathbb{E}_N \int_0^T dt \sum_{k, \ell \in I_q} \alpha(m_k, m_\ell) V_\varepsilon(x_k - x_\ell) \frac{m_\ell}{m_k + m_\ell} \varepsilon^{2(d-2)} \\ &\quad \times \sum_{i \in I_q} |u^\varepsilon(x_\ell - x_i + z; m_k + m_\ell, m_i) - u^\varepsilon(x_\ell - x_i; m_k + m_\ell, m_i)| \\ &\quad \times |\hat{J}(x_\ell, m_k + m_\ell, x_i, m_i, t)|, \end{aligned}$$

$$\begin{aligned} D_3 &= \mathbb{E}_N \int_0^T dt \sum_{k, \ell \in I_q} \alpha(m_k, m_\ell) V_\varepsilon(x_k - x_\ell) \varepsilon^{2(d-2)} \\ &\quad \times \sum_{i \in I_q} |u^\varepsilon(x_k - x_i + z; m_k, m_i) - u^\varepsilon(x_k - x_i; m_k, m_i)| |\hat{J}(x_k, m_k, x_i, m_i, t)|, \end{aligned}$$

and

$$\begin{aligned} D_4 &= \mathbb{E}_N \int_0^T dt \sum_{k, \ell \in I_q} \alpha(m_k, m_\ell) V_\varepsilon(x_k - x_\ell) \varepsilon^{2(d-2)} \\ &\quad \times \sum_{i \in I_q} |u^\varepsilon(x_\ell - x_i + z; m_\ell, m_i) - u^\varepsilon(x_\ell - x_i; m_\ell, m_i)| |\hat{J}(x_\ell, m_\ell, x_i, m_i)|. \end{aligned}$$

The other four terms each take the form of one of the above terms, the particle indices that appear in the arguments of the functions u^ε and \hat{J} being switched, along with the mass pair labels for these functions.

The estimates involved for each of the eight cases are in essence identical. We will examine the case of D_3 . We write $D_3 = D^1 + D^2$, decomposing the inner i -indexed sum according to the

respective index sets

$$\{i \in I_{\mathbf{q}}, i \neq k, \ell, |x_k - x_i| > \rho\} \quad \text{and} \quad \{i \in I_{\mathbf{q}}, i \neq k, \ell, |x_k - x_i| \leq \rho\}.$$

Here, ρ is a positive parameter that satisfies the bound $\rho \geq \max\{2|z| + C_0\varepsilon, 2C_0\varepsilon\}$. By the second part of Lemma 4.2, we have that

$$D^1 \leq \frac{c_0|z|\varepsilon^{d-2}}{\rho^{d-1}} \mathbb{E}_N \int_0^T dt \sum_{k, \ell \in I_{\mathbf{q}}} \alpha(m_k, m_\ell) V_\varepsilon(x_k - x_\ell),$$

where we have also used the fact that the test function \hat{J} is of compact support, and the fact that the total number of particles living at any given time is bounded above by $Z\varepsilon^{2-d}$. From the bound on the collision that is provided by Lemma 4.1 it follows that

$$D^1 \leq \frac{c_1|z|}{\rho^{d-1}}.$$

To bound the term D^2 , note that by Lemma 4.2, the term D^2 is bounded above by

$$\begin{aligned} & \mathbb{E}_N \int_0^T \varepsilon^{2(d-2)} \sum_{k, \ell \in I_{\mathbf{q}}} \alpha(m_k, m_\ell) V_\varepsilon(x_k - x_\ell) \\ & \quad \times \sum_{i \in I_{\mathbf{q}}} \mathbb{1}\{|x_i - x_k| \leq \rho\} \left[|u^\varepsilon(x_k - x_i + z; m_k, m_i)| + |u^\varepsilon(x_k - x_i; m_k, m_i)| \right] \\ & \quad \times |\hat{J}(x_i, m_i, x_k, m_k, t)| dt \\ & \leq c_1 \mathbb{E}_N \int_0^T \varepsilon^{3(d-2)} \sum_{k, \ell \in I_{\mathbf{q}}} \alpha(m_k, m_\ell) V^\varepsilon(x_k - x_\ell) \\ & \quad \times \sum_{i \in I_{\mathbf{q}}} \mathbb{1}\{|x_i - x_k| \leq \rho, \max\{m_k, m_i, |x_k|, |x_i|\} \leq L, m_k + m_i \geq L^{-1}\} \\ & \quad \alpha'(m_k, m_i) \left[|x_k - x_i + z|^{2-d} + |x_k - x_i|^{2-d} \right] dt, \end{aligned}$$

where $V^\varepsilon = \varepsilon^{2-d} V_\varepsilon$ and L is chosen so that $\hat{J}(x, m, y, n) = 0$ if any of the conditions

$$m + n \geq L^{-1}, \quad \max(m, n) \leq L, \quad \max(|x|, |y|) \leq L,$$

does not hold. We note that if $m_k + m_i \geq L^{-1}$, then $\alpha'(m_k, m_i) \leq c_2\alpha(m_k, m_i)$, for a constant c_2 that depends on L . On the other hand, the conditions

$$m_k \leq L, \quad m_i \leq L, \quad m_k \text{ or } m_i \geq \frac{1}{2}L^{-1},$$

imply that for a constant $c_3 = c_3(L)$,

$$\alpha(m_k, m_\ell)\alpha(m_k, m_i) \leq c_3\gamma_3(m_i)\gamma_3(m_\ell)\gamma_3(m_k),$$

where we have used second part of Hypothesis 1.1. We are now in a position to apply Corollary 3.1. For this we choose $k = 3$ and

$$\begin{aligned} K(x_1, x_2, x_3) &= V^\varepsilon(x_1 - x_2) \mathbb{1}\{|x_2 - x_3| \leq \rho, |x_2|, |x_3| \leq L\} \\ & \quad \times \left[|x_2 - x_3 + z|^{2-d} + |x_2 - x_3|^{2-d} \right]. \end{aligned}$$

Note that K is discontinuous and Corollary 3.1 cannot be applied directly. We can readily replace the indicator function with an appropriate continuous function and obtain a new function \tilde{K} which is continuous off the set of points with $x_2 = x_3$. We can apply Corollary 3.1 to \tilde{K} as we explained in Remark 3.1. As a result, $D^2 \leq D(z) + D(0)$ where $D(z)$ is given by

$$\begin{aligned} c_4 \int V^\varepsilon(x_1 - x_2) \mathbb{1}\{|x_2 - x_3| \leq \rho, |x_2|, |x_3| \leq L\} |x_2 - x_3 + z|^{2-d} \prod_1^3 (\bar{h}_3 * \lambda_3)(x_r) dx_r \\ \leq c_5 \int V^\varepsilon(x_1 - x_2) \mathbb{1}\{|x_2 - x_3| \leq \rho, |x_2|, |x_3| \leq L\} |x_2 - x_3 + z|^{2-d} dx_1 dx_2 dx_3 \\ \leq c_6 \int_{|a| \leq \rho} |a + z|^{2-d} da \leq c_7(\rho + |z|)^2, \end{aligned}$$

where, for the first inequality, we used Hypothesis 1.2(ii). Combining these estimates yields

$$D_3 = D^1 + D^2 \leq c_1 \frac{|z|}{\rho^{d-1}} + c_7(\rho + |z|)^2.$$

Making the choice $\rho = |z|^{\frac{1}{d+1}}$ leads to the inequality $D_3 \leq c_8 |z|^{\frac{2}{d+1}}$. Since each of the cases of $\{D_i : i \in \{1, \dots, 8\}\}$ may be treated by a nearly verbatim proof, we are done. \square

Proof of (2.15). Setting $\mathbb{L} = \mathbb{A}_0 + \mathbb{A}_c$, the process

$$M_z(T) = X_z(\mathbf{q}(T), T) - X_z(\mathbf{q}(0), 0) - \int_0^T \left(\frac{\partial}{\partial t} + \mathbb{L} \right) X_z(\mathbf{q}(t), t) dt$$

is a martingale which satisfies

$$\mathbb{E}_N[M_z(T)^2] = \mathbb{E}_N \int_0^T \left(\mathbb{L} X_z^2 - 2X_z \mathbb{L} X_z \right) (\mathbf{q}(t), t) dt = \sum_{i=1}^3 \mathbb{E}_N \int_0^T A_i(\mathbf{q}(t), t) dt,$$

where

$$A_1(\mathbf{q}, t) = 2\varepsilon^{4(d-2)} \sum_{i \in I_{\mathbf{q}}} d(m_i) \left[\nabla_{x_i} \sum_{j \in I_{\mathbf{q}}} u^\varepsilon(x_i - x_j + z; m_i, m_j) \hat{J}(x_i, m_i, x_j, m_j, t) \right]^2,$$

and

$$A_2(\mathbf{q}, t) = 2\varepsilon^{4(d-2)} \sum_{j \in I_{\mathbf{q}}} d(m_j) \left[\nabla_{x_j} \sum_{i \in I_{\mathbf{q}}} u^\varepsilon(x_i - x_j + z; m_i, m_j) \hat{J}(x_i, m_i, x_j, m_j, t) \right]^2,$$

while $A_3(\mathbf{q}, t)$ is given by

$$\begin{aligned} \frac{1}{2} \varepsilon^{4(d-2)} \sum_{i, j \in I_{\mathbf{q}}} \alpha(m_i, m_j) \varepsilon^{-2} V_\varepsilon(x_i - x_j) \\ \times \left\{ \sum_{k \in I_{\mathbf{q}}} \left[\frac{m_i}{m_i + m_j} u^\varepsilon(x_i - x_k + z; m_i + m_j, m_k) \hat{J}(x_i, m_i + m_j, x_k, m_k, t) \right. \right. \\ \left. \left. + \frac{m_i}{m_i + m_j} u^\varepsilon(x_k - x_i + z; m_k, m_i + m_j) \hat{J}(x_k, m_k, x_i, m_i + m_j, t) \right. \right. \\ \left. \left. + \frac{m_j}{m_i + m_j} u^\varepsilon(x_j - x_k + z; m_i + m_j, m_k) \hat{J}(x_j, m_i + m_j, x_k, m_k, t) \right. \right. \end{aligned}$$

$$\begin{aligned}
 & + \frac{m_j}{m_i + m_j} u^\varepsilon(x_k - x_j + z; m_k, m_i + m_j) \hat{J}(x_k, m_k, x_j, m_i + m_j, t) \\
 & - u^\varepsilon(x_i - x_k + z; m_i, m_k) \hat{J}(x_i, m_i, x_k, m_k, t) \\
 & - u^\varepsilon(x_k - x_i + z; m_k, m_i) \hat{J}(x_k, m_k, x_i, m_i, t) \\
 & - u^\varepsilon(x_j - x_k + z; m_j, m_k) \hat{J}(x_j, m_j, x_k, m_k, t) \\
 & - u^\varepsilon(x_k - x_j + z; m_k, m_i) \hat{J}(x_k, m_k, x_j, m_j, t) \Big] \\
 & - u^\varepsilon(x_i - x_j + z; m_i, m_j) \hat{J}(x_i, m_i, x_j, m_j, t) \Big\}^2. \tag{4.11}
 \end{aligned}$$

We now bound the three terms. Of the first two, we treat only A_1 , the other being bounded by an identical argument. By multiplying out the brackets appearing in the definition of A_1 , and using $\sup_{m \in (0, \infty)} d(m) < \infty$, (which is assumed by Hypothesis 1.1), we obtain that $A_1 \leq A_{11} + A_{12}$ with

$$\begin{aligned}
 A_{11} &= c_0 \varepsilon^{4(d-2)} \sum_{i,j,k \in I_q} |u_x^\varepsilon(x_i - x_j + z; m_i, m_j)| |u_x^\varepsilon(x_i - x_k + z; m_i, m_k)| \\
 &\quad \times |\hat{J}(x_i, m_i, x_j, m_j, t)| |\hat{J}(x_i, m_i, x_k, m_k, t)| \\
 A_{12} &= c_0 \varepsilon^{4(d-2)} \sum_{i,j,k \in I_q} |u^\varepsilon(x_i - x_j + z; m_i, m_j)| |u^\varepsilon(x_i - x_k + z; m_i, m_k)| \\
 &\quad \times |\hat{J}_x(x_i, m_i, x_j, m_j, t)| |\hat{J}_x(x_i, m_i, x_k, m_k, t)|.
 \end{aligned}$$

Let us assume that $z = 0$ because this will not affect our arguments. We bound the term A_{11} with the aid of Corollary 3.1 and Lemma 4.2. The term A_{12} can be treated likewise. To bound A_{11} , first observe that even though i and j are distinct, k and j can coincide. Because of this, let us write $A_{11} = A_{111} + A_{112}$ where A_{111} represents the case of distinct i, j and k . We only show how to bound A_{111} where the correlation bound in the case of $k = 3$ is used. The term A_{112} can be treated in a similar fashion with the aid of Corollary 3.1 when $k = 2$. Since $\hat{J}(x, m, y, n) \neq 0$ implies that $m, n, |x|, |y| \leq L$ and $m + n \geq L^{-1}$. Using second part of Hypothesis 1.1, we can find a constant $c_1 = c_1(L)$ such that

$$\alpha(m_i, m_j) \alpha(m_i, m_k) \leq c_2 \gamma_3(m_i) \gamma_3(m_j) \gamma_3(m_k),$$

whenever

$$m_i, m_j, m_k \leq L, \quad m_i + m_j, m_i + m_k \geq L^{-1}.$$

As a result, we may apply Corollary 3.1 with $k = 3$ and

$$K(x_1, x_2, x_3) = \varepsilon^{d-2} |x_1 - x_2|^{1-d} |x_1 - x_3|^{1-d} \mathbb{1}(|x_1|, |x_2|, |x_3| \leq L),$$

to deduce

$$A_{111} \leq c_2 \varepsilon^{d-2} \int |x_1 - x_2|^{1-d} |x_1 - x_3|^{1-d} \mathbb{1}(|x_1|, |x_2|, |x_3| \leq L) \prod_{r=1}^3 (\bar{h}_3 * \lambda_3)(x_r) dx_r.$$

As before, we may approximate K with a continuous function \tilde{K} and we apply Corollary 3.1 as we explained in Remark 3.1. From this and Hypothesis 1.2, we deduce

$$A_{11} \leq c_3 \varepsilon^{d-2} \int |x_1 - x_2|^{1-d} |x_1 - x_3|^{1-d} \mathbb{1}(|x_1|, |x_2|, |x_3| \leq L) dx_1 dx_2 dx_3 = c_4 \varepsilon^{d-2}.$$

This and an analogous argument that treats the terms A_{112} , A_{12} and A_2 lead to the conclusion that

$$A_1 + A_2 \leq c_4 \varepsilon^{d-2}. \tag{4.12}$$

We must treat the third term, A_3 . An application of the inequality

$$(a_1 + \dots + a_n)^2 \leq n(a_1^2 + \dots + a_n^2)$$

to A_3 , given in (4.11), implies that

$$A_3(\mathbf{q}, t) \leq \frac{9}{2} \varepsilon^{4(d-2)} \sum_{i,j \in I_q} \alpha(m_i, m_j) V_\varepsilon(x_i - x_j) \left[\sum_{n=1}^8 \left(\sum_{k \in I_q} Y_n \right)^2 + Y_9^2 \right] =: \frac{9}{2} \sum_{i=1}^9 A_{3i}, \tag{4.13}$$

where Y_1 is given by

$$\frac{m_i}{m_i + m_j} u^\varepsilon(x_i - x_k + z; m_i + m_j, m_k) \hat{J}(x_i, m_i + m_j, x_k, m_k, t),$$

and where $\{Y_i : i \in \{2, \dots, 8\}\}$ denote the other seven expressions in (4.11) that appear in a sum over $k \in I_q$, while Y_9 denotes the last term in (4.11) that does not appear in this sum. There are nine cases to consider. The first eight are practically identical, and we treat only the fifth. Let us again assume that $z = 0$ because this will not affect our arguments. Note that

$$\begin{aligned} A_{35} &= \varepsilon^{4(d-2)} \sum_{i,j \in I_q} \alpha(m_i, m_j) V_\varepsilon(x_i - x_j) \left(\sum_{k \in I_q} Y_5 \right)^2 \\ &= \varepsilon^{5(d-2)} \sum_{i,j \in I_q} \alpha(m_i, m_j) V^\varepsilon(x_i - x_j) \left[\sum_{k,l \in I_q} u^\varepsilon(x_i - x_k; m_i, m_k) \right. \\ &\quad \left. \times u^\varepsilon(x_i - x_l; m_i, m_l) \hat{J}(x_i, m_i, x_k, m_k, t) \hat{J}(x_i, m_i, x_l, m_l, t) \right]. \end{aligned}$$

In the sum with indices involving $k, l \in I_q$, we permit the possibility that these two may be equal, though they must be distinct from each of i and j (which of course must themselves be distinct by the overall convention). Let us write $A_{35} = A_{351} + A_{352}$, where A_{351} corresponds to the case when all the indices i, j, k and l are distinct and A_{352} corresponds to the remaining cases. Again, our assumption on α as in Hypothesis 1.2 would allow us to treat the term A_{351} with the aid of Corollary 3.1. This time $k = 4$ and our bound on u given in the first part of Lemma 4.2 suggests the following choice for K :

$$K(x_1, \dots, x_4) = \varepsilon^{d-2} V^\varepsilon(x_1 - x_2) |x_1 - x_3|^{2-d} |x_1 - x_4|^{2-d} \mathbb{1}(|x_1|, |x_2|, |x_3|, |x_4| \leq L).$$

Again K can be approximated by a continuous function \tilde{K} and apply Corollary 3.1 as we explained in Remark 3.1. From Corollary 3.1 and Hypothesis 1.1 on the initial data we deduce that the expression $\int_0^T A_{351} dt$ is bounded above by

$$\begin{aligned} c_5 \varepsilon^{d-2} \int V^\varepsilon(x_1 - x_2) |x_1 - x_3|^{2-d} |x_1 - x_4|^{2-d} \mathbb{1}(|x_1|, |x_2|, |x_3|, |x_4| \leq L) dx_1 \dots dx_4 \\ = c_6 \varepsilon^{d-2}. \end{aligned}$$

A similar reasoning applies to A_{352} , except that [Corollary 3.1](#) in the case of $k = 3$ would be employed. Hence,

$$\sum_{i=1}^8 A_{3i} \leq c_7 \varepsilon^{d-2}. \tag{4.14}$$

We now treat the ninth term, as they are classified in [\(4.13\)](#). It takes the form

$$\varepsilon^{4d-8} \sum_{i,j \in I_q} \alpha(m_i, m_j) V_\varepsilon(x_i - x_j) u^\varepsilon(x_i - x_j + z; m_i, m_j)^2 \hat{J}(x_i, m_i, x_j, m_j, t)^2.$$

This is bounded above by

$$c_8 \varepsilon^{2d-4} \sum_{i,j \in I_q} \alpha(m_i, m_j) V_\varepsilon(x_i - x_j),$$

because $u^\varepsilon \leq c_9 \varepsilon^{2-d}$ by the first part of [Lemma 4.2](#). The expected value of the integral on the interval of time $[0, T]$ of this last expression is bounded above by

$$c_7 \varepsilon^{2d-4} \mathbb{E}_N \int_0^T \sum_{i,j \in I_q} \alpha(m_i, m_j) V_\varepsilon(x_i - x_j) dt \leq c_{10} \varepsilon^{d-2}.$$

where we used [Lemma 4.1](#) for the last inequality. This and [\(4.12\)–\(4.14\)](#) complete the proof of [\(2.15\)](#). \square

5. Bounds on the macroscopic densities

In this section we show how [Corollary 3.1](#) can be used to obtain certain bounds on the macroscopic densities. These bounds will be used for the derivation of the macroscopic equation. Recall that

$$g^\varepsilon(dx, dn, t) = \varepsilon^{d-2} \sum_i \delta_{(x_i(t), m_i(t))}(dx, dn),$$

and that the law of

$$\mathbf{q} \mapsto g^\varepsilon(dx, dn, t)$$

induces a probability measure \mathcal{P}^ε on the space \mathcal{X} . The main result is [Theorem 5.1](#).

Theorem 5.1. *Let \mathcal{P} be a limit point of \mathcal{P}^ε . The following statements are true:*

- 1. For every positive L_1 , and $k \in \{2, 3, 4\}$,

$$\sup_\delta \int_{\mathcal{X}} \int_0^\infty \int_{|x| \leq L_1} \left[\int_0^\infty \int \xi^\delta(x - y) \gamma_k(n) g(dy, dn, t) \right]^k dx dt d\mathcal{P} < \infty, \tag{5.1}$$

where $\xi^\delta(x) = \delta^{-d} \xi(\frac{x}{\delta})$, with ξ a non-negative smooth function of compact support satisfying $\int \xi = 1$.

- 2. We have $g(dx, dn, t) = f(x, t, dn)dx$ for almost all g with respect to the probability measure \mathcal{P} .

- 3. For every continuous R of compact support and positive L ,

$$\lim_{\delta \rightarrow 0} \int \left| \int_0^T \int_{L^{-1}}^L \int_{L^{-1}}^L \int R(x, m, n, t) f^\delta(x, t, dm) f^\delta(x, t, dn) dx dt - \int_0^T \int_{L^{-1}}^L \int_{L^{-1}}^L \int R(x, m, n, t) f(x, t, dm) f(x, t, dn) dx dt \right| d\mathcal{P} = 0, \tag{5.2}$$

where

$$f^\delta(x, t, dn) = \int \xi^\delta(x - y) g(dy, dn, t). \tag{5.3}$$

Proof. Fix $x \in \mathbb{R}^d$ and choose

$$K(y_1, \dots, y_k) = \prod_{r=1}^k \xi^\delta(x - y_r),$$

in Corollary 3.1. The right-hand side of (3.7) equals

$$\int \prod_{r=1}^k \xi^\delta(x - x_r) \bar{h}_k * \lambda_k(x_r) dx_r,$$

which, by the second part of Hypothesis 1.2, is bounded by a constant $c_1(L_1)$ when $k = 2, 3, 4$, and $|x| \leq L_1$. As a result,

$$\mathbb{E}_N \int_0^\infty \int_{|x| \leq L_1} \varepsilon^{k(d-2)} \sum_{i_1, \dots, i_k} \prod_{r=1}^k \xi^\delta(x - x_{i_r}(t)) \gamma_k(m_{i_r}(t)) dx dt \leq c_1(L_1) \tag{5.4}$$

for a constant $c_1(L)$ which is independent of δ and ε . Here we are assuming that the indices i_1, \dots, i_k are distinct. Note that if we allow non-distinct indices in the summation, then the difference would go to 0 as $\varepsilon \rightarrow 0$ because the summation is multiplied by $\varepsilon^{k(d-2)}$ while the number of additional terms is of order $O(\varepsilon^{(k-1)(2-d)})$. As a consequence, we can use (5.4) to deduce (5.1).

Recall that the function γ_k is a positive continuous function. From this and (5.1), one can readily deduce part 2.

It remains to establish part 3. First observe that by (5.1) and the positivity of γ_4 ,

$$\sup_{\delta} \int \int_0^T \int_{|x| \leq L_1} \left[\int_{L^{-1}}^L f^\delta(x, t, dn) \right]^4 dx dt \mathcal{P}(dg) \leq c_2(L_1, L). \tag{5.5}$$

Because of this, it suffices to prove that

$$\begin{aligned} & \lim_{\delta \rightarrow 0} \int_0^T \int \int_{L^{-1}}^L \int_{L^{-1}}^L R_p(x, m, n, t) f^\delta(x, t, dm) f^\delta(x, t, dn) dx dt \\ & = \int_0^T \int_{L^{-1}}^L \int_{L^{-1}}^L \int R_p(x, m, n, t) f(x, t, dm) f(x, t, dn) dx \end{aligned}$$

for each p , provided that $\lim_{p \rightarrow \infty} R_p(x, m, n, t) = R(x, m, n, t)$, uniformly for $m, n \in [L^{-1}, L]$, $|x| \leq L_1$ and $t \leq T$. By approximation, we may assume that R is of the form

$R(x, m, n, t) = \sum_{i=1}^{\ell} J_1^{\ell}(x, t) J_2^{\ell}(m) J_3^{\ell}(n)$. Hence it suffices to establish (5.2) for R of the form $R(x, m, n, t) = J_1(x, t) J_2(m) J_3(n)$. But now the left-hand side of (5.2) equals

$$\lim_{\delta \rightarrow 0} \int_0^T \int \left[\int_{L^{-1}}^L J_2(m) f^{\delta}(x, t, dm) \right] \left[\int_{L^{-1}}^L J_3(n) f^{\delta}(x, t, dn) \right] J_1(x, t) dx dt.$$

We note that

$$\int_{L^{-1}}^L J_2(m) f^{\delta}(x, t, dm) = \left(\int_{L^{-1}}^L J_2(m) f(\cdot, t, dm) \right) *_x \xi^{\delta}(x).$$

converges almost everywhere to

$$\int_{L^{-1}}^L J_2(m) f(x, t, dm).$$

The same comment applies to $\int_{L^{-1}}^L J_3(n) f_n^{\delta}(x, t) dn$. From this and (5.5) we deduce (5.2). \square

6. Deriving the PDE

We wish to derive (1.6) from the identity (2.1). There is a technical issue that we need to settle first: in (2.2), the function $\hat{J}(x, m, y, n, t)$ does not have a compact support with respect to (m, n) , even if J is of compact support. Recall that in Theorem 1.1 we have assumed that \hat{J} is of compact support. Lemma 6.1 settles this issue.

Lemma 6.1. *There exists a constant C_4 independent of ε such that*

$$\mathbb{E}_N \int_0^T \varepsilon^{2(d-2)} \sum_{i,j \in I_q} \alpha(m_i(t), m_j(t)) V_{\varepsilon}(x_i(t) - x_j(t)) m_i(t) m_j(t) dt \leq C_4. \tag{6.1}$$

Moreover,

$$\begin{aligned} \lim_{L \rightarrow \infty} \sup_{\varepsilon} \mathbb{E}_N \int_0^T \varepsilon^{2(d-2)} \sum_{i,j \in I_q} \alpha(m_i(t), m_j(t)) V_{\varepsilon}(x_i(t) - x_j(t)) \mathbb{1} \\ \times (\min\{m_i(t), m_j(t)\} \leq L^{-1}) dt = 0. \end{aligned} \tag{6.2}$$

Proof. Let us take a smooth function $J : \mathbb{R}^d \rightarrow [0, \infty)$ and set

$$H(x) = c_0(d) \int \frac{J(y)}{|x - y|^{d-2}} dy \tag{6.3}$$

with $c_0(d) = (d - 2)^{-1} \omega_d^{-1}$ with ω_d denoting the surface area of the unit sphere in \mathbb{R}^d . Note that $H \geq 0$ and $-\Delta H = J$. Let $\psi : (0, \infty) \times (0, \infty) \rightarrow [0, \infty)$ be a continuous symmetric function and set

$$X_N(\mathbf{q}) = \varepsilon^{2(d-2)} \sum_{i,j \in I_q} H(x_i - x_j) \psi(m_i, m_j). \tag{6.4}$$

We have

$$\begin{aligned} - \mathbb{E}_N \int_0^T \mathcal{A}_c X_N(\mathbf{q}(s)) ds - \mathbb{E}_N \int_0^T \mathcal{A}_0 X_N(\mathbf{q}(s)) ds &= \mathbb{E}_N X_N(\mathbf{q}(0)) - \mathbb{E}_N X_N(\mathbf{q}(T)) \\ &\leq \mathbb{E}_N X_N(\mathbf{q}(0)), \end{aligned} \tag{6.5}$$

where

$$\mathcal{A}_0 X_N(\mathbf{q}) = -\varepsilon^{2(d-2)} \sum_{i,j \in I_{\mathbf{q}}} J(x_i - x_j) \psi(m_i, m_j) (d(m_i) + d(m_j)),$$

and $\mathcal{A}_c X_N(\mathbf{q}) = Y_1(\mathbf{q}) + Y_2(\mathbf{q})$, with

$$Y_1(\mathbf{q}) = -\frac{1}{2} \varepsilon^{2(d-2)} \sum_{i,j \in I_{\mathbf{q}}} \alpha(m_i, m_j) V_\varepsilon(x_i - x_j) \psi(m_i, m_j) H(x_i - x_j)$$

$$Y_2(\mathbf{q}) = \frac{1}{2} \varepsilon^{2(d-2)} \sum_{i,j,k \in I_{\mathbf{q}}} \alpha(m_i, m_j) V_\varepsilon(x_i - x_j) \Gamma(x_i, x_j, x_k, m_i, m_j, m_k),$$

where

$$\begin{aligned} \Gamma(x_i, x_j, x_k, m_i, m_j, m_k) = & \left[\frac{m_i}{m_i + m_j} \psi(m_i + m_j, m_k) - \psi(m_i, m_k) \right] H(x_i - x_k) \\ & + \left[\frac{m_j}{m_i + m_j} \psi(m_i + m_j, m_k) - \psi(m_j, m_k) \right] H(x_j - x_k) \\ & + \left[\frac{m_i}{m_i + m_j} \psi(m_k, m_i + m_j) - \psi(m_k, m_i) \right] H(x_k - x_i) \\ & + \left[\frac{m_j}{m_i + m_j} \psi(m_k, m_i + m_j) - \psi(m_k, m_j) \right] H(x_k - x_j). \end{aligned}$$

We consider two examples for ψ . As the first example, we choose $\psi(m, n) = mn$. This yields $Y_2 = 0$. We find that

$$\sup_N \mathbb{E}_N \int_0^T Y_1(\mathbf{q}(s)) \, ds \leq \mathbb{E}_N X_N(\mathbf{q}(0)). \tag{6.6}$$

The hope is that a suitable choice of J would yield the desired assertion (6.1). For this, we simply choose $J(x) = \varepsilon^{-d} A\left(\frac{x}{\varepsilon}\right)$ where A is a smooth non-negative function of compact support. We then have that $H(x) = \varepsilon^{2-d} B\left(\frac{x}{\varepsilon}\right)$ where $\Delta B = -A$. As a result,

$$Y_1(\mathbf{q}) = \frac{1}{2} \varepsilon^{d-2} \sum_{i,j \in I_{\mathbf{q}}} V_\varepsilon(x_i - x_j) B\left(\frac{x_i - x_j}{\varepsilon}\right) m_i m_j \alpha(m_i, m_j) \tag{6.7}$$

with

$$B(x) = c_0(d) \int \frac{A(y)}{|x - y|^{d-2}} \, dy.$$

Recall that the support of V is contained in the set y with $|y| \leq C_0$. If we choose A so that

$$\mathbb{1}(|y| \leq 3C_0) \leq A(y) \leq \mathbb{1}(|y| \leq 4C_0),$$

then, for $|x| \leq C_0$,

$$B(x) \geq c_0(d) \int_{3C_0 \geq |y| \geq 2C_0} \frac{dy}{|x - y|^{d-2}} \leq c_0(d) C_0^{2-d} \int_{3C_0 \geq |y| \geq 2C_0} dy =: \tau_0 > 0.$$

On the other hand, if $|x| \leq 5C_0$, then

$$B(x) \leq c_0(d) \int_{|x-y| \leq 9C_0} \frac{dy}{|x - y|^{d-2}} = \frac{1}{2} c_0(d) \omega_d (9C_0)^2 \tag{6.8}$$

and if $|x| \geq 5C_0$, then

$$B(x) \leq c_0(d) \left| \frac{4x}{5} \right|^{2-d} \int_{C_0 \geq |y|} dy = c_1 |x|^{2-d}.$$

From this, (6.8) and the third part of Hypothesis 1.2, we learn that the right-hand side of (6.6) is uniformly bounded in ε . This completes the proof of (6.1). \square

As for (6.2), we choose $\psi(m, n) = \mathbb{1}(m \leq \delta) + \mathbb{1}(n \leq \delta)$. This time we have that $Y_2 \leq 0$. Such a function ψ is not continuous. But by a simple approximation procedure we can readily see that (6.5) is valid for such a choice. By the third part of Hypothesis 1.2 on the initial data, we know that

$$\int_0^\infty \int h_n(x) \hat{h}(y) |x - y|^{2-d} dx dy dn < \infty.$$

From this we learn that

$$\lim_{\delta \rightarrow 0} \int_0^\delta \int h_n(x) \hat{h}(y) |x - y|^{2-d} dx dy dn = 0,$$

whence

$$\limsup_{\delta \rightarrow 0} \mathbb{E}_N X_N(\mathbf{q}(0)) = 0.$$

This and (6.5) imply (6.2).

Proof of Theorem 1.1. *Step 1.* We take a smooth test function J of compact support in $\mathbb{R}^d \times (0, \infty) \times [0, \infty)$ and study the decomposition (2.1). Firstly, we show that the martingale term goes to 0. The term M is a martingale satisfying

$$\begin{aligned} \mathbb{E}_N[M_T^2] &= \mathbb{E}_N \int_0^T (\mathbb{L}Y^2 - 2Y\mathbb{L}Y)(\mathbf{q}(t), t) dt \\ &= \mathbb{E}_N \int_0^T A_1(\mathbf{q}(t), t) dt + \mathbb{E}_N \int_0^T A_2(\mathbf{q}(t), t) dt, \end{aligned}$$

where $A_1(\mathbf{q}, t)$ and $A_2(\mathbf{q}, t)$ are respectively set equal to

$$A_1(\mathbf{q}, t) = \varepsilon^{2(d-2)} \sum_{i \in I_{\mathbf{q}}} d(m_i) |J_x(x_i, m_i, t)|^2,$$

and

$$A_2(\mathbf{q}, t) = \frac{1}{2} \varepsilon^{2(d-2)} \sum_{i, j \in I_{\mathbf{q}}} \alpha(m_i, m_j) V_\varepsilon(x_i - x_j) \hat{J}(x_i, m_i, x_j, m_j, t)^2.$$

We can readily show

$$A_1(\mathbf{q}, t) \leq c_1 \varepsilon^{2(d-2)} \sum_{i \in I_{\mathbf{q}}} d(m_i) \leq c_2 \varepsilon^{d-2}, \tag{6.9}$$

$$\mathbb{E}_N \int_0^T A_2(\mathbf{q}(t), t) dt \leq c_3 \mathbb{E}_N \int_0^T \varepsilon^{2(d-2)} \sum_{i, j \in I_{\mathbf{q}}} \alpha(m_i, m_j) V_\varepsilon(x_i - x_j) dt \leq c_4 \varepsilon^{d-2}, \tag{6.10}$$

where we have Lemma 4.1 in the last inequality. From these inequalities, we deduce that the martingale tends to zero, in the $\varepsilon \downarrow 0$ limit.

Step 2. We rewrite the terms of (2.1) in terms of the empirical measures. We have that

$$Y(\mathbf{q}(t), t) = \int_0^\infty \int_{\mathbb{R}^d} J(x, n, t)g(dx, dn, t), \tag{6.11}$$

and that

$$\int_0^T \left(\frac{\partial}{\partial t} + \mathbb{A}_0 \right) Y(\mathbf{q}(t), t) dt = \int_0^T \int_0^\infty \int_{\mathbb{R}^d} \left(\frac{\partial}{\partial t} + d(n)\Delta_x \right) J(x, n, t)g(dx, dn, t). \tag{6.12}$$

Furthermore, by Theorem 1.2 and Lemma 6.1,

$$\int_0^T \mathbb{A}_c Y(\mathbf{q}(t), t) dt = \frac{1}{2} \int_0^T \Gamma_L^\delta(\mathbf{q}(t), t) dt + Err^1(\varepsilon, L) + Err^2(\varepsilon, \delta, L), \tag{6.13}$$

where T is large enough that $J(\cdot, \cdot, t) = 0$ for $t \geq T$, the expression $\Gamma_L^\delta(\mathbf{q}, t)$ is given by

$$\begin{aligned} & \iint \int_{L^{-1}}^L \int_{L^{-1}}^L \alpha(m, n)U_{n,m}^\varepsilon(w_1 - w_2) f^\delta(w_1, dm; \mathbf{q}) \\ & \times f^\delta(w_2, dn; \mathbf{q}) \hat{J}(w_1, m, w_2, n, t) dw_1 dw_2, \end{aligned}$$

and

$$\lim_{L \rightarrow \infty} \sup_\varepsilon \mathbb{E}_N |Err^1(\varepsilon, L)| = 0, \quad \lim_{\delta \rightarrow 0} \limsup_{\varepsilon \rightarrow 0} \mathbb{E}_N |Err^2(\varepsilon, \delta, L)| = 0.$$

We note that if we replace $f^\delta(w_2, dn; \mathbf{q}) \hat{J}(w_1, m, w_2, n, t)$ with $f^\delta(w_1, dn; \mathbf{q}) \hat{J}(w_1, m, w_1, n, t)$, then we produce an error which is of order $O(L\delta^{-\delta-1}\varepsilon)$, which goes to 0 because we send $\varepsilon \rightarrow 0$ first. As a result, (6.13) equals

$$\begin{aligned} & \frac{1}{2} \int_0^\infty \int_{\mathbb{R}^d} \int_{L^{-1}}^L \int_{L^{-1}}^L \beta(m, n)(g *_x \xi^\delta)(x, t, dm)(g *_x \xi^\delta)(x, t, dn) \tilde{J}(x, m, n, t) dx dt \\ & + Err^1(\varepsilon, L) + Err^3(\varepsilon, \delta, L), \end{aligned}$$

where

$$\lim_{\delta \rightarrow 0} \limsup_{\varepsilon \rightarrow 0} \mathbb{E}_N |Err^3(\varepsilon, \delta, L)| = 0.$$

By passing to the limit in low ε , we find that any weak limit \mathcal{P} is concentrated on the space of measures $g(dx, dn, t) dt$ such that

$$\begin{aligned} & \int_0^\infty \int_{\mathbb{R}^d} h_n(x)J(x, n, 0) dx dn + \int_0^\infty \int_0^\infty \int_{\mathbb{R}^d} g(dx, dn, t) \left(\frac{\partial}{\partial t} + d(n)\Delta_x \right) J(x, n, t) dt \\ & + \frac{1}{2} \int_0^\infty \int_{\mathbb{R}^d} \int_{L^{-1}}^L \int_{L^{-1}}^L \beta(m, n)(g *_x \xi^\delta)(x, t, dm)(g *_x \xi^\delta)(x, t, dn) \tilde{J}(x, m, n, t) dx dt \\ & + Err^4(L) + Err^5(\delta) = 0, \end{aligned} \tag{6.14}$$

where the \mathcal{P} -expectation of $|Err^5(\delta)|$ goes to zero as $\delta \downarrow 0$, and the \mathcal{P} -expectation of $|Err^4(L)|$ goes to zero as $L \rightarrow \infty$. From Theorem 5.1 we know that $g(dx, dn, t) = f(x, t, dn)dx$, \mathcal{P} -almost surely and that by (5.2) we can replace $g *_x \xi$ with f . Hence

$$\begin{aligned} & \int_0^\infty \int_{\mathbb{R}^d} h_n(x) J(x, n, 0) dx dn \\ & + \int_0^\infty \int_0^\infty dt \int_{\mathbb{R}^d} f(x, t, dn) \left(\frac{\partial}{\partial t} + d(n)\Delta_x \right) J(x, n, t) \\ & + \frac{1}{2} \int_0^\infty \int_{L^{-1}}^L \int_{L^{-1}}^L \int_{\mathbb{R}^d} \beta(m, n) f(x, t, dm) f(x, t, dn) \tilde{J}(x, m, n, t) dx dt \\ & + Err^4(L) = 0. \end{aligned} \tag{6.15}$$

It remains to replace L^{-1} and L with 0 and ∞ respectively. For this, recall that by assumption, there exists ℓ such that $J(x, m, t) = 0$ if $m \notin (\ell^{-1}, \ell)$. Hence, when $\tilde{J}(x, m, n, t) \neq 0$, we must have that $m + n > \ell^{-1}$ and $\min\{m, n\} < \ell$. By the first remark that we made after the statement of Theorem 1.1, we know that $\beta \leq \alpha$. From the second part of Hypothesis 1.1 we deduce that there exists a constant $c_5 = c_5(\ell)$ such that $\beta(m, n) \leq \alpha(m, n) \leq c_5 \gamma_2(m) \gamma_2(n)$ provided that $m + n > \ell^{-1}$ and $\min\{m, n\} < \ell$. (Here we are using the fact that $d(m)^{d/2} \phi^{d-1}$ is uniformly positive and bounded over the interval $[\ell^{-1}/2, \ell]$.) On the other hand, we know by part 1 of Theorem 5.1 that

$$\int_0^T \int_{|x| \leq L_1} \int_0^\infty \int_0^\infty \gamma_2(n) \gamma_2(m) f(x, t, dm) f(x, t, dn) dx dt < \infty,$$

\mathcal{P} -almost surely, where L_1 is chosen so that the set $\{|x| \leq L_1\}$ contains the support of \tilde{J} in the spatial variable. From this we deduce

$$\begin{aligned} & \lim_{L \rightarrow \infty} \int_0^T \int \int_0^\infty \int_0^\infty \beta(m, n) f(x, t, dm) f(x, t, dn) \\ & \mathbb{1} \left(\max\{m, n\} \geq L \quad \text{or} \quad \min\{m, n\} \leq L^{-1} \right) \tilde{J}(x, m, n, t) dx dt = 0. \end{aligned}$$

This allows us to replace L^{-1} and L with 0 and ∞ respectively in (6.15), concluding that $f(x, t, dn)$ solves (1.1) weakly in the sense of (1.6). \square

As we stated in Section 1, the family \mathcal{P}^ε is defined on a compact metric space \mathcal{X} which consists of measures $\mu(dx, dn, dt)$ which are absolutely continuous with respect to the time variable. This can be proved by standard arguments.

Lemma 6.2. *Every measure $\mu \in \mathcal{X}$ is of the form $\mu(dx, dn, dt) = g(dx, dn, t)dt$.*

Proof. Let $J_k : \mathbb{R}^d \times [0, \infty) \rightarrow \mathbb{R}$, $k \in \mathbb{N}$, be a sequence of linearly independent continuous functions of compact support such that $J_1 = 1$ and the linear span Y of this sequence is dense in the space of continuous functions of compact support. Given $\mu \in \mathcal{X}$, it is not hard to show that for each k , there exists a measurable function $G_{J_k} : [0, T] \rightarrow \mathbb{R}$ such that $\|G_{J_k}\|_{L^\infty} \leq Z \sup_{x,n} |J_k(x, n)|$, and

$$\int_{\mathbb{R}^d} \int_0^\infty J_k(x, n) \mu(dx, dn, dt) = G_{J_k}(t) dt.$$

We wish to define G_J for every continuous J of compact support. Note that each G_{J_k} is defined almost everywhere in the interval $[0, \infty)$. For our purposes, we need to construct G_J in such a way that for almost all t , the operator $J \mapsto G_J(t)$ is linear. For this, let us set $G_J = r_1 G_{J_1} + \dots + r_l G_{J_l}$ when $J = r_1 J_1 + \dots + r_l J_l$ with r_1, \dots, r_l rational. The set of such J is denoted by Y' . Since Y' is countable, There exists a set $A \subset [0, \infty)$ of 0 Lebesgue measure such that for $t \notin A$, the operator $J \mapsto G_J(t)$ from Y' to \mathbb{R} is linear over rationals. By denseness of rationals, we can extend $J \mapsto G_J(t)$ for $J \in Y$ and $t \notin A$. For such (J, t) ,

$$\int_{\mathbb{R}^d} \int_0^\infty J(x, n) \mu(dx, dn, dt) = G_J(t) dt.$$

We then take a point in $[0, \infty) \setminus A$ and use Riesz Representation Theorem to find a measure $g(dx, dn, t)$ such that

$$G_J(t) = \int_{\mathbb{R}^d} \int_0^\infty J(x, n) g(dx, dn, t),$$

for every $J \in Y$. Hence

$$\int_{\mathbb{R}^d} \int_0^\infty J(x, n) \mu(dx, dn, dt) = \int_{\mathbb{R}^d} \int_0^\infty J(x, n) g(dx, dn, t) dt.$$

for every $J \in Y$. This completes the proof. \square

7. Entropy

In this section, we establish entropy-like inequalities to show that the macroscopic density g is absolutely continuous with respect to Lebesgue measure.

Proof of Theorem 1.2. *Step 1.* Recall that initially we have \mathcal{N} particles. We choose $I_{\mathbf{q}(0)} = \{1, \dots, \mathcal{N}\}$, and label the initial particles as $(x_1, m_1), \dots, (x_{\mathcal{N}}, m_{\mathcal{N}})$. If a coagulation occurs at time t , one of the coagulating particles disappears from the system, and $I_{\mathbf{q}} \subseteq \{1, \dots, \mathcal{N}\}$ satisfies $|I_{\mathbf{q}(t^+)}| = |I_{\mathbf{q}(t)}| - 1$. We write $\mathcal{N}(\mathbf{q}) = |I_{\mathbf{q}}|$ for the number of particles of the configuration \mathbf{q} . Note that $\mathcal{N}(\mathbf{q})$ takes values in the set $\{1, \dots, \mathcal{N}\}$. We write $F(\mathbf{q}, t) \nu_N(d\mathbf{q})$ for the law of $\mathbf{q}(t)$, and define

$$H_N(t) = \int F(\mathbf{q}, t) \log F(\mathbf{q}, t) \nu_N(d\mathbf{q}).$$

By standard arguments,

$$\frac{\partial H_N}{\partial t}(t) = \int \left(\mathbb{L}(\log F)(\mathbf{q}, t) \right) F(\mathbf{q}, t) \nu_N(d\mathbf{q}) = \Omega_1 + \Omega_2, \tag{7.1}$$

where

$$\Omega_1 = \int \left(\mathbb{A}_0(\log F)(\mathbf{q}, t) \right) F(\mathbf{q}, t) \nu_N(d\mathbf{q}),$$

$$\Omega_2 = \int \left(\mathbb{A}_c(\log F)(\mathbf{q}, t) \right) F(\mathbf{q}, t) \nu_N(d\mathbf{q}).$$

We have

$$\Omega_1 = \int \sum_{i \in I_{\mathbf{q}}} d(m_i) (\Delta_{x_i} F) \log F \, d\nu_N$$

$$\begin{aligned}
 &= - \int \sum_{i \in I_{\mathbf{q}}} d(m_i) \frac{|\nabla_{x_i} F|^2}{F} \, d\nu_N + \int \sum_{i \in I_{\mathbf{q}}} d(m_i) \nabla_{x_i} F \cdot x_i \, d\nu_N \\
 &= - \int \sum_{i \in I_{\mathbf{q}}} d(m_i) \frac{|\nabla_{x_i} F|^2}{F} \, d\nu_N - \int \sum_{i \in I_{\mathbf{q}}} d(m_i) (d - |x_i|^2) F \, d\nu_N \\
 &\leq D \int \sum_{i \in I_{\mathbf{q}}} |x_i|^2 F \, d\nu_N,
 \end{aligned}$$

where we integrated by parts for the second and third equality, and D is an upper bound for the function $d(\cdot)$. To bound the right-hand side, we use the Markov property of the process $\mathbf{q}(t)$ to write

$$\begin{aligned}
 \mathbb{E}_N \sum_{i \in I_{\mathbf{q}(t)}} |x_i(t)|^2 &\leq \mathbb{E}_N \sum_{i \in I_{\mathbf{q}(0)}} |x_i(0)|^2 + 2d \int_0^t \mathbb{E}_N \sum_{i \in I_{\mathbf{q}(s)}} d(m_i(s)) \, ds \\
 &\leq c\varepsilon^{2-d} + 2dt D Z \varepsilon^{2-d},
 \end{aligned}$$

where, in the first inequality, we used that the coagulation is non-positive, which follows from our assumption that a particle, newly born in a coagulation event, is placed in the location of one of the departing particles. The second inequality is due to our assumption that D is a uniform upper bound on $d : (0, \infty) \rightarrow (0, \infty)$ and to the hypothesis we make on the initial condition. We learn that

$$\Omega_1 \leq c_1(t + 1)\varepsilon^{2-d}. \tag{7.2}$$

We now concentrate on the contribution coming from coagulations, namely the expression Ω_2 . This expression equals

$$\begin{aligned}
 &\frac{1}{2} \int \sum_{i,j \in I_{\mathbf{q}}} V_\varepsilon(x_i - x_j) \alpha(m_i, m_j) \left[\frac{m_i}{m_i + m_j} \log \frac{F(S_{i,j}^1 \mathbf{q}, t)}{F(\mathbf{q}, t)} + \frac{m_j}{m_i + m_j} \log \frac{F(S_{i,j}^2 \mathbf{q}, t)}{F(\mathbf{q}, t)} \right] \\
 &\quad \times F(\mathbf{q}, t) \, \nu_N(d\mathbf{q}) \\
 &\leq \frac{1}{2} \int \sum_{i,j \in I_{\mathbf{q}}} V_\varepsilon(x_i - x_j) \alpha(m_i, m_j) \left[\frac{m_i}{m_i + m_j} F(S_{i,j}^1 \mathbf{q}, t) + \frac{m_j}{m_i + m_j} F(S_{i,j}^2 \mathbf{q}, t) \right] \\
 &\quad \times \nu_N(d\mathbf{q}) \\
 &= \frac{1}{2} \int \sum_{i,j \in I_{\mathbf{q}}} V_\varepsilon(x_i - x_j) \alpha(m_i, m_j) F(S_{i,j}^1 \mathbf{q}, t) \, \nu_N(d\mathbf{q}),
 \end{aligned}$$

where we used the elementary inequality $\log x \leq x$ for the second line. To bound this, we first observe

$$\int V_\varepsilon(x_i - x_j) (2\pi)^{-d/2} \exp\left(-\frac{|x_j|^2}{2}\right) \, dx_i \leq (2\pi)^{-d/2} \int V_\varepsilon(x_i - x_j) \, dx_i \leq C\varepsilon^{d-2}.$$

We then make a change of variables $m_i + m_j \mapsto m_i$. As a result, Ω_2 is bounded above by

$$\varepsilon^{d-2} \int \sum_{i \in I_{\mathbf{q}}} \rho(m_i) F(\mathbf{q}, t) \, d\nu_N(d\mathbf{q}),$$

where the function ρ is defined (1.8).

From the second part of Hypothesis 1.3, we deduce that Ω_2 is bounded by a constant multiple of ε^{d-2} . This, the first part of Hypothesis 1.3, and (7.2) yield

$$H_N(t) \leq c_2(t + 1)\varepsilon^{d-2}. \tag{7.3}$$

Step 2. Note that by Sanov’s theorem, the empirical measure $\varepsilon^{d-2} \sum_i \delta_{(x_i, m_i)}$ satisfies a large deviation principle with respect to the measure ν_N as $\varepsilon \rightarrow 0$. The large deviation rate function $\mathcal{I}(g) = \infty$ unless $g(dx, dn) = f(x, n)r(x, n) dx dn$ and if such a function f exists, then

$$\mathcal{I}(g) = \int_0^\infty \int (f \log f - f + 1)r \, dx \, dn.$$

By an argument similar to the proof of Lemma 6.3 of [3], we can use (7.3) to deduce that if \mathcal{P} is any limit point of the sequence \mathcal{P}^ε , then

$$\int \mathcal{I}(g(\cdot, t)) \mathcal{P}(dg) < \infty,$$

for every t . This completes the proof of Theorem 1.2. \square

Acknowledgments

The work of the second author is supported in part by NSF grant DMS-0707890. We thank James Colliander and Pierre Germain for valuable comments that relate to the discussion in the appendix. We also thank anonymous referees for a number of useful suggestions and comments.

Appendix. Scaling of the continuous Smoluchowski equation

We comment on the scaling satisfied by the system (1.1), under the assumptions that

$$d(n) = n^{-\phi}$$

and

$$\beta(n, m) = n^\eta + m^\eta, \tag{A.1}$$

with $\phi, \eta \in [0, \infty)$. Rescaling the equations,

$$g_n(x, t) = \lambda^\alpha f_{n\lambda^\gamma}(\lambda^\tau x, \lambda t), \tag{A.2}$$

we note that g_n satisfies (1.1) provided that

$$1 - \gamma\phi - 2\tau = 0 \tag{A.3}$$

and

$$-\alpha + \gamma(1 + \eta) + 1 = 0, \tag{A.4}$$

(A.3) ensuring that the free motion term is preserved, (A.4) the interaction term. The mass

$$h_f(t) = \int_0^\infty n \int_{\mathbb{R}^d} f_n(x, t) \, dx \, dn,$$

which, formally at least, is conserved in time, is mapped by the rescaling to

$$h_g(t) = \lambda^{\alpha - \tau d - 2\gamma} h_f(\lambda t). \tag{A.5}$$

The mass, then, is conserved by the rescaling provided that

$$\alpha - \tau d - 2\gamma = 0. \tag{A.6}$$

In the critical case, where each of (A.3), (A.4) and (A.6) is satisfied, we have that

$$\begin{aligned} \gamma &= \frac{d/2 - 1}{\eta + \phi d/2 - 1}, \\ \alpha &= \frac{d/2(\phi + \eta + 1) - 2}{\eta + \phi d/2 - 1} \end{aligned}$$

and

$$\tau = \frac{\eta + \phi - 1}{2(\eta + \phi d/2 - 1)}. \tag{A.7}$$

In the case where the dimension $d = 2$, the values $\gamma = 0$, $\alpha = 1$ and $\tau = 1/2$ are adopted, whatever the values taken for the input parameters ϕ and η . The only critical scaling, then, leaves the mass unchanged and performs a diffusive rescaling of space–time.

Regarding the critical scaling, we recall from Remark 1.2 of [6] that the condition $\eta + \phi = 1$, which is a natural transition for the rescaling g_n (as is apparent from (A.7)), represents the limit of the parameter range for which uniqueness and mass conservation of the solution of (1.1) are proved: indeed, the condition required by [6] is $\eta + \phi < 1$, along with some hypothesis on the initial data.

Do we expect the complementary condition $\eta + \phi \geq 1$ to have physical meaning? To consider this question, we take positive and fixed ϕ and η , and consider the rescaling (A.2) under the constraints (A.3) and (A.4). Seeking to understand the formation of massive particles, rather than spatial blow-up, we fix $\tau = 0$. We are led to

$$\gamma = \phi^{-1} \tag{A.8}$$

and

$$\alpha = 1 + \frac{1 + \eta}{\phi}. \tag{A.9}$$

Returning to (A.2), a self-similar blow-up profile is consistent with the scaling

$$t^{-\alpha} f_{nt^{-\gamma}}(x, 1)$$

given by $\lambda = t^{-1}$ provided that its mass (A.5) does not grow to infinity as $\lambda \rightarrow 0$. We have set $\tau = 0$: as such, the condition that ensures this is $\alpha - 2\gamma \geq 0$, which, by (A.8) and (A.9), amounts to the inequality $\phi + \eta \geq 1$.

We conclude that considerations of scaling would in principle permit a blow-up in the equations in the mass variable under the condition that $\eta + \phi \geq 1$. The blow-up that we considered is in a low λ limit, which corresponds to heavy mass at late times: as such, it should be considered not as a gelation, in which particles of infinite mass develop in finite time, but rather as the appearance of populations of arbitrarily heavy particles at correspondingly high time-scales. Expressed more precisely, the weak form of blow-up considered is the statement that, for each K strictly less than the total initial mass $\int_0^\infty \int_{\mathbb{R}^d} m f_m(x, 0) dx dm$ and any $m_0 \in \mathbb{R}^+$, there exists

$t \in [0, \infty)$,

$$\int_{m_0}^{\infty} \int_{\mathbb{R}^d} m f_m(x, t) \, dx \, dm > K. \tag{A.10}$$

(This condition is correct in the absence of gelation. Gelation would remove mass from all finite levels. Note also that the absence of fragmentation in (1.1) means that, in fact, (A.10) implies the stronger statement that most of the mass accumulates in arbitrarily high levels at all sufficiently late times.) For dimension $d \geq 3$, (1.11) of Theorem 1.1 in [6] shows that the discrete analogue of (A.10) fails if $\eta + \phi < 1$.

A parallel may be drawn between the Smoluchowski PDE and the non-linear Schrödinger equation. Consider, for example, a solution of cubic defocusing NLS, $u : \mathbb{R}^d \times \mathbb{R}^+ \rightarrow \mathbb{C}$ of

$$i \frac{\partial}{\partial t} u - \Delta u = -|u|^2 u, \tag{A.11}$$

which may be written in Fourier space as

$$i \frac{\partial}{\partial t} \hat{u} - |\xi|^2 \hat{u} = - \int \int \hat{u}(\xi - \eta) \hat{u}(\sigma) \hat{u}(\eta - \sigma) \, d\eta \, d\sigma. \tag{A.12}$$

We see that the mass variable in (1.1) may be viewed as analogous to the frequency variable in (A.12): the non-linear interaction term in each case is a type of convolution. Pursuing the analogy, the quantity $\frac{1}{2} \|\nabla u\|_2^2 + \frac{1}{4} \|u\|_4^4$ is formally conserved in NLS, as is the mass $\int_0^\infty \int_{\mathbb{R}^d} m f_m \, dx \, dm$ for the Smoluchowski PDE. For NLS, the term weak turbulence refers to the growth to infinity in time of the H^s norm

$$\|u\|_{H^s} = \int |\hat{u}(\xi)|^2 |\xi|^{2s} \, d\xi,$$

for some $s > 1$, a circumstance that is anticipated in (A.11) in a periodic domain. (See Section II.2 of [1] for a discussion.) The counterpart of weak turbulence for the system (1.1) is

$$\int_0^\infty \int_{\mathbb{R}^d} m^r f_m(x, t) \, dx \, dm \rightarrow \infty \quad \text{as } t \rightarrow \infty,$$

for some $r > 1$. (Note that (A.10) implies this statement for every $r > 1$ on a subsequence of times.)

Comparing the system (1.1) to its spatially homogeneous counterpart, given by $\{f_n : [0, \infty) \rightarrow [0, \infty) : n \in (0, \infty)\}$ satisfying

$$\frac{d}{dt} f_n(t) = \frac{1}{2} \int_0^n \beta(m, n - m) f_m(t) f_{n-m}(t) \, dm - 2 \int_0^\infty \beta(m, n) f_m(t) f_n(t) \, dm, \tag{A.13}$$

we see the stabilizing role of diffusion: for example, it is easy to see that taking $\beta(n, m)$ identically equal to a constant in (A.13) ensures the analogue of (A.10), while we have seen in the spatial case that scaling arguments do not disallow (A.10) under the condition that $\eta + \phi \geq 1$.

Regarding the prospect of proving mass conservation for at least some part of the parameter space where $\phi + \eta \geq 1$, we comment that, in [6], hypotheses of the form $\beta(n, m) \leq n^\eta + m^\eta$ were used. It may be that, if $\beta(n, m) \leq n^{1+\varepsilon} + m^{1+\varepsilon}$ or $\beta(n, m) \leq n^{1/2+\varepsilon} m^{1/2+\varepsilon}$ (with $\varepsilon > 0$ a small constant), but β is permitted to have space–time dependence subject to such a bound, then gelation is more liable to occur. Hence, an argument for mass conservation would have to exploit the assumption that $\beta(n, m)$ is constant in space–time in a way that those in [6] did not.

References

- [1] J. Bourgain, Global solutions of nonlinear Schrödinger equations, American Mathematical Society Colloquium Publications, 1999.
- [2] S. Großkinsky, C. Klingenberg, K. Oelschläger, A rigorous derivation of Smoluchowski's equation in the moderate limit, *Stoch. Anal. Appl.* 22 (2004) 113–141.
- [3] M.Z. Guo, G.C. Papanicolaou, S.R.S. Varadhan, Nonlinear diffusion limit for a system with nearest neighbor interactions, *Commun. Math. Phys.* 118 (1988) 31–59.
- [4] A. Hammond, F. Rezakhanlou, The kinetic limit of a system of coagulating Brownian particles, *Arch. Ration. Mech. Anal.* 185 (2007) 1–67.
- [5] A. Hammond, F. Rezakhanlou, Kinetic limit for a system of coagulating planar Brownian particles, *J. Stat. Phys.* 124 (2006) 997–1040.
- [6] A. Hammond, F. Rezakhanlou, Moment bounds for the Smoluchowski equation and their consequences, *Commun. Math. Phys.* 276 (2007) 645–670.
- [7] R. Lang, X.-X. Nyugen, Smoluchowski's theory of coagulation in colloids holds rigorously in the Boltzmann–Grad limit, *Z. Wahrsch. Verw. Gebiete* 54 (1980) 227–280.
- [8] P. Laurencot, S. Mischler, The continuous coagulation–fragmentation equations with diffusion, *Arch. Ration. Mech. Anal.* 162 (2002) 45–99.
- [9] J. Norris, Brownian coagulation, *Commun. Math. Sci.* 2 (Suppl. 1) (2004) 93–101.
- [10] F. Rezakhanlou, The coagulating Brownian particles and Smoluchowski's equation, *Markov Process. Related Fields* 12 (2006) 425–445.
- [11] F. Rezakhanlou, Moment bounds for the solutions of the Smoluchowski equation with coagulation and fragmentation. Preprint.