

# A Bernstein Type $L^p$ Inequality for a Certain Class of Polynomials

Robert Gardner\*

*Department of Mathematics, East Tennessee State University, Johnson City.*

metadata, citation and similar papers at [core.ac.uk](http://core.ac.uk)

and

Amy Weems

*Department of Mathematics, University of Colorado, Boulder, Colorado 80309-0395*

*Submitted by Robert A. Gustafson*

Received October 21, 1996

Bernstein's classical theorem states that for a polynomial  $P$  of degree at most  $n$ ,  $\max_{|z|=1} |P'(z)| \leq n \max_{|z|=1} |P(z)|$ . We give related results for polynomials  $P$  satisfying the conditions  $P'(0) = P''(0) = \dots = P^{(m-1)}(0) = 0$  and  $P(z) \neq 0$  for  $|z| < K$ , where  $K \geq 1$ . We give  $L^p$  inequalities valid for  $0 \leq p \leq \infty$ . © 1998 Academic Press

## 1. INTRODUCTION AND HISTORY

Let  $\mathcal{P}_n$  be the linear space of all polynomials over the complex field of degree less than or equal to  $n$ . For  $P \in \mathcal{P}_n$ , define

$$\|P\|_0 = \exp\left(\frac{1}{2\pi} \int_0^{2\pi} \log |P(e^{i\theta})| d\theta\right),$$
$$\|P\|_p = \left(\frac{1}{2\pi} \int_0^{2\pi} |P(e^{i\theta})|^p d\theta\right)^{1/p} \quad \text{for } 0 < p < \infty,$$

\* E-mail: [gardnerr@etsuarts.etsu.edu](mailto:gardnerr@etsuarts.etsu.edu).

and

$$\|P\|_\infty = \max_{|z|=1} |P(z)|.$$

Notice that  $\|P\|_0 = \lim_{p \rightarrow 0^+} \|P\|_p$  and  $\|P\|_\infty = \lim_{p \rightarrow \infty} \|P\|_p$ . For  $1 \leq p \leq \infty$ ,  $\|\cdot\|_p$  is a norm (and therefore  $\mathcal{P}_n$  is a normed linear space under  $\|\cdot\|_p$ ). However, for  $0 \leq p < 1$ ,  $\|\cdot\|_p$  does not satisfy the triangle inequality and is therefore not a norm (this follows from Minkowski's inequality—see [10] for details).

Bernstein's well known result relating the supremum norm of a polynomial and its derivative states that if  $P \in \mathcal{P}_n$  then  $\|P'\|_\infty \leq n\|P\|_\infty$  [2]. This inequality reduces to equality if and only if  $P(z) = \alpha z^n$  for some complex constant  $\alpha$ . Erdős conjectured and Lax proved [6]:

**THEOREM 1.1.** *If  $P \in \mathcal{P}_n$  and  $P(z) \neq 0$  for  $|z| < 1$ , then*

$$\|P'\|_\infty \leq \frac{n}{2} \|P\|_\infty.$$

Malik generalized Theorem 1.1 and proved [7]:

**THEOREM 1.2.** *If  $P \in \mathcal{P}_n$  and  $P(z) \neq 0$  for  $|z| < K$  where  $K \geq 1$ , then*

$$\|P'\|_\infty \leq \frac{n}{1+K} \|P\|_\infty.$$

Of course, Theorem 1.1 follows from Theorem 1.2 when  $K = 1$ . Chan and Malik [3] introduced the class of polynomials of the form  $P(z) = a_0 + \sum_{v=m}^n a_v z^v$ . We denote the linear space of all such polynomials as  $\mathcal{P}_{n,m}$ . Notice that  $\mathcal{P}_{n,1} = \mathcal{P}_n$ . Chan and Malik presented the following result [3]:

**THEOREM 1.3.** *If  $P \in \mathcal{P}_{n,m}$  and  $P(z) \neq 0$  for  $|z| < K$  where  $K \geq 1$ , then*

$$\|P'\|_\infty \leq \frac{n}{1+K^m} \|P\|_\infty.$$

Qazi, independently of Chan and Malik, presented the following result which includes Theorem 1.3 [8]:

**THEOREM 1.4.** *If  $P(z) = a_0 + \sum_{v=m}^n a_v z^v \in \mathcal{P}_{n,m}$  and  $P(z) \neq 0$  for  $|z| < K$  where  $K \geq 1$ , then*

$$\|P'\|_\infty \leq \frac{n}{1+s_0} \|P\|_\infty,$$

where

$$s_0 = K^{m+1} \left( \frac{m|a_m|K^{m-1} + n|a_0|}{n|a_0| + m|a_m|K^{m+1}} \right).$$

Since  $m|a_m|K^m \leq n|a_0|$ , Theorem 1.4 implies Theorem 1.3 (see [8] for details).

Zygmund [11] extended Bernstein's result to  $L^p$  norms. DeBruijn [4] extended Theorem 1.1 to  $L^p$  norms by showing:

**THEOREM 1.5.** *If  $P \in \mathcal{P}_n$  and  $P(z) \neq 0$  for  $|z| < 1$ , then for  $1 \leq p \leq \infty$*

$$\|P'\|_p \leq \frac{n}{\|1+z\|_p} \|P\|_p.$$

Of course, Theorem 1.5 reduces to Theorem 1.1 with  $p = \infty$ . Rahman and Schmeisser [9] proved that Theorem 1.5 in fact holds for  $0 \leq p \leq \infty$ . The purpose of this paper is to show that Theorems 1.3 and 1.4 can be extended to  $L^p$  inequalities where  $0 \leq p \leq \infty$ .

## 2. STATEMENT OF RESULTS

Our main result is:

**THEOREM 2.1.** *If  $P(z) = a_0 + \sum_{v=m}^n a_v z^v \in \mathcal{P}_{n,m}$  and  $P(z) \neq 0$  for  $|z| < K$  where  $K \geq 1$ , then for  $0 \leq p \leq \infty$*

$$\|P'\|_p \leq \frac{n}{\|s_0+z\|_p} \|P\|_p,$$

where  $s_0$  is as given in Theorem 1.4.

With  $p = \infty$ , Theorem 2.1 reduces to Theorem 1.4. As mentioned in Section 1, we can deduce:

**COROLLARY 2.2.** *If  $P \in \mathcal{P}_{n,m}$  and  $P(z) \neq 0$  for  $|z| < K$  where  $K \geq 1$ , then for  $0 \leq p \leq \infty$*

$$\|P'\|_p \leq \frac{n}{\|K^m+z\|_p} \|P\|_p.$$

With  $p = \infty$ , Corollary 2.2 reduces to Theorem 1.3.

Of special interest, is the fact that Theorem 2.1 and Corollary 2.2 hold for  $L^p$  norms for all  $1 \leq p \leq \infty$ . In particular, we have:

**COROLLARY 2.3.** *If  $P \in \mathcal{P}_{n,m}$  and  $P(z) \neq 0$  for  $|z| < K$  where  $K \geq 1$ , then for  $1 \leq p \leq \infty$*

$$\|P'\|_p \leq \frac{n}{\|K^m + z\|_p} \|P\|_p.$$

With  $m = 1$ , Corollary 2.3 yields an  $L^p$  version of Theorem 1.2. With  $p = \infty$ , Corollary 2.3 reduces to Theorem 1.3. With  $m = 1$  and  $p = \infty$ , Corollary 2.3 reduces to Theorem 1.2. Finally, with  $m = 1$ ,  $p = \infty$ , and  $K = 1$ , Corollary 2.3 reduces to Theorem 1.1.

### 3. LEMMAS

We need the following lemmas for the proof of our theorem.

**LEMMA 3.1.** *If the polynomial  $P(z)$  of degree  $n$  has no roots in the circular domain  $C$  and if  $\zeta \in C$  then  $(\zeta - z)P'(z) + nP(z) \neq 0$  for  $z \in C$ .*

Lemma 3.1 is due to Laguerre [5].

**DEFINITION 3.2.** For  $\gamma = (\gamma_0, \dots, \gamma_n) \in \mathbf{C}^{n+1}$  and  $P(z) = \sum_{v=0}^n c_v z^v$ , define

$$\Lambda_\gamma P(z) = \sum_{v=0}^n \gamma_v c_v z^v.$$

The operator  $\Lambda_\gamma$  is said to be *admissible* if it preserves one of the following properties:

- (a)  $P(z)$  has all its zeros in  $\{z \in \mathbf{C} : |z| \leq 1\}$ ,
- (b)  $P(z)$  has all its zeros in  $\{z \in \mathbf{C} : |z| \geq 1\}$ .

The proof of Lemma 3.3 was given by Arestov [1]:

**LEMMA 3.3.** *Let  $\phi(x) = \psi(\log x)$  where  $\psi$  is a convex non-decreasing function on  $\mathbf{R}$ . Then for all  $P(z) \in \mathcal{P}_n$  and each admissible operator  $\Lambda_\gamma$*

$$\int_0^{2\pi} \phi(|\Lambda_\gamma P(e^{i\theta})|) d\theta \leq \int_0^{2\pi} \phi(c(\gamma, n)|P(e^{i\theta})|) d\theta,$$

where  $c(\gamma, n) = \max(|\gamma_0|, |\gamma_n|)$ .

Qazi proved [8]:

LEMMA 3.4. *If  $P(z) = c_0 + \sum_{v=m}^n c_v z^v$  has no zeros in  $|z| < K$ ,  $K \geq 1$  then for  $|z| = 1$*

$$K^m |P'(z)| \leq s_0 |P'(z)| \leq |Q'(z)|,$$

where  $Q(z) = z^n \overline{P(1/\bar{z})}$  and  $s_0$  is as defined in Theorem 1.4.

#### 4. PROOF OF THEOREM 2.1

By Lemma 3.1 we have  $nP(z) - (z - \zeta)P'(z) \neq 0$  for  $|z| \leq 1$ ,  $|\zeta| \leq 1$ . Therefore, setting  $\zeta = -ze^{-i\alpha}$ ,  $\alpha \in \mathbf{R}$ , the operator  $\Lambda$  defined by

$$\Lambda P(z) = (e^{i\alpha} + 1)zP'(z) - ne^{i\alpha}p(z)$$

is admissible and so by Lemma 3.3 with  $\psi(x) = e^{px}$ ,

$$\int_0^{2\pi} \left| (e^{i\alpha} + 1) \frac{dP(e^{i\theta})}{d\theta} - ine^{i\alpha}P(e^{i\alpha}) \right|^p d\theta \leq n^p \int_0^{2\pi} |P(e^{i\theta})|^p d\theta$$

for  $p > 0$ . Then

$$\int_0^{2\pi} \left| \frac{dP(e^{i\theta})}{d\theta} + e^{i\alpha} \left\{ \frac{dP(e^{i\theta})}{d\theta} - inP(e^{i\theta}) \right\} \right|^p d\theta \leq n^p \int_0^{2\pi} |P(e^{i\theta})|^p d\theta.$$

This gives

$$\begin{aligned} & \int_0^{2\pi} \int_0^{2\pi} \left| \frac{dP(e^{i\theta})}{d\theta} + e^{i\alpha} \left\{ \frac{dP(e^{i\theta})}{d\theta} - inP(e^{i\theta}) \right\} \right|^p d\theta d\alpha \\ & \leq 2\pi n^p \int_0^{2\pi} |p(e^{i\theta})|^p d\theta. \end{aligned} \tag{4.1}$$

Now

$$\begin{aligned}
 & \int_0^{2\pi} \int_0^{2\pi} \left| \frac{dP(e^{i\theta})}{d\theta} + e^{i\alpha} \left\{ \frac{dP(e^{i\theta})}{d\theta} - inP(e^{i\theta}) \right\} \right|^p d\theta d\alpha \\
 &= \int_0^{2\pi} \left| \frac{dP(e^{i\theta})}{d\theta} \right|^p \int_0^{2\pi} \left| 1 + e^{i\alpha} \left\{ \frac{dP(e^{i\theta})/d\theta - inP(e^{i\theta})}{dP(e^{i\theta})/d\theta} \right\} \right|^p d\alpha d\theta \\
 &= \int_0^{2\pi} \left| \frac{dP(e^{i\theta})}{d\theta} \right|^p \int_0^{2\pi} \left| e^{i\alpha} + \left| \frac{dP(e^{i\theta})/d\theta - inP(e^{i\theta})}{dP(e^{i\theta})/d\theta} \right| \right|^p d\alpha d\theta \\
 &= \int_0^{2\pi} \left| \frac{dP(e^{i\theta})}{d\theta} \right|^p \int_0^{2\pi} \left| e^{i\alpha} + \left| \frac{Q'(e^{i\theta})}{P'(e^{i\theta})} \right| \right|^p d\alpha d\theta \\
 &\geq \int_0^{2\pi} \left| \frac{dP(e^{i\theta})}{d\theta} \right|^p \int_0^{2\pi} |e^{i\alpha} + s_0|^p d\alpha d\theta \quad \text{by Lemma 3.4} \quad (4.2)
 \end{aligned}$$

by the fact that  $|e^{i\alpha} + r|$  is an increasing function of  $r$  for  $r \geq 1$ . Thus combining (4.1) and (4.2) we see that

$$\left( \int_0^{2\pi} \left| \frac{dP(e^{i\alpha})}{d\theta} \right|^p d\theta \right) \left( \int_0^{2\pi} |e^{i\alpha} + s_0|^p d\alpha \right) \leq 2\pi n^p \int_0^{2\pi} |P(e^{i\theta})|^p d\theta$$

from which the theorem follows for  $0 < p < \infty$ . The result holds for  $p = 0$  and  $p = \infty$  by letting  $p \rightarrow 0^+$  and  $p \rightarrow \infty$ , respectively.

## REFERENCES

1. V. Arestov, On integral inequalities for trigonometric polynomials and their derivatives, *Math. USSR-Izv.* **18** (1982), 1–17.
2. S. Bernstein, "Leçons sur les propriétés extrémales et la meilleure approximation des fonctions analytiques d'une variable réelle," Collection Borel, Paris, 1926.
3. T. Chan and M. Malik, On Erdős–Lax theorem, *Proc. Indian Acad. Sci.* **92** (1983), 191–193.
4. N. DeBruijn, Inequalities concerning polynomials in the complex domain, *Nederl. Akad. Wetensch. Proc.* **50**, (1947), 1265–1272.
5. E. Laguerre, "Oeuvres," Vol. 1; *Nouvelles Ann. Math.* **17**, No. 2 (1878).
6. P. Lax, Proof of a conjecture of P. Erdős on the derivative of a polynomial, *Bull. Amer. Math. Soc.* **50** (1944), 509–513.

7. M. Malik, On the derivative of a polynomial, *J. London Math. Soc.* **1**, No. 2 (1969), 57–60.
8. M. Qazi, On the maximum modulus of polynomials, *Proc. Amer. Math. Soc.* **115** (1992), 337–343.
9. Q. Rahman and G. Schmeisser,  $L^p$  inequalities for polynomials, *J. Approx. Theory* **53** (1988), 26–32.
10. H. Royden, “Real Analysis,” 3rd ed., Macmillan Co., New York, 1988.
11. A. Zygmund, “Trigonometric Series,” 2nd rev. ed., Vols. I, II, Cambridge Univ. Press, Cambridge, UK, 1959.