

Enhanced Goblet Cell Hyperplasia in HDC Knockout Mice with Allergic Airway Inflammation

Kohei Yamauchi¹, Hong Mei Piao^{1,5}, Toshihide Nakadate¹, Toshiki Shikanai¹, Yutaka Nakamura¹, Harumasa Ito¹, Takashi Mouri¹, Hitoshi Kobayashi¹, Chihaya Maesawa³, Takashi Sawai², Hiroshi Ohtsu⁴ and Hiroshi Inoue¹

ABSTRACT

Background: Histamine is known to have immunoregulatory roles in allergic reactions through histamine receptor 1 (H1R), H2R, H3R and H4R. However, its role in goblet cell hyperplasia in the airways of asthma patients is yet to be clarified.

Objective: This study was designed to examine the role of histamine in goblet cell hyperplasia using histamine-deficient mice (*Hdc*^{-/-} mice) with allergic airway inflammation.

Methods: Wild-type and *Hdc*^{-/-} C57BL/6 mice were sensitized with ovalbumin (OVA). After a 2-week exposure to OVA, goblet cell hyperplasia was evaluated. Cell differentials and cytokines in BALF were analyzed. The mRNA levels of MUC5AC and Gob-5 gene were determined quantitatively.

Results: The number of eosinophils in BALF increased in both the sensitized wild-type mice and *Hdc*^{-/-} mice with OVA inhalation. In addition, the numbers of alveolar macrophages and lymphocytes in BALF increased significantly in the sensitized *Hdc*^{-/-} mice with OVA inhalation compared to the wild-type mice under the same conditions. The concentrations of Interleukin-4 (IL-4), IL-5, IL-13, Interferon- γ (IFN- γ), tumor necrosis factor- α (TNF- α) and IL-2 in the BALF all increased significantly in both groups compared to those exposed to saline. In particular, the concentration of TNF- α in the *Hdc*^{-/-} mice exposed to OVA was significantly higher than that in the wild-type mice under the same conditions. The mRNA levels of Gob-5 and MUC5AC, and the ratio of the goblet cells in the airway epithelium significantly increased in *Hdc*^{-/-} mice exposed to OVA compared to wild-type mice.

Conclusions: These results suggested that histamine may play a regulatory role in goblet cell hyperplasia in allergic airway inflammation.

KEY WORDS

airway remodeling, asthma, Gob-5, MUC5AC, TNF- α

INTRODUCTION

Goblet cell hyperplasia and mucus overproduction are important features of bronchial asthma.^{1,2} Although goblet cell hyperplasia with mucus hypersecretion has been reported to be associated with the development of airway hyperresponsiveness and the increase of severity and mortality in bronchial asthma¹⁻⁴, the mechanisms responsible for goblet cell

hyperplasia are not completely understood.

Histamine is known to be a strong chemical mediator that stimulates secretion from goblet cells. Bryce *et al.* recently reported that the airway inflammatory response and goblet cell hyperplasia to allergic stimuli diminished in H1 receptor (H1R) knock-out mice, suggesting that histamine may be involved in the immunomodulation in airway allergic reaction including Th1 and Th2 cytokine production and goblet cell hy-

¹Third Department of Internal Medicine, ²First Department of Pathology, ³Second Department of Pathology, Iwate Medical University School of Medicine, Iwate, ⁴Tohoku University Graduate School of Engineering, Miyagi, Japan and ⁵Department of Respiratory Medicine, Affiliated Hospital of Yanbian University College of Medicine, Yanji, China.

Correspondence: Kohei Yamauchi, M.D., Associate Professor,

Department of Internal Medicine, Division of Pulmonary Medicine, Allergy and Rheumatology, Iwate Medical University School of Medicine, 19-1 Uchimarui, Morioka, Iwate 020-8505, Japan.

Email: kyamauch@iwate-med.ac.jp

Received 25 February 2008. Accepted for publication 1 September 2008.

©2009 Japanese Society of Allergology

perplasia.⁵

It has been reported that histamine can modulate T-cell-mediated immune responses.⁶ Mast cells and basophils, which have been considered mainly to be effector cells for IgE-mediated reactions, can modulate the immune response through secreted histamine. Jutel *et al.* demonstrated, using H1R and histamine H2 receptor (H2R) knock-out mice, that histamine augmented TH1 response T lymphocytes through H1R and suppressed both TH1 and TH2 responses through H2R.⁷ In addition, histamine H4 receptor (H4R) has recently been identified and an H4R inhibitor has been demonstrated to have a suppressive effect on allergic airway inflammation, including eosinophil and mast cell infiltration in murine asthma models.⁸⁻¹¹ In this context, histamine may have various roles in allergic airway inflammation through H1R, H2R and H4R in immune cells including T lymphocytes and dendritic cells.

To date, goblet cell hyperplasia has been thought to be induced after epithelial damage by numerous stimuli, such as endogenous oxidants induced by neutrophil elastase,¹² or exogenous oxidants produced by cigarette smoke¹³ or diesel engine emissions.¹⁴ On the other hand, it has been demonstrated that the TH2 lymphocyte-derived cytokines interleukin-4 (IL-4), IL-5, IL-9, and IL-13 induce goblet cell hyperplasia in animal models and *in vitro* studies.¹⁵⁻¹⁹

According to previous reports, several genes are up-regulated in association with goblet cell hyperplasia. Among them, MUC genes encode various mucin glycoproteins which are primary secretory proteins consisting of a family of genes (ie, MUC-1, MUC-2 and MUC4).^{20,21} MUC5AC is the predominant MUC gene expressed in the airway.²¹ It has been demonstrated that the level of MUC5AC gene expression increases in both animal models of asthma and tissue samples from patients with asthma.²²⁻²⁴

In addition, the overexpression of the Gob-5 gene has been reported in airway epithelia with exacerbated AHR, goblet cell hyperplasia, mucus overproduction, and eosinophil infiltration in a murine asthma model.²⁵ Gob-5 gene in the mouse corresponds to CLCA1 gene in humans, which regulates calcium activated chloride conductance. It has been shown to induce mucin gene expression (MUC5AC) in the human mucoepidermoid cell line NCI-H292 and its expression was demonstrated to increase in patients with asthma.²⁵⁻²⁷

Histamine-deficient mice are now available by disrupting *L*-histidine decarboxylase gene (HDC knockout mice: *Hdc*^{-/-}).²⁸ Histamine exerts its various biological actions via H1, H2, H3 and H4. Due to the overlapping, and sometimes antagonistic function of the receptors in the presence of endogenous histamine, receptor blocking alone cannot achieve complete elimination of the histamine system. It is, there-

fore, reasonable to evaluate the total role of histamine with *Hdc*^{-/-} mice.

An analysis of cytokine production and cell differentials in BALF in sensitized *Hdc*^{-/-} mice with allergic airway inflammation has been previously reported, however, these investigated only the acute airway responses in murine asthma models with a short exposure to allergens.^{29,30} There have been few data about airway remodeling that included goblet cell hyperplasia in histamine-deficient mice.

In this study, we designed an experimental asthma model with a relatively long duration of allergen exposure to *Hdc*^{-/-} mice to elucidate the roles of histamine in airway remodeling, focusing especially on goblet cell hyperplasia.

METHODS

ANIMALS

HDC knockout (*Hdc*^{-/-}) mice were generated by Ohtsu *et al.*²⁸ In the present study, we used 6 to 8-week-old female *Hdc*^{-/-} mice backcrossed for six generations with C57BL/6 mice, and used age/gender-matched C57BL/6N mice (Japan SLC, Shizuoka, Japan) as controls. The mice used for this study were generated by breeding homozygous *Hdc*^{-/-} mice in the facilities of Tohoku University (Sendai, Japan). Both *HDC*^{-/-} and wild-type mice were kept on a normal diet containing <0.3 mg of histamine/g of food. All experiments described in this study were performed according to the guidelines for the care and use of experimental animals as determined by the Japanese Association for Laboratory Animals Science in 1987.

IMMUNIZATION AND AEROSOLIZATION PROTOCOL

The mice were sensitized according to the methods described in a previous study.³¹ In brief, mice were sensitized on days 0 and 5 of the protocol by an intraperitoneal injection of 0.5 ml aluminum hydroxide-precipitated antigen containing 8 µg OVA (Sigma Chemical Co., St. Louis, MO) adsorbed overnight at 4°C to 4 mg of aluminium hydroxide (Wako Chemical Co., Tokyo, Japan) in phosphate-buffered saline (PBS). Twelve days after the second immunization, the wild-type mice and *Hdc*^{-/-} mice were divided into 3 groups respectively, each consisting of 6 animals. Each of the wild-type mice groups and one of the *Hdc*^{-/-} mice groups were killed for analysis as controls without inhalation. To perform chronic exposure of OVA, 2 groups of each of the wild mice and *Hdc*^{-/-} mice were placed in a plastic chamber (10 cm × 15 cm × 25 cm) and exposed to aerosolized OVA (5 mg/ml in 0.9% saline) for 1 hour every other day for 14 days. The other 2 groups of the wild-type mice and *Hdc*^{-/-} mice were exposed only to 0.9% saline every other day for 14 days. The aerosolized OVA was produced by a Pulmo-Aide Compressor/Nebulizer

(Devilbiss) (Sunrise Medical HHG, Inc., Somerset, PA, USA) at a flow rate of 5–7 liter/minute.

COLLECTION AND MEASUREMENT OF SPECIMENS

After being exposed to aerosolized saline or OVA every other day for 2 weeks, each group of mice was killed on the 14th day, 24 hours after final inhalation, and bronchoalveolar lavage fluid (BALF) and lung tissues were collected. To collect BALF, the lungs were dissected and the trachea was cannulated with a polyethylene tube (Becton Dickinson, Sparks, MD, USA). The lungs were lavaged twice with 0.5 ml PBS, and up to 0.8 ml of the instilled fluid was consistently recovered. The recovered fluid was centrifuged ($300 \times g$ for 6 minutes) and the cells were resuspended in 0.5 ml PBS. The total number of cells was counted using an improved Neubauer hemocytometer chamber. An air-dried slide preparation was made from each sample containing 10,000 cells by cytopspin (Cytocentrifuge, Sakura Seiki, Tokyo, Japan) and stained with May-Grunwald-Giemsa stain. Differential counts of at least 500 cells were made according to standard morphologic criteria. The numbers of cells recovered per mouse were then expressed as the mean and standard error of the mean (SEM) for each treated group.

After centrifugation, supernatants were stored at -80°C for the measurement of cytokines. After harvesting BALF, lungs were fixed with formaldehyde and were embedded in paraffin. These 3- μm -thick sections were stained with hematoxylin eosin (HE) and periodic acid-Schiff (PAS). In addition, other parts of the left lungs were stored at -80°C for RNA extraction.

MEASUREMENT OF SERUM CONCENTRATION OF IgE

Serum IgE levels were determined using the commercially available ELISA kit (Yamasa, Chiba, Japan).

HISTOLOGICAL MEASUREMENT OF GOBLET CELL NUMBER

The 3 mm paraffin sections of the lungs described above were stained with PAS for evaluation of goblet cells. Goblet cell hyperplasia was determined by counting the number of PAS positive cells in more than 5 large (diameter $>200 \mu\text{m}$) and more than 10 small bronchi (diameter $<150 \mu\text{m}$) per lung under microscopy. The results were expressed as percentages of PAS positive cells per total epithelial cells. In addition, lung sections were stained with HE.

ISOLATION OF TOTAL RNA AND REAL-TIME QUANTITATIVE PCR

Total RNA from the whole lobes of the left lungs was obtained using ISOGEN (Wako Pure Chemicals, Osaka, Japan), quantified by spectrophotometry. The quality of the obtained RNA was confirmed by aga-

rose gel electrophoresis.

To quantify the mRNA of Gob-5, MUC5AC, and glyceraldehyde-3-phosphatase dehydrogenase (GAPDH) expression in the murine lungs, quantitative PCR was carried out using an ABI Prism 7700 Sequence Detector (Perkin-Elmer Applied Biosystems, Foster City, CA, USA) as previously described.³²

Oligonucleotide PCR primer pairs and fluorogenic probes for murine MUC5AC and Gob-5 were designed from the published sequences using Primer Express software (Perkin-Elmer) (sense primer: 5'-ACTGTTACTATGCGATGTGTAGCCA-3', antisense primer: 5'-GAGGAAACACATTGCACCGA-3'; Taqman probe: 5'-[FAM] ACTGCCACCTGTCACTGGGCGG [TAMRA]-3') for MUC5AC and (sense primer: 5'-AGGGCATCGTCATCGCC-3'; antisense primer: 5'-TCCTTTATGTGTTGAATGAGGGC-3'; Taqman probe: 5'-[FAM] AGACCACGACGTGCCGGAAGATG [TAMRA]-3') for Gob-5.³³ Primers and the labeled probe (VIC) for rodent GAPDH were purchased from Perkin-Elmer Applied Biosystems. 100 ng of RNA dissolved in 10 μl of water from each aliquot of murine lung tissue was denatured at 90°C for 90 seconds. Each RNA sample (100 ng/10 μl of water) was mixed in 40 μl of buffer containing the following reagents for the one-step RT-PCR reaction: 50 mM KCl, 10 mM Tris-HCl (pH 8.3), 0.01 mM EDTA, 60 nM Passive Reference 1 (Applied Biosystems), 5 mM MgCl_2 , 100 nM sense primer, 100 nM antisense primer, 0.3 mM deoxynucleoside triphosphate (Boehringer), 0.4 U/ μl RNase inhibitor (Promega), 0.4 U/ μl Moloney murine leukemia virus RT (Perkin Elmer), 0.0025 U/ μl Taq Gold Polymerase (Perkin Elmer), and 100 nM Taqman probe, as described above. The fragment of mRNA for Gob-5, MUC5AC and GAPDH was reversely transcribed into cDNA (30 minutes at 48°C) and amplified by PCR for 40 cycles (15 seconds at 95°C and 1 minute at 60°C). Whole reactions of the RT-PCR and detection of the fluorescence emission signal for every PCR cycle were performed at the same time in a single tube in a sequence detector (ABI 7700). The minimum PCR cycle to detect the fluorescent signal was defined as the cycle threshold (C), which is predictive of the quantity of an input target fragment.³⁴ The standard curve was obtained between the fluorescence emission signals and C by means of duplicated serial dilutions of each quantified cDNA fragment which contained the targeted sequences. The expression of MUC5AC and Gob-5 mRNA was normalized to the constitutive expression of GAPDH mRNA.

CYTOKINE ASSAY

Concentrations of IL-4, IL-5, Interferon- γ (IFN- γ), tumor necrosis factor- α (TNF- α) and IL-2 in BALF were determined with cytokine Bead Array inflammatory kits using flow cytometry according to the manufacturer's instructions (BD PharMingen, San Diego, CA,

Table 1 Total cell number and cell differential in BALF of the wild-type and *Hdc*^{-/-} mice after saline and OVA inhalation

	C57BL/6: wild saline, 14 th day	C57BL/6: wild OVA, 14 th day	C57BL/6: <i>Hdc</i> ^{-/-} saline, 14 th day	C57BL/6: <i>Hdc</i> ^{-/-} OVA, 14 th day
Total cells (×10 ⁴ /ml)	2.29 ± 0.64 (n = 6)	28.2 ± 5.1 (n = 6)	3.60 ± 0.21 (n = 6)	62.1 ± 7.0* (n = 6)
Alveolar macrophages	2.16 ± 0.06 (94.2 ± 2.5%)	2.09 ± 0.56 (7.40 ± 2.0%)	3.43 ± 0.11 (95.2 ± 3.0%)	23.5 ± 3.73* (37.9 ± 6.0%)
Lymphocytes	0.13 ± 0.06 (5.75 ± 2.5%)	0.45 ± 0.25 (1.6 ± 0.9%)	0.15 ± 0.09 (4.20 ± 2.6%)	10.8 ± 4.04* (17.4 ± 6.5%)
Eosinophils	< 0.001 (< 0.1%)	25.7 ± 0.50 (91.0 ± 1.8%)	0.22 ± 0.01 (0.6 ± 0.4%)	27.8 ± 6.40 (44.8 ± 10.3%)
Neutrophils	< 0.01 (< 0.1%)	< 0.01 (< 0.1%)	< 0.01 (< 0.1%)	< 0.01 (< 0.1%)
Epithelial cells	< 0.01 (< 0.1%)	< 0.01 (< 0.1%)	< 0.01 (< 0.1%)	< 0.01 (< 0.1%)

The absolute number of cells in BALF on 14th day after saline and OVA inhalation in mice sensitized with OVA was calculated. The numbers in parentheses indicated % in the total cells in BALF. BALF was collected as described in "Methods". Values were expressed as the means ± SEM of 6 mice. * *p* < 0.01 relative to the respective value of wild-type mice with OVA inhalation.

USA). The concentration of IL-13 in BALF was measured using ELISA (RayBio, Norcross, GA, USA).

STATISTICAL ANALYSIS

Data were expressed as mean ± SEM. Multiple comparisons of mean data among the groups were analyzed by the Mann-Whitney U test. Probability values of less than 0.05 were considered to indicate a statistically significant difference.

RESULTS

CELL DIFFERENTIALS IN BALF OF *Hdc*^{-/-} MICE

Repetitive exposure to OVA for 2 weeks induced a marked increase of the total cell numbers in BALF in both wild-type and *Hdc*^{-/-} mice sensitized with OVA, compared with those from the wild-type mice group and mice sensitized with OVA and exposed to saline (Table 1). The total cell numbers in BALF from the *Hdc*^{-/-} mice sensitized with OVA and then exposed to OVA increased significantly more than in wild-type mice under the same conditions. The absolute numbers of alveolar macrophages, lymphocytes and eosinophils in *Hdc*^{-/-} mice sensitized with and exposed to OVA increased significantly compared to mice sensitized with OVA and exposed to saline. In particular, the absolute numbers of alveolar macrophages and lymphocytes in *Hdc*^{-/-} mice sensitized with and exposed to OVA were significantly higher than those of wild-type mice under the same condition. Eosinophils were the predominant cells in BALF from the wild-type mice exposed to OVA and the ratio of eosinophils in BALF was higher compared with that of *Hdc*^{-/-} mice. However, there was no significant difference in the absolute cell numbers of eosinophils in these 2 groups.

CYTOKINES IN BALF OF *Hdc*^{-/-} MICE

TNF- α , IL-4, IL-5, IL-13, INF- γ and IL-2 in BALF were

measured. The concentrations of TNF- α in BALF of both wild (15.08 ± 0.58 pg/ml: mean ± SEM) and *Hdc*^{-/-} mice (25.30 ± 4.23 pg/ml) exposed to OVA increased markedly compared to those of wild-type (2.28 ± 0.40 pg/ml) and *Hdc*^{-/-} mice (2.27 ± 0.40 pg/ml) exposed to saline (Fig. 1A). The concentration of TNF- α in the *Hdc*^{-/-} mice exposed to OVA was significantly higher than that of the wild-type mice under the same conditions.

The concentrations of IL-4 in BALF from both wild-type and *Hdc*^{-/-} mice sensitized with and exposed to OVA increased significantly compared to wild and *Hdc*^{-/-} mice sensitized with OVA and exposed to saline. The concentration of IL-4 in the *Hdc*^{-/-} mice exposed to OVA was significantly lower than that of wild mice under the same conditions (Fig. 1B).

The concentrations of IL-2, IL-5, IL-13 and INF- γ in BALF from both wild-type and *Hdc*^{-/-} mice sensitized with and exposed to OVA increased significantly compared with that of wild-type and *Hdc*^{-/-} mice sensitized with OVA and exposed to saline. No statistically significant difference in these cytokines between the wild-type and *Hdc*^{-/-} mice exposed to OVA (Fig. 1C, D, E, F).

SERUM IgE OF *Hdc*^{-/-} MICE

The concentrations of serum IgE in both wild-type (2133.6 ± 370.3 ng/ml) and *Hdc*^{-/-} mice (998.9 ± 254.0 ng/ml) exposed to OVA were markedly higher than those exposed to saline (wild-type exposed to saline: 416.1 ± 25.1 ng/ml, *Hdc*^{-/-} mice exposed to saline: 371.7 ± 17.2 ng/ml) and without exposure (wild-type: 390.1 ± 32.5; *Hdc*^{-/-}: 328.1 ± 14.8) (Fig. 2). The concentration of serum IgE in the *Hdc*^{-/-} mice exposed to OVA was significantly lower than that of the wild-type mice under the same conditions.




Fig. 1 A, B, C, D, E, F: Outlined column: wild-type mice; Solid column: *Hdc*^{-/-} mice; Column a and c: saline inhalation; Column b and d: OVA inhalation, **A**: TNF- α , **B**: IL-4, **C**: IL-5, **D**: IL-13, **E**: INF- γ , **F**: IL-2 concentration in BALF. * $p < 0.01$, ** $p < 0.05$.

Gob-5 GENE EXPRESSION IN THE LUNG OF *Hdc*^{-/-} MICE

Levels of Gob-5 mRNA in lung tissue of both wild-type and *Hdc*^{-/-} mice sensitized with and exposed to OVA increased significantly compared with those of wild-type and *Hdc*^{-/-} mice sensitized with OVA and exposed to saline. The level of Gob-5 mRNA in the *Hdc*^{-/-} mice exposed to OVA was significantly higher than that of wild-type mice under the same conditions (Fig. 3).

MUC5AC GENE EXPRESSION IN THE LUNG OF *Hdc*^{-/-} MICE

Levels of MUC5AC mRNA in lung tissue of both wild-type and *Hdc*^{-/-} mice sensitized with OVA and exposed to OVA increased markedly compared with those of wild and *Hdc*^{-/-} mice sensitized with OVA exposed to saline. The level of MUC5AC mRNA in the *Hdc*^{-/-} mice exposed to OVA was significantly higher than that of wild-type mice under the same conditions (Fig. 4).

GOBLET CELLS OF LARGE AND SMALL AIRWAYS

PAS staining demonstrated an increase in the number of goblet cells in the epithelium in large and small airways of both wild-type and *Hdc*^{-/-} mice sensitized with OVA (Fig. 5A a-f, B a-f). Goblet cell hyperplasia was more prominent in the *Hdc*^{-/-} mice exposed to OVA than in the wild-type mice under the same conditions (Fig. 5A c, A f, B c, B f). Repetitive saline inhalation induced a mild increase in the number of goblet cells in large and small airways of both wild-type and *Hdc*^{-/-} mice (Fig. 5A b, A e, B b, B e).

QUANTIFICATION OF GOBLET CELLS IN LARGE AND SMALL AIRWAYS

The ratios of goblet cells in the epithelium in large and small airways, of both wild-type and *Hdc*^{-/-} mice sensitized with and exposed to OVA, increased markedly compared with those of the wild-type and *Hdc*^{-/-} mice sensitized with OVA and exposed to saline (Fig. 6A, B). The ratios of goblet cells in the epithelium in large and small airways of the *Hdc*^{-/-} mice exposed to OVA were significantly higher than that of wild-type mice under the same condition.




Fig. 2 IgE concentration in serum. Outlined column: wild-type mice; Solid column: *Hdc*^{-/-} mice; Column A and D: before inhalation; Column B and E: saline inhalation; Column C and F: OVA inhalation. * $p < 0.01$, ** $p < 0.05$, ns: not significant.




Fig. 3 Gob-5 gene expression in the lung. Outlined column: wild-type mice; Solid column: *Hdc*^{-/-} mice; Column A and C: saline inhalation; Column B and D: OVA inhalation. * $p < 0.01$.

DISCUSSION

In this study, we demonstrated that goblet cell hyperplasia was enhanced in *Hdc*^{-/-} mice sensitized with OVA after repetitive OVA exposure for 14 days. We




Fig. 4 MUC5AC gene expression in the lung. Outlined column: wild-type mice; Solid column: *Hdc*^{-/-} mice; Column A and C: saline inhalation; Column B and D: OVA inhalation. * $p < 0.01$.

also demonstrated a striking difference in differential cytological features in BALF between 2 murine asthma models with *Hdc*^{-/-} mice, representing the acute allergic airway response with a short period of OVA exposure previously reported^{29,30} and a relatively chronic allergic airway response shown in the present study. We found significant increases of alveolar macrophages and lymphocytes in BALF of *Hdc*^{-/-} mice with repetitive exposure to OVA for 2 weeks compared with the wild-type mice under the same conditions. The concentration of TNF- α in BALF in the *Hdc*^{-/-} mice after repetitive OVA exposure was significantly higher than that of the wild-type mice under the same conditions.

Histamine has been recognized as a chemical mediator playing a central role in allergic reactions including mucosal edema, mucous gland secretion, smooth muscle contraction, etc. Recent studies have demonstrated that histamine plays a critical role in immunomodulation by acting on different types of histamine receptors such as H1R, H2R and H4R on the surface of immune cells and inflammatory cells.⁶⁻¹¹ Using mutant mice lacking H1R and H2R, Jutel *et al.* demonstrated that histamine enhances TH1-type responses by triggering H1R, whereas both TH1- and TH2-type responses were negatively regulated by H2R. Indeed, the deletion of H1R resulted in suppression of IFN- γ , IL-4 and IL-13, and the deletion of H2R showed an up-regulation of both TH1 and TH2 cytokines (IL-4 and IL-13).⁷ Bryce *et al.* recently demonstrated the interesting opposite effects of H1R inhibition on cultured T cells *in vitro* and the airway response with allergic inflammation *in vivo* using a murine asthma model.⁵ In their report, allergen-stimulated splenic T cells from sensitized H1R knock-




Fig. 5 **A:** Goblet cells in the large airways, **B:** Goblet cells in the small airways. The goblet cells in the epithelium were stained with PAS. **a, b** and **c:** wild-type mice; **d, e** and **f:** *Hdc*^{-/-} mice; **a** and **d:** before inhalation; **b** and **e:** saline inhalation; **c** and **f:** OVA inhalation. black bars indicate 50µm.

out mice exhibited enhanced TH2 cytokine production. However, in contrast, allergen-challenged H1R knockout mice exhibited diminished lung TH2 cytokine mRNA levels, airway inflammation, goblet cell metaplasia, and airway hyperresponsiveness (AHR).

The present study demonstrated that the concentration of IL-4 in BALF was significantly lower in the *Hdc*^{-/-} mice than in the wild-type mice after exposure to OVA, which was consistent with the experiment of the *Hdc*^{-/-} mice by Kozma *et al.*,³⁰ despite differences in the duration of OVA exposure. The concentrations of IL-2, IL-5, IL-13 and INF- γ showed no significant differences between the *Hdc*^{-/-} mice and the wild-

type mice. The lower level of IL-4 in the *Hdc*^{-/-} mice was consistent with the lower level of serum IgE in the *Hdc*^{-/-} mice after the exposure to OVA. These changes may be explained at least in part by the suppressed recruitment of activated TH2 cells into the lung due to the blockade of H1R in T lymphocytes, as suggested by Bryce *et al.*⁵

H1R, H2R and H4R are expressed in monocytes, macrophages and dendritic cells, and histamine plays a regulatory role in expression of cytokines and adhesion molecules in these cells.³⁵ Mazzoni *et al.* demonstrated that histamine inhibited INF- α and released TNF- α from activated plasmacytoid dendritic cells




Fig. 6 **A:** Goblet cell ratio in the large airways, **B:** Goblet cell ratio in the small airways. Outlined column: wild-type mice; Solid column: *Hdc*^{-/-} mice; Column **a** and **c**: saline inhalation, Column **b** and **d**: OVA inhalation. * $p < 0.01$.

through H2R.³⁶ Histamine also inhibited lipopolysaccharide-induced TNF- α production by down-regulating intercellular adhesion molecules (ICAM)-1 in human monocytes through H2R.³⁷ Alveolar macrophages are known to be a major source of TNF- α in the lung, and the increased production of TNF- α by alveolar macrophages in patients with asthma has been reported.^{38,39} The present study demonstrated an increase in the TNF- α concentration in BALF in the *Hdc*^{-/-} mice exposed to OVA compared to the wild-type mice under the same condition. This result suggested that histamine may play a role in suppressing TNF- α production through H2R in allergic airway inflammation in mice.

Busse *et al.* have recently demonstrated that chronic exposure of TNF- α to the airway induced goblet cell hyperplasia.⁴⁰ Taking these findings into consideration, we speculate that the depletion of histamine in the *Hdc*^{-/-} mice induced the up-regulation of TNF- α production, resulting in the increase of gob-

let cells in the airways with allergic inflammation.

In association with the increase of goblet cells, higher levels of Gob-5 and MUC5ac mRNA were found in the *Hdc*^{-/-} mice exposed to OVA. Previous reports have suggested that the expression of the Gob-5 gene, which corresponds to hCLCA1 in human beings, would be one of the first steps in mucus cell metaplasia and hyperplasia by inducing mucin gene expression.²⁵ MUC5ac gene, a member of the mucin gene family, has been shown to be expressed exclusively in the airway and associated with the hyperplasia of goblet cells. In addition to IL-4, IL-5 and IL-13, TNF- α was reported to increase the expression of the Gob-5 and MUC5ac genes.⁴¹ These findings suggested that the increased production of TNF- α in the BALF of the *Hdc*^{-/-} mice exposed to OVA may be, at least in part, involved in the increase of mRNA levels of the Gob-5 and MUC5ac genes, leading to the increased number of goblet cells in the airway.

Antigen challenge of sensitized mice or human asthmatic subjects results in increased TNF- α expression in BALF, peripheral blood, and tissue biopsy specimens.⁴²⁻⁴⁶ TNF- α production is partly regulated by histamine via H2R in macrophages and dendritic cells, as described above. Although we had no direct evidence to evaluate the role of TNF- α in goblet cell hyperplasia, the results of our study suggested that the increased level of TNF- α might contribute to enhance goblet cell hyperplasia and that histamine may be involved in the negative regulation of TNF- α production in allergic airway inflammation *via* the histamine receptors of inflammatory cells, including macrophages. To confirm the above, further studies are needed, such as experiments using mice in which the TNF- α activity is blocked with antibody or gene disruption.

There was a discrepancy between the TNF- α levels and the macrophage counts. The concentration of TNF- α in the knock-out mice was twice that in the wild-type mice, and the number of alveolar macrophages in *Hdc*^{-/-} mice was ten times that in the wild-type mice. We have no data to explain this, however, to answer this question, it would be necessary to evaluate the capacity of TNF- α production by alveolar macrophages in *Hdc*^{-/-} mice and wild mice. It would also be necessary to evaluate the time course of the concentration of TNF- α and the number of alveolar macrophages to understand their relationship.

We demonstrated the increased number of lymphocytes in BALF of *Hdc*^{-/-} mice. It has been reported by Jutel *et al.* that histamine stimulated the Th1 response through the H1 receptor. On the other hand, the activities of both Th1 and Th2 lymphocytes were inhibited by histamine through the H2 receptor.⁷ Elimination of the inhibitory actions on lymphocytes by histamine through the H2 receptor in *Hdc*^{-/-} mice may contribute to the increased number of lymphocytes in BALF.

The number of lymphocytes in the BALF of *Hdc*^{-/-} mice was twenty times that of wild-type mice, although no significant difference in the concentration of IL-2 in the BALF was found between knock-out and wild-type mice. The number of cells and cytokine levels in BALF are not always concordant in the time courses of *in vivo* experiments. The level of IL-2 in BALF peaked within 1 week during exposure to OVA (unpublished data). However, in this study we examined the cytokine levels and the cell differentials in BALF only on the 14th day. To explain the discrepancy between the IL-2 level and the number of lymphocytes in BALF in the study, it would have been helpful to evaluate the time course of IL-2 concentration and the number of lymphocytes in BALF.

Although we designed a 2-week exposure program to evaluate the extent of goblet cell hyperplasia in the *Hdc*^{-/-} mice, we did not examine airway hyperresponsiveness (AHR). However, we found no difference in AHR between the *Hdc*^{-/-} and the wild mice in our preliminary study as was reported in the previous acute model (unpublished data).²⁹

In the present study, we did not measure the anti-OVA specific IgE in serum. Although the levels of serum OVA-specific IgE and total IgE are not always consistent, the procedures for sensitization with OVA and exposure to OVA were directly associated with the increase of IgE in serum. Despite the fact that the levels of serum OVA-specific IgE were different from the level of total IgE in serum, the level of total IgE in serum reflected strongly the levels of serum OVA-specific IgE in our study.

Anti-histamine drugs available for therapy consist of H1 and H2 blockers. H1 blockers have not been recommended for therapy for asthma, but their anti-allergic actions have been recognized. According to the results of this study, the H2 blocker may have the possibility of stimulating goblet cell hyperplasia in patients with allergic asthma. It will be necessary to evaluate the effects of H2 blockers in a clinical study in the future.

In conclusion, we demonstrated enhanced goblet cell hyperplasia in the airway of OVA sensitized histamine-deficient mice. The results of our study suggested that histamine may play a significant role in goblet cell hyperplasia in the airway with allergic inflammation.

ACKNOWLEDGEMENTS

The authors would like to thank Miss M. Niisato and Miss M. Shibanaï for technical assistance, and are grateful to Mr. Brent Bell for his critical reading of the manuscript.

This study was supported by the Ministry of Education, Culture, Sports, Science and Technology, Japan.

REFERENCES

- Laitinen A, Laitinen LA. Pathology of asthma. *Allergy Proc* 1994;**15**:323-8.
- Shimura S, Andoh Y, Haraguchi M, Shirato K. Continuity of airway goblet cells and intraluminal mucus in the airways of patients with bronchial asthma. *Eur Respir J* 1996;**9**:1395-401.
- Carroll N, Carello S, Cooke C, James A. Airway structure and inflammatory cells in fatal attacks of asthma. *Eur Respir J* 1996;**9**:709-15.
- Aikawa T, Shimura S, Sasaki H *et al*. Marked goblet cell hyperplasia with mucus accumulation in the airways of patients who died of severe acute asthma attack. *Chest* 1992;**101**:916-21.
- Bryce PJ, Mathias CB, Harrison KL *et al*. The H1 histamine receptor regulates allergic lung responses. *J Clin Invest* 2006;**116**:1624-32.
- Siegel JN, Schwartz A, Askenase PW, Gershon RK. T-cell suppression and contrasuppression induced by histamine H2 and H1 receptor agonists, respectively. *Proc Natl Acad Sci U S A* 1982;**79**:5052-6.
- Jutel M, Watanabe T, Klunker S *et al*. Histamine regulates T-cell and antibody responses by differential expression of H1 and H2 receptors. *Nature* 2001;**413**:420-5.
- Oda T, Matsumoto S, Masuho Y *et al*. cDNA cloning and characterization of porcine histamine H4 receptor. *Biochim Biophys Acta* 2002;**1575**:135-8.
- Nakayama T, Kato Y, Hieshima K *et al*. Liver-expressed chemokine/CC chemokine ligand 16 attracts eosinophils by interacting with histamine H4 receptor. *J Immunol* 2004;**173**:2078-83.
- Ling P, Ngo K, Nguyen S *et al*. Histamine H4 receptor mediates eosinophil chemotaxis with cell shape change and adhesion molecule upregulation. *Br J Pharmacol* 2004;**142**:161-71.
- Hofstra CL, Desai PJ, Thurmond RL, Fung-Leung WP. Histamine H4 receptor mediates chemotaxis and calcium mobilization of mast cells. *J Pharmacol Exp Ther* 2003;**305**:1212-21.
- Voynow JA, Fischer BM, Malarkey DE *et al*. Neutrophil elastase induces mucus cell metaplasia in mouse lung. *Am J Physiol Lung Cell Mol Physiol* 2004;**287**:L1293-302.
- Harkema JR, Hotchkiss JA. Ozone- and endotoxin-induced mucous cell metaplasias in rat airway epithelium: novel animal models to study toxicant-induced epithelial transformation in airways. *Toxicol Lett* 1993;**68**:251-63.
- Ichinose T, Takano H, Miyabara Y *et al*. Enhancement of antigen-induced eosinophilic inflammation in the airways of mast-cell deficient mice by diesel exhaust particles. *Toxicology* 2002;**180**:293-301.
- Dabbagh K, Takeyama K, Lee HM *et al*. IL-4 induces mucin gene expression and goblet cell metaplasia *in vitro* and *in vivo*. *J Immunol* 1999;**162**:6233-7.
- Longphre M, Li D, Gallup M *et al*. Allergen-induced IL-9 directly stimulates mucin transcription in respiratory epithelial cells. *J Clin Invest* 1999;**104**:1375-82.
- Temann UA, Prasad B, Gallup MW *et al*. A novel role for murine IL-4 *in vivo*: induction of MUC5AC gene expression and mucin hypersecretion. *Am J Respir Cell Mol Biol* 1997;**16**:471-8.
- Zhu Z, Homer RJ, Wang Z *et al*. Pulmonary expression of interleukin-13 causes inflammation, mucus hypersecretion, subepithelial fibrosis, physiologic abnormalities, and eotaxin production. *J Clin Invest* 1999;**103**:779-88.
- Trifilieff A, Fujitani Y, Coyle AJ *et al*. IL-5 deficiency abol-

- ishes aspects of airway remodelling in a murine model of lung inflammation. *Clin Exp Allergy* 2001;**31**:934-42.
20. Rose MC, Voynow JA. Respiratory tract mucin genes and mucin glycoproteins in health and disease. *Physiol Rev* 2006;**86**:245-78.
 21. Fahy JV. Goblet cell and mucin gene abnormalities in asthma. *Chest* 2002;**122**:320S-6.
 22. Groneberg DA, Eynott PR, Lim S *et al*. Expression of respiratory mucins in fatal status asthmaticus and mild asthma. *Histopathology* 2002;**40**:367-73.
 23. Ordonez CL, Khashayar R, Wong HH *et al*. Mild and moderate asthma is associated with airway goblet cell hyperplasia and abnormalities in mucin gene expression. *Am J Respir Crit Care Med* 2001;**163**:517-23.
 24. Zuhdi Alimam M, Piazza FM, Selby DM *et al*. Muc-5/5ac mucin messenger RNA and protein expression is a marker of goblet cell metaplasia in murine airways. *Am J Respir Cell Mol Biol* 2000;**22**:253-60.
 25. Nakanishi A, Morita S, Iwashita H *et al*. Role of gob-5 in mucus overproduction and airway hyperresponsiveness in asthma. *Proc Natl Acad Sci U S A* 2001;**98**:5175-80.
 26. O'Neill L. Gob genes, mucus and asthma. *Trends Immunol* 2001;**22**:353.
 27. Zhou Y, Shapiro M, Dong Q *et al*. A calcium-activated chloride channel blocker inhibits goblet cell metaplasia and mucus overproduction. *Novartis Found Symp* 2002;**248**: 150-65; discussion 165-70, 277-82.
 28. Ohtsu H, Tanaka S, Terui T *et al*. Mice lacking histidine decarboxylase exhibit abnormal mast cells. *FEBS Lett* 2001;**502**:53-6.
 29. Koarai A, Ichinose M, Ishigaki-Suzuki S *et al*. Disruption of L-histidine decarboxylase reduces airway eosinophilia but not hyperresponsiveness. *Am J Respir Crit Care Med* 2003;**167**:758-63.
 30. Kozma GT, Losonczy G, Keszei M *et al*. Histamine deficiency in gene-targeted mice strongly reduces antigen-induced airway hyper-responsiveness, eosinophilia and allergen-specific IgE. *Int Immunol* 2003;**15**:963-73.
 31. Ohkawara Y, Lei XF, Stampfli MR *et al*. Cytokine and eosinophil responses in the lung, peripheral blood and bone marrow compartments in a murine model of allergen-induced airways inflammation. *Am J Respir Cell Mol Biol* 1997;**16**:510-20.
 32. Martell M, Gomez J, Esteban JI *et al*. High-throughput real-time reverse transcription-PCR quantitation of hepatitis C virus RNA. *J Clin Microbiol* 1999;**37**:327-32.
 33. Ryseck RP, Macdonald-Bravo H, Mattei MG, Bravo R. Structure, mapping, and expression of fisp-12, a growth factor-inducible gene encoding a secreted cysteine-rich protein. *J Cell Growth Differ* 1991;**2**:225-33.
 34. Heid CA, Stevens J, Livak KJ, Williams PM. Real time quantitative PCR. *Genome Res* 1996;**6**:986-94.
 35. Gutzmer R, Diestel C, Mommert S *et al*. Histamine H4 receptor stimulation suppresses IL-12p70 production and mediates chemotaxis in human monocyte-derived dendritic cells. *J Immunol* 2005;**174**:5224-32.
 36. Mazzoni A, Leifer CA, Mullen GE *et al*. Cutting edge: histamine inhibits IFN- α release from plasmacytoid dendritic cells. *J Immunol* 2003;**170**:2269-73.
 37. Morichika T, Takahashi HK, Iwagaki H *et al*. Histamine inhibits lipopolysaccharide-induced tumor necrosis factor- α production in an intercellular adhesion molecule-1- and B7.1-dependent manner. *J Pharmacol Exp Ther* 2003;**304**: 624-33.
 38. Azevedo I, de Blic J, Dumarey CH *et al*. Increased spontaneous release of tumour necrosis factor- α by alveolar macrophages from wheezy infants. *Eur Respir J* 1997;**10**: 1767-73.
 39. Broide DH, Lotz M, Cuomo AJ *et al*. Cytokines in symptomatic asthma airways. *J Allergy Clin Immunol* 1992;**89**: 958-67.
 40. Busse PJ, Zhang TF, Srivastava K *et al*. Chronic exposure to TNF- α increases airway mucus gene expression *in vivo*. *J Allergy Clin Immunol* 2005;**116**:1256-63.
 41. Hauber HP, Daigneault P, Frenkiel S *et al*. Niflumic acid and MSI-2216 reduce TNF- α -induced mucin expression in human airway mucosa. *J Allergy Clin Immunol* 2005;**115**: 266-71.
 42. Gosset P, Tscopoulos A, Wallaert B *et al*. Increased secretion of tumor necrosis factor alpha and interleukin-6 by alveolar macrophages consecutive to the development of the late asthmatic reaction. *J Allergy Clin Immunol* 1991;**88**:561-71.
 43. Gosset P, Tscopoulos A, Wallaert B *et al*. Tumor necrosis factor- α and interleukin-6 production by human mononuclear phagocytes from allergic asthmatics after IgE-dependent stimulation. *Am Rev Respir Dis* 1992;**146**:768-74.
 44. Cembrzynska-Nowak M, Szklarz E, Inglot AD, Teodorczyk-Injeyan JA. Elevated release of tumor necrosis factor- α and interferon- γ by bronchoalveolar leukocytes from patients with bronchial asthma. *Am Rev Respir Dis* 1993;**147**:291-5.
 45. Lukacs NW, Strieter RM, Chensue SW *et al*. TNF- α mediates recruitment of neutrophils and eosinophils during airway inflammation. *J Immunol* 1995;**154**:5411-7.
 46. Krishnaswamy G, Liu MC, Su SN *et al*. Analysis of cytokine transcripts in the bronchoalveolar lavage cells of patients with asthma. *Am J Respir Cell Mol Biol* 1993;**9**: 279-86.